



Lithofacies control on deformation bands

Oliver Wakefield^{*1}, Ed Hough¹, Jan Hennissen¹, Dan Parkes¹

¹ - British Geological Survey, Nottingham. oliverw@bgs.ac.uk




**British
Geological Survey**



NATURAL ENVIRONMENT RESEARCH COUNCIL

Overview

- Secondary strain induced localisation features (mostly planar) that typically develop in porous sandstones (Fossen et al., 2007).
- The development of deformation bands results in combinations of grain re-organisation, grain-dissolution, cataclasis (grain-breaking) and precipitation of new cements.
- Commonly have reduced permeability (orders of magnitude less) than the host rock.
- Classification schemes based on mechanism of deformation or kinematically.





Overview



British Geological Survey


NATURAL ENVIRONMENT RESEARCH COUNCIL

Overview



Thurstaston
Common

100 m



Data collected:
Dimensions (x, y, z)
Orientation
Permeability
Host facies
Type

Def. band
3 – 30 mD



Host SDST
80 – 2500 mD











British
Geological Survey

NATURAL ENVIRONMENT RESEARCH COUNCIL



Classification Scheme

- Current classification schemes don't make assessments for deformation geometry.
- The following proposed scheme can be used in conjunction with others.



Facies Control? *cont'd.*

Factors known linked to the formation of deformation:

- grain size
- grain sorting
- grain/clast composition (mineralogy)
- grain/clast roundness
- porosity

**Primary
Sedimentological**

- burial depth
- lithification
- amount and duration of stress
- pressure exerted by porefluids during deformation

**Secondary
'Stuff'**

What about variations facies....?



**British
Geological Survey**

NATURAL ENVIRONMENT RESEARCH COUNCIL

Facies Control?

Reference	Host succession	Depositional system
Antonellini et al. (1994)	Chinle Formation (Utah, USA) Wingate Sandstone Formation (Utah, USA) Kayenta Formations (Utah, USA)	Mixed fluvial, aeolian, lacustrine & palustrine Aeolian dunefield Aeolian dunefield
Aydin (1978)	Entrada Sandstone Formation (Utah, USA) Navajo Sandstone Formation (Utah, USA)	Aeolian; sabkha & dunefield Aeolian sand dune system
Cashman & Cashman (2000)		
Draganits et al. (2005)		
Edwards et al. (1993)	Hopeman Sandstone Formation (Moray Firth, UK)	Aeolian dunefield (with minor fluvial incursions)
Fossen (2010)	Entrada Sandstone Formation (Utah, USA) Navajo Sandstone Formation (Utah, USA)	Aeolian; sabkha & dunefield Aeolian dunefield
Fossen & Bale (2007)	Entrada Sandstone Formation (Utah, USA) Navajo Sandstone Formation (Utah, USA)	Aeolian; sabkha & dunefield Aeolian dunefield
Fowles & Burley (1994)	Penrith Sandstone Formation (Vale of Eden & Dumfries, UK)	Aeolian dunefield with fringing alluvial fans
Griffiths et al. (2016)	Sherwood Sandstone (Thurstaston, UK)	Mixed aeolian-fluvial (semi-arid)
Hodson et al. (2016)	Moab Tongue Member [Curtis Formation] (Utah, USA)	Aeolian dunefield
Johansen & Fossen (2008)	Carmel Formation (Utah, USA)	Aeolian; sabkha & dunefield
Johansen et al. (2005)	Moab Member [Entrada Sandstone Formation] (Utah, USA)	Aeolian; sabkha & dunefield
Main et al. (2000)	Hopeman Sandstone (Lossiemouth, Scotland)	Aeolian system (possibly wet)*
Mollema & Antonellini (1996)	Navajo Sandstone Formation (Utah, USA)	Aeolian dunefield
Parnell et al. (2004)	Upper Old Red Sandstone (Caithness, Scotland)	Aeolian dunefield
Parry et al. (2004)	Navajo Sandstone Formation (Utah, USA)	Aeolian dunefield
Raduha et al. (2016)	Entrada Sandstone Formation (Utah, USA)	Aeolian; sabkha & dunefield
Rotevatn et al. (2013)		
Schuessler et al. (2013)	Entrada Sandstone Formation (Utah, USA) Navajo Sandstone Formation (Utah, USA)	Aeolian; sabkha & dunefield Aeolian dunefield
Schuessler et al. (2013)	Aztec Sandstone (Nevada, USA) Hopeman Sandstone (Lossiemouth, Scotland)	Aeolian dunefield Aeolian system (possibly wet)*
Mollema & Antonellini (1996)	Navajo Sandstone Formation (Utah, USA)	Aeolian dunefield
Parnell et al. (2004)	Upper Old Red Sandstone (Caithness, Scotland)	Aeolian dunefield
Parry et al. (2004)	Navajo Sandstone Formation (Utah, USA)	Aeolian dunefield
Raduha et al. (2016)	Entrada Sandstone Formation (Utah, USA)	Aeolian; sabkha & dunefield
Rotevatn et al. (2013)		
Schuessler et al. (2013)	Entrada Sandstone Formation (Utah, USA) Navajo Sandstone Formation (Utah, USA)	Aeolian; sabkha & dunefield Aeolian dunefield
Schuessler et al. (2013)	Aztec Sandstone (Nevada, USA)	Aeolian dunefield
Skurtveit et al. (2015)	Navajo Sandstone Formation (Utah, USA) Page Formations (Utah, USA)	Aeolian dunefield Aeolian dunefield
Sternlof et al. (2006)	Aztec Sandstone (Nevada, USA)	Aeolian dunefield
Taylor & Pollard (2000)	Aztec Sandstone (Nevada, USA)	Aeolian dunefield
Tindall & Davis (2003)	Navajo Sandstone Formation (Utah, USA)	Aeolian dunefield
Torabi & Fossen (2009)	Entrada Sandstone Formation (Utah, USA)	Aeolian; sabkha & dunefield
Torabi et al. (2008)	Navajo Sandstone Formation (Utah, USA) Entrada Sandstone Formation (Utah, USA)	Aeolian dunefield Aeolian; sabkha & dunefield
Zuluaga et al. (2014)	Navajo Sandstone Formation (Utah, USA)	Aeolian dunefield



31 of 42 the stratigraphic units are aeolian (exclusively or mixed).



British Geological Survey

NATURAL ENVIRONMENT RESEARCH COUNCIL


Facies Control? *cont'd.*



British Geological Survey

NATURAL ENVIRONMENT RESEARCH COUNCIL

Facies Control? *cont'd.*




British Geological Survey

NATURAL ENVIRONMENT RESEARCH COUNCIL

Transect data

- Total transect length of ~1.7 km (T1 – T8), with 812 deformation bands counted.



Facies	Code	Total segment length %	Total number of bands %	Concentration factor *
Grainfall & grainflow	Sgl & Sgw	83.6	97.3	1.32
Trough xbedded sandstone	St	15.1	2.3	0.18
Wind-ripple facies	Sw	1.1	0.4	0.39
Massive sandstone	Sm	0.2	0	0
Deformed sandstone	Sa	>0.1	0	0
Upperplain beds	Sh	>0.1	0	0
Pebbly sandstone	Ss	>0.1	0	0
Low-angle inclined SDST	Sl	>0.1	0	0
Laminated sandstone	Sr	>0.1	0	0
Planas xbedded sandstone	Sp	>0.1	0	0

* - measure of def. band frequency per metre



British Geological Survey

NATURAL ENVIRONMENT RESEARCH COUNCIL



Conclusions?

Question: *Do facies exert a control on deformation band frequency or style?*


Answer: *Yes... and.....no!.....but for a good reason*



British
Geological Survey

NATURAL ENVIRONMENT RESEARCH COUNCIL

Conclusions...?



British Geological Survey

NATURAL ENVIRONMENT RESEARCH COUNCIL

Conclusions...?

Aeolian transport

- Discriminating grain selection.
- Dominant saltation transport method.
- Density of air ($\sim 1.3\text{kg/m}^2$).



Entrada SDST, Utah




Wilmslow SDST, UK



British Geological Survey

NATURAL ENVIRONMENT RESEARCH COUNCIL



Questions?



British Geological Survey

NATURAL ENVIRONMENT RESEARCH COUNCIL