

This is a post-peer-review, pre-copyedit version of an article published in *Cellular and Molecular Neurobiology* published by Springer. The final authenticated version is available online at: <http://dx.doi.org/10.1007/s10571-017-0571-4>

# Cellular and Molecular Neurobiology

## Effects of acute alcohol exposure on layer 5 pyramidal neurons of juvenile mice --Manuscript Draft--

<b>Manuscript Number:</b>	CEMN-D-17-00264R3	
<b>Full Title:</b>	Effects of acute alcohol exposure on layer 5 pyramidal neurons of juvenile mice	
<b>Article Type:</b>	Original Article	
<b>Keywords:</b>	Ethanol; cerebral cortex; electrophysiology; pyramidal neurons; acute slice; mIPSC	
<b>Corresponding Author:</b>	Alberto Granato, M.D. Catholic University Milan, ITALY	
<b>Corresponding Author Secondary Information:</b>		
<b>Corresponding Author's Institution:</b>	Catholic University	
<b>Corresponding Author's Secondary Institution:</b>		
<b>First Author:</b>	Francesco Ferrini, D.V.M.	
<b>First Author Secondary Information:</b>		
<b>Order of Authors:</b>	Francesco Ferrini, D.V.M.	
	Benjamin Dering, Ph.D.	
	Andrea De Giorgio, Ph.D.	
	Laura Lossi, D.V.M.	
	Alberto Granato, M.D.	
<b>Order of Authors Secondary Information:</b>		
<b>Funding Information:</b>	Università degli Studi di Torino	Dr. Francesco Ferrini Dr. Laura Lossi
	Università Cattolica del Sacro Cuore	Dr. Alberto Granato
<b>Abstract:</b>	<p>Early-onset drinking during childhood or preadolescence is a serious social problem. Yet, most of the basic neurobiological research on the acute effects of ethanol has been carried out on adult or early postnatal animals. We studied the effect of alcohol exposure on the basic electrophysiological properties and cell viability of layer 5 pyramidal neurons from the somatosensory cortex of juvenile (P21-P23) C57BL/6N mice. After bath application of 50 mM ethanol to acute slices of the somatosensory cortex, no adverse effects were detected on cells survival, whereas the input resistance and firing rate of layer 5 neurons were significantly reduced. While the effect on the input resistance was reversible, the depressing effect on cell firing remained stable after 6 minutes of alcohol exposure. Ethanol application did not result in any significant change of mIPSC frequency, amplitude, and rise time. A slight increase of mIPSC decay time was observed after 6 minutes of ethanol exposure. The molecular mechanisms leading to these alterations and their significance for the physiology of the cerebral cortex are briefly discussed.</p>	
<b>Response to Reviewers:</b>	<p>Reviewer's comments and our answers are interleaved. Changes to the text are highlighted in red in the manuscript file.</p> <p>REVIEWER: The additional references provided are helpful. However, there are several canonical reviews devoted to this area explicitly that would be hugely beneficial to readers less familiar with the nuances of this research area. Dr. LP Spear et al. has authored many such reviews for example.</p>	

Response:

As suggested by the Reviewer, the following text has been added to Introduction: "Indeed, ethanol affects different cortical regions and induces important ontogenetic alterations during adolescence, which critically influence subsequent drug self-administration in adulthood (Spear 2016)".

REVIEWER:

I understand referencing work using very similar methods. However, it seems relevant to at least briefly discuss how the current work fits in the context of developmental patch work in other brain areas/rodents strains. For example, findings from Brodie and Appel (2000) suggest that ethanol may have very different effects as a function of age and brain region examined, as well as in other rodent strains.

Response:

To point out that different alcohol effects might be observed, depending on the rodent strain and/or developmental stage, the following text has been added to Discussion: "As a more general remark on experimental alcohol studies, it should be taken into account that different effects of ethanol can be observed not only in different brain regions, but also as a function of age and rodent strains (Brodie and Appel 2000; Ikonomidou et al. 2000; Dou et al. 2013)."

REVIEWER:

Related to this last point, this reviewer was only able to find information pertaining to the /6N substrain of the C7BL mice when Charles River (Italy) was queried. Differences between these 2 substrains (/6J and /6N) in ethanolrelated behaviors have been reported, and it seems reasonable that these may be due to differences in ethanolinduced brain function. Please confirm which of these substrains was used. Please also include details on the number of generations from the founding stock mice were that were used for experiments. Were littermates used for these studies or was there any effort to test mice from different litters to control for this potential factor?

Response:

We thank the Reviewer for raising this point. Both substrains of C57BL/6 mice were ordered in the past by the University of Turin. The ones used in the present study were purchased from Charles River Italy, as indicated in the manuscript, but they were actually C57BL/6N, not C57BL/6J, as mistakenly indicated in the previous version of the manuscript. We have now corrected the substrain indication. In addition, the phrase of Material and Methods: "All animals used were bred in house" has been modified as follows: "All animals used were bred in house for four generations. Experiments were conducted on mice obtained from different litters."

Finally, according to the instructions for authors, we have slightly modified Figure 4 (part d), in order to remove reference to colors from the caption.

[Click here to view linked References](#)

# Effects of acute alcohol exposure on layer 5 pyramidal neurons of juvenile mice

Francesco Ferrini<sup>1</sup>, Benjamin Dering<sup>2</sup>, Andrea De Giorgio<sup>3</sup>, Laura Lossi<sup>1</sup>, Alberto Granato<sup>4</sup>

1. Department of Veterinary Sciences, University of Turin, Largo Braccini 2, 10095 Grugliasco, Italy.

2. Faculty of Natural Sciences, University of Stirling, Stirling, United Kingdom.

3. Faculty of Psychology, University eCampus, Novedrate, Italy.

4. Department of Psychology, Catholic University, Milan, Italy

Corresponding Author:

Alberto Granato

Department of Psychology, Catholic University

Largo A. Gemelli 1, 20123, Milan, Italy

Ph.: +39 02 72348588

alberto.granato@unicatt.it

1  
2 **Abstract**  
3

4 Early-onset drinking during childhood or preadolescence is a serious social problem. Yet, most of  
5 the basic neurobiological research on the acute effects of ethanol has been carried out on adult or  
6 early postnatal animals. We studied the effect of alcohol exposure on the basic electrophysiological  
7 properties and cell viability of layer 5 pyramidal neurons from the somatosensory cortex of juvenile  
8 (P21-P23) C57BL/6N mice. After bath application of 50 mM ethanol to acute slices of the  
9 somatosensory cortex, no adverse effects were detected on cells survival, whereas the input  
10 resistance and firing rate of layer 5 neurons were significantly reduced. While the effect on the  
11 input resistance was reversible, the depressing effect on cell firing remained stable after 6 minutes  
12 of alcohol exposure. Ethanol application did not result in any significant change of mIPSC  
13 frequency, amplitude, and rise time. A slight increase of mIPSC decay time was observed after 6  
14 minutes of ethanol exposure. The molecular mechanisms leading to these alterations and their  
15 significance for the physiology of the cerebral cortex are briefly discussed.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

27 **Keywords**  
28

29 Ethanol; cerebral cortex; electrophysiology; pyramidal neurons; acute slice; mIPSC.  
30  
31  
32  
33  
34

35 **Author Contributions**  
36

37 Francesco Ferrini designed and performed the experiments, analyzed the data and wrote the  
38 manuscript. Benjamin Dering designed and performed the experiments and analyzed the data.  
39 Andrea De Giorgio performed the experiments and analyzed the data. Laura Lossi performed the  
40 experiments and analyzed the data. Alberto Granato designed and performed the experiments,  
41 analyzed the data and wrote the manuscript. All the authors have read and approved the manuscript.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Introduction

Alcohol abuse is one of the most serious problems faced by our society, with a tremendous impact on public health and economy (Rehm et al. 2009). The issues raised by childhood- and preadolescence-onset drinking (Donovan 2013) are even more worrisome, since early-onset drinking is positively associated with the development of alcohol dependence later in life (Hingson et al. 2006). In particular, alcohol consumption during adolescence has been associated with disruptions of many normal developmental processes, potentially disturbing the maturation of higher-order executive functions (McMurray et al. 2016). **Indeed, ethanol affects different cortical regions and induces important ontogenetic alterations during adolescence, which critically influence subsequent drug self-administration in adulthood (Spear 2016).** Surprisingly, despite the huge amount of work devoted to experimental research on alcohol dependence, little is known about the neurobiology of alcohol effects in juvenile lab animals.

Pyramidal neurons of layer 5 (L5), the most complex cells among those providing the output from the cerebral cortex (Ramaswamy and Markram 2015), are highly sensitive to the long-term deleterious effects of ethanol. Early exposure to alcohol during the first stages of development can permanently modify the electrophysiological properties of these neurons (Granato et al. 2012). As to the acute effects of ethanol, there are reports describing the alterations of the electrophysiological parameters of L5 neurons after alcohol application on acute slices from adult rodents (Proctor et al. 1992; Sessler et al. 1998). Other studies, instead, focused on effects of ethanol in cortical slices from postnatal rodents showing not only effects on neuronal activity, as in the adult, but also on cell survival (Sanderson et al. 2009). Conversely, little is known about the effects of ethanol on layer 5 neurons of juvenile rodents (P21-23), a developmental stage considered to correspond to “early adolescence” in humans. To address this issue and fill the gap of knowledge at such a critical age, we designed *in vitro* electrophysiological experiments and cell viability tests on P21-P23 C57BL/6 mice, the strain most commonly adopted in alcohol research as genetic background for the generation of transgenic mice (Heit et al. 2015).

## Materials and Methods

**Animals.** Juvenile male (P21-P23) C57BL/6N mice were used in the present study. The breeding colony was purchased from Charles River Laboratories (Italy). All animals used were bred in house **for four generations. Experiments were conducted on mice obtained from different litters.** To avoid exposing mice to unwanted stressful events prior to the experimental procedures, adolescent mice were not weaned until sacrifice.

1 All the experiments were conducted in accordance with the Society for Neuroscience Policies on  
2 the Use of Animals and Humans in Neuroscience Research, as well as current Italian and EU  
3 regulation on animal experimentation and welfare.  
4

5 **Electrophysiology.** Mice were anaesthetized (Pentothal 50mg/Kg i.p.) and, after decapitation, the  
6 brain was quickly removed. Coronal slices (300  $\mu\text{m}$ ) of the primary somatosensory cortex were cut  
7 on a vibratome in ice-cold, oxygenated solution containing (in mM): 250 sucrose, 25  $\text{NaHCO}_3$ , 2.5  
8 KCl, 1.25  $\text{NaH}_2\text{PO}_4$ , 1  $\text{MgCl}_2$ , 25 glucose, 2  $\text{CaCl}_2$ . The composition of the ACSF used for all the  
9 subsequent procedures was the same as the cutting solution, with the exception that the sucrose was  
10 replaced by NaCl 125 mM. Slices were kept at 34°C in oxygenated ACSF for 1h and then at room  
11 temperature until use. Recording pipettes (4-8  $\text{M}\Omega$ ) were made from borosilicate glass capillaries.  
12 Whole-cell current clamp recordings were obtained from the soma of visually identified pyramidal  
13 neurons of layer 5 using IR-DIC optics. Recordings were carried out using a Multiclamp 700B  
14 amplifier (Molecular Devices, Sunnyvale, CA) and acquired with a Digidata digitizer (Molecular  
15 Devices). Series resistance ( $R_s$ ) was monitored throughout the recordings and compensated using  
16 the bridge balance circuit of the amplifier. Recordings were discarded if  $R_s$  changed by more than  
17 20% during the experimental procedure.  
18

19 Current clamp recordings were obtained with an internal solution containing (in mM): 135 K  
20 gluconate, 5 KCl, 10 HEPES, 2  $\text{MgCl}_2$ , 4 $\text{Na}_2\text{ATP}$ , 0.4 NaGTP, pH 7.2. For some recordings, the  
21 intracellular solution contained 0.1% Lucifer Yellow. Several steps of hyperpolarizing and  
22 depolarizing current were injected into the soma (increments of 50 pA; duration: 800 ms). Once  
23 stable recordings were obtained in normal ACSF (preEt), the slice was perfused with ACSF + 50  
24 mM ethanol and additional recordings were obtained from the same cell 3 and 6 minutes after the  
25 beginning of ethanol application (3mEt and 6mEt, respectively). Recordings were also performed 5  
26 and 10 minutes after the beginning of alcohol washout; cells were discarded if the number of spikes  
27 did not return to at least 40% of baseline during washout.  
28

29 Active and passive membrane properties were analyzed using Igor Pro (Wavemetrics, Lake  
30 Oswego, OR). The following subthreshold parameters were measured: resting membrane potential,  
31 input resistance, membrane time constant, depolarizing voltage sag (calculated as  $(V_{\text{max}} - V_{\text{ss}}) /$   
32  $V_{\text{max}}$ , where  $V_{\text{max}}$  is the transient voltage peak reached soon after a hyperpolarizing current step and  
33  $V_{\text{ss}}$  is the steady state voltage). Measured suprathreshold parameters were: spike threshold, spike  
34 amplitude, spike half-width (i.e. the spike width at half amplitude), afterhyperpolarization,  
35 interspike interval (ISI), and ratio between the last and 1<sup>st</sup> ISI of a spike train.  
36

37 For voltage clamp recordings K gluconate and KCl in the intracellular solution were substituted  
38 with CsCl (140 mM). Miniature inhibitory post-synaptic currents (mIPSCs) were isolated in  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 presence of TTX (1  $\mu$ M; Tocris Cookson, Bristol, UK), APV (40  $\mu$ M; Sigma, St. Louis, MO, USA),  
2 NBQX (10  $\mu$ M; Sigma) at a holding potential of -60 mV. Under these experimental conditions  $E_{Cl}$  is  
3 near 0 mV, thus IPSCs are inwardly directed.  
4

5 mIPSCs were analyzed by Mini Analysis software (Synaptosoft Inc., Decatur, GA). mIPSC  
6 frequency and amplitude were sampled for periods of 100 s. Rise time and decay time kinetics were  
7 analyzed by averaging over 100 synaptic events/cell per each experimental condition. Rise time was  
8 defined as the duration of the rise from 10 to 90% of the peak. Decay time was calculated by fitting  
9 the 10-90% decay phase with a monoexponential function and expressed as time constant  $\tau$ .  
10

11 **Cell viability assay.** After recovery in ACSF, three slices for each experimental condition (control,  
12 3mEt, and 6mEt) were transferred into ACSF containing 1.5 mM propidium iodide (PI) and  
13 incubated for 10 min at room temperature. They were then washed three times in plain ACSF,  
14 followed by 30 min fixation in 4% paraformaldehyde in 0.1 M phosphate-buffered saline (PBS) (pH  
15 7.4-7.6). Slices were then washed in PBS (2 x 10 min) and double distilled water (2 x 5 min) and  
16 mounted in fluorescent free medium (Vectashield® Antifade Mounting Medium, Vector  
17 Laboratories, Burlingame, CA). They were then photographed using a Leica DM6000 wide-field  
18 fluorescence microscope (Leica Microsystems, Wetzlar, Germany) with a 20x lens. For each slice,  
19 10 randomly selected microscope fields were photographed at a resolution of 1392x1040 pixels  
20 (0.3084 mm<sup>2</sup>) and PI stained nuclei were counted with the "Count Particles" function of the ImageJ  
21 software (NIH, Bethesda, MD, USA) in an interval of area size between 12.56 and 78.50  
22  $\mu$ m<sup>2</sup> (diameter 4-10  $\mu$ m). Values were expressed as number of PI stained nuclei/mm<sup>2</sup>.  
23

24 **Statistical analysis.** Statistical differences of all grouped electrophysiological data, but ISI, were  
25 evaluated using the ANOVA for repeated measures test followed by Bonferroni post-hoc. As ISI  
26 data distribution is not normal, differences were analyzed through the non-parametric Wilcoxon  
27 signed-rank test. Differences in the number of dead cells in the cell viability assay were tested by a  
28 one-way ANOVA. All values are expressed as mean  $\pm$  standard error of the mean (s.e.m.).  
29 Differences were considered to be significant at  $P < 0.05$ .  
30

## 31 Results

32 Acute effects of ethanol on neuronal activity in L5 pyramidal neurons were assessed after 3 minutes  
33 (3mEt) and 6 minutes (6mEt) of slice exposure to 50 mM ethanol, a concentration previously  
34 adopted in several functional studies *in vitro* and *in vivo* (e.g.: Criswell et al. 2008; Ehlich et al.  
35 2012, Huang et al. 2012) and known to mimic ethanol levels in the blood of heavy drinkers (Han et  
36 al. 1998; Ehlich et al. 2012). This brain concentration of ethanol causes cognitive dysfunction in  
37 humans and animal models, as well as altered synaptic function (reviewed in Tipps et al., 2014;  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 White, 2003; Zorumski et al., 2014). Moreover, levels of ethanol in the central nervous system have  
2 been shown to match with blood levels in animal models (Gilpin et al., 2009; Smolen and Smolen,  
3 1989).  
4

5 To exclude the possibility that the effects of ethanol on neuronal activity could be linked to its pro-  
6 apoptotic effects (Olney, 2014) we have evaluated the incidence of cell death with PI in any of the  
7 three experimental conditions of the present study (Fig. 1). Dead cells are easily spotted as their  
8 nuclei are intensely fluorescent following PI uptake. PI stained nuclei are scattered throughout the  
9 slice and there are no obvious differences in the number of positive nuclei between the three  
10 experimental conditions (Fig. 1a-c) Fig. 1d shows the results of statistical analysis in graph form.  
11 After one-way ANOVA there are no statistically significant differences among the three  
12 experimental groups ( $F_{2,30} = 0.467$ ;  $P = 0.63$ ).  
13  
14  
15  
16  
17  
18  
19

20 All the cells recorded in current clamp and used for the quantitative evaluation ( $n = 8$ ) were regular  
21 spiking L5 pyramidal neurons with a moderate spike frequency adaptation and input resistance  
22 ranging from 92 to 262 M $\Omega$  (mean  $\pm$  s.e.m. =  $161.53 \pm 20.75$  M $\Omega$ ). The morphological control  
23 performed on neurons filled with Lucifer yellow confirmed the presence of basal dendrites and of  
24 an apical dendrite reaching the most superficial layers of the cortex. Figure 2 summarizes the main  
25 subthreshold parameters before ethanol exposure and 3-6 minutes after the beginning of ethanol  
26 bath application. The mean resting membrane potential was hyperpolarized by about 5 mV after  
27 ethanol application, with a statistically significant difference between preEt and 3mEt, and between  
28 preEt and 6mEt (Fig. 2a). The input resistance and the membrane time constant showed a similar  
29 trend (Fig. 2b-c): they were significantly lower than preEt at 3mEt, while they returned to baseline  
30 levels at 6mEt. No changes were observed in depolarizing voltage sag (Fig. 2d). Suprathreshold  
31 parameters obtained upon somatic injections of depolarizing current are shown in Figure 3. As  
32 demonstrated by the ISI, the firing frequency was significantly reduced at both 3mEt and 6mEt, as  
33 compared to preEt (Fig. 3a). This behavior was substantially maintained for all the levels of injected  
34 current (rheobase, 50 pA, and 100 pA above rheobase; Fig. 3b-d). For current injections 50 pA  
35 above rheobase, there was a significantly higher frequency adaptation at 6mEt, as compared to  
36 preEt (Fig. 3c - *inset*). Other action potential parameters (threshold, amplitude, half-width, and  
37 afterhyperpolarization; Fig. 3e-h) were not significantly different before and after ethanol exposure,  
38 except threshold values that were significantly higher at 6mEt than at 3mEt.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

54 As ethanol is known to enhance GABAergic inhibition (Lobo and Harris 2008; Förster et al.  
55 2016), we tested whether alterations in active and passive membrane properties were also paralleled  
56 by increased miniature inhibitory postsynaptic transmission (mIPSCs; Fig. 4). Ethanol application  
57 did not significantly alter mIPSC frequency, suggesting a lack of pre-synaptic effects (Fig. 4b).  
58  
59  
60  
61  
62  
63  
64  
65

1 Also, no differences were observed in mIPSC amplitude and rise time (Fig. 4c-e). However a slight  
2 but significant increase in decay time was observed at 6mEt as compared to preEt level (Fig. 4f).  
3  
4

## 5 **Discussion**

6  
7 The present study provides a comprehensive description of the effects of alcohol exposure on the  
8 electrophysiological properties of neocortical pyramidal neurons in juvenile rodents. The  
9 deleterious actions of ethanol in the immature brain mainly involve a direct depression of neuronal  
10 activity (Lotfullina and Khazipov 2017). The inhibitory effects of ethanol decrease with age and, in  
11 adult animals, ethanol only mildly depresses neuronal firing. In newborn P3-P9 rats, the overall  
12 suppression of neuronal activity is mediated by both the depression of NMDA receptors and the  
13 potentiation of GABAergic activity (Galindo et al. 2005, Sanderson et al., 2009). This combined  
14 effect has been associated with ethanol-induced apoptosis in the developing brain (Lotfullina and  
15 Khazipov 2017). Although the suppressive effects of ethanol on cortical activity and the consequent  
16 apoptosis decrease with age, yet in adult cortical neurons ethanol still decreases NMDA currents,  
17 spike firing, and input resistance with no effects on GABAergic transmission (Badanich et al. 2013  
18 Sessler et al. 1998). Our data confirm that ethanol has suppressive effects on firing activity of  
19 pyramidal neurons from juvenile adolescent mice, without causing detectable cell death. Unlike  
20 adult neurons, however, it also causes a mild post-synaptic enhancement of GABAergic  
21 transmission, which may further amplify the suppressive effect on firing activity.  
22  
23

24 In particular, we have shown a significant reduction of the input resistance and membrane time  
25 constant after 3 minutes of 50 mM ethanol bath application, accompanied by a hyperpolarization of  
26 resting membrane potential. Consistently, firing rate was also significantly reduced after both 3 and  
27 6 minutes of alcohol exposure. Our data are in line with previous observations made on acute slices  
28 of adult rodents, although some discrepancies among different works can be found. Sessler et al.  
29 (1998) reported a decreased firing rate of L5 neurons of young (125-200 gr) rats after bath  
30 application of ethanol. A reduction in input resistance was also observed, while the authors did not  
31 find consistent changes of membrane potential (Sessler et al. 1998). Conversely, the  
32 hyperpolarization, in presence of a small, non-significant reduction of input resistance, was  
33 observed in rats by Proctor and coworkers (1992). In large regular spiking neurons of the seven-  
34 week-old mouse orbitofrontal cortex, the application of ethanol led to a reduction of spike firing  
35 and input resistance, accompanied by hyperpolarization (Badanich et al. 2013). These effects seem  
36 to be region-specific, since in hippocampal slices the acute exposure to alcohol was more frequently  
37 accompanied by depolarization rather than hyperpolarization (Siggins et al. 1987). **As a more  
38 general remark on experimental alcohol studies, it should be taken into account that different effects**  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 of ethanol can be observed not only in different brain regions, but also as a function of age and  
2 rodent strains (Brodie and Appel 2000; Ikonomidou et al. 2000; Dou et al. 2013).

3 There are several possible explanations for the decrease in intrinsic excitability and input resistance  
4 observed in the present study. The potentiation of GABAergic transmission by ethanol has been  
5 repeatedly observed in brain slices (reviewed in Weiner and Valenzuela 2006). Our data on  
6 mIPSCs, however, show no significant difference of amplitude and frequency after exposure to  
7 ethanol. These findings are in agreement with those obtained by Fleming et al. (2009) on cultured  
8 cortical neurons. In other studies, a change of miniature or spontaneous IPSCs has been observed  
9 after exposure of dissociated cortical neurons to 100 mM ethanol (Marszalec et al. 1998; Moriguchi  
10 et al. 2007) and increased presynaptic release of GABA has been reported in different immature  
11 rodent models (Galindo et al. 2005, Sanderson et al., 2009). Even in absence of  
12 frequency/amplitude modifications, the slightly longer duration of mIPSCs observed in our study  
13 after 6 minutes of ethanol application, and therefore the increased GABA-mediated Cl<sup>-</sup> charge  
14 transfer, may partly explain the decreased excitability of pyramidal neurons. However, as no effects  
15 on decay time were detected after 3 minutes, the increased inhibitory synaptic transmission is  
16 unlikely to underlie early ethanol-induced changes in cell activity. On the other hand, we cannot  
17 exclude that ethanol may also potentiate extrasynaptic GABA<sub>A</sub> receptors, thus challenging tonic  
18 GABAergic inhibition. Indeed, ethanol-mediated effects on tonic inhibition have been previously  
19 reported in hippocampal neurons (Wei et al. 2004). Further investigations are thus required to  
20 specifically address this point in cortical neurons.

21 Another effect potentially accounting for a reduction of input resistance is the potentiation of  
22 hyperpolarization-activated currents (I<sub>h</sub>). It is known that ethanol augments I<sub>h</sub> in neuronal and non-  
23 neuronal cell types (e.g. Okamoto et al. 2006; Chen et al. 2012). The alcohol-mediated potentiation  
24 of I<sub>h</sub> on hippocampal interneurons is more effective in adolescent than in adult rats (Yan et al.  
25 2009). However, the expression of HCN channels mediating I<sub>h</sub> at somatic locations of L5 pyramidal  
26 neurons is constantly low during development (Atkinson and Williams 2009). Furthermore, in the  
27 present study we did not observe a significant change of the depolarizing sag after injection of  
28 hyperpolarizing current. Therefore, the effect of ethanol on I<sub>h</sub>, if any, is unlikely to be the main  
29 factor responsible for the decrease of input resistance. We cannot rule out that ethanol can affect  
30 other ion channels, as the electrophysiological properties of neurons are the result of a combined  
31 effect of several conductances (e.g., Day et al. 2005).

32 Interestingly, we have also found that some of the ethanol-related changes are time-dependent. In  
33 fact, while the decreased spike firing is maintained throughout the alcohol superfusion, the input  
34 resistance and membrane time constant display a more complex time course, showing a significant  
35

1 decrease after 3 minutes of alcohol application, followed by a return to control values after 6  
2 minutes. This biphasic action of ethanol might involve slow-onset processes such as ion channel  
3 phosphorylation (reviewed in Trudell et al., 2014).  
4

5 Whatever the mechanism accounting for the alterations observed in the present study, they can  
6 impair the function of juvenile neurons acutely exposed to alcohol. The changes of membrane time  
7 constant, besides affecting synaptic integration, can also modify the induction of spike-timing  
8 dependent plasticity (Fuenzalida et al. 2007). The reduced intrinsic excitability of pyramidal  
9 neurons can have deep consequences on network activity and on plastic adjustments required for  
10 learning and memory processes (see Cohen et al. 2017, for a discussion on the interplay among  
11 neuron excitability, plasticity, and network remodeling). These alterations are expected to have a  
12 dramatic effect on cortical activity of young drinkers, since the blood ethanol concentrations in  
13 adolescents rise more rapidly as compared to adults. According to the NIAAA (National Institute on  
14 Alcohol Abuse and Alcoholism), binge drinking in adults corresponds to 5 drinks within a 2 hour  
15 period, which lead to a blood concentration above 80 mg/dl. In the adolescent population, three  
16 drinks only are sufficient to pass the threshold (Donovan, 2009), thus increasing the likelihood that  
17 a concentration close to that used in our experiments is reached. Therefore, our data contribute to  
18 clarify that even single, acute exposures to alcohol can have dramatic and potential long-term  
19 consequences on brain electrical activity and behavior.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

- Atkinson SE, Williams SR (2009) Postnatal development of dendritic synaptic integration in rat neocortical pyramidal neurons. *J Neurophysiol* 102:735-751. doi: 10.1152/jn.00083.2009.
- Badanich KA, Mulholland PJ, Beckley JT, Trantham-Davidson H, Woodward JJ (2013) Ethanol reduces neuronal excitability of lateral orbitofrontal cortex neurons via a glycine receptor dependent mechanism. *Neuropsychopharmacology* 38:1176-1188. doi: 10.1038/npp.2013.12.
- Brodie MS, Appel SB (2000) Dopaminergic neurons in the ventral tegmental area of C57BL/6J and DBA/2J mice differ in sensitivity to ethanol excitation. *Alcohol Clin Exp Res* 24:1120-1124.**
- Chen Y, Wu P, Fan X, Chen H, Yang J, Song T, Huang C (2012) Ethanol enhances human hyperpolarization-activated cyclic nucleotide-gated currents. *Alcohol Clin Exp Res* 36:2036-2046. doi: 10.1111/j.1530-0277.2012.01826.x.
- Cohen EJ, Quarta E, Bravi R, Granato A, Minciacchi D (2017) Neural plasticity and network remodeling: From concepts to pathology. *Neuroscience*. 344:326-345. doi: 10.1016/j.neuroscience.2016.12.048.
- Criswell HE, Ming Z, Kelm MK, Breese GR (2008) Brain regional differences in the effect of ethanol on GABA release from presynaptic terminals. *J Pharmacol Exp Ther*. 326: 596-603. doi: 10.1124/jpet.107.135418.
- Day M, Carr DB, Ulrich S, Ilijic E, Tkatch T, Surmeier DJ (2005) Dendritic excitability of mouse frontal cortex pyramidal neurons is shaped by the interaction among HCN, Kir2, and K<sub>leak</sub> channels. *J Neurosci* 25: 8776-8787.
- Donovan JE (2009) Estimated blood alcohol concentrations for child and adolescent drinking and their implications for screening instruments. *Pediatrics*. 123:e975-981. doi: 10.1542/peds.2008-0027.
- Donovan JE (2013) The burden of alcohol use: focus on children and preadolescents. *Alcohol Res* 35:186-192.
- Dou X, Wilkemeyer MF, Menkari CE, Parnell SE, Sulik KK, Charness ME (2013) Mitogen-activated protein kinase modulates ethanol inhibition of cell adhesion mediated by the L1 neural cell adhesion molecule. *Proc Natl Acad Sci U S A* 110: 5683-5688. doi: 10.1073/pnas.1221386110.**
- Ehrlich D, Pirchl M, Humpel C (2012) Ethanol transiently suppresses choline-acetyltransferase in basal nucleus of Meynert slices. *Brain Res* 1459: 35-42. doi: 10.1016/j.brainres.2012.04.020.
- Fleming RL, Manis PB, Morrow AL (2009) The effects of acute and chronic ethanol exposure on presynaptic and postsynaptic gamma-aminobutyric acid (GABA) neurotransmission in cultured cortical and hippocampal neurons. *Alcohol* 43: 603-618. doi: 10.1016/j.alcohol.2009.10.006.
- Förstera B, Castro PA, Moraga-Cid G, Aguayo LG (2016) Potentiation of Gamma Aminobutyric Acid Receptors (GABAAR) by Ethanol: How Are Inhibitory Receptors Affected? *Front Cell Neurosci*. 10: 114. doi: 10.3389/fncel.2016.00114.

1 Fuenzalida M, Fernandez de Sevilla D, Buño W (2007) Changes of the EPSP waveform regulate  
2 the temporal window for spike-timing-dependent plasticity. *J Neurosci* 27:11940-11948.  
3

4 Galindo R, Zamudio PA, Valenzuela CF (2005) Alcohol is a potent stimulant of immature neuronal  
5 networks: implications for fetal alcohol spectrum disorder. *J Neurochem* 94:1500-1511.  
6

7  
8 Gilpin NW, Smith AD, Cole M, Weiss F, Koob GF, Richardson HN (2009) Operant behavior and  
9 alcohol levels in blood and brain of alcohol-dependent rats. *Alcohol Clin Exp Res* 33:2113–2123.  
10 doi: 10.1111/j.1530-0277.2009.01051.x  
11

12  
13 Granato A, Palmer LM, De Giorgio A, Tavian D, Larkum ME (2012) Early exposure to alcohol  
14 leads to permanent impairment of dendritic excitability in neocortical pyramidal neurons. *J*  
15 *Neurosci* 32:1377-1382. doi: 10.1523/JNEUROSCI.5520-11.2012.  
16

17  
18 Han CL, Liao CS, Wu CW, Hwong CL, Lee AR, Yin SJ (1998) Contribution to first-pass  
19 metabolism of ethanol and inhibition by ethanol for retinol oxidation in human alcohol  
20 dehydrogenase family--implications for etiology of fetal alcohol syndrome and alcohol-related  
21 diseases. *Eur J Biochem* 254: 25-31.  
22

23  
24 Heit C, Dong H, Chen Y, Shah YM, Thompson DC, Vasiliou V (2015) Transgenic mouse models  
25 for alcohol metabolism, toxicity, and cancer. *Adv Exp Med Biol* 815:375-87. doi: 10.1007/978-3-  
26 319-09614-8\_22.  
27

28  
29 Hingson RW, Heeren T, Winter MR (2006) Age at drinking onset and alcohol dependence: age at  
30 onset, duration, and severity. *Arch Pediatr Adolesc Med* 160:739-746.  
31

32  
33 Huang JJ, Yen CT, Tsai ML, Valenzuela CF, Huang C (2012) Acute ethanol exposure increases  
34 firing and induces oscillations in cerebellar Golgi cells of freely moving rats. *Alcohol Clin Exp Res*  
35 36:2110-2116. doi: 10.1111/j.1530-0277.2012.01818.x.  
36

37  
38 Ikonomidou C, Bittigau P, Ishimaru MJ, Wozniak DF, Koch C, Genz K, Price MT, Stefovská V,  
39 Hörster F, Tenkova T, Dikranian K, Olney JW (2000) Ethanol-induced apoptotic neurodegeneration  
40 and fetal alcohol syndrome. *Science* 287: 1056-1060.  
41

42  
43 Lobo IA, Harris RA (2008) GABA(A) receptors and alcohol. *Pharmacol Biochem Behav.* 90:90-94.  
44 doi: 10.1016/j.pbb.2008.03.006.  
45

46  
47 Lotfullina N, Khazipov R (2017) Ethanol and the Developing Brain: Inhibition of Neuronal  
48 Activity and Neuroapoptosis. *Neuroscientist*. doi: 10.1177/1073858417712667.  
49

50  
51 Marszalec W, Aistrup GL, Narahashi T (1998) Ethanol modulation of excitatory and inhibitory  
52 synaptic interactions in cultured cortical neurons. *Alcohol Clin Exp Res.* 22:1516-1524.  
53

54  
55 McMurray MS, Amodeo LR, Roitman JD (2016) Consequences of Adolescent Ethanol  
56 Consumption on Risk Preference and Orbitofrontal Cortex Encoding of Reward.  
57 *Neuropsychopharmacology* 41:1366-1375. doi: 10.1038/npp.2015.288.  
58

59  
60 Moriguchi S, Zhao X, Marszalec W, Yeh JZ, Narahashi T (2007) Effects of ethanol on excitatory  
61 and inhibitory synaptic transmission in rat cortical neurons. *Alcohol Clin Exp Res* 31:89-99.  
62  
63  
64  
65

1 Okamoto T, Harnett MT, Morikawa H (2006) Hyperpolarization-activated cation current (I<sub>h</sub>) is an  
2 ethanol target in midbrain dopamine neurons of mice. *J Neurophysiol* 95:619-626.

3 Olney JW (2014) Focus on apoptosis to decipher how alcohol and many other drugs disrupt brain  
4 development. *Front Pediatr* 2:81. doi: 10.3389/fped.2014.00081.

5  
6  
7 Proctor WR, Soldo BL, Allan AM, Dunwiddie TV (1992) Ethanol enhances synaptically evoked  
8 GABAA receptor-mediated responses in cerebral cortical neurons in rat brain slices. *Brain Res*  
9 595:220-227.

10  
11  
12 Ramaswamy S, Markram H (2015) Anatomy and physiology of the thick-tufted layer 5 pyramidal  
13 neuron. *Front Cell Neurosci* 9:233. doi: 10.3389/fncel.2015.00233

14  
15  
16 Rehm J, Mathers C, Popova S, Thavorncharoensap M, Teerawattananon Y, Patra J (2009) Global  
17 burden of disease and injury and economic cost attributable to alcohol use and alcohol-use  
18 disorders. *Lancet*. 373:2223-2233. doi: 10.1016/S0140-6736(09)60746-7.

19  
20  
21 Sanderson JL, Donald Partridge L, Valenzuela CF (2009) Modulation of GABAergic and  
22 glutamatergic transmission by ethanol in the developing neocortex: an in vitro test of the excessive  
23 inhibition hypothesis of fetal alcohol spectrum disorder. *Neuropharmacology* 56:541-555. doi:  
24 10.1016/j.neuropharm.2008.10.012

25  
26  
27 Sessler FM, Hsu FC, Felder TN, Zhai J, Lin RC, Wieland SJ, Kosobud AE (1998) Effects of  
28 ethanol on rat somatosensory cortical neurons. *Brain Res* 804:266-274.

29  
30  
31 Siggins GR, Pittman QJ, French ED (1987) Effects of ethanol on CA1 and CA3 pyramidal cells in  
32 the hippocampal slice preparation: an intracellular study. *Brain Res* 414:22-34.

33  
34  
35 Smolen TN, Smolen A (1989) Blood and brain ethanol concentrations during absorption and  
36 distribution in long-sleep and short-sleep mice. *Alcohol* 6:33-38.

37  
38  
39 **Spear LP (2016) Consequences of adolescent use of alcohol and other drugs: Studies using rodent  
40 models. *Neurosci Biobehav Rev* 70: 228-243. doi: 10.1016/j.neubiorev.2016.07.026.**

41  
42  
43 Tipps ME, Raybuck JD, Lattal KM (2014). Substance abuse, memory, and post-traumatic stress  
44 disorder. *Neurobiol Learn Mem* 112:87-100. doi: 10.1016/j.nlm.2013.12.002.

45  
46  
47 Trudell JR, Messing RO, Mayfield J, Harris RA (2014) Alcohol dependence: molecular and  
48 behavioral evidence. *Trends Pharmacol Sci* 35:317-3723. doi: 10.1016/j.tips.2014.04.009.

49  
50  
51 Wei W, Faria LC, Mody I (2004) Low ethanol concentrations selectively augment the tonic  
52 inhibition mediated by delta subunit-containing GABAA receptors in hippocampal neurons. *J*  
53 *Neurosci* 24:8379-8382.

54  
55  
56 Weiner JL, Valenzuela CF (2006) Ethanol modulation of GABAergic transmission: the view from  
57 the slice. *Pharmacol Ther* 111:533-354.

58  
59  
60 White AM (2003) What happened? Alcohol, memory blackouts, and the brain. *Alcohol Res Health*  
61 27:186-196.

1 Yan H, Li Q, Fleming R, Madison RD, Wilson WA, Swartzwelder HS (2009) Developmental  
2 sensitivity of hippocampal interneurons to ethanol: involvement of the hyperpolarization-activated  
3 current, *Int J Neurophysiol* 101:67-83. doi: 10.1152/jn.90557.2008.

4 Zorumski CF, Mennerick S, Izumi Y (2014) Acute and chronic effects of ethanol on learning-  
5 related synaptic plasticity. *Alcohol* 48:1–17. doi: 10.1016/j.alcohol.2013.09.045.  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



## Figure legends

1  
2 **Fig. 1.** Staining of dead cells in mouse cortical slices after propidium iodide uptake.  
3 **a-c:** representative microscopy fields showing the incidence of dead cells in control slices (**a**) and  
4 slices treated with ethanol for 3 or 6 min (**b, c**). Note that the nuclei of dead cells are intensely  
5 fluorescent and stand out very neatly over tissue background. Scale bar = 100 $\mu$ m. **d:** graph showing  
6 the result of quantitative analysis of the density (positive nuclei/mm<sup>2</sup>) of dead cells in control and  
7 ethanol-treated slices.  
8  
9  
10  
11  
12  
13

14  
15 **Fig. 2 a-d.** Subthreshold parameters measured before ethanol exposure (preEt), 3 minutes, and 6  
16 minutes after the beginning of ethanol superfusion (3mEt, 6 mEt). RMP: resting membrane  
17 potential; n = 8 cells. \*: P < 0.05. \*\*: P < 0.01. Broken lines beside each graph show the values  
18 recorded from single neurons for each experimental condition.  
19  
20  
21  
22  
23

24 **Fig. 3** The main firing properties measured before ethanol exposure (preEt), 3 minutes, and 6  
25 minutes after the beginning of alcohol superfusion (3mEt, 6mEt). **a:** representative recordings from  
26 the same L5 pyramidal neuron. For all the recordings, both the hyperpolarizing and the depolarizing  
27 current were 100 pA. **b-d:** mean interspike interval (ISI) at rheobase, 50 pA, and 100 pA above  
28 rheobase (where rheobase is that found for each neuron before ethanol application). The symbols  
29 (x) indicate the number of neurons with less than two spikes / train. In these cases the ISI was  
30 approximated to 801 ms (0 spike / train) or to the longest interval between the single spike and the  
31 extreme points of the depolarization envelope. The broken lines beside b show the ISI recorded  
32 from single neurons at rheobase, for each experimental condition. The symbols (#) indicate  
33 recordings with 0 spike / train in two neurons. Inset of **c:** ratio between the last and the first ISI of  
34 the spike train 50 pA above rheobase. **e-h:** the main parameters regarding the action potentials.  
35 AHP = afterhyperpolarization; n = 8 cells. \*: P < 0.05. \*\*: P < 0.01  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 **Fig. 4** Effects of ethanol on mIPSCs. **a:** Representative voltage clamp recordings of mIPSCs from a  
49 L5 pyramidal neuron before ethanol exposure (preEt), and after 3 (3mEt) and 6 minutes (6mEt) of  
50 exposure. **b-c:** Pooled data of mIPSC frequency (n = 4) and amplitude (n = 4). **d:** Averaged mIPSCs  
51 obtained at preEt and at 6mEt from the neuron in **a** and scaled at the peak amplitude for comparing  
52 time courses. **e-f** Pooled data of mIPSC rise time (n = 4) and decay time (n = 4). Broken lines  
53 beside b-c and e-f show the values recorded from single neurons for each experimental condition \*:  
54 P < 0.05.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65





Figure 2

[Click here to download Figure Figure2.eps](#)

