

1 **HABITAT CHARACTERISTICS OF WINTERING WOOD WARBLER (*Phylloscopus***
2 ***sibilatrix*) IN CENTRE REGION OF CAMEROON: CONSERVATION**
3 **IMPLICATIONS**

4
5 **Taku Awa II^{1*}, Tsi A. Evaristus², Robin C. Whytock³, Tsetagho Guilain¹, John Mallord⁴**

6 ¹ University of Dschang, Faculty of Science, Laboratory of Applied Biology and Ecology, P.
7 O. Box 67, Dschang, West Region, Cameroon.

8 ² Department of Fundamental Science, Higher Technical Teacher Training College, P.O. Box
9 39, Bambili, University of Bamenda, North West Region, Cameroon.

10 ³ School of Biological and Environmental Sciences, Natural Sciences, University of Stirling,
11 Stirling FK9 4LA, UK.

12 ⁴ Royal Society for the Protection of Birds, Centre for Conservation Science, RSPB, The
13 Lodge Sandy Beds, SG19 2DL, UK.

14 *Corresponding author: takuawa@yahoo.co.uk

15
16 **This is an Accepted Manuscript of an article published by Taylor & Francis Group in**
17 ***Ostrich: Journal of African Ornithology* on 15 Nov 2017, available online:**

18 <http://www.tandfonline.com/10.2989/00306525.2017.1368037>

19
20 **ABSTRACT**

21
22 Populations of many Afro-Palaearctic birds have declined, with those wintering in sub-Saharan
23 Africa, such as Wood Warbler *Phylloscopus sibilatrix*, particularly affected. In this study we
24 investigated the relationship between habitat characteristics on Wood Warbler
25 presence/absence in Centre Region of Cameroon. A total of six transects were established in
26 three habitat types (forest, forest-savannah transitional zone and savannah). Call playback
27 surveys were conducted monthly from November 2015 to April 2016 to determine Wood
28 Warbler presence / absence. Detailed habitat measurements were also recorded in each
29 transect. A total of 86 responses were recorded: 33 (mean 6.6 ± 2.3) in forest habitat, 47
30 (mean 9.4 ± 3.36) in forest-savannah transitional zone, and 6 (mean 2 ± 1.1) in savannah
31 habitat. Wood Warbler presence increased significantly with the number of trees between 3 -
32 7 m in height, and decreased significantly with the number of shrubs between 0.5 - 3 m in
33 height. Anthropogenic disturbance such as agriculture cycle and burning were not found to
34 have an effect on Wood Warblers presence / absence. We conclude that Wood Warblers

35 overwinter in forested habitats with a relatively low canopy and an open understorey,
36 supporting a similar study in West Africa. Forest clearance in sub-Saharan Africa potentially
37 threatens wintering habitat for Wood Warblers.

38 **Keywords:** Wood Warbler, Wintering habitat, Habitat characteristics

39 INTRODUCTION

40

41 Biodiversity conservation and environmental sustainability have become global priorities that
42 shape international policy (United Nation et al. 2000). Biodiversity faces multiple threats
43 ranging from agricultural expansion, overexploitation, forest fires and habitat fragmentation
44 (Laurance et al. 2009). In many cases, species are particularly vulnerable if their geographic
45 range spans multiple habitats or even continents (Grendelmeier 2011), as is the case with
46 migrant birds.

47

48 There is increasing evidence from Europe suggesting that many migrant birds from the
49 African - Palearctic region (Afro-Palearctic migrants) are undergoing population declines
50 (Evans et al. 2012). This decline is more marked in long distance migrants that overwinter in
51 sub-Saharan Africa and breed in Eurasia than short distance and sedentary species (Heldbjerg
52 and Fox 2008). However, the mechanisms leading to the decline of these species are poorly
53 understood (Evans et al. 2012). Moreover, almost nothing is known of the wintering ecology
54 of many species (Hobson et al. 2014).

55

56 The Wood Warbler (*Phylloscopus sibilatrix*) is one of such migrants that have undergone a
57 striking population decline (PECBMS 2011). Wood warblers breed in Europe and Central
58 Asia, and winter in Equatorial Africa (Urban et al. 2007). Natural and semi-natural
59 broadleaved as well as coniferous woodland is the preferred breeding habitat, with densities
60 highest where Oak *Quercus* is dominant (Wesołowski and Bazaars 2009; Mallord et al.
61 2012a). Wood Warbler populations declined significantly in Europe and between 1990 and
62 2006 (Voříšek et al. 2008), but it remains obscure whether this pattern of change is due to
63 events in the breeding areas or on wintering grounds, or both. Preliminary data from Great
64 Britain (Mallord et al. 2012a, 2012b) and Switzerland (Gerber 2011; Grendelmeier 2011), do
65 not indicate that changes on the breeding grounds are implicated in declines. Thus, events
66 occurring outside the breeding grounds are likely more important (Hobson 2014). Despite
67 this, the information about their ecology on their wintering ground is poorly known,
68 particularly in Central Africa (Evans et al. 2012; Mallord et al. 2016). In a review of the

69 potential factors that are driving migrant bird declines, Vickery et al. (2014) highlighted the
70 urgent need for detailed studies on the wintering grounds to inform conservation management,
71 where there is a paucity of ecological information for most species. Such is the case with
72 Wood Warbler wintering habitat use which is poorly understood, making it difficult to
73 identify potential threats and hindering conservation efforts. In particular, no studies to date
74 have quantified habitat use throughout the entire wintering period, which could change in
75 response to resource availability and environmental cues.

76

77 This paper presents the first systematic study carried out to investigate the relationship
78 between habitat characteristics on Wood Warbler presence/absence in Central Africa. Based
79 on work from West Africa (Mallord et al. 2016), we expected that (1) Wood Warblers would
80 prefer the forest savannah transitional zone, and (2) that anthropogenic activities such as
81 agricultural practices and burning would negatively influence Wood Warblers presence.

82

83 **MATERIALS AND METHODS**

84

85 **Study area**

86 The study was conducted in the Centre Region of Cameroon, at three sites that differed in
87 broad vegetation characteristics identified from satellite imagery and assessments on the
88 ground: Batschenga (4°16'60" N 11°39'0" E) characterized by its forest cover, Nachtigal
89 (4°20'48.01" N 11°38'5.99" E) characterized by its forest savannah transitional zone cover
90 and Ntui (4°27'0"N 11°37'60" E) characterized by its savannah cover (Figure 1). The study
91 area holds a high diversity of floral species such as Ebony (*Diopyros spp*), Sapelli
92 (*Entandrophragma cylindricum*), Moabi (*Baillonella toxisperma*), White Doussie (*Afzelia*
93 *pachyloba*), Ayous (*Triplochyton sceroxylon*) and Fraké (*Terminalia superba*). The area has
94 no legal protection status. Two transects of 2 km of length were established in each of the
95 study sites (forest, forest savannah transitional zone and savannah) by non - random sampling
96 based on homogeneity of habitat.

97

98 **Wood Warblers presence / absence**

99 The data was collected in repeated manner from November 2015 to April 2016 and a day per
100 month was devoted to the visit of each transect. Surveys began at 06:20 am and were
101 completed at 11:00 am; since birds are more active during this period and their detectability is
102 high (Bibby 2000). Along each transect we established 20 listening points at intervals of 100

103 m measured using a handheld GPS. Given that Wood Warbler have a small size (Hobson et
104 al. 2014) and are easily overlooked when not singing or calling, we used call playback
105 (Sutherland et al. 2004) to increase Wood Warbler detectability. At each station along the
106 transect, the advertising call of Wood Warbler (recording taken from Chappuis 2001) was
107 played using an mp3 player connected to a “Radio Shack” mini Amplifier-Speaker,
108 alternating in all directions during 30 seconds (short period to minimize disturbance) and 10
109 seconds (to avoid attracting a bird from outside afar). Presence/absence was recorded on a
110 data sheet alongside the GPS position. This method is described by RSPB / BTO in
111 collaboration with Naturama and Ghana wildlife society, in their migrant project
112 (*unpublished*).

113

114 **Habitat characteristics**

115 Habitat characteristics within 50 m of each point were described with variables referring to
116 habitat structure and anthropogenic activities. Data were collected at all bird survey points,
117 and additionally at 200 m intervals between points. Table 1 below shows the different
118 variables recorded.

119

120 **Statistical analysis**

121 The effects of habitat structure and land management practices on Wood Warbler presence /
122 absence at a listening point were modelled using a Generalised Linear Mixed effects Model
123 (GLMM) with a binomial error structure. There was no multicollinearity between predictors
124 ($r = < 0.6$). Continuous fixed effects in the model were n trees 3-7 m, n trees 7-14 m, n trees >
125 14 m, n shrubs 0.5-3 m, n shrubs 3 – 5 m, vegetation density (mean of four measurements
126 taken at the cardinal points). Burning period, wood removal and agricultural cycle were
127 included in the model as categorical factors. Continuous fixed effects were mean centred and
128 scaled by 1 standard deviation to compare coefficient effect sizes (β). Point ID nested in
129 transect was included as a random intercept to account for pseudo replication. The minimum
130 adequate model was selected using likelihood ratio tests, and 95% confidence intervals were
131 bootstrapped.

132 After selecting the minimum adequate model for habitat structure and land management
133 practices, we then tested if there was a seasonal effect on wood warbler presence/absence.
134 This was done by re-running the minimum adequate model with the addition of month as a
135 fixed effect, and also including a two-way interaction between month and each of the

136 remaining habitat structure / land management variables. All analyses were conducted using R
137 (R Core Team 2014) and the lme4 package (Bates et al. 2014).

138

139 **RESULTS**

140

141 During the six month survey, a total of 86 responses were recorded. Respectively, 33 (mean
142 6.6 ± 2.3) responses were recorded in forest habitat, 47 (mean 9.4 ± 3.36) responses were
143 recorded in the forest-savannah transitional zone, and 6 (mean 2 ± 1.1) responses were
144 recorded in savannah habitat (Figure 2).

145

146 The maximum number of responses were recorded in January, and by April no birds were
147 detected (Figure 3). In other words, few responses were recorded in November and this
148 gradually peaked over time until the maximum number of records in January, with no
149 responses recorded in April.

150

151 **Habitat Preferences**

152 The best minimum adequate GLMM (Table 2) for Wood Warbler presence included only the
153 fixed effects of n trees 3-7 m and n shrubs 0.5-3 m. The probability of Wood Warbler
154 presence was positively correlated with n trees 3-7 m ($\beta = 0.45$, $p = 0.014$; Figure 4), and
155 negatively correlated with n shrubs at 0.5-3 m ($\beta = -0.66$, $p = 0.019$; Figure 5). Based on β
156 effect sizes, n shrubs at 0.5-3 m had a greater influence on Wood Warbler presence. Marginal
157 R^2 for the minimum adequate model was 13%. The number of responses did not vary
158 significantly during the season, and there was no evidence of seasonal changes in habitat
159 preferences based on the interactions tested ($P > 0.05$)

160

161 Conditional modes of the intercept for the random effect of study site (Batschenga, forest;
162 Nachtigal, forest-savannah transition; Ntui, savannah) showed that the probability of detecting
163 Wood Warblers was, on average, greatest in the forest-savannah transition zone (Figure 6).

164

165 The various anthropogenic activities recorded as potential threat on Wood Warbler habitat in
166 the study area are agricultural activities dominated by perennial cultures (100% in forest,
167 33,33% in forest-savannah transitional zone and 83,33% in savannah), pasture (5% present
168 only in forest-savannah transitional), bush fires (55% in forest-savannah transitional zone,

169 45% in savannah and absent in forest), wood removal (40% in forest, 29% in forest-savannah
170 transitional zone, 47% in savannah).

171

172 **DISCUSSION**

173

174 This study is the first to investigate Wood Warbler habitat preferences in Central Africa. The
175 majority of responses were recorded in forest-savannah transitional zone with a greater
176 probability of Wood Warbler detection in its two transects (NAT1 and NAT2) than those of
177 others habitats. Our results are consistent with those of Mallord et al. (2016) in similar study
178 in Burkina Faso and Ghana, and Evans et al. (2012) using stable isotope for declining avian
179 migrants wintering habitat investigation. These could be due to the proximity of our forest-
180 savannah transitional zone (in Nachtigal) to Sanaga River, which provide a micro climate,
181 suitable for abundance of a broad insect's diversity, principal food resources of Wood
182 Warbler. Miguel and Aide (2008) found in Puerto Rico that, Migratory species were mostly
183 abundant in habitats situated near a river. This abundance could be due to the availability of
184 nutritive resources (Lefebvre et al. 1994; Lefebvre and Poulin 1996), characterized by a broad
185 specific insects richness and abundance (Meades et al. 2002), considered as the main food
186 resources of a large number of these migratory species (Russell, 1980; Lefebvre et al. 1994).
187 Also, bioavailability of insects in this habitat could reduce competition between Wood
188 Warbler and others insectivores of the same or different taxa (Moreau, 1952; Miguel and Aide
189 2008). Furthermore, more than one response was recorded at some listening points, mostly in
190 forest and forest-savannah transitional zone habitats. This observation is similar with those
191 found by Mallord et al. (2016) and suggests that, Wood Warblers are not territorial on the
192 wintering grounds, and forms ephemeral intra- and inter-specific feeding flocks (Mallord et
193 al. 2016).

194

195 **Effect of wintering period on Wood Warbler presence/absence**

196 A great number of responses were recorded in January, while no response in April. This
197 means that Wood Warbler wintering population is completely installed in January, and return
198 to Europe by end of March and early April. Also we found no seasonal change in habitat
199 preferences of Wood Warbler during its wintering period. Then we could hypothesize that,
200 this Afro Palearctic birds arrive very quickly in Cameroon, setup a non-breeding territory,
201 stay for five months then quickly depart. This fast departure could be justified by the fact that

202 they should synchronize their reproductive period with the time of prey availability on the
203 breeding grounds (Wesolowski and Maziarz 2009).

204

205 **Effect of Habitat characteristics on Wood Warbler presence/absence**

206 Probability of Wood Warbler detection or presence increased significantly (estimate = 0.45; p
207 = 0.014) with number of trees between 3 - 7 m in height (which were dominant in forest-
208 savannah transitional zone). This result is consistent with those of Mallord et al. (2016) in
209 Burkina Faso and Ghana. This could be due to the fact that Wood Warbler preferred wooded
210 habitat with relative widely-spaced branches to allow movement and foraging (Gerber 2011).
211 Also, spacing between branches requires a certain vegetation height (Glutz von Blotzheim and
212 Bauer 1991). In addition, these trees could provide adequate cover and camouflage,
213 particularly when roosting. Preference for these trees could suggest that Wood Warblers like
214 landscape with open canopy structure which probably increased the physical availability of
215 prey (Mallord et al. 2016). Probability of Wood Warbler detection or presence decreased
216 significantly (estimate = -0.66; p = 0.019) with number of shrubs between 0.5 – 3m in height
217 (which were dominant in savannah habitat). Based on absolute effect sizes, the *n* shrubs 0.5 –
218 3 m has a greater effect than *n* trees 3- 7 m. This result could be justified by the fact that, a
219 high shrub density obstructs Wood Warbler moving or foraging. In addition, Quelle and
220 Lemke (1988); Delahaye and Vandevyvre (2008); Marti (2007); Hillig (2009); Reinhardt and
221 Bauer (2009) found in Wood Warbler breeding ground that high shrubs density limit the
222 establishment of a dense herb layer, which is necessary to conceal Wood Warbler nests from
223 predators. Furthermore, herb cover and herb height were found to have significant influences
224 on Wood Warbler territory choice (Gerber 2011), yet we found that they weren't have
225 significant influence on Wood Warbler presence/absence in this study. This could be due to
226 contrasting habitat requirements on the breeding grounds and during the overwintering period.

227

228 Marginal R^2 for the minimum adequate model was 13%. This means that only 13% of
229 variation in Wood Warbler presence/absence was explained by the most parsimonious model.
230 Suggesting Wood Warbler presence could dependent on other factors that we did not look at.
231 These might include, for example, the availability of food resources or specific tree species.
232 Since Mallord et al. (2016) found a significant correlation between Wood Warbler presence
233 and tree species like *Albizia sp* and *Anogeissus leiocarpus*, this could play an important part in
234 determining Wood Warbler presence that was overlooked during this study.

235

236 Anthropogenic activities such as agriculture are amongst the main causes of deforestation in
237 Africa and greater threats on avifauna (Bobo 2007). Agriculture was found in all sites and was
238 dominated by cocoa land, mostly in Forest habitat in Batchenga following by savannah in
239 Ntui. But we found that activities like agriculture and bush fire were not having significant
240 effect on Wood Warbler presence/absence. These results are similar with those found by
241 Mallord et al. (2016). This suggests that species tolerate farm land holding high density of
242 trees. Nonetheless, Wood Warblers, like other Afro-Paleartic migrants could be vulnerable to
243 woodland loss due to the land-use change in tropical Africa (Mallord et al. 2016).

244

245 Wood Warblers were mostly found in forest-savannah transitional zone habitat, and prefer
246 forested habitats with a relatively low canopy and an open understory which probably favors
247 the species' foraging strategy. Wood Warbler shows fine-scale selection in terms of tree
248 height, and land-use change and forest clearance in sub-Saharan Africa could therefore be
249 contributing to declines by deforestation of wooded land. We suggest future conservation
250 research to focus on investigating changes in Wood Warbler habitat. This study encourages
251 farmers to retain trees on farmland to increase suitable tree cover on farmland required by
252 Wood Warbler. Also regenerate degraded woodland with a diversity of trees which could help
253 the species and others birds. Furthermore, managing these regenerated forests by
254 incorporating crop rotation, edge type forest, crop activity, given that Wood Warbler avoid
255 close forest and may be consider as species that require adoption of wildlife-friendly farming
256 practices that integrate the needs of birds and people.

257

258 **ACKNOWLEDGEMENTS**

259

260 We are grateful to IDEA WILD for equipment grant provided for field work. We are also very
261 thankful to the chiefs of Dji village (at Batchenga), and Nachtigal village for their hospitality
262 and for allowing us to carry out this study in their respective villages. We are also indebted to
263 our field guide, Toa Gregoire, for unmatched field assistance during the survey.

264

265 **REFERENCES**

266

267 Bates D, Maechler M, Bolker B, Walker S. 2014. lme4: Linear mixed-effects models using
268 Eigen and S4. R package version 1.1-7, <URL: [http://CRAN.R-project.org/package =](http://CRAN.R-project.org/package=lme4)
269 [lme4](http://CRAN.R-project.org/package=lme4)> dernière consultation 10-01-2016.

- 270 Bibby C, Jones M, Stuart M (eds). 2000. *Expedition Field Techniques: Bird Surveys*. BirdLife
271 International. Cambridge. 139p
- 272 Bobo KS. 2007. From forest to farmland: Effects of land use on understory birds of
273 Afrotropical rainforests. PhD thesis, University of Göttingen. 195 p.
- 274 Chappuis C. 2001. African Bird Sounds. CD set. Claude Chappuis & British Library,
- 275 Delahaye L, Vandevyvre X. 2008. Le Pouillot siffleur (*Phylloscopus sibilatrix*) est-il une
276 espèce indicatrice de la qualité des forêts feuillues ardennaises ?. *Aves* 45: 3-14.
- 277 Evans KL, Newton J, Mallord JW, Markman S. 2012. Stable isotope analysis provides new
278 information on winter habitat use of declining avian migrants that is relevant to their
279 conservation. *Plos one* 7: e34542. doi:10.1371/journal.pone.0034542
- 280 Gerber M. 2011. Territory choice of the Wood Warbler *Phylloscopus sibilatrix* in Switzerland
281 in relation to habitat structure and rodent density. Master thesis, Institute of evolutionary
282 biology and environmental studies, University of Zürich, Switzerland. 27 p
- 283 Glutz von Blotzheim UN, Bauer KM. 1991. Handbuch der Vögel Mitteleuropas
284 Passeriformes (3. Teil): Sylviidae (Grasmücken, Laubsänger, Goldhähnchen). AULA-
285 Verlag GmbH.
- 286 Gregory RD, van Strien A, Vorisek P, Gmelig Meyling AW, Noble DG, Foppen RP, Gibbons
287 DW. 2005. Developing indicators for European birds. *Philosophical Transactions of the
288 Royal Society B* 360: 269-288.
- 289 Grendelmeier A. 2011. Breeding success and nest predation in the declining Wood Warbler
290 *Phylloscopus sibilatrix*. Master thesis, University of Bern, Switzerland. 108 p.
- 291 Heldbjerg H, Fox TAD. 2008. Long-term population declines in Danish trans-Saharan
292 migrant birds. *Bird Study* 55: 267–279.
- 293 Hillig F. 2009. Verursachen Veränderungen im Brutgebiet den Bestandsrückgang des
294 Waldlaubsänger *Phylloscopus sibilatrix*? Eine Untersuchung im Schwalm-Eder Kreis
295 (Hessen) unter Berücksichtigung von Bruterfolg und Habitatveränderung. Diploma /
296 Master, Fachhochschule Osnabrück.
- 297 Hobson KA, Van Wilgenburg SL, Wesolowski T, Maziarz M, Bijlsma RG, Grendelmeier A,
298 Mallord JW. 2014. A multi-isotope ($\delta^2\text{H}$, $\delta^{13}\text{C}$, $\delta^{15}\text{N}$) approach to establishing
299 migratory connectivity in Palearctic-Afrotropical migrants: An example using Wood
300 Warblers *Phylloscopus sibilatrix*. *Acta Ornithologica* 49: 57–69.
- 301 Laurance WF, Goosem M, Laurance SGW. 2009. Impacts of roads and linear clearings on
302 tropical forests. *Trends in Ecology and Evolution* 24: 659–669.
- 303 Lefebvre G, Poulin B. 1996. Seasonal abundance of migrant birds and food resources in
304 Panama-nian mangrove forests. *Wilson Bulletin* 108: 748-759.
- 305 Lefebvre GB, Poulin McNeil R. 1994. Temporal dynamics of mangrove bird communities in
306 Venezuela with special reference to migrant warblers. *The Auk* 111: 405-415.
- 307 Mallord JW, Orsman CJ, Roberts JT, Skeen R, Sheehan DK, Vickery JA. 2016: Habitat use
308 and tree selection of a declining Afro-Palearctic migrant at sub-Saharan staging and
309 wintering sites. *Bird Study* 63: 459-469.
- 310 Mallord JW, Charman EC, Cristinacce A, Orsman CJ. 2012a. Habitat associations of Wood
311 Warblers *Phylloscopus sibilatrix* breeding in Welsh oakwoods. *Bird Study* 59: 403–415.

312 Mallord JW, Orsma CJ, Cristinacce A, Butcher N, Stowe TJ, Charman EC. 2012b. Mortality
313 of Wood Warbler *Phylloscopus sibilatrix* nests in Welsh Oakwoods: predation rates and
314 the identification of nest predators using miniature nest cameras. *Bird Study* 59: 286–
315 295.

316 Meades LL, Rodgerson AY, French K. 2002. Assessment of the diversity and abundance of
317 terrestrial mangrove arthropods in southern New South Wales, Australia. *Austral*
318 *Ecology* 27: 451-458.

319 Miguel AA, Mitchell AT. 2008. Bird community dynamics and habitat associations in karst,
320 mangrove and pterocarpus forest fragments in an urban zone in Puerto Rico. *Caribbean*
321 *Journal of Science* 44: 402-416.

322 Moreau RE. 1952. The place of Africa in the Palearctic migration system. *Journal of Animal*
323 *Ecology* 21: 250-271.

324 PECBMS 2011. Trends of common birds in Europe, 2011 update. European Bird Census
325 Council, Prague. <http://www.ebcc.info/index.php?ID=457> (accessed 24 January 2012).

326 Quelle M, Lemke W. 1988. Strukturanalyse von Waldlaubsängerrevieren. *Charadrius* 24: 196
327 – 213.

328 R Core Team 2014. R: A language and environment for statistical computing. R Foundation
329 for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.

330 Russell SM. 1980. Distribution and abundance of North American migrants in lowlands of
331 Northern Colombia. In Keast A, Morton ES (eds), *Migrant birds in the Neotropics:*
332 *Ecology, behavior, distribution and conservation*. Smithsonian Institution Press,
333 Washington, DC. pp 249-252.

334 Sutherland WJ, Newton I, Green RE (eds). 2004. *Bird Ecology and Conservation: A*
335 *Handbook of Techniques (Techniques in Ecology and Conservation)*. Oxford University
336 Press.

337 UN, OECD, IMF, WBG. 2000. A better world for all: International Development Goals.
338 *Washington DC , US*.

339 Urban EK, Fry CH, Keith S (eds). 1997. *The Birds of Africa*, vol. 5. London: Academic Press.

340 Vickery JA, Ewing SR, Smith KW, Pain DJ, Bairlein F, Škorpiłová J, Gregory RD. 2014. The
341 decline of Afro-Palearctic migrants and an assessment of potential causes. *Ibis* 156: 1–
342 22.

343 Voříšek P, Škorpiłová J, Klvaňová A. 2008. Trends of common birds in Europe, 2008 update.
344 Downloaded from [www.ebcc.info/index.php? ID=358](http://www.ebcc.info/index.php?ID=358) on 2. 11. 2008, 6:00.

345 Wesołowski T, Maziarz M. 2009. Changes in breeding phenology and performance of Wood
346 Warbler *Phylloscopus sibilatrix* in a primeval forest: a thirty-year perspective. *Acta*
347 *Ornithologica* 44: 69–80.

348
349
350
351
352
353
354

355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372

TABLES

Table 1: Variable names and descriptions recorded in each of the three habitats (forest, forest-savannah transitional zone and savannah).

Variable	Description
Dominant land cover	Dominant vegetation: dense forest, open forest, agricultural/natural mosaic, others (arable, bare ground, grassland, plantation, shrub land)
Tree cover	Tree cover: 1 = 0%, 2 = 1 – 4%, 3 = 4 – 15%, 4 = 15 – 40%, 5 = 40 – 65%, 6 = >65%
Tree 3 – 7 m	Number of trees
Tree 7 – 14 m	Number of trees
Tree > 14 m	Number of trees
Shrub cover	Shrub cover: 1 = 0%, 2 = 1 – 4%, 3 = 4 – 15%, 4 = 15 – 40%, 5 = 40 – 65%, 6 = >65%
Shrub 0.5 – 3 m	Number of shrubs
Shrub 3 – 5 m	Number of shrubs
Grass cover	1 = 0%, 2 = 1 – 4%, 3 = 4 – 15%, 4 = 15 – 40%, 5 = 40 – 65%, 6 = >65%
Grass 0.03 – 0.3 m	Grass with height of 0.03 – 0.3 m
Grass 0.3 – 1 m	Grass with height of 0.3 – 1 m
Grass 1 – 3 m	Grass with height of 1 – 3 m
Wood	Evidence of wood removal (whole trees, branches or for charcoal): yes/no

Burn	Evidence of burning, either in this year, the previous year or not at all: yes/no
Agriculture	Evidence of agriculture cycle: 1 = annual, 2 = perennial, 3 = 2 crops / year.
Vegetation density	Mean of four measurements taken at the cardinal points.

373
374
375
376
377
378
379

380 **Table 2:** Habitat characteristics explaining Wood Warbler presence/absence from the best
381 GLMM. β coefficients are given on the link scale (odds-ratios)

Variable	β	-95% CI	+95% CI	P
Intercept	-2.57	-3.45	-1.97	< 0.001
n trees 3-7m	0.45	0.04	0.84	0.014
n shrubs 0.5-3 m	-0.66	-1.31	-0.13	0.019

386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404

405
406
407
408
409
410
411
412
413
414
415
416
417
418
419

420 **FIGURE LEGENDS**

421

422 **Figure 1:** Map of study area and study site in Centre Region of Cameroon.

423 **Figure 2:** Average number of responses per habitat

424 **Figure 3:** Number of responses recorded per month surveyed

425 **Figure 4:** Fitted values (red line +/- 95% CI) from the top GLMM explaining Wood Warbler
426 presence showing the effect of n trees 3-7 m on Wood Warbler presence at a survey point.


427 The histograms show the frequency of presence/absence for a given value on the x-axis.

428 **Figure 5:** Fitted values (red line +/- 95% CI) from the top GLMM explaining Wood Warbler
429 presence showing the effect of n shrubs 0.5 - 3 m on Wood Warbler presence at a survey
430 point. The histograms show the frequency of presence/absence for a given value on the x-axis.

431 **Figure 6:** Conditional modes of the intercept (+/- 95% CI) for the random effect of transect
432 nested in study site extracted from the minimum adequate GLMM explaining Wood Warbler
433 presence. Study site abbreviations are NAT: Nacthigal, BAT: Batschenga and NTUI: Ntui.

434


435 **FIGURES**



436

437 **Figure 1:** Map of study area and study site in Centre Region of Cameroon.


438



439

440 **Figure 2:** Average number of responses per habitat


441



442

443 **Figure 3:** Number of responses recorded per month surveyed

444




445

446 **Figure 4:** Fitted values (red line +/- 95% CI) from the top GLMM explaining Wood Warbler


447 presence showing the effect of n trees 3-7 m on Wood Warbler presence at a survey point.

448 The histograms show the frequency of presence/absence for a given value on the x-axis.



449
450
451
452
453

Figure 5: Fitted values (red line +/- 95% CI) from the top GLMM explaining Wood Warbler presence showing the effect of n shrubs 0.5 - 3 m on Wood Warbler presence at a survey point. The histograms show the frequency of presence/absence for a given value on the x-axis.



454
455
456
457
458

Figure 6: Conditional modes of the intercept (+/- 95% CI) for the random effect of transect nested in study site extracted from the minimum adequate GLMM explaining Wood Warbler presence. Study site abbreviations are NAT: Nacthigal, BAT: Batschenga and NTUI: Ntui.