

Age associated blood pressure distributions in pediatric intensive care units differ from healthy children.

Samer Abdelrazeq¹, Samiran Ray^{1,2}, Libby Rogers³, David P Noren⁴, Mark J Peters^{1,2}, David P Inwald⁵

Author affiliations:

¹ Paediatric Intensive Care Unit, Great Ormond Street Hospital NHS Trust, London, UK

² Respiratory, Critical Care and Anaesthesia Section, UCL GOS Institute of Child Health, London, UK

³ UCL Clinical Operations Research Unit, University College London, London, UK

⁴ Philips Research North America, Cambridge, MA 02141, USA

⁵ Paediatric Intensive Care Unit, St Mary's Hospital, Imperial College Healthcare NHS Trust, London, UK

Corresponding author:

Samiran Ray

Respiratory, Critical Care & Anaesthesia Section, UCL GOS Institute of Child Health, 30 Guildford Street, London, UK WC1N 1EH

Phone: 02074059200 ext 0032

Fax: 02078138206

Reprints not requested

Funding information: No direct funding was received for this study. Part of this work was undertaken at UCL GOS Institute of Child Health, which received a proportion of funding from the Department of Health's NIHR Biomedical Research Centre's funding scheme.

Keywords: blood pressure; children; monitoring; critical care

Normal blood pressure (BP) centiles for age and height in children have been derived from healthy children by the US NIH Task Force (1,2). Restoration of BP towards healthy normal values in the resuscitation phase of childhood critical illness has been associated with an outcome benefit (3). However, paediatric intensive care unit (PICU) admission systolic BP has a U-shaped association with mortality (4). Current studies in critically-ill adults are exploring a more permissive approach to BP (5). Description of the distribution of BP in critically-ill children is an essential step towards similar work.

We aimed to describe the systolic and mean BP distribution according to sex and age in children admitted to two PICUs in London, UK. Electronic health records were interrogated to yield all non-invasive oscillometric blood pressure measurements (Intellivue, Philips, Netherlands) in children admitted to the general (non-cardiac) PICUs at Great Ormond Street Hospital (between 2012-2016) and St Mary's Hospital (between 2009-2015). Only non-invasive measurements were considered because (a) all children underwent non-invasive measurement, and (b) the NIH normal values were derived from non-invasive measurements. Systolic measurements <30 mmHg and >250 mmHg, and mean measurements <16 mmHg and >216 mmHg were excluded as likely spurious.

To account for multiple measurements in each patient, multi-level quantile regression analysis was used, with blood pressure as the dependent variable, age in years as the fixed effect variable and the unique patient identifier as the random effects variable. The 50th centile for each year of age were compared with the 50th centile from the US Task Force NIH data: as height data were not routinely available for children in our cohort, the 50th height centile values were used for comparison. The values were considered significantly different ($p < 0.05$) if the 50th centile value from the Task Force NIH data lay outwith the 95% confidence interval for the 50th centile value from our cohort. Data were analysed using r (www.cran.r-project.org).

Systolic BP centiles were generated from 309520 measurements from 2415 boys and 251877 measurements from 1946 girls; mean BP centiles from 313492 measurements from 2413 boys and 256662 measurements from 1944 girls ~~(Figure)~~. Following multi-level quantile regression for the 50th centile, the PICU 50th centile values for systolic BP were significantly higher for boys between 1 and 9 years, and girls between 2 and 8 years compared to the ~~Task Force~~NIH 50th centile ~~(Figure)~~e. The PICU 50th centile values for mean BP were significantly higher for boys between 1 and 3 years, and girls aged 2 years and 4 years (Electronic Supplementary Material [1, 2](#)).

Compared to population-based normal values, younger children in our cohort had higher observed BP measurements. ~~It is not clear if this is due to therapeutic intervention, varying levels of sedation or individual disease pathophysiology. However, the therapeutic benefit of such supra-normal values has not been demonstrated~~This may not be therapeutically driven, but a consequence of levels of sedation, or disease pathophysiology (Electronic Supplementary Material 3). The frequency of measurements is likely to be biased by haemodynamic status. Our data suggestRegardless, we demonstrate uncertainty regarding optimal BP targets in ~~PICU~~critically-ill children. Prospective trials examining the effect of different BP targets on outcome in children in intensive care, similar to the adult SEPSISPAM trial (5), are needed to guide practice.

Ethical approval and consent to participate: Ethical approval was not sought as we used routinely collected clinical data retrospectively. Consent to participate was not sought as this is a retrospective observational study and no patient identifiable data are presented.

Consent to publish: Consent was not sought as no patient identifiable data are reported.

Availability of data and materials: The datasets used and/or analysed during the current study available from the corresponding author on reasonable request.

Competing interest: The authors declare that they have no conflict of interest.

Author's contributions: SA, SR, MJP, and DPI conceived and designed the study; SA, LR, SR and DPN collected and verified the data; SR and LR analysed the data. SA and SR drafted the manuscript, all authors contributed to and approved the final version of the manuscript.

Acknowledgements: Not applicable

Authors' information: DPI is the chief investigator and MJP is a co-investigator on the UK NIHR funded Fluids in Shock (FiSh) external pilot trial (IRAS 195544)

References

1. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. National Institutes of Health, National Heart, Lung, and Blood Institute, NIH Publication No. 05-5267, May 2005
2. Haque IU, Zaritsky AL. Analysis of the evidence for the lower limit of systolic and mean arterial pressure in children. *Pediatr Crit Care Med.* 2007;8:138–144
3. Han YY, Carcillo JA, Dragotta MA, Bills DM, Watson RS, Westerman ME, Orr RA. Early reversal of pediatric-neonatal septic shock by community physicians is associated with improved outcome. *Pediatrics.* 2003 Oct;112(4):793-9.
4. Straney L, Clements A, Parslow RC, Pearson G, Shann F, Alexander J, Slater A; ANZICS Paediatric Study Group and the Paediatric Intensive Care Audit Network. Paediatric index of mortality 3: an updated model for predicting mortality in pediatric intensive care. *Pediatr Crit Care Med.* 2013 Sep;14(7):673-81.
5. Asfar P, Meziani F, Hamel JF, Grelon F, Megarbane B, Anguel N, Mira JP, Dequin PF, Gergaud S, Weiss N, Legay F, Le Tulzo Y, Conrad M, Robert R, Gonzalez F, Guitton C, Tamion F, Tonnelier JM, Guezennec P, Van Der Linden T, Vieillard-Baron A, Mariotte E, Pradel G, Lesieur O, Ricard JD, Hervé F, du Cheyron D, Guerin C, Mercat A, Teboul JL, Radermacher P; SEPSISPAM Investigators: High versus low blood-pressure target in patients with septic shock. *N Engl J Med.* 370(17):1583-93, 2014

Figure: Systolic non-invasive blood pressure distributions of children in 2 PICUs. The blue line in the top left panel shows the 50th centile of 309520 systolic blood pressure measurements from 2415 boys according to age. The blue shaded area represents the area between the 5th and 95th centiles. The red lines show the population 5th, 50th and 95th centile values for boys according to age, for the 50th centile for height (as described by the NIH Task Force, and Haque and Zaritsky). The green line in the top right panel shows the 50th centile of 251877 systolic blood pressure measurements from 1946 girls according to age. The green shaded area represents the area between the 5th and 95th centiles. The red lines show the population 5th, 50th and 95th centile values for girls according to age, for the 50th centile for height (as described by the NIH Task Force, and Haque and Zaritsky).

The bottom panels show the 50th centile systolic blood pressure values for each age group following multi-level quantile regression, with 95% confidence intervals for boys (blue) and girls (green). Systolic blood pressure in our cohort is significantly higher than NIH Task Force 50th centiles in boys between 1 and 9 years and girls 2 and 8 years of age.

~~**Figure:** Systolic and mean non-invasive blood pressure distributions of children in 2 PICUs. The blue line in the top left panel shows the 50th centile of 309520 systolic blood pressure measurements from 2415 boys according to age. The blue shaded area represents the area between the 5th and 95th centiles. The red lines show the population 5th, 50th and 95th centile values for boys according to age, for the 50th centile for height (as described by the NIH Task Force, and Haque and Zaritsky). The green line in the top right panel shows the 50th centile of 251877 systolic blood pressure measurements from 1946 girls according to age. The green shaded area represents the area between the 5th and 95th centiles. The red lines show the population 5th, 50th and 95th centile values for girls according to age, for the 50th centile for height (as described by the NIH Task Force, and Haque and Zaritsky).~~

~~The bottom panels show the corresponding mean blood pressure distributions, based on 313492 measurements from 2413 boys (left), and 256662 measurements from 1944 girls (right). The red~~

lines show the population 5th, 50th and 95th centile values according to age for 50th centile for height (as described by Haque and Zaritsky).