

# Review of South African research on volcanic and related rocks and mantle-derived materials: 1999–2002

J.S. Marsh\*

---

**This report reviews South African research relating to the scientific interests of the International Association of Volcanology and Chemistry of the Earth's Interior (IAVCEI) and which was published between 1999 and 2002. The focus is on published work and does not include conference presentations and abstract volumes or other informal documents.**

---

As a previous National Correspondent<sup>1</sup> has noted, it is not easy to determine what precisely is covered by 'South African research' and the 'scientific interests of IAVCEI'. In compiling this report, one approach I could have adopted was to include all research of any igneous or volcanic flavour. Another was to aim at comprehensive coverage but to select on the basis of research results which I perceived to be of interest to the broader IAVCEI membership. As an example, consider an Archaean pyroclastic deposit. A paper focused on the description and emplacement of the deposit would clearly fall within the interests of IAVCEI. If the deposit has been tectonized and metamorphosed and is mentioned briefly in a paper whose main focus is the dating of zircons recovered from the unit, then it is legitimate to consider the paper to have little relevance to global IAVCEI membership. However, the timing of volcanism and associated igneous activity in relation to tectonism is of considerable interest to our understanding of temporal trends in global volcanism and tectonics, a topic falling within the broad interests of IAVCEI members and, on balance, might be included in the report to IAVCEI. In this context, this review does report on papers concerned with dating of volcanic and igneous rocks.

Apart from research on volcanic rocks and associated intrusions, this article also reports on inclusions of deep origin contained in kimberlite and other rare alkaline rocks, as they are a rich source of information on the chemistry of the Earth's interior. Work carried out by South African-based scientists in contiguous countries as well as in the oceanic environment surrounding South Africa is also covered. I do not report research on South African rocks carried out exclusively by scientists based in other countries, as this work is expected to be reflected by the reports from the National Correspondent of those countries. Much of such work is, however, published in collaboration with South African scientists and is included here.

Most of the material in the report is organized according to geological time periods starting with the Archaean. The exceptions are carbonatitic and kimberlitic rocks and associated materials, which are discussed together regardless of age. Some published research has probably been inadvertently omitted, but for some the omission is on the basis of a judgement of

relevance. Such omissions are minor and do not detract from the overall impact of this review as a statement of the research achievements of South African scientists from 1999 to 2002.

## Archaean greenstones, komatiites, and granitoid rocks

These are complex entities involving volcanism and plutonism and allied structural deformation and metamorphism. Much of the research carried out is by necessity of a multidisciplinary nature, including age dating.

The 3.48 Gyr Komati Formation at Barberton was mapped by Dann<sup>2</sup> on a scale of 1:5000, with special attention to detailed mapping of chill-bounded cooling units and to defining volcanic, intrusive and tectonic contacts. This mapping allowed the magmatic architecture of the Komati Formation to be established. The formation consists of a lower sequence of komatiitic sheet flows emplaced in a lava-plain setting, followed by pillowed komatiites emplaced in a more irregular topographic environment, possibly caused by faulting. The sequence is intruded by pyroxene-spinifex-textured sills, but the wherlite dykes are younger than the Komati and overlying Hoogenoeg Formations. Thus the komatiites do not have a feeder dyke system analogous to post-Archaean ophiolites. Cloete<sup>3</sup> also gives a detailed account of the volcanology and geochemistry of the komatiitic eruptives and concluded that they have attributes akin to modern oceanic plateaux rather than typical mid-ocean ridge crust. His work is also concerned with primary sea-floor alteration of the komatiites.

Anhaeusser<sup>4</sup> argues that the more metamorphosed komatiites and komatiitic basalts of the Nelshoogte Schist Belt and associated intrusions represent a younger (*c.* 3250 Myr) ultramafic volcanic event and he rejects previous proposals that these rocks are similar to Phanerozoic ophiolites.

A number of studies report single zircon ages for a variety of granitic rocks in the greenstone belts. These ages are important in providing a temporal framework for the evolution of the belts. Hence, Poujol and Robb<sup>5</sup> demonstrate that the granitic magmatism was contemporaneous with the deposition of the 3.09–2.97 Gyr Murchison greenstone belt and extended to younger ages of 2.85 Gyr. This was subsequently extended to an even younger age of 2680 Myr.<sup>6</sup> In the Pietersburg greenstone terrane felsic volcanism is dated at  $2949.7 \pm 0.2$  Myr,<sup>7</sup> indicating that it is coeval with felsic volcanism in the Murchison belt to the east. A granite intruding the greenstone is dated at  $2853 \pm 18$  Myr. In addition, the associated Turfloop Batholith was emplaced at about 2.78 Gyr on the basis of U-Pb, Sm-Nd and Rb-Sr dating<sup>8</sup> and has an origin in the lower Archaean crust. In the Amalia-Kraaipan greenstone belt in the western Kaapvaal craton, the earliest tonalites and trondhjemites was reported as varying in age from  $3162 \pm 8$  to  $3070 \pm 7$  Myr and the youngest pluton, the Mosita adamellite, was dated at  $2749 \pm 3$  Myr.<sup>9</sup> In the same area, Poujol *et al.*<sup>10</sup> dated granite magmatism varying from *c.* 3008 Myr (trondhjemitic) to 2791 Myr (granodioritic). These new results suggest a temporal correlation of the youngest

\*The author is South African National Correspondent for the International Association of Volcanology and Chemistry of the Earth's Interior, and is in the Department of Geology, Rhodes University, Grahamstown 6140, South Africa. E-mail: goonie.marsh@ru.ac.za

granitoid activity with the emplacement of the Gaborone Granite Complex in Botswana. In the Giyani greenstone belt, meta andesites have yielded an age of  $3203.3 \pm 0.2$  Myr and younger intrusive quartz porphyries have an age of  $2874.1 \pm 0.2$  Myr.<sup>7</sup> In the Johannesburg Dome a variety of trondhjemitic and granitic rocks yield single zircon U/Pb ages from  $3340 \pm 3$  Myr in the north to  $3114 \pm 2.3$  Myr in the south.<sup>11</sup> Similar age ranges were obtained using a variety of dating techniques by Barton *et al.*<sup>12</sup>

### Archaean supracrustal sequences

The oldest supracrustal sequence in the Kaapvaal craton is the Pongola Supergroup with a lower Nsuze Group of volcanic and sedimentary rocks and an overlying siliclastic Mozaan Group. Gutzmer *et al.*<sup>13</sup> report an age of  $2837 \pm 5$  Myr for quartz porphyry sill intruded into these sediments and deformed with them.

The Late Archaean Ventersdorp Supergroup continues to attract attention. It consists of basal formations of mafic lava overlain by a clastic wedge deposit, the Kameeldoorns Formation, which formed in grabens. Intermediate to felsic ash-flow deposits of the Makwassie and Goedgenoeg formations overlie these sediments. Large spherical structures and clasts have been described at the base of the Makwassie Formation at T'kuip in the Northern Cape Province<sup>14</sup> and are ascribed to eruption of the hot ash-flow onto water-saturated sediments. De Bruijn *et al.*<sup>15</sup> also report on the geochemistry of alteration of andesitic and basaltic andesite lavas of the uppermost Allanridge Formation. Such studies are important for effective use of geochemistry in petrogenetic studies on these old sequences. Hall and Els<sup>16</sup> report on lava-soft sediment interactions at the base of the Ventersdorp Supergroup. Features such as sediment injection into lava, 'ball and pillow' structures in lava, and soft sediment deformation are described. These features developed during dewatering of the sediments in response to their differential loading by the overlying lava.

### Palaeoproterozoic

Oberholzer and Eriksson<sup>17</sup> described the volcanic sequence in the Palaeoproterozoic Hekpoort Formation of the Transvaal Supergroup as consisting of equal volumes of basaltic-andesite lava flows and intercalated with volcanoclastic rocks emplaced in a subaerial intracratonic setting. The volcanoclastic rocks are thought to represent a variety of pyroclastic flow and lahatic deposits. In a paper which is significant for correlation of the Kheis and Magondi mobile belts as components of an extensive Palaeoproterozoic orogen in southern Africa, McCourt *et al.*<sup>18</sup> report an age of  $1997.5 \pm 2.6$  Myr for the syn-kinematic Hurungwe granite in Zimbabwe. This age invalidates the Kheis-Magondi correlation.

### The Bushveld Magmatic Province

The Bushveld Complex is purported to be the largest continental layered mafic-ultramafic intrusion (Rustenberg Layered Suite) and is associated with coeval mafic and silicic volcanic suites (the Rooiberg Group) and intrusive granites (Lebowa Granite Suite). Collectively these suites form the Bushveld Magmatic Province. The enormous metal reserves [platinum-group elements (PGEs), Cr, Ti and Fe] in the complex has ensured that it is the target of much research. The *South African Journal of Geology* published two special issues largely relating to the Bushveld Complex. The first, edited by Cawthorn,<sup>19</sup> commemorated the 75th anniversary of the discovery of the Merensky Reef, and the second, edited by Maier,<sup>20</sup> was directed at platinum-group minerals and elements. Not all of the papers in these two

volumes fall within the interests of IAVCEI

Bushveld magmatism was initiated by eruption of basaltic andesites and felsic volcanic rocks of the Dullstroom Formation, the lowest formation of the Rooiberg Group. Buchanan *et al.*<sup>21</sup> demonstrate that the mafic rocks can be divided into a high- and a low-Ti lineage with strong compositional similarities to the Mesozoic Karoo flood basalt geochemical types. They suggest that a mantle plume was responsible for the mafic volcanism and present petrogenetic models involving magma mixing and assimilation to account for the compositional variability. Associated high-Mg felsic rocks in the Dullstroom Formation, as well as siliceous to intermediate rocks in the upper part of the Rooiberg Group, are derived from the low-Ti mafic magmas by assimilation of crustal material and fractional crystallization in shallow magma chambers.<sup>22</sup>

Maier *et al.*<sup>23</sup> use the PGE geochemistry of the ultramafic Pyroxenite Marker within the mafic Main Zone of the Bushveld Complex to constrain a model for the formation of this zone. In a study using mineral composition reversal and whole-rock compositional trends within the vicinity of the Pyroxenite Marker, Nex *et al.*<sup>24</sup> examined the processes of magma addition to the Main Zone. The transition from the Critical to Main zones is believed to exist in the vicinity of the Giant Mottled Anorthosite but has not been pinpointed precisely. Mitchell and Manthre<sup>25</sup> investigated this problem and developed a model for the emplacement of the Main Zone.

The Merensky Reef in the top of the Critical Zone is arguably the most famous horizon within the Bushveld Complex. A number of papers<sup>26-32</sup> address problems of geology, mineralogy, and geochemistry of the reef and discuss implications for its origin. The development of cyclic units in the Critical Zone below the Merensky Reef is ascribed by Cawthorn<sup>33</sup> to crystal sorting during settling.

In broader studies, Maier and Barnes<sup>34</sup> analysed for PGEs in a wide range of silicate rocks through the complex in an attempt to constrain ore-forming processes within the intrusion. Eales<sup>35</sup> examined the Cr budget in the western Bushveld and demonstrated that the amount of chromite in the rocks exceeds the Cr solubility of mafic magmas and suggests that the magmas emplaced into the Bushveld chamber were carrying up to 3% chromite as microphenocrysts. Willmore *et al.*<sup>36,37</sup> have investigated halogen geochemistry of the Bushveld and identify correlations of Cl/F ratio with a number of geochemical and mineralogical trends through the complex. Bushveld magmas appear to have been unusually enriched in Cl, and all evidence points to the halogens being a primary magmatic component and not derived by assimilation of, or infiltration from, country rocks. Separation of Cl-rich fluids have played a role in mineralization in the Lower and Critical zones. This is supported by chlorine isotopes. Maier *et al.*<sup>38</sup> report on a wide-ranging Nd isotope study of the Bushveld Complex, which supports older Sr isotope data indicating a large crustal component in the upper part of the intrusion. The isotope data are decoupled from highly incompatible element concentrations. This is interpreted in terms of changes in the nature of the crustal assimilant with evolution of the complex. Contamination by crust, including unusual compositions such as dolomite, are also indicated in the work of Harris and Chaumba<sup>39</sup> on the Platreef in the Northern Limb of the Bushveld Complex.

A structural study relating to mechanisms of intrusion of the Rustenberg Layered Suite along its southwestern margin (Spruitfontein Inlier) was documented by Clarke *et al.*<sup>40</sup>

Discordant ultramafic iron-rich pegmatoidal bodies are common in the Rustenberg Layered Suite and their origin and petrogen-

esis is controversial. A detailed study<sup>41</sup> on the Tweefontein pipe indicated that it was magmatically intruded when the layered rocks were still extremely hot and that it is a magmatic as opposed to a metasomatic feature. The pipe magmas are not residual liquids derived from adjacent layered rock, but distinct magma batches in their own right. Reid and Basson<sup>42</sup> focused on similar bodies replacing the Merensky Reef at Northam Platinum Mine. They concluded that the pegmatoid bodies result from replacement of pre-existing rocks by residual melts migrating from the upper Critical Zone. Scoon and Eales<sup>43</sup> showed that spinels in the pegmatoids can be divided into three types and that there is a relationship between spinel type and stratigraphic height. The composition of the Fe-Ti-Cr spinels is not duplicated by cumulus spinels in the layered rocks but the disseminated Ti-magnetites are very similar to that found in the Upper Zone.

Minor intrusions associated with the Bushveld Complex include the Uitkomst Complex and numerous mafic-ultramafic sills in the footwall of the Bushveld Complex. Using olivine compositions and sulphur isotopes, Li *et al.*<sup>44</sup> present a model for multiple intrusion emplacement of the Uitkomst Complex which is consistent with its being a conduit to the Bushveld Complex as proposed by Gauert.<sup>45</sup> Age and geochemical similarities between the Uitkomst Complex and a diorite intrusive into the Marble Hall fragment suggest that there is a genetic link between the two and led De Waal and Armstrong<sup>46</sup> to define a new magma type (Bu-type) which preceded the emplacement of the Bushveld B1-type magma and which may also have been significant in a number of sub-Rustenberg Layered Suite intrusions such as the Lindequesdrift, Roodekraal, and Rietfontein complexes. The nature of the Bushveld magmas is also discussed by Eales.<sup>47</sup> Maier *et al.*<sup>48</sup> discuss PGE mineralization of ultramafic footwall sills exposed around the eastern margin of the complex.

Maier and Barnes<sup>49</sup> investigated the origin of the Cu sulphide deposits associated with mafic-ultramafic intrusive bodies, 2.0–2.3 Gyr in age, in the Curaçá Valley, Brazil. These intrusions have been emplaced in a high-grade metamorphic terrane and as such resemble the O'okiep deposits in western South Africa.

Although it is strictly an Archaean intrusion, research on the Great Dyke of Zimbabwe is reported here because of its similarities to the Bushveld Complex. Wilson and Prendergast<sup>50</sup> reviewed the magma evolution and magma chamber structure of the Great Dyke with emphasis on implications for PGE mineralization. In more detail Wilson<sup>51</sup> and Wilson *et al.*<sup>52</sup> reported on the geology, mineralogy and geochemistry of the Selukwe subchamber of the Great Dyke. The zone of PGE enrichment is associated with sharp compositional changes in orthopyroxene, and the layered subzones, characterized by different PGE contents, may reflect original liquid layering in the chamber.

### Mesoproterozoic

Research results are of two types; they relate either to petrogenetic studies on igneous intrusions and volcanic sequences and dating of intrusive and extrusive rocks with the aim to constrain correlations, or to the tectonic and metamorphic evolution of mobile belts. In the latter category fall the work of Gutzmer *et al.*<sup>53</sup> on the Koras bimodal volcanic suite and of Mendonidis *et al.*<sup>54</sup> on the Glenmore granite, KwaZulu-Natal south coast. Petrogenetic studies include that of Kruger *et al.*,<sup>55</sup> who described the petrology and geochemistry of the 1.1 Gyr Oranjekom Complex of layered gabbro-norites and anorthosite. The Oranjekom magmas were derived from a depleted mantle source and were Al-rich. Differentiation took place by sorting of

mafic phases with plagioclase remaining largely suspended. Evans *et al.*<sup>56</sup> investigated the petrogenesis of the 1.0 Gyr Tete complex in northwestern Mozambique. Magmas forming this complex were also derived from depleted mantle sources with little evidence of crustal contamination and were emplaced at shallow levels in the upper crust with differentiation to form gabbroic, pyroxenitic and anorthositic rocks. Maier *et al.*<sup>57</sup> reported on PGE mineralization in this complex.

South African-based scientists have also been involved in research related to the 1.33 Gyr Voisey's Bay troctolite-gabbro intrusion and associated Ni-Cu-Co sulphide deposits in Canada. These involve Nd-Sr-Pb isotopes and crustal assimilation,<sup>58</sup> melting of gneiss inclusions in the ore-associated breccia,<sup>59</sup> comparison of Voisey's Bay and the Mushuau intrusion,<sup>60</sup> the oxygen isotope geochemistry of Voisey's Bay intrusion,<sup>61</sup> the oxygen fugacity during sulphide segregation,<sup>62</sup> and the Re-Os isotope systematics of the intrusion with implications for parental magma chemistry and ore genesis.<sup>63</sup>

### Neoproterozoic

Research on Neoproterozoic rocks essentially concerns granite magmatism, especially the precise dating of magmatic activity. Frimmel *et al.*<sup>64</sup> report ages ranging from  $833 \pm 2$  Myr through to  $741 \pm 6$  Myr for various components of the felsic extrusive and intrusive Richtersveld Igneous Complex, northwestern South Africa. This magmatism developed in accordance with crustal thinning over a mantle plume. Also in this area are the post-orogenic alkaline granites of the Kuboos-Bremen line of intrusions trending NE from northwest South Africa into southern Namibia. One of the largest plutons is the Kuboos pluton, which intrudes the Pan-African Gariep belt whose main phase of deformation occurred at  $545 \pm 2$  Myr. Frimmel *et al.*<sup>65</sup> report a U-Pb age of  $507 \pm 6$  Myr for the youngest intrusive phase of this pluton. Scheepers and co-workers focused on the 515–552 Myr Cape granite suite intrusive in the Pan-African mobile belt in the southwestern part of the Western Cape Province. They reported broadly synchronous SHRIMP ages of  $547 \pm 6$  Myr for the early syntectonic Darling granite and  $536 \pm 5$  Myr for the post-tectonic Robertson granite.<sup>66</sup> Nd isotopes suggest that both granites were derived from Mesoproterozoic crustal sources. This range of ages for granite activity was extended by subsequent SHRIMP age determinations on three granites comprising the Saldanha batholith, believed to be amongst the oldest of the Cape Granite Suite.<sup>67</sup> Ages obtained range from  $552 \pm 4$  Myr to  $539 \pm 4$  Myr. Scheepers and Nortje<sup>68</sup> described rhyolitic ignimbrites at Postberg, associated with the Saldanha batholith, and Scheepers and Pojoul<sup>69</sup> discussed their geochemistry and petrogenesis and demonstrated by a single zircon age of  $515 \pm 3$  Myr that they represent the final phase of Cape Granite Suite magmatism. A detailed petrogenetic study of the Malmesbury batholith was published by Siegfried.<sup>70</sup> He showed that the batholith is built of seven granitoid intrusions derived largely from a mafic igneous source by fractional crystallization with crustal assimilation.

Further afield, Handke *et al.*<sup>71</sup> report U-Pb zircon and baddeleyite ages of 804–779 Myr for 11 coeval gabbroic and granitoid intrusions along a 450-km belt in Madagascar. Yibas *et al.*<sup>72</sup> report U-Pb ages of 880–526 Myr granitoids in Ethiopia. In a petrogenetic study, Harris and Ashwal<sup>73</sup> report results of an oxygen isotope study on the 750 Myr Seychelles granites, indicating that the Mahé type was derived from juvenile mafic to intermediate crust, whereas the Praslin type was derived from a source that was a mixture of this crust and older crust which acquired its low  $\delta^{18}\text{O}$  significantly before granite genesis.

## Mesozoic volcanism

### Karoo Flood Basalt Province

Most publications focus on the intrusive element of this classic continental flood basalt province (CFB). Chevallier and Woodford<sup>74</sup> analysed a large amount of field data for three sill-ring complexes in the western Karoo basin, South Africa. They showed that these structures are complex and built of stacked saucer-like intrusions (inclined sheets) interlinked with flat inner and outer sills and arcuate dykes. Their work shows that emplacement of these complexes was initiated from dykes which fed the inclined sheets. These then propagated into the outer sill and then into the inner sill. The implications are that the sheets are an integral part of the feeding system to the subaerial lavas.

In an analysis of magma flow directions at the mineralized Insizwa intrusion using the AMS technique, Ferre *et al.*<sup>75</sup> show that magnetic and mineral lineations coincide and represent magma flow directions. Furthermore, the intrusion is built by multiple injection events from a lower level located to the SE of the Insizwa massif. In a paper focused on the Ni-Cu-PGE potential of Insizwa, Maier *et al.*<sup>76</sup> show that the magmas feeding the intrusion were depleted in PGE before emplacement and contained no entrained sulphide. They indicate little potential for an economic deposit associated with Insizwa. Expanding this work to the lavas of the Karoo province and other CFB sequence of all ages in southern Africa, these authors<sup>77</sup> found that all the flood basalts are depleted in PGE relative to Cu, except the oldest (Dominion, Ventersdorp). They conclude that the data are consistent with minor sulphide segregation during ascent and that these CFBs have to be considered poor exploration targets.

The behaviour of lava flows is addressed by De Bruijn *et al.*,<sup>78</sup> who present evidence of flow concentration of olivine phenocrysts in a basalt flow high in the sequence in Lesotho to produce a picritic core to the flow. Such evidence is important in accounting for olivine-rich layers within large differentiated intrusions such as Insizwa. In the Sabie River Formation in the northern Lebombo, the occurrence of peperite in the upper margin of a basalt sheet is interpreted as a shallow sheet that has burrowed into unconsolidated water-saturated sediment.<sup>79</sup> It is suggested that sediment-impregnated blocky flow tops that occur elsewhere in the Karoo volcanic sequence might have a similar origin. Weinert and Dunlevey<sup>80</sup> report the first recorded occurrence of the zeolite yugawaralite in Karoo basalts of the southern Lebombo.

### Etendeka Igneous Province

Marsh *et al.*<sup>81</sup> reviewed decades of research on the Etendeka Group, Namibia, which is equivalent to the Paraná Province in South America. On the basis of geochemistry they define eight mafic and seventeen silicic magma types and describe their areal and stratigraphic distribution. There is a marked but not exclusive geochemical provinciality within the Etendeka Province with incompatible element-enriched (high-Ti) mafic and silicic rocks having a close geographic association as do the low-Ti mafic and silicic rocks. The Doros complex is shown to be the eruptive site of the early Tafelkop basalts, which have affinities to the Tristan Plume. Comparisons with the Paraná indicate that all the important silicic types in the Paraná have geochemical equivalents in the Etendeka, hence extending the previous transatlantic correlations of the two provinces. These correlations indicate that volumetrically giant silicic systems developed with pronounced lithospheric thinning and rifting in continental flood basalt provinces.

Associated with the Etendeka flood volcanism are a large number of central volcanic complexes. One of these, Messum, has been subjected to detailed studies. Ewart *et al.*<sup>82</sup> showed that the complex igneous geology at Messum is best interpreted as a down-sag cauldron subsidence which is distinctly different from the classic cauldron subsidence structures found in the western United States. Focusing on the alkaline rocks in the core of the complex, Harris *et al.*<sup>83</sup> showed that trends towards silica-oversaturated syenites reflect crustal contamination of nepheline-bearing parental magmas. Basanite dykes are isotopically similar to the modern Tristan lavas and may reflect an input from the Tristan plume into the complex.

### Other Mesozoic igneous suites

As part of the South African Earth Sciences Research Programme in Antarctica, Harris *et al.*<sup>84</sup> discussed the petrogenesis of the largest Mesozoic intrusion in Dronning Maud Land — the Siste fjell Syenite, which is associated with the 'Karoo' CFB igneous event in that continent. Major, trace, radiogenic and stable isotope data indicate that the composition variation in the syenite is consistent with fractional crystallization with some contamination by the metamorphic basement and the low  $\delta^{18}\text{O}$  Sistenup lavas of unknown age. Harris also contributed to the understanding of the large-volume rhyolitic volcanism in Patagonia and the Antarctic Peninsula,<sup>85</sup> which parallels the palaeo-Pacific margin of Gondwana.

## Cenozoic to Recent volcanism

### Ocean island volcanism

Harris *et al.*<sup>86</sup> discuss oxygen isotopes measured on phenocryst in lavas from those Tristan da Cunha and Gough Island basalts which have enriched isotopic signatures. For Gough Island the phenocrysts are in isotopic equilibrium and suggest that the least evolved magmas had the same oxygen isotopic composition as MORB and that more-evolved lavas evolved by closed-system fractional crystallization. At Tristan, oxygen isotope ratios are lower, suggesting that material from the volcanic edifice contaminated the parental mafic magmas. The most primitive lavas evolved by fractional crystallization involving crystal accumulation, but the evolved lavas formed by closed-system fractional crystallization.

### Mid-ocean ridge volcanism

In a series of papers, Le Roux *et al.*<sup>87-89</sup> describe the geochemistry and petrogenesis of MORB erupted along the southern Mid-Atlantic Ridge (MAR) between 40 and 55 degrees. This part of the MAR is moderately slow-spreading and lies in the vicinity of the Discovery and Shona mantle plumes. There are systematic distributions of E-MORB and N-MORB on the ridge, with the former related to the proximity of the mantle plumes. The basalts also exhibit mild geochemical signatures of subduction. A model is proposed that involves interaction in the melting regime of the mantle plumes, delaminated continental lithosphere from Gondwana break-up, and subduction-contaminated asthenospheric mantle. Along this segment of the ridge the source in the vicinity of the Discovery plume has greater proportions of clinopyroxene. Partial melting (15–17%) at 18 kbar produces N-MORB, whereas slightly higher degrees of melting at slightly higher pressures characterize melting in the vicinity of the plumes. Subsequent evolution of melts involve crystallization of olivine, plagioclase and clinopyroxene. Crystallization of N-MORB happened at pressures of 3–6 kbar, whereas that in the plume-influenced melts occurred over a greater pressure range

(1 atm–7 kbar, but predominantly at 1 atm–3 kbar). These differences are ascribed to higher temperatures, more constant magma fluxes and increased longevity of subaxial magma chambers in the vicinity of mantle plumes.

#### Continental rift volcanism

Spath *et al.*<sup>90,91</sup> and Le Roex *et al.*<sup>92</sup> reported on geochemical studies of volcanoes on the flanks of the southern Kenya rift (Chyulu Hills, 10 km E of the rift) and from the rift walls and floor (Lake Magadie area). The Chyulu Hills volcanoes range in composition from nepheline-normative nephelinites, basanite, and hawaiites to orthopyroxene-normative subalkali basalts. Compositions are consistent with olivine-dominated differentiation. Spatial and temporal variations in degree of silica-undersaturation is explained in terms of variation in depth and degree of melting. The striking geochemical feature in these lavas is their depletion in K in mantle-normalized trace element plots; this is explained in terms of residual amphibole in the melting regime. This in turn implies a source in the lithosphere which is metasomatized by a rising mantle plume to be followed later by melting. Similarly, the mildly nepheline-normative within-rift alkali basalts and basanite are generated by variable amounts of melting to depths extending into the garnet stability field and in the presence of residual amphibole, again suggesting melting in the subcontinental lithosphere rather than in the asthenosphere or rising mantle plume. The lithosphere must extend to depths of 75 km beneath the Magadie area.

#### Anorogenic alkaline volcanism

Using the southwestern Cape melilitite-alkali basalt province (76–58 Myr), Janney *et al.*<sup>93</sup> demonstrate that strong trace element and isotopic variations correlate with lithospheric thickness. Eruptives on the continental shelf have strong HIMU affinities, whereas those on thick Proterozoic lithosphere have EM 1 isotopic features and kimberlite-like trace element patterns. A complex two-stage mixing process is used to account for the data and it is proposed that, unlike oceanic areas where the HIMU component is supplied from a conventional mantle plume, under Africa the HIMU component is in pods of recycled oceanic crust incorporated in a laterally broad, long-lived upwelling.

#### Carbonatites

Reviewing Sr and Nd isotope composition of carbonatites, Harmer<sup>94</sup> compared these to kimberlites and argued that CO<sub>2</sub>-rich residues forming from ponded kimberlites rise from the base of the lithosphere and are trapped at the peridotite-CO<sub>2</sub> thermal maximum at 2 GPa and enrich the lithospheric peridotite there. This enriched peridotite forms the source for subsequent carbonatitic magmas. Ionov and Harmer<sup>95</sup> report results of laser-ablation ICPMS trace element determinations in calcite-dolomite carbonatites with well-preserved igneous textures from Spitskop. Calcite phenocrysts have low rare-earth element (REE) contents and flat patterns in comparison with interstitial late-crystallizing calcite, which is strongly enriched in LREE. This suggests that REE are incompatible during fractional crystallization of carbonatite liquids. The phenocryst abundances are similar to carbonates in mantle xenoliths, indicating that the latter are crystal cumulates rather than quenched carbonatite liquids.

In a paper focusing on potassic trachytes, the only significant silicate rock of the Dicker Willem carbonatite, Namibia, Cooper and Reid<sup>96</sup> show that the trachytes are isotopically similar to the highest grade of potassic fenites associated with the carbonatite. They propose that the trachytes are partially melted fenites.

#### Kimberlites and mantle materials

In 1998, the 7th International Kimberlite Conference took place in Cape Town and a two-volume set of papers relating to that meeting was published in 1999.<sup>97</sup> Other publications on this topic have appeared in numerous journals.

#### Kimberlites

Field and Scott-Smith<sup>98</sup> presented a comprehensive survey of the geology of kimberlite pipes, reviewing the standard model which was established from observations on South African pipes and comparing it to others in southern Africa and the newly discovered pipes in Canada. They review and redefine terminology and demonstrate the variability of pipes in terms of pipe shape and internal geology. They demonstrate relationships between pipe type and the nature of the country rocks and discuss implications for emplacement mechanisms. Emplacement mechanism is also a concern of Rice,<sup>99</sup> who applied engineering studies on blasting to understanding kimberlite eruption mechanisms.

Skinner *et al.*<sup>100</sup> addressed the controversial topic of the presence of melilitite in kimberlite and showed that it is present in most Group II kimberlites and in some phlogopite-rich Group I kimberlites. Crystallization is favoured in the diatreme facies as a consequence of CO<sub>2</sub> loss. Recognition of late-crystallizing phases which formed subsequent to melt degassing is important for Ar-Ar dating of kimberlite, as shown by Phillips *et al.*,<sup>101</sup> who successfully applied laser probe step-heating analysis to single groundmass phlogopite and K-richrichterite grains in eleven kimberlites in South Africa.

In more general studies on the origin of kimberlite magma, Sweeney and Winter<sup>102</sup> integrated results from high-pressure experimental petrology and major element composition of kimberlite to constrain the depth of melting and the residual mineralogy of the source. Nowell *et al.*<sup>103</sup> adopted an isotopic approach, offering an Hf isotope perspective on components contributing to both Group I and II kimberlite magmas. They recognize a negative  $\Delta\epsilon_{\text{Hf}}$  component, which they propose derives from a deep sub-lithospheric source. Kimberlite magmas are generated in a plume derived from this deep source and attain additional isotopic variability by assimilating variably enriched lithospheric mantle.

Several descriptions of individual kimberlite pipes in southern Africa<sup>104,105</sup> and Australia<sup>106,107</sup> are also reported.

#### Xenocrysts

Xenocrysts, including diamond and its inclusions, have proved to be fertile sources for information about the Earth's interior. The 240 Myr Jwaneng pipe in Botswana is one of the few where diamonds with an eclogitic inclusion assemblage (eclogitic diamonds) occur.<sup>108</sup> These inclusions yield a Mesoproterozoic (1580 Myr) Sm-Nd isotope age, similar to that obtained from Finsch eclogitic diamonds, and initial Nd isotopic composition indicative of derivation from a depleted mantle source. The Jwaneng and Finsch results point to a regional eclogitic diamond formation event in the Mesoproterozoic Proterozoic related to interaction between subducted lithosphere with the stable Archaean lithosphere.

Viljoen *et al.*<sup>109</sup> and Aulbach *et al.*<sup>110</sup> reviewed the inclusions in diamonds from the Venetia pipes, which are important as they have an unusual and somewhat anomalous location in the Limpopo Mobile Belt at the junction of the Kaapvaal and Zimbabwe cratons. The most common inclusion type is sulphide, with peridotitic oxide and silicate inclusions and minor eclogitic and websteritic types. Mineral chemistry

suggests diamond crystallized at 900–1400°C at 55–70 kbar in a thick, ancient cratonic root dominated by highly depleted magnesian peridotite. Results for websteritic and eclogitic inclusions are consistent with eclogite representing subducted ocean crust, whereas the websterites represent the product of reaction of slab-derived melts with mantle peridotite. Viljoen<sup>111</sup> reports on infrared studies on diamonds from the Venetia kimberlite. Nitrogen contents and nitrogen aggregation states are highly variable with unusually high aggregation in the majority of diamonds — making it unusual compared to the cratonic kimberlite localities in southern Africa. High aggregation is a feature of other craton margin localities worldwide. Deformation in the mantle is thought to accelerate nitrogen aggregation in the diamonds.

Descriptions of inclusions in diamond from a craton-margin kimberlite, Dokolwayo, Swaziland, are reported by Daniels and Gurney.<sup>112</sup> They describe a variety of inclusions (peridotitic assemblages predominating) with variable composition recording macro- and micro-scale geochemical heterogeneity in the mantle where the diamonds formed. Among the inclusions is staurolite, which has implications for the high-*P* stability of this mineral. They also discuss the carbon isotope composition of a suite of 88 diamonds from Dokolwayo,<sup>113</sup> the majority of which lie within the normal range, regardless of their eclogitic or peridotitic character. The carbon source is considered to be methane which degassed from the lower mantle or core.

Grutter *et al.*<sup>114</sup> surveyed the composition of xenocrystic peridotitic garnets occurring on a regional scale on the Slave craton, Canada. Using the Cr<sub>2</sub>O<sub>3</sub>–CaO relationships as an indicator of depletion, they show that there are marked differences across at least three distinct northeast-trending lithospheric domains which are not reflected in isotopic compositions of Archaean crustal rock, suggesting isotopic decoupling between crust and mantle. For South Africa, these authors demonstrate significant compositional differences in the lithospheric mantle on either side of the wrench-fault system that defines the SW margin of the Kaapvaal craton. Towards the centre of the craton, lithospheric thickness is correlated with geochemical signatures of melt depletion in peridotite, which imparts stability to thickened cratonic roots.

### Xenoliths

Xenolith studies encompass thermobarometry, whole rock major and trace element geochemistry, and isotope geochemistry. Girnis *et al.*<sup>115</sup> carried out experiments in the FMASCr system to derive internally consistent thermobarometers for spinel and garnet harzburgites and applied the results to touching and non-touching inclusions in diamond. Tainton *et al.*<sup>116</sup> applied garnet thermobarometry to garnets recovered from a number of intrusions in the Central Tanzanian Craton and define heatflow patterns within this craton. In a detailed description of the xenolith suite from Venetia, South Africa, Stiefenhofer *et al.*<sup>117</sup> show that the suite is dominated by peridotites with pyroxenite and demonstrate that they have been affected by melt-metasomatism. Thermobarometry reveals the presence of a high-temperature inflection in the Venetia geotherm. In Namibia non-kimberlitic mantle xenoliths have been recovered from a c. 75 Myr nephelinite<sup>118</sup> at Swakopmund and from a lamprophyre pipe at Okenyenya,<sup>119</sup> both sited on the Damara mobile belt between the Congo and Kaapvaal cratons. Refractory and fertile peridotite xenoliths are present, the latter representing a variety of mantle not previously documented in southern Africa. They define a hot geotherm which probably relates to regional Cretaceous magmatism associated with the opening of the

South Atlantic Ocean.

Van Achtebergh *et al.*<sup>120</sup> have estimated the element fluxes during metasomatism of a suite garnet-bearing mantle xenoliths from the Letlhakane kimberlite, Botswana. They show that modally metasomatized rocks become enriched in Sr, Na, K, LREE and HFSE (Ti, Zr and Nb), with the removal of Al, Cr, and Fe and garnet-compatible trace elements. The proposed depletion in Al challenges a previous view that the metasomatic environment was merely Al-poor.

Gregoire *et al.*<sup>121</sup> re-examined the occurrence of the two phlogopite-bearing xenolith suites, MARID and PIC, and demonstrated that both are clearly distinguished in terms of major and trace element compositions of common minerals and Sr and Nd isotopes. Moreover, geochemical data indicate a genetic relationship of the xenolith types to Group I kimberlites (PIC) and Group II kimberlites (MARID). In a detailed textural compositional and oxygen isotope study, Zhang *et al.*<sup>122</sup> interpret the ilmenite-rich pliomict mantle xenoliths from Bultfontein as precipitates from a Fe-Ti-Cr-rich melt, which in turn could have separated through immiscibility of a migrating high-Ti silicate melt. Carlson *et al.*<sup>123</sup> present new, and review previously published, Re-Os isotope data for peridotite xenoliths in southern Africa. Menzies *et al.*<sup>124</sup> report additional data for the Newlands peridotites. Overall, the Kaapvaal craton peridotites have similar early Proterozoic to late Archaean re-depletion ages, and have the lowest <sup>187</sup>Os/<sup>186</sup>Os ratio of any terrestrial rock.

- Duncan A.R. (1995). A review of South African research on volcanic rocks, related intrusive rocks, and mantle derived materials: 1991–1995. *S. Afr. J. Sci.* **91**, 255–264.
- Dann J.C. (2000). The 3.5 Ga Komati Formation, Barberton Greenstone Belt, South Africa, Part I: new maps and magmatic architecture. *S. Afr. J. Geol.* **103**(1), 47–68.
- Cloete M. (1999). Aspects of volcanism and metamorphism on the Onverwacht Group Lavas in the southwestern portion of the Barberton Greenstone Belt. *Geological Survey of South Africa Memoir* **84**.
- Anhaeusser C.R. (2001). The anatomy of an extrusive-intrusive Archaean mafic-ultramafic sequence: the Nelshoogte Schist Belt and Stolzberg Layered Ultramafic Complex, Barberton Greenstone Belt, South Africa. *S. Afr. J. Geol.* **104**, 167–204.
- Poujol M. and Robb L.J. (1999). New U-Pb zircon ages on gneisses and pegmatite from south of the Murchison greenstone belt, South Africa. *S. Afr. J. Geol.* **102**(2), 93–97.
- Poujol M. (2001). U-Pb isotopic evidence for episodic granitoid emplacement in the Murchison greenstone belt, South Africa. *J. Afr. Earth Sci.* **33**(1), 155–163.
- Kröner A., Jaekel P. and Brandl G. (2000). Single zircon ages for felsic to intermediate rocks from the Pietersburg and Giyani greenstone belts and bordering granitoid orthogneisses, northern Kaapvaal Craton, South Africa. *J. Afr. Earth Sci.* **30**(4), 773–793.
- Henderson D.R., Long L.E. and Barton J.M. (2000). Isotopic ages and chemical and isotopic composition of the Archaean Turfloop Batholith, Pietersburg granite-greenstone terrane, Kaapvaal Craton, South Africa. *S. Afr. J. Geol.* **103**(1), 38–46.
- Anhaeusser C.R. and Walraven F. (1999). Episodic granitoid emplacement in the Western Kaapvaal Craton: evidence from the Archaean Kraaipan granite-greenstone terrane, South Africa. *J. Afr. Earth Sci.* **28**(2), 289–309.
- Poujol M., Anhaeusser C.R. and Armstrong R.A. (2002). Episodic granitoid emplacement in the Archaean Amalia–Kraaipan terrane, South Africa: confirmation from single zircon U-Pb geochronology. *J. Afr. Earth Sci.* **35**, 147–161.
- Poujol M. and Anhaeusser C.R. (2001). The Johannesburg Dome, South Africa: new single zircon U-Pb isotopic evidence for early Archaean granite-greenstone development within the central Kaapvaal Craton. *Precambrian Res.* **108**, 139–157.
- Barton J.M., Barton E.S. and Kröner A. (1999). Age and isotopic evidence for the origin of the Archaean granitoid intrusives of the Johannesburg Dome, South Africa. *J. Afr. Earth Sci.* **28**(3), 693–702.
- Gutzmer J., Nhleko N., Beukes N.J., Pickard A. and Barley M.E. (1999). Geochemistry and ion microprobe (SHRIMP) age of a quartz porphyry sill in the Mozaan Group of the Pongola Supergroup: implications for the Pongola and Witwatersrand Supergroups. *S. Afr. J. Geol.* **102**(2), 139–146.
- van der Westhuizen W.A. and de Bruijn H. (2000). High temperature ash flow–wet sediment interaction in the Makwassie Formation, Ventersdorp Supergroup, South Africa. *Precambrian Res.* **101**, 341–351.
- de Bruijn H., Schoch A.E., Whitelaw H.T. and van der Westhuizen W.A. (2002). Alteration of the Allanridge Formation of the Ventersdorp Supergroup near Douglas, Northern Cape Province. *S. Afr. J. Geol.* **105**, 75–92.
- Hall R.C.B. and Els B.G. (2002). The origin and significance of load-induced deformation structures in soft-sediment and lava at the base of the Archaean Ventersdorp Supergroup, South Africa. *J. Afr. Earth Sci.* **35**, 135–145.

17. Oberholzer J.D. and Eriksson P.G. (2000). Subaerial volcanism in the Palaeoproterozoic Hekpoort Formation (Transvaal Supergroup), Kaapvaal Craton. *Precambrian Res.* **101**, 193–210.
18. McCourt S., Hilliard P., Armstrong R.A. and Munyanyiwa H. (2001). SHRIMP U-Pb zircon geochronology of the Hurungwe granite north-west Zimbabwe: age constraints on the timing of the Magondi orogeny and implications for the correlation between the Kheis and Magondi Belts. *S. Afr. J. Geol.* **104**, 39–46.
19. Cawthorn R.G. (Ed.) (1999). Special Issue commemorating the 75th anniversary of the discovery of the Merensky Reef. *S. Afr. J. Geol.* **102**, 176–302.
20. Maier W.D. (Ed.) (2001). Special issue on platinum-group elements and minerals in southern African rocks. *S. Afr. J. Geol.* **104**, 273–364.
21. Buchanan P.C., Koeberl C. and Reimold W.U. (1999). Petrogenesis of the Dullstroom Formation, Bushveld Magmatic Province, South Africa. *Contrib. Mineral. Petrol.* **137**, 133–146.
22. Buchanan P.C., Reimold W.U., Koeberl C. and Kruger F.J. (2002). Geochemistry of intermediate to siliceous volcanic rocks of the Rooiberg Group, Bushveld Magmatic Province, South Africa. *Contrib. Mineral. Petrol.* **144**, 131–143.
23. Maier W.D., Barnes S.-J. and van der Merwe M.J. (2001). Platinum-group elements in the pyroxenite marker, Bushveld Complex: implications for the formation of the Main Zone. *S. Afr. J. Geol.* **104**, 301–308.
24. Nex P.A.M., Cawthorn R.G. and Kinnaird J.A. (2002). Geochemical effects of magma addition: compositional reversals and decoupling of trends in the Main Zone of the western Bushveld Complex. *Min. Mag.* **66**(6), 833–856.
25. Mitchell A.A. and Manthorpe R. (2002). The Giant Mottled Anorthosite: a transitional sequence at the top of the Upper Critical Zone of the Bushveld Complex. *S. Afr. J. Geol.* **105**, 15–24.
26. Viring R.G. and Cowell M.W. (1999). The Merensky Reef on Northam Platinum Limited. *S. Afr. J. Geol.* **102**(3), 192–208.
27. Lomberg K.G., Martin E.S., Patterson M.A. and Venter J.E. (1999). The morphology of potholes in the UG2 chromitite layer and Merensky Reef (pothole reef facies) at Union Section, Rustenburg Platinum Mines. *S. Afr. J. Geol.* **102**(3), 209–220.
28. Viljoen M.J. (1999). The nature and origin of the Merensky Reef of the western Bushveld Complex based on geological facies and geophysical data. *S. Afr. J. Geol.* **102**(3), 221–239.
29. Cawthorn R.G. (1999). Platinum-group element mineralisation in the Bushveld Complex — a critical reassessment of geochemical models. *S. Afr. J. Geol.* **102**(3), 268–281.
30. Scoon R.N. (2002). A new occurrence of Merensky Reef on the flanks of the Zaaikloof Dome, northeastern Bushveld Complex: relationship between diaspirism and magma replenishment. *Econ. Geol.* **97**, 1037–1049.
31. Carr H.W., Kruger F.J., Groves D.I. and Cawthorn R.G. (1999). The petrogenesis of Merensky Reef potholes at the Western Platinum Mine, Bushveld Complex: Sr-isotopic evidence for synmagmatic deformation. *Mineralium Deposita* **34**, 335–347.
32. Barnes S.-J. and Maier W.D. (2002). Platinum-group elements and microstructures of normal Merensky Reef from Impala Platinum Mines, Bushveld Complex. *J. Petrol.* **43**(1), 103–128.
33. Cawthorn R.G. (2002). Delayed accumulation of plagioclase in the Bushveld Complex. *Min. Mag.* **66**(6), 881–893.
34. Maier W.D. and Barnes S.-J. (1999). Platinum-group elements in silicate rocks of the Lower, Critical and Main Zones at Union Section, Western Bushveld Complex. *J. Petrol.* **40**(11), 1647–1671.
35. Eales H.V. (2000). Implications of the chromium budget of the Western Limb of the Bushveld Complex. *S. Afr. J. Geol.* **103**(2), 141–150.
36. Willmore C.C., Boudreau A.E. and Kruger F.J. (2000). The halogen geochemistry of the Bushveld Complex, Republic of South Africa: implications for chalcophile element distribution in the Lower and Critical Zones. *J. Petrol.* **41**, 1517–1539.
37. Willmore C.C., Boudreau A.E., Spivack A. and Kruger F.J. (2002). Halogens of Bushveld Complex, South Africa: <sup>37</sup>Cl and Cl/F evidence for hydration melting of the source region in a back-arc setting. *Chem. Geol.* **182**, 503–511.
38. Maier W.D., Arndt N.T. and Curl E.A. (2000). Progressive crustal contamination of the Bushveld Complex: evidence from Nd isotopic analyses of the cumulate rocks. *Contrib. Mineral. Petrol.* **140**, 316–327.
39. Harris C. and Chaumba J.B. (2001). Crustal Contamination and fluid-rock interaction during the Formation of the Platreef, Northern Limb of the Bushveld Complex, South Africa. *J. Petrol.* **42**(7), 1321–1347.
40. Clarke B.M., Uken R. and Watkeys M.K. (2000). Intrusion mechanisms of the southwestern Rustenburg Layered Suite as deduced from the Spruitfontein inlier. *S. Afr. J. Geol.* **103**(2), 120–127.
41. Cawthorn R.G., Harris C. and Kruger F.J. (2000). Discordant ultramafic pegmatoidal pipes in the Bushveld Complex. *Contrib. Mineral. Petrol.* **140**, 119–133.
42. Reid D.L. and Basson I.J. (2002). Iron-rich ultramafic pegmatite replacement bodies within the Upper Critical Zone, Rustenburg Layered Suite, Northam Platinum Mine, South Africa. *Min. Mag.* **66**(6), 895–914.
43. Scoon R.N. and Eales H.V. (2002). Unusual Fe-Ti-Cr spinels from discordant bodies of iron-rich ultramafic pegmatite at the Amandelbult Platinum mine, northwestern Bushveld Complex. *Min. Mag.* **66**(6), 857–880.
44. Li C., Ripley E.M., Maier W.D. and Gomwe T.E.S. (2002). Olivine and sulfur isotope compositions of the Uitkomst Ni-Cu sulfide ore-bearing complex, South Africa: evidence for sulfur contamination and multiple magma emplacements. *Chem. Geol.* **188**, 149–159.
45. Gauert C. (2001). Sulphide and oxide mineralisation in the Uitkomst Complex, South Africa: origin in a magma conduit. *J. Afr. Earth Sci.* **32**(2), 149–161.
46. de Waal S.A. and Armstrong R.A. (2000). The age of the Marble Hall diorite, its relationship to the Uitkomst Complex, and evidence for a new magma type associated with the Bushveld igneous event. *S. Afr. J. Geol.* **103**(2), 128–140.
47. Eales H.V. (2002). Caveats in defining the magmas parental to the mafic rocks of the Bushveld Complex, and the manner of their emplacement: review and commentary. *Min. Mag.* **66**(6), 815–832.
48. Maier W.D., Slipe J., Barnes S.-J., de Waal S.A. and Li C. (2001). PGE-bearing mafic-ultramafic sills in the floor of the eastern Bushveld Complex on the farms Blaauwboschkraal, Zwartkopje, and Waterval. *S. Afr. J. Geol.* **104**, 343–354.
49. Maier W.D. and Barnes S.-J. (1999). The origin of Cu sulfide deposits in the Curaçá Valley, Bahia, Brazil: evidence from Cu, Ni, Se, and platinum-group element concentrations. *Econ. Geol.* **94**(2), 165–184.
50. Wilson A.H. and Prendergast M.D. (2001). Platinum-group element mineralisation in the Great Dyke, Zimbabwe, and its relationship to magma evolution and magma chamber structure. *S. Afr. J. Geol.* **104**, 319–342.
51. Wilson A.H. (2001). Compositional and lithological controls on the PGE-bearing sulphide zones in the Selukwe subchamber, Great Dyke: a combined equilibrium-Rayleigh fractionation model. *J. Petrol.* **42**(10), 1845–1867.
52. Wilson A.H., Murahwi C.Z. and Coghill B. (2000). Stratigraphy, geochemistry and platinum group element mineralisation of the central zone of the Selukwe Subchamber of the Great Dyke, Zimbabwe. *J. Afr. Earth Sci.* **30**(4), 833–853.
53. Gutzmer J., Beukes N.J., Pickard A. and Barley M.E. (2000). 1170 Ma SHRIMP age for Koras Group bimodal volcanism, Northern Cape Province. *S. Afr. J. Geol.* **103**(1), 32–37.
54. Mendonidis P., Armstrong R.A., Eglinton B.M., Grantham G.H. and Thomas R.J. (2002). Metamorphic history and U-Pb Zircon (SHRIMP) geochronology of the Glenmore granite: implications for the tectonic evolution of the Natal Metamorphic Province. *S. Afr. J. Geol.* **105**, 325–336.
55. Kruger F.J., Geringer G.J. and Havenga A.T. (2000). The geology, petrology, geochronology and source region character of the layered gabbronoritic Oranjekom Complex in the Kibaran Namaqua mobile belt, South Africa. *J. Afr. Earth Sci.* **30**(3), 675–687.
56. Evans R.J., Ashwal L.D. and Hamilton M.A. (1999). Mafic, ultramafic, and anorthositic rocks of the Tete Complex, Mozambique: petrology, age, and significance. *S. Afr. J. Geol.* **102**(2), 153–166.
57. Maier W.D., Barnes S.-J., Ashwal L.D. and Li C. (2001). A reconnaissance study on the Magmatic Cu-Ni-PGE sulphide potential of the Tete Complex, Mozambique. *S. Afr. J. Geol.* **104**, 355–364.
58. Amelin Y., Li C., Valeyev O. and Naldrett A.J. (2000). Nd-Pb-Sr isotope systematics of crustal assimilation in the Voisey's Bay and Mashuau intrusions, Labrador, Canada. *Econ. Geol.* **95**, 815–830.
59. Li C. and Naldrett A.J. (2000). Melting reactions of gneissic inclusions with enclosing magma at Voisey's Bay, Labrador, Canada: implications with respect to ore genesis. *Econ. Geol.* **95**, 801–814.
60. Li C., Lightfoot P.C., Amelin Y. and Naldrett A.J. (2000). Contrasting petrological and geochemical relationships in the Voisey's Bay and Mashuau intrusions, Labrador, Canada: implications for ore genesis. *Econ. Geol.* **95**, 771–799.
61. Ripley E.M., Park Y.-R., Li C. and Naldrett A.J. (2000). Oxygen isotope studies of the Voisey's Bay Ni-Cu-Co deposit, Labrador, Canada. *Econ. Geol.* **95**, 831–844.
62. Brennan J.M. and Li C. (2000). Constraints on oxygen fugacity during sulfide segregation in the Voisey's Bay intrusion, Labrador, Canada. *Econ. Geol.* **95**, 901–915.
63. Lambert D.D., Frick L.R., Foster J.G., Li C. and Naldrett A.J. (2000). Re-Os isotope systematics of the Voisey's Bay Ni-Cu-Co magmatic sulfide system, Labrador, Canada: II. Implications for parental magma chemistry, ore genesis, and metal redistribution. *Econ. Geol.* **95**, 867–888.
64. Frimmel H.E., Zartman R.E. and Späth A. (2001). The Richtersveld igneous complex, South Africa: U-Pb zircon and geochemical evidence for the beginning of Neoproterozoic continental breakup. *J. Geol.* **109**, 493–508.
65. Frimmel H.E. (2000). New U-Pb zircon ages for the Kuboos pluton in the Pan-African Gariep belt, South Africa: Cambrian mantle plume or far field collision effect? *S. Afr. J. Geol.* **103**, 207–214.
66. Da Silva L.C., Gresse P.G., Scheepers R., McNaughton N.J., Hartmann L.A. and Fletcher I. (2000). U-Pb SHRIMP and Sm-Nd age constraints on the timing and sources of the Pan-African Cape Granite Suite, South Africa. *J. Afr. Earth Sci.* **30**(4), 795–815.
67. Scheepers R. and Armstrong R. (2002). New U-Pb SHRIMP zircon ages of the Cape Granitic Suite: implications for the magmatic evolution of the Saldanha Belt. *S. Afr. J. Geol.* **105**, 241–256.
68. Scheepers R. and Nortjé A.N. (2000). Rhyolitic ignimbrites of the Cape Granite Suite, southwestern Cape Province, South Africa. *J. Afr. Earth Sci.* **31**(3/4), 647–656.
69. Scheepers R. and Poujol M. (2002). U-Pb zircon ages of Cape Granite Suite ignimbrites: characteristics of the last phases of the Saldanian magmatism. *S. Afr. J. Geol.* **105**, 163–178.
70. Siegfried H.P. (1999). The Malmesbury Batholith and its relationship to granitic plutons in the Swartland Tectonic Domain. *S. Afr. Council for Geoscience Bull.* **125**, 122 pp.
71. Handke M.J., Tucker R.D. and Ashwal L.D. (1999). Neoproterozoic continental arc magmatism in west-central Madagascar. *Geology* **27**(4), 351–354.
72. Yibas B., Reimold W.U., Armstrong R., Koeberl C., Anhaeusser C.R. and Phillips D. (2002). The tectonostratigraphy, granulite geochronology and geological evolution of the Precambrian of southern Ethiopia. *J. Afr. Earth Sci.* **34**, 57–84.
73. Harris C. and Ashwal L.D. (2002). The origin of low  $\delta^{18}\text{O}$  granites and related rocks from the Seychelles. *Contrib. Mineral. Petrol.* **143**, 366–376.
74. Chevallier L. and Woodford A. (1999). Morpho-tectonics and mechanism of emplacement of the dolerite rings and sills of the western Karoo, South Africa. *S. Afr. J. Geol.* **102**(1), 43–54.
75. Ferré E.C., Bordarier C. and Marsh J.S. (2002). Magma flow inferred from AMS fabrics in a layered mafic sill, Insizwa, South Africa. *Tectonophysics* **354**, 1–23.
76. Maier W.D., Marsh J.S., Barnes S.-J. and Dodd D.C. (2002). The distribution of platinum group elements in the Insizwa Lobe, Mount Ayliff Complex, South Africa: implications for Ni-Cu-PGE sulfide exploration in the Karoo Igneous

- Province. *Econ. Geol.* **97**, 1293–1306.
77. Maier W.D., Barnes S.-J. and Marsh J.S. (2001). In *Mineral Deposits at the Beginning of the 21st Century*, eds Piestrzynski et al., pp. 665–668. Swets & Zeitlinger, Lisse.
  78. de Bruijn H., Schoch A.E., van der Westhuizen W.A. and Myburgh C.A. (2000). Picrite from the Katse area, Lesotho: evidence for flow differentiation. *J. Afr. Earth Sci.* **31**(3–4), 657–668.
  79. Rawlings D.J., Watkeys M.K. and Sweeney R.J. (1999). Peperitic upper margin of an invasive flow, Karoo flood basalt province, northern Lebombo. *S. Afr. J. Geol.* **102**(4), 377–383.
  80. Weinert C.H.-S.W. and Dunlevey J.N. (2000). Yugawaralite in the Letaba Formation, northeastern Kwazulu-Natal, South Africa. *S. Afr. J. Geol.* **103**(1), 69–73.
  81. Marsh J.S., Ewart A., Milner S.C., Duncan A.R. and Miller R.McG. (2001). The Etendeka Igneous Province: magma types and their stratigraphic distribution with implications for the evolution of the Paraná-Etendeka flood basalt province. *Bull. Volcanol.* **62**, 464–486.
  82. Ewart A., Milner S.C., Duncan A.R. and Bailey M. (2002). The Cretaceous Messum igneous complex, S.W. Etendeka, Namibia: reinterpretation in terms of a downsag-cauldron subsidence model. *J. Volcanol. Geothermal Res.* **114**, 251–273.
  83. Harris C., Marsh J.S. and Milner S.C. (1999). Petrology of the alkaline core of the Messum Igneous Complex, Namibia: evidence for the progressively decreasing effect of crustal contamination. *J. Petrol.* **40**(9), 1377–1397.
  84. Harris C., Johnstone W.P. and Phillips D. (2002). Petrogenesis of the Mesozoic Sistejell syenite intrusion, Dronning Maud Land, Antarctica and surrounding low  $\delta^{18}\text{O}$  lavas. *S. Afr. J. Geol.* **105**, 205–226.
  85. Riley T.R., Leat P.T., Pankhurst R.J. and Harris C. (2001). Origins of large volume rhyolitic volcanism in the Antarctic Peninsula and Patagonia by crustal melting. *J. Petrol.* **42**(6), 1043–1065.
  86. Harris C., Smith H.S. and le Roex A.P. (2000). Oxygen isotope composition by phenocrysts from Tristan da Cunha and Gough Island lavas: variation with fractional crystallisation and evidence for assimilation. *Contrib. Mineral. Petrol.* **138**, 164–175.
  87. le Roux P.J., le Roex A.P. and Schilling J.-G. (2002). Crystallisation processes beneath the southern Mid-Atlantic Ridge (40–55°S), evidence for high-pressure initiation of crystallisation. *Contrib. Mineral. Petrol.* **142**, 582–602.
  88. le Roux P.J., le Roex A.P., Schilling J.-G., Shimizu N., Perkins W.W. and Pearce N.J.G. (2002). Mantle heterogeneity beneath the southern Mid-Atlantic Ridge: trace element evidence for contamination of ambient asthenospheric mantle. *Earth Planet. Sci. Lett.* **203**, 479–498.
  89. le Roux P.J., le Roex A.P. and Schilling J.-G. (2002). MORB melting processes beneath the southern Mid-Atlantic Ridge (40–55°S): a role for mantle plume-derived pyroxenite. *Contrib. Mineral. Petrol.* **144**, 206–229.
  90. Späth A., le Roex A.P. and Opiyo-Akech N. (2000). The petrology of the Chyulu Hills Volcanic Province, southern Kenya. *J. Afr. Earth Sci.* **31**(2), 337–358.
  91. Späth A., le Roex A.P. and Opiyo-Akech N. (2001). Plume-lithosphere interaction and the origin of continental rift-related alkaline volcanism — the Chyulu Hills Volcanic Province, Southern Kenya. *J. Petrol.* **42**(4), 765–787.
  92. le Roex A.P., Späth A. and Zartman R.E. (2001). Lithospheric thickness beneath the southern Kenya Rift: implications from basalt geochemistry. *Contrib. Mineral. Petrol.* **142**, 89–106.
  93. Janney P.E., le Roex A.P., Carlson R.W. and Viljoen K.S. (2002). A chemical and multi-isotope study of the Western Cape Olivine Melilitite Province, South Africa: implications for the sources of kimberlites and the origin of the HIMU signature in Africa. *J. Petrol.* **43**(12), 2339–2370.
  94. Harmer R.E. (1999). A common source for carbonatites, kimberlites and megacrysts? In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 332–340. Red Roof Design, Cape Town.
  95. Ionov D. and Harmer R.E. (2002). Trace element distribution in calcite-dolomite carbonatites from Spitskop: inferences for differentiation of carbonatite magmas and the origin of carbonates in mantle xenoliths. *Earth Planet. Sci. Lett.* **198**, 495–510.
  96. Cooper A.F. and Reid D.L. (2000). The association of potassic trachytes and carbonatites at the Dicker Willem Complex, southwest Namibia: coexisting, immiscible, but not cogenetic magmas. *Contrib. Mineral. Petrol.* **139**, 570–583.
  97. Gurney J.J., Gurney J.L., Pascoe M.D. and Richardson S.H. (Eds) (1999). *Proceedings of the VIIIth International Kimberlite Conference*. Red Roof Design, Cape Town.
  98. Field M. and Scott Smith B.H. (1999). Contrasting geology and near-surface emplacement of kimberlite pipes in southern Africa and Canada. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 214–237. Red Roof Design, Cape Town.
  99. Rice A. (1999). Can the blasting excavation engineering services provide insight into the processes of kimberlite emplacement and eruption? In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 699–708. Red Roof Design, Cape Town.
  100. Skinner E.M.W., Mahotkin I.L. and Grutter H.S. (1999). Melilitite in kimberlites. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 788–794. Red Roof Design, Cape Town.
  101. Phillips D., Kiviets G.B., Barton E.S., Smith C.B., Viljoen K.S. and Fourie L.F. (1999).  $^{40}\text{Ar}/^{39}\text{Ar}$  dating of kimberlites and related rocks: problems and solutions. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 677–688. Red Roof Design, Cape Town.
  102. Sweeney R.J. and Minter F. (1999). Kimberlite as high-pressure melts: the determination of segregation depth from major element chemistry. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 846–851. Red Roof Design, Cape Town.
  103. Nowell G.M., Pearson D.G., Kempton P.D., Noble S.R. and Smith C.B. (1999). Origins of kimberlites: a Hf isotope perspective. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 616–624. Red Roof Design, Cape Town.
  104. Seggie A.G., Hannweg G.W., Colgan E.A. and Smith C.B. (1999). The geology and geochemistry of the Venetia Kimberlite Cluster, Northern Province, South Africa. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 750–756. Red Roof Design, Cape Town.
  105. Williams C.M. and Robey J.V.A. (1999). Petrography and mineral chemistry of the Mwenzi-01 Kimberlite, Zimbabwe. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 896–903. Red Roof Design, Cape Town.
  106. Wyatt B.A., Sumpton J.D.H., Stiefenhofer J., Shee S.R. and Smith T.W. (1999). Kimberlites in the Forrest River area, Kimberley Region, Western Australia. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 912–922. Red Roof Design, Cape Town.
  107. Berryman A.K., Stiefenhofer J., Shee S.R., Wyatt B.A. and Belousova E.A. (1999). The discovery and geology of the Timber Creek kimberlites, Northern Territory, Australia. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 30–39. Red Roof Design, Cape Town.
  108. Richardson S.H., Chinn I.L. and Harris J.W. (1999). Age and origin of eclogitic diamonds from the Jwaneng kimberlite, Botswana. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 709–713. Red Roof Design, Cape Town.
  109. Viljoen K.S., Phillips D., Harris J.W. and Robinson D.N. (1999). Mineral inclusions in diamonds from the Venetia kimberlites, Northern Province, South Africa. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 888–895. Red Roof Design, Cape Town.
  110. Aulbach S., Stachel T., Viljoen K.S., Brey G.P. and Harris J.W. (2002). Eclogitic and websteritic diamond sources beneath the Limpopo Belt — is slab-melting the link? *Contrib. Mineral. Petrol.* **143**, 56–70.
  111. Viljoen K.S. (2002). An infrared investigation of inclusion-bearing diamonds from the Venetia kimberlite, Northern Province, South Africa: implications for diamonds from craton-margin settings. *Contrib. Mineral. Petrol.* **144**, 98–108.
  112. Daniels L.R.M. and Gurney J.J. (1999). Diamond inclusions from the Dokolwayo kimberlite, Swaziland. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 134–142. Red Roof Design, Cape Town.
  113. Daniels L.R.M. and Gurney J.J. (1999). Dokolwayo diamond carbon isotope. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 143–147. Red Roof Design, Cape Town.
  114. Grütter H.S., Apter D.B. and Kong J. (1999). Crust-mantle coupling: evidence from mantle-derived xenocrystic garnets. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 307–313. Red Roof Design, Cape Town.
  115. Giris A.V., Stachel T., Brey C.P., Harris J.W. and Phillips D. (1999). Internally consistent geothermobarometers for garnet harzburgites: model refinement and application. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 247–256. Red Roof Design, Cape Town.
  116. Tainton K.M., Seggie A.G., Bayly B.A., Tomlinson I. and Quadling K.E. (1999). Garnet thermobarometry: implications for mantle heat flow within the Tanzanian Craton. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 852–860. Red Roof Design, Cape Town.
  117. Stiefenhofer J., Viljoen K.S., Tainton K.M., Dobbe R. and Hannweg G.W. (1999). The petrology of a mantle xenolith suite from Venetia, South Africa. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 836–845. Red Roof Design, Cape Town.
  118. Whitehead K., le Roex A., Class C. and Bell D. (2002). Composition and Cretaceous thermal structure of the upper mantle beneath the Damara Mobile Belt: evidence from nepheline-hosted peridotite xenoliths, Swakopmund. *Namibia. J. Geol. Soc.* **159**, 307–321.
  119. Baumgartner M.C., le Roex A.P. and Gurney J.J. (2000). Mantle and crustal xenoliths from the Okenyena lamprophyre diatreme: constraints on the upper mantle and lower crust beneath the Damara Belt, northwestern Namibia. *Comm. Geol. Suro. Namibia* **12**, 279–290.
  120. van Acherbergh E., Griffin W.L. and Stiefenhofer J. (2001). Metasomatism in mantle xenoliths from the Lethakane kimberlites: estimation of element fluxes. *Contrib. Mineral. Petrol.* **141**, 397–414.
  121. Grégoire M., Bell D.R. and le Roex A.P. (2002). Trace element geochemistry of phlogopite-rich mafic mantle xenoliths: their classification and their relationship to phlogopite-bearing peridotites and kimberlites revisited. *Contrib. Mineral. Petrol.* **142**, 603–625.
  122. Zhang H.-F., Menzies M.A., Matthey D.P., Hinton R.W. and Gurney J.J. (2001). Petrology, mineralogy and geochemistry of oxide minerals in polymict xenoliths from the Bultfontein kimberlites, South Africa: implication for low bulk-rock oxygen isotopic ratios. *Contrib. Mineral. Petrol.* **141**, 367–379.
  123. Carlson R.W., Pearson D.G., Boyd F.R., Shirey S.B., Irvine G., Menzies A.H. and Gurney J.J. (1999). Re-Os systematics of lithospheric peridotites: implications for lithosphere formation and preservation. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 99–108. Red Roof Design, Cape Town.
  124. Menzies A.H., Carlson R.W., Shirey S.B. and Gurney J.J. (1999). Re-Os systematics of Newlands peridotite xenoliths: implications for diamond and lithosphere formation. In *Proc. VIIIth International Kimberlite Conference*, eds J.J. Gurney, J.L. Gurney, M.D. Pascoe and S.H. Richardson, pp. 566–573. Red Roof Design, Cape Town.