

Acta Biologica Plantarum Agriensis 5(1): 27 (2017) DOI:10.21406/abpa.2017.5.1.27
ISSN 2061-6716 (Print), 2063-6725 (Online) 4th CC 2017 Abstract
<http://abpa.ektf.hu/> Plenary lecture

RECOVERY OF ACETONE RINSED LICHEN THALLI OF *CLADONIA FOLIACEA*
Cladonia foliacea zuzmótelepek regenerálódása acetonos kezelést követően

Edit FARKAS¹, Rita ENGEL¹, Bernadett BIRÓ¹, Krisztina SZABÓ¹,
Katalin VERES¹ & Zsolt CSINTALAN²

¹Institute of Ecology and Botany, MTA Centre for Ecological Research, Hungarian Academy of Sciences, H-2163 Vácrátót, Alkotmány u. 2-4., Hungary; ²Institute of Botany and Ecophysiology, Szent István University, H-2100 Gödöllő, Páter Károly u. 1., Hungary; e-mail: farkas.edit@okologia.mta.hu

The optimal period for applying acetone (best for extraction and shortest possible for avoiding detrimental effects), was established for dry lichen thalli rinsed in acetone following the methods by Solhaug and Gauslaa applied in 2001. Secondary lichen substances (usnic and fumarprotocetraric acids) were analysed by HPTLC and HPLC, the detrimental effect of acetone was tested by chlorophyll fluorescence measurement. The optimal duration for acetone rinsing proved to be around 1–2(–5) days in the samples collected in summer. The determined Fv/Fm values of *Cladonia foliacea* remained relatively high even after 1024 hours compared to any of the 12 species studied previously. A seasonal difference between summer and winter collected thalli was found. According to our results the thalli collected in winter are more sensitive to acetone rinsing. Therefore the summer period – when lichens are more frequently in dry condition and therefore photosynthetically active for a shorter period – is more advantageous for collecting samples for transplantation experiments combined with acetone rinsing. Higher concentrations of usnic acid and fumarprotocetraric acid were found in winter samples than in summer ones. The considerable variability is possibly due to a natural variability among the samples (substance content among and within thalli) and partly to the preparation methods. However, homogenised (lyophilised, then ground) samples showed no difference in usnic acid content if extracted for 15, 30, 40 or 50 minutes.

The thalli re-placed in controlled field condition after acetone rinsing experiments were observed monthly. The digital images showed serious damage already after 1–2 months at most of the treated thalli. Samples were investigated after a 6 months recovery period. The results confirm the advantage of summer collecting. Though both winter and summer collected samples rinsed in acetone less than 16/32 hours showed similar, good vitality based on Fv/Fm measurements, however photosynthetically active parts of thalli were smaller in case of winter collected samples.

Supported by the Hungarian Scientific Research Fund OTKA 81232, 101713, the Research Centre of Excellence – 11476-3/2016/FEKUT and the project NKFI K 124341 financed by the National Research Development and Innovation Fund.