

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(Biphenyl-2-yl)bromidobis(2-methyltetrahydrofuran- κ O)magnesium(II)Simon Nordschild,^a D. Wohlgemuth^a and Michael Bolte^{b*}^aChemetall GmbH, Lithium Division, Trakehner Strasse 3, 60487 Frankfurt am Main, Germany, and ^bInstitut für Anorganische Chemie, J. W. Goethe-Universität Frankfurt, Max-von-Laue-Strasse 7, 60438 Frankfurt/Main, Germany

Correspondence e-mail: bolte@chemie.uni-frankfurt.de


Received 23 March 2009; accepted 1 April 2009

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.045; wR factor = 0.108; data-to-parameter ratio = 17.4.

In the title Grignard reagent, $[\text{MgBr}(\text{C}_{12}\text{H}_9)(\text{C}_5\text{H}_{10}\text{O})_2]$, the Mg centre adopts a distorted tetrahedral MgCO_2Br arrangement. The dihedral angle between the two aromatic rings of the biphenyl residue is $44.00(14)^\circ$. Each molecule incorporates one R - and one S -configured 2-methyltetrahydrofuran molecule.

Related literature

For background to Grignard-type compounds, see Elschenbroich (2008); Schwetlick (1996); Silverman & Rakita (1996).



Experimental

Crystal data

 $[\text{MgBr}(\text{C}_{12}\text{H}_9)(\text{C}_5\text{H}_{10}\text{O})_2]$ $M_r = 429.67$

Monoclinic, $P2_1/n$
 $a = 11.6887(5)$ Å
 $b = 16.8061(9)$ Å
 $c = 11.7888(5)$ Å
 $\beta = 103.757(4)^\circ$
 $V = 2249.38(18)$ Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.87$ mm⁻¹
 $T = 173$ K
 $0.29 \times 0.28 \times 0.26$ mm

Data collection

Stoe IPDSII two-circle diffractometer
 Absorption correction: multi-scan (MULABS; Spek, 2003; Blessing, 1995)
 $T_{\min} = 0.614$, $T_{\max} = 0.642$

54470 measured reflections
 4107 independent reflections
 3484 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.069$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.108$
 $S = 1.05$
 4107 reflections

236 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.76$ e Å⁻³
 $\Delta\rho_{\min} = -0.41$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Mg1—O1	2.022 (2)	Mg1—C21	2.143 (3)
Mg1—O11	2.030 (2)	Mg1—Br1	2.4750 (10)
O1—Mg1—O11	97.02 (10)	O1—Mg1—Br1	109.97 (9)
O1—Mg1—C21	114.41 (11)	O11—Mg1—Br1	104.75 (7)
O11—Mg1—C21	106.87 (11)	C21—Mg1—Br1	120.53 (8)

Data collection: *X-AREA* (Stoe & Cie, 2001); cell refinement: *X-AREA*; data reduction: *X-AREA*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL-Plus* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2933).

References

- Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
 Elschenbroich, C. (2008). In *Organometallchemie*. Stuttgart: Teubner BG.
 Schwetlick, K. (1996). In *Organikum*. Heidelberg: Johann Ambrosius Barth.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Silverman, G. S. & Rakita, P. E. (1996). In *Handbook of Grignard Reagents*. New York: Dekker.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
 Stoe & Cie (2001). *X-AREA*. Stoe & Cie, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2009). E65, m493 [doi:10.1107/S1600536809012185]

(Biphenyl-2-yl)bromidobis(2-methyltetrahydrofuran- κ O)magnesium(II)

S. Nordschild, D. Wohlgemuth and M. Bolte

Comment

The title compound is a Grignard reagent, a kind of compounds which is widely used for C—C bond formation or as base in organic chemistry. The Mg centre is four coordinate in a distorted tetrahedral fashion. The bond angles range from 97.02 (1)° for O—Mg—O to 120.53 (8)° for C—Mg—Br. The bond lengths are 2.4750 (10)Å for Mg—Br, 2.143 (3)Å for Mg—C and 2.022 (2)° and 2.030 (2)Å for the Mg—O bonds. The dihedral angles between the two aromatic rings of the biphenyl residue is 44.00 (14)°.

Experimental

This compound is commercially available from Chemetall GmbH (CAS 82214–69-5, Product No. 408562). Colourless blocks of (I) were obtained from a solution due to long term storage at ambient temperature.

Refinement

The H atoms bonded were geometrically positioned and refined with fixed individual displacement parameters [$U(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ or $U(\text{H}) = 1.5 U_{\text{eq}}(\text{C}_{\text{methyl}})$] using a riding model with $C_{\text{aromatic}}\text{—H} = 0.95 \text{ \AA}$, $C_{\text{methyl}}\text{—H} = 0.98 \text{ \AA}$, $C_{\text{methylene}}\text{—H} = 0.99 \text{ \AA}$ and $C_{\text{tertiary}}\text{—H} = 1.00 \text{ \AA}$.

Figures




Fig. 1. Perspective view of (I) with displacement ellipsoids drawn at the 50% probability level; H atoms omitted for clarity.

(Biphenyl-2-yl)bromidobis(2-methyltetrahydrofuran- κ O)magnesium(II)

Crystal data

[MgBr(C₁₂H₉)(C₅H₁₀O)₂]

$M_r = 429.67$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 11.6887 (5) \text{ \AA}$

$F_{000} = 896$

$D_x = 1.269 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 40381 reflections

$\theta = 3.6\text{--}26.6^\circ$

supplementary materials

$b = 16.8061 (9) \text{ \AA}$
 $c = 11.7888 (5) \text{ \AA}$
 $\beta = 103.757 (4)^\circ$
 $V = 2249.38 (18) \text{ \AA}^3$
 $Z = 4$

$\mu = 1.87 \text{ mm}^{-1}$
 $T = 173 \text{ K}$
Block, colourless
 $0.29 \times 0.28 \times 0.26 \text{ mm}$

Data collection

Stoe IPDSII two-circle diffractometer
Radiation source: fine-focus sealed tube
Monochromator: graphite
 $T = 173 \text{ K}$
 ω scans
Absorption correction: multi-scan (MULABS; Spek, 2003; Blessing, 1995)
 $T_{\min} = 0.614$, $T_{\max} = 0.642$
54470 measured reflections

4107 independent reflections
3484 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.069$
 $\theta_{\max} = 25.4^\circ$
 $\theta_{\min} = 3.6^\circ$
 $h = -14 \rightarrow 14$
 $k = -20 \rightarrow 20$
 $l = -14 \rightarrow 14$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.108$
 $S = 1.05$
4107 reflections
236 parameters

Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0462P)^2 + 2.3133P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.76 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.41 \text{ e \AA}^{-3}$
Extinction correction: SHELXL97 (Sheldrick, 2008),
 $F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.0043 (6)

Primary atom site location: structure-invariant direct methods
Secondary atom site location: difference Fourier map

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Mg1	0.56815 (8)	0.31486 (6)	0.70780 (8)	0.0349 (2)
Br1	0.60093 (3)	0.24360 (2)	0.89652 (3)	0.05398 (15)
O1	0.4214 (2)	0.38260 (15)	0.6841 (2)	0.0572 (6)
C2	0.4112 (4)	0.4578 (4)	0.6045 (5)	0.0967 (17)
H2A	0.4898	0.4781	0.6011	0.116*
H2B	0.3649	0.4465	0.5243	0.116*
C3	0.3515 (7)	0.5122 (4)	0.6625 (8)	0.159 (4)
H3A	0.3114	0.5542	0.6085	0.190*
H3B	0.4067	0.5376	0.7296	0.190*
C4	0.2618 (6)	0.4595 (5)	0.7036 (7)	0.145 (3)
H4A	0.2312	0.4860	0.7653	0.173*
H4B	0.1951	0.4450	0.6379	0.173*
C5	0.3367 (5)	0.3866 (6)	0.7512 (5)	0.143 (3)
H5	0.3763	0.3948	0.8354	0.172*
C6	0.2702 (6)	0.3147 (6)	0.7372 (7)	0.170 (4)
H6A	0.3212	0.2705	0.7720	0.256*
H6B	0.2383	0.3044	0.6539	0.256*
H6C	0.2052	0.3199	0.7762	0.256*
O11	0.50159 (19)	0.23129 (11)	0.58553 (18)	0.0412 (5)
C12	0.4587 (3)	0.2531 (2)	0.4632 (3)	0.0536 (8)
H12A	0.4940	0.3040	0.4464	0.064*
H12B	0.3718	0.2587	0.4430	0.064*
C13	0.4955 (5)	0.1870 (3)	0.3968 (4)	0.0852 (15)
H13A	0.4355	0.1777	0.3231	0.102*
H13B	0.5713	0.1998	0.3774	0.102*
C14	0.5080 (4)	0.1158 (2)	0.4719 (4)	0.0672 (11)
H14A	0.5833	0.0884	0.4732	0.081*
H14B	0.4426	0.0782	0.4422	0.081*
C15	0.5056 (3)	0.14444 (18)	0.5944 (3)	0.0521 (9)
H15	0.5799	0.1279	0.6508	0.063*
C16	0.4023 (4)	0.1148 (3)	0.6359 (4)	0.0790 (13)
H16A	0.4065	0.1351	0.7147	0.119*
H16B	0.4032	0.0565	0.6373	0.119*
H16C	0.3294	0.1334	0.5829	0.119*
C21	0.7121 (2)	0.37048 (16)	0.6550 (2)	0.0341 (6)
C22	0.7636 (2)	0.44648 (17)	0.6866 (3)	0.0395 (7)
C23	0.8611 (3)	0.4719 (2)	0.6460 (3)	0.0549 (9)
H23	0.8928	0.5234	0.6666	0.066*
C24	0.9121 (3)	0.4234 (3)	0.5767 (4)	0.0627 (11)
H24	0.9794	0.4411	0.5518	0.075*
C25	0.8648 (3)	0.3495 (2)	0.5439 (3)	0.0542 (9)
H25	0.8986	0.3159	0.4959	0.065*
C26	0.7668 (3)	0.32490 (18)	0.5821 (3)	0.0419 (7)
H26	0.7345	0.2741	0.5577	0.050*
C31	0.6804 (3)	0.4711 (2)	0.8611 (3)	0.0483 (8)

supplementary materials

H31	0.6914	0.4163	0.8802	0.058*
C32	0.7136 (3)	0.50025 (17)	0.7624 (3)	0.0442 (8)
C33	0.6971 (4)	0.5819 (2)	0.7384 (3)	0.0638 (11)
H33	0.7205	0.6039	0.6732	0.077*
C34	0.6474 (5)	0.6307 (2)	0.8081 (4)	0.0854 (16)
H34	0.6354	0.6855	0.7890	0.102*
C35	0.6154 (4)	0.6014 (3)	0.9041 (4)	0.0853 (16)
H35	0.5821	0.6356	0.9517	0.102*
C36	0.6318 (4)	0.5206 (2)	0.9319 (3)	0.0656 (11)
H36	0.6099	0.4997	0.9986	0.079*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mg1	0.0308 (5)	0.0364 (5)	0.0377 (5)	-0.0051 (4)	0.0085 (4)	-0.0023 (4)
Br1	0.0496 (2)	0.0716 (3)	0.0401 (2)	-0.01271 (17)	0.00923 (14)	0.00894 (16)
O1	0.0412 (13)	0.0673 (16)	0.0586 (15)	0.0132 (11)	0.0029 (11)	-0.0154 (12)
C2	0.059 (3)	0.116 (4)	0.106 (4)	0.027 (3)	0.002 (3)	0.014 (3)
C3	0.129 (6)	0.114 (5)	0.174 (8)	0.051 (5)	-0.082 (6)	-0.039 (5)
C4	0.103 (5)	0.204 (8)	0.133 (6)	0.102 (6)	0.039 (4)	-0.001 (5)
C5	0.080 (4)	0.269 (10)	0.082 (4)	0.081 (5)	0.023 (3)	-0.001 (5)
C6	0.086 (5)	0.277 (12)	0.145 (7)	-0.072 (6)	0.022 (4)	0.027 (7)
O11	0.0477 (12)	0.0308 (11)	0.0408 (11)	-0.0064 (9)	0.0017 (9)	0.0021 (8)
C12	0.065 (2)	0.0477 (19)	0.0407 (17)	-0.0002 (17)	-0.0019 (15)	0.0043 (15)
C13	0.137 (5)	0.065 (3)	0.054 (2)	-0.002 (3)	0.024 (3)	-0.013 (2)
C14	0.072 (3)	0.050 (2)	0.075 (3)	0.0072 (19)	0.009 (2)	-0.0172 (19)
C15	0.054 (2)	0.0317 (16)	0.061 (2)	-0.0068 (14)	-0.0061 (16)	0.0057 (15)
C16	0.090 (3)	0.069 (3)	0.073 (3)	-0.043 (2)	0.010 (2)	0.003 (2)
C21	0.0299 (13)	0.0292 (14)	0.0409 (15)	0.0013 (11)	0.0038 (11)	0.0036 (12)
C22	0.0312 (14)	0.0343 (15)	0.0451 (17)	-0.0061 (12)	-0.0063 (12)	0.0082 (13)
C23	0.0436 (19)	0.054 (2)	0.058 (2)	-0.0204 (16)	-0.0055 (16)	0.0150 (17)
C24	0.0345 (17)	0.086 (3)	0.067 (2)	-0.0100 (18)	0.0113 (16)	0.025 (2)
C25	0.0435 (18)	0.066 (2)	0.057 (2)	0.0104 (17)	0.0198 (16)	0.0174 (18)
C26	0.0397 (16)	0.0364 (16)	0.0507 (18)	0.0063 (13)	0.0129 (14)	0.0064 (13)
C31	0.0481 (18)	0.0392 (17)	0.0478 (19)	0.0046 (14)	-0.0079 (14)	-0.0072 (14)
C32	0.0417 (16)	0.0341 (16)	0.0450 (18)	-0.0045 (13)	-0.0132 (13)	-0.0052 (13)
C33	0.082 (3)	0.0333 (18)	0.055 (2)	-0.0038 (17)	-0.0242 (19)	-0.0040 (16)
C34	0.124 (4)	0.038 (2)	0.065 (3)	0.015 (2)	-0.035 (3)	-0.017 (2)
C35	0.103 (4)	0.064 (3)	0.066 (3)	0.033 (2)	-0.025 (3)	-0.034 (2)
C36	0.069 (2)	0.068 (2)	0.049 (2)	0.013 (2)	-0.0076 (18)	-0.0172 (18)

Geometric parameters (\AA , $^\circ$)

Mg1—O1	2.022 (2)	C14—H14A	0.9900
Mg1—O11	2.030 (2)	C14—H14B	0.9900
Mg1—C21	2.143 (3)	C15—C16	1.492 (5)
Mg1—Br1	2.4750 (10)	C15—H15	1.0000
O1—C5	1.408 (6)	C16—H16A	0.9800
O1—C2	1.562 (6)	C16—H16B	0.9800

C2—C3	1.421 (9)	C16—H16C	0.9800
C2—H2A	0.9900	C21—C26	1.412 (4)
C2—H2B	0.9900	C21—C22	1.424 (4)
C3—C4	1.536 (11)	C22—C23	1.403 (5)
C3—H3A	0.9900	C22—C32	1.485 (5)
C3—H3B	0.9900	C23—C24	1.385 (6)
C4—C5	1.533 (9)	C23—H23	0.9500
C4—H4A	0.9900	C24—C25	1.377 (6)
C4—H4B	0.9900	C24—H24	0.9500
C5—C6	1.425 (11)	C25—C26	1.390 (4)
C5—H5	1.0000	C25—H25	0.9500
C6—H6A	0.9800	C26—H26	0.9500
C6—H6B	0.9800	C31—C36	1.393 (5)
C6—H6C	0.9800	C31—C32	1.399 (5)
O11—C12	1.456 (4)	C31—H31	0.9500
O11—C15	1.463 (4)	C32—C33	1.404 (4)
C12—C13	1.481 (5)	C33—C34	1.384 (6)
C12—H12A	0.9900	C33—H33	0.9500
C12—H12B	0.9900	C34—C35	1.366 (7)
C13—C14	1.475 (6)	C34—H34	0.9500
C13—H13A	0.9900	C35—C36	1.398 (6)
C13—H13B	0.9900	C35—H35	0.9500
C14—C15	1.529 (5)	C36—H36	0.9500
O1—Mg1—O11	97.02 (10)	C13—C14—C15	106.8 (3)
O1—Mg1—C21	114.41 (11)	C13—C14—H14A	110.4
O11—Mg1—C21	106.87 (11)	C15—C14—H14A	110.4
O1—Mg1—Br1	109.97 (9)	C13—C14—H14B	110.4
O11—Mg1—Br1	104.75 (7)	C15—C14—H14B	110.4
C21—Mg1—Br1	120.53 (8)	H14A—C14—H14B	108.6
C5—O1—C2	109.6 (4)	O11—C15—C16	110.0 (3)
C5—O1—Mg1	129.5 (3)	O11—C15—C14	104.7 (3)
C2—O1—Mg1	118.7 (2)	C16—C15—C14	113.8 (3)
C3—C2—O1	102.0 (5)	O11—C15—H15	109.4
C3—C2—H2A	111.4	C16—C15—H15	109.4
O1—C2—H2A	111.4	C14—C15—H15	109.4
C3—C2—H2B	111.4	C15—C16—H16A	109.5
O1—C2—H2B	111.4	C15—C16—H16B	109.5
H2A—C2—H2B	109.2	H16A—C16—H16B	109.5
C2—C3—C4	103.3 (6)	C15—C16—H16C	109.5
C2—C3—H3A	111.1	H16A—C16—H16C	109.5
C4—C3—H3A	111.1	H16B—C16—H16C	109.5
C2—C3—H3B	111.1	C26—C21—C22	114.9 (3)
C4—C3—H3B	111.1	C26—C21—Mg1	116.1 (2)
H3A—C3—H3B	109.1	C22—C21—Mg1	128.9 (2)
C5—C4—C3	101.7 (5)	C23—C22—C21	120.8 (3)
C5—C4—H4A	111.4	C23—C22—C32	119.2 (3)
C3—C4—H4A	111.4	C21—C22—C32	120.0 (3)
C5—C4—H4B	111.4	C24—C23—C22	121.3 (3)
C3—C4—H4B	111.4	C24—C23—H23	119.3

supplementary materials

H4A—C4—H4B	109.3	C22—C23—H23	119.3
O1—C5—C6	109.4 (6)	C25—C24—C23	119.7 (3)
O1—C5—C4	104.4 (6)	C25—C24—H24	120.1
C6—C5—C4	112.8 (7)	C23—C24—H24	120.1
O1—C5—H5	110.0	C24—C25—C26	119.0 (3)
C6—C5—H5	110.0	C24—C25—H25	120.5
C4—C5—H5	110.0	C26—C25—H25	120.5
C5—C6—H6A	109.5	C25—C26—C21	124.2 (3)
C5—C6—H6B	109.5	C25—C26—H26	117.9
H6A—C6—H6B	109.5	C21—C26—H26	117.9
C5—C6—H6C	109.5	C36—C31—C32	121.6 (3)
H6A—C6—H6C	109.5	C36—C31—H31	119.2
H6B—C6—H6C	109.5	C32—C31—H31	119.2
C12—O11—C15	108.6 (2)	C31—C32—C33	117.2 (3)
C12—O11—Mg1	120.80 (18)	C31—C32—C22	121.0 (3)
C15—O11—Mg1	129.73 (19)	C33—C32—C22	121.8 (3)
O11—C12—C13	105.0 (3)	C34—C33—C32	121.1 (4)
O11—C12—H12A	110.7	C34—C33—H33	119.5
C13—C12—H12A	110.7	C32—C33—H33	119.5
O11—C12—H12B	110.7	C35—C34—C33	121.0 (4)
C13—C12—H12B	110.7	C35—C34—H34	119.5
H12A—C12—H12B	108.8	C33—C34—H34	119.5
C14—C13—C12	106.9 (3)	C34—C35—C36	119.7 (4)
C14—C13—H13A	110.3	C34—C35—H35	120.1
C12—C13—H13A	110.3	C36—C35—H35	120.1
C14—C13—H13B	110.3	C31—C36—C35	119.4 (4)
C12—C13—H13B	110.3	C31—C36—H36	120.3
H13A—C13—H13B	108.6	C35—C36—H36	120.3
O11—Mg1—O1—C5	101.5 (5)	C13—C14—C15—C16	115.6 (4)
C21—Mg1—O1—C5	-146.4 (5)	O1—Mg1—C21—C26	-130.5 (2)
Br1—Mg1—O1—C5	-7.1 (5)	O11—Mg1—C21—C26	-24.4 (2)
O11—Mg1—O1—C2	-97.5 (3)	Br1—Mg1—C21—C26	94.8 (2)
C21—Mg1—O1—C2	14.7 (3)	O1—Mg1—C21—C22	52.0 (3)
Br1—Mg1—O1—C2	154.0 (3)	O11—Mg1—C21—C22	158.2 (2)
C5—O1—C2—C3	20.6 (6)	Br1—Mg1—C21—C22	-82.6 (3)
Mg1—O1—C2—C3	-144.0 (4)	C26—C21—C22—C23	0.3 (4)
O1—C2—C3—C4	-38.6 (5)	Mg1—C21—C22—C23	177.7 (2)
C2—C3—C4—C5	43.7 (7)	C26—C21—C22—C32	179.6 (3)
C2—O1—C5—C6	127.7 (6)	Mg1—C21—C22—C32	-2.9 (4)
Mg1—O1—C5—C6	-69.9 (7)	C21—C22—C23—C24	-1.6 (5)
C2—O1—C5—C4	6.8 (7)	C32—C22—C23—C24	179.0 (3)
Mg1—O1—C5—C4	169.2 (4)	C22—C23—C24—C25	1.7 (5)
C3—C4—C5—O1	-29.7 (8)	C23—C24—C25—C26	-0.4 (5)
C3—C4—C5—C6	-148.4 (7)	C24—C25—C26—C21	-1.0 (5)
O1—Mg1—O11—C12	60.3 (3)	C22—C21—C26—C25	1.1 (4)
C21—Mg1—O11—C12	-57.9 (3)	Mg1—C21—C26—C25	-176.7 (2)
Br1—Mg1—O11—C12	173.1 (2)	C36—C31—C32—C33	0.8 (5)
O1—Mg1—O11—C15	-131.4 (3)	C36—C31—C32—C22	-179.0 (3)
C21—Mg1—O11—C15	110.4 (3)	C23—C22—C32—C31	-136.9 (3)

Br1—Mg1—O11—C15	-18.6 (3)	C21—C22—C32—C31	43.7 (4)
C15—O11—C12—C13	-28.8 (4)	C23—C22—C32—C33	43.3 (4)
Mg1—O11—C12—C13	141.7 (3)	C21—C22—C32—C33	-136.1 (3)
O11—C12—C13—C14	25.2 (5)	C31—C32—C33—C34	-1.6 (5)
C12—C13—C14—C15	-12.7 (5)	C22—C32—C33—C34	178.2 (3)
C12—O11—C15—C16	-102.0 (3)	C32—C33—C34—C35	1.6 (6)
Mg1—O11—C15—C16	88.6 (3)	C33—C34—C35—C36	-0.7 (7)
C12—O11—C15—C14	20.7 (4)	C32—C31—C36—C35	0.1 (5)
Mg1—O11—C15—C14	-148.7 (2)	C34—C35—C36—C31	-0.1 (6)
C13—C14—C15—O11	-4.6 (4)		

Fig. 1

