

# A new species of the basal araneomorph spider genus *Ectatosticta* (Araneae, Hypochilidae) from China

Norman I. Platnick<sup>1,†</sup>, Peter Jäger<sup>2‡</sup>

**1** Division of Invertebrate Zoology, American Museum of Natural History, New York, New York USA **2** Forschungsinstitut Senckenberg, Frankfurt am Main, Germany

† [urn:lsid:zoobank.org:author:3DA57711-13BB-4F42-AE8D-03CD8BE779C5](https://doi.org/urn:lsid:zoobank.org:author:3DA57711-13BB-4F42-AE8D-03CD8BE779C5)

‡ [urn:lsid:zoobank.org:author:A79CD670-BC87-4A1B-8EDF-B391514CCB2A](https://doi.org/urn:lsid:zoobank.org:author:A79CD670-BC87-4A1B-8EDF-B391514CCB2A)

Corresponding author: Norman I. Platnick ([platnick@amnh.org](mailto:platnick@amnh.org))

---

Academic editor: Pavel Stoev | Received 17 February 2009 | Accepted 16 April 2009 | Published 29 July 2009

---

[urn:lsid:zoobank.org:pub:1B2DBE14-0767-4362-B814-6A4D94ABAC30](https://doi.org/urn:lsid:zoobank.org:pub:1B2DBE14-0767-4362-B814-6A4D94ABAC30)

---

**Citation:** Platnick NI, Jäger P (2009) A new species of the basal araneomorph spider genus *Ectatosticta* (Araneae, Hypochilidae) from China. In: Stoev P, Dunlop J, Lazarov S (Eds) A life caught in a spider's web. Papers in arachnology in honour of Christo Deltchev. ZooKeys 16: 209-215. doi: 10.3897/zookeys.16.231

---

## Abstract

The hypochilid spider *Ectatosticta davidi* (Simon) is redescribed on the basis of adults from Mt. Taibaishan in Shaanxi Province, China; the specimens from Qinghai Province previously identified as *E. davidi* by most modern authors belong to a new species described as *E. deltshevi*.

## Keywords

Araneae, Araneomorphae, Hypochilidae, *Ectatosticta*, China

## Introduction

The spiders of the family Hypochilidae have long been recognized as the most primitive of all araneomorph, or true, spiders. In modern cladistic analyses, they appear as the sister group to all other araneomorphs (Platnick 1977; Forster et al. 1987; Griswold et al. 2005).

Only two genera of hypochilids are known, *Hypochilus* Marx from the United States, and *Ectatosticta* Simon from China. Although 10 species of *Hypochilus* are now

recognized, for *Ectatosticta* only the type species, *E. davidi* (Simon, 1888), has been described, and that species has been known from only a few specimens.

Even the type locality of *E. davidi* is problematic. In the original description, Simon (1888) indicated that the male and female syntypes were collected by A. David from “Montagnes au nord de Péking”. Those specimens are now labeled only “Sina”, but Simon (1892) later listed the species as being from “China merid.” (southern China), which hardly describes areas north of Beijing. Many decades ago, one of Simon’s female specimens from “Inkiaphou” in Shaanxi Province, China, was donated to Dr Willis Gertsch; the label with that specimen reads “Inkiaphou, Chine méridionale, Mus. Nat. d’Hist. Nat. coll. E. Simon”. According to Schenkel (1963: 8), A. David collected at “Inkia-fu, Süd Schensi” in 1873. Repeated attempts by Chinese colleagues to recollect the species in the mountains north of Beijing have all been unsuccessful; Forster et al. (1987: 24) therefore concluded that the syntypes were probably collected at Inkiaphou. The exact location of Inkiaphou (in French) or Inkia-fu (in German) is also uncertain, but those names might refer to Inkiapo, now known as Yinjiapo (on Mt. Qinling) in Shaanxi (near 34°00’N, 108°00’E).

The redescription and illustrations of *E. davidi* provided by Forster et al. (1987) and other modern authors were based primarily on specimens taken in Qinghai Province, to the west of Inkiaphou. The collection of a good series of specimens, by the second author and colleagues, at Mt. Taibaishan in Shaanxi Province, has led us to reexamine the available material. We conclude that there are at least two species of *Ectatosticta*, that the specimens from Mt. Taibaishan and Inkiaphou in Shaanxi Province belong to *E. davidi*, and that the specimens from Qinghai Province described and illustrated by Forster et al. (1987) and other modern authors belong instead to a different species, here described as *E. deltshevi*.

The format of the descriptions follows that of Forster et al. (1987). It is with great pleasure that we dedicate this paper, and the new species, to our Bulgarian colleague, Dr Christo Deltshev, in honor of his many contributions to arachnology. The material examined is housed in the following collections:


- AMNH** American Museum of Natural History, New York  
**IZCAS** Institute of Zoology, Chinese Academy of Sciences, Beijing  
**MNHN** Muséum National d’Histoire Naturelle, Paris  
**SMF** Forschungsinstitut Senckenberg, Frankfurt am Main

### *Ectatosticta davidi* (Simon)

Figs 1-4

*Hypochilus davidi* Simon, 1888: ccviii (one male and one female syntypes, supposedly from mountains north of Beijing but probably from Inkiaphou, Shaanxi Province, China, in MNHN, examined).

*Ectatosticta davidi*: Simon 1892: 204, figs. 143-146, 148-149. Gertsch 1958: 13, figs. 10, 19, 22-31. Lehtinen 1967: 298, fig. 15.



**Figures 1-4.** *Ectatosticta davidi* (Simon): **1** left male palp, prolateral view **2** same, ventral view **3** same, retrolateral view **4** female spermathecae, dorsal view.

**Diagnosis.** Males of *E. davidi* can be distinguished from those of *E. deltshevi* by the details of the distal modifications on the retrolateral side of the palpal tibia. In *E. davidi*, there is a small triangular lobe followed apically by a large ledge bearing 5-7 thickened setae, all closely appressed (Fig. 3); in *E. deltshevi*, there are only four thickened setae, and the most dorsal of those is smaller and well separated from the main group of three (Forster et al. 1987: fig. 81). Females of *E. davidi* (Fig. 4) have the inner pair of spermathecae relatively larger and more robust than is the case in *E. deltshevi* (Forster et al. 1987: fig. 82).

**Male.** Total length 9.57. Carapace 4.61 long, 3.78 wide, yellow, with margins of pars cephalica and pair of broad marginal bands mottled with purplish brown, similar mottling along midline, with additional pair of purple lines originating anterior of thoracic groove, extending to posterior median eyes; clypeus at middle about twice AME diameter in height. Eye sizes and interdistances: AME 0.09, ALE 0.18, PME 0.21, PLE 0.15; AME-AME 0.15, AME-ALE 0.26, PME-PME 0.22, PME-PLE 0.15, ALE-PLE 0.06; MOQ length 0.37, front width 0.33, back width 0.63. Chelicerae yellow, unmarked; endites light brown, darkened distally; labium light brown, darkened distally; sternum light brown, with three broad, transverse bands of purple pigment covering most of surface. Legs light brown, with femora,

patellae, and tibiae darkened distally. Femur I more than twice as long as carapace. Leg spination (only surfaces bearing spines listed): femora: I d3-0-0, p0-4-1, r1-3-1; II d3-0-0, p0-3-2, r3-3-1; III d3-0-0, p1-2-1, r3-2-1; IV d4-0-0, p1-3-1, r0-2-2; tibiae: I d1-0-1, p2-2-1, v4-4-3, r2-3-1; II d1-0-1, p2-2-1, v4-4-3, r2-2-1; III d1-0-1, p1-2-1, v2-2-2, r1-2-1; IV d1-0-1, p1-2-1, v2-2-1, r1-2-1; metatarsi: I d1-0-0, p1-2-2, v2-4-4, r1-1-0; II, III d1-0-0, p1-1-0, v4-4-2, r1-1-0; IV p1-0-1, v2-2-0, r1-0-1. Dorsum of abdomen white with about six purple chevrons, posterior ones stronger than anterior ones, covering most of posterior one-third, sides and venter mottled except on lung covers. Retrolateral side of palpal tibia distally with small triangular lobe followed distally by large lobe bearing 5-7 modified setae, all closely appressed (Fig. 3); embolus rounded, more smoothly curved than in *E. deltshevi* (Figs 1-3).

**Female.** As in male, except as noted. Total length 10.98. Carapace 4.20 long, 2.97 wide. Eye sizes and interdistances: AME 0.11, ALE 0.20, PME 0.20, PLE 0.18, AME-AME 0.14, AME-ALE 0.25, PME-PME 0.30, PME-PLE 0.15, ALE-PLE 0.09; MOQ length 0.37, front width 0.36, back width 0.70. Chelicerae light brown; sternum with only one anterior transverse light band interrupting purple mottling. Femora darkened medially and distally, tibiae and metatarsi with basal, median, and distal darkenings. Leg spination (only surfaces bearing spines listed): femora: I d2-0-0, p1-2-3, r2-4-2; II d2-1-0, p1-1-2, r2-3-1; III d2-3-0, p1-1-1, r3-3-2; IV d3-1-0, p0-2-1, r0-1-2; tibiae: I d1-0-1, p2-1-1, v2-5-2, r1-2-1; II d1-0-1, p1-2-1, v4-5-2, r1-2-1; III d1-0-1, p1-2-1, v2-2-2, r1-1-1; IV d1-0-1, p1-1-0, v2-3-2, r1-1-0; metatarsi: I d1-0-0, p1-1-0, v2-4-4, r1-1-0; II d1-0-0, p1-1-0, v4-4-4, r1-1-0; III d1-0-0, p1-1-0, v4-2-2, r1-1-0; IV p0-1-0, v1p-1p-1p, r0-1-0. Inner pair of spermathecae relatively long, robust throughout their length (Fig. 4).

**Material Examined.** CHINA: **Shaanxi:** Inkiaphou (AMNH), 1♀; presumably Inkiaphou, label reads only "Sina" (MNHN 7285), 1♂, 1♀ (syntypes); Mt. Taibais-han, S flanks, above Houshenzi, 33°54'43.98"N, 107°46'44.06"E, June 12-15, 1997, tree line, scattered mixed coniferous/*Rhododendron* forest, elev. 3050 m (P. Jäger, C. Fischer, AMNH), 1♂, 2♀, same (IZCAS), 1♂, 1♀, same (SMF), 1♂, 8♀ (not all dissected, some may be juvenile), June 25, 1997, primary broad-leaved forest, elev. 2500-2600 m (J. Martens, P. Jäger, SMF), 1♀.

**Natural History.** According to observations by the second author at Mt. Taibais-han, *E. davidi* occurs in stony debris in open, semi-open, and forest-covered habitats. One part of the elongate sheet-web (Fig. 5) is situated at the surface and connected to a stone or rock and surrounding vegetation. The apparently larger part of the web leads, as a narrow band, deep into the debris. Spiders cannot be seen or lured out of their retreat during the day. When it gets dark, the spiders sit close to the surface part of the web, ready to escape to their retreat. One male was found walking around, apparently searching for a female. Another male was observed during courtship, at the margin of a female's surface web; that male touched the web only with its first pair of legs. The third male was taken in its own web.

**Distribution.** Known with certainty only from Shaanxi Province, China (Fig. 6).



Figure 5. Web of *Ectatosticta davidi* (Simon) from Mt. Taibaishan, Shaanxi, China.




Figure 6. Map of China, showing records of *Ectatosticta*.

***Ectatosticta deltshevi* sp. n.**

urn:lsid:zoobank.org:act:FF9F3FC8-8CB0-4912-9D8E-75EA18F2E57E

*Ectatosticta davidi* (misidentification): Li and Zhu 1984: 510, figs. a-g. Forster et al. 1987: 23, figs. 6-16, 18-20, 23-24, 31-36, 78-82. Song et al. 1999: 41, figs. 11D, 17Q-T. Hu 2001: 69, figs. 1.1-6. Song et al. 2001: 64, figs. 24A-E.

**Type. Male holotype** from Huangyuan County, Qinghai Province, China, 15.IX.1984, Z.S. Li leg., deposited in IZCAS.

**Etymology.** The specific name is a patronym in honor of Dr Christo Deltshev.

**Diagnosis.** Males of *E. deltshevi* can be distinguished by the form of the specialized setae situated distally on the retrolateral ledge of the palpal tibia, as detailed above, and by the distally more sharply angled embolus (Forster et al. 1987: fig. 79). Females have the inner pair of spermathecae shorter, less robust, and distally less sclerotized than in *E. davidi*.

**Male.** Described by Forster et al. (1987: 23), as *E. davidi*.

**Female.** Described by Forster et al. (1987: 23), as *E. davidi*.

**Other material examined.** CHINA: **Qinghai:** Huangyuan County, Sept. 10, 1983 (Z.S. Li, AMNH), 1♂, 1♀, Sept. 15, 1984 (Z.S. Li, IZCAS), 2♂, 3♀.

**Distribution.** Known with certainty only from Huangyuan County, Qinghai Province, China. A juvenile taken at an elevation of 2300-2700 m in Beishan National Park, 120 km NE Xining City, Qinghai Province, China, 36°56'N, 102°29'E, 23.V – 8.VI.1996, J. Martens leg. (SMF) may also belong to this species (Fig. 6).

## Acknowledgements

We thank Dr Christine Rollard (MNHN) for loaning the syntypes, the late Prof. Song Daxiang and Dr Li Shuqiang (IZCAS) for specimens and information, Dr Mohammad Shadab (AMNH) for providing the illustrations, and Liu Jie (IZCAS) for constructing the map. The expedition of Jochen Martens and Peter Jäger to China in 1997 was partly supported by the Stifterverband für die Deutsche Wissenschaft.

## References

- Forster RR, Platnick NI, Gray MR (1987) A review of the spider superfamilies Hypochiloidea and Austrochiloidea (Araneae, Araneomorphae). *Bulletin of the American Museum of Natural History* 185: 1-116.
- Gertsch WJ (1958) The spider family Hypochilidae. *American Museum Novitates* 1912: 1-28.
- Griswold CE, Ramírez MJ, Coddington JA, Platnick NI (2005) Atlas of phylogenetic data for entelegyne spiders (Araneae: Araneomorphae: Entelegynae) with comments on their phylogeny. *Proceedings of the California Academy of Sciences ser. 4, 56 (Supplement II)*: 1-324.
- Hu JL (2001) Spiders in Qinghai-Tibet Plateau of China. Henan Science and Technology Publishing House, Henan, 658 pp.

- Lehtinen PT (1967) Classification of the cribellate spiders and some allied families. *Annales Zoologici Fennici* 4: 199-468.
- Li ZS, Zhu CD (1984) *Ectatosticta davidi* (Simon, 1888) of China (Araneae: Hypochilidae). *Journal of Norman Bethune Medical University* 10: 510.
- Platnick NI (1977) The hypochiloid spiders: a cladistic analysis, with notes on the Atypoidea (Arachnida, Araneae). *American Museum Novitates* 2627: 1-23.
- Schenkel E (1963) Ostasiatische Spinnen aus dem Muséum d'Histoire Naturelle de Paris. *Mémoires du Muséum National d'Histoire Naturelle, ser. A, Zoologie* 25: 1-481.
- Simon E (1888) [without title]. *Annales de la Société Entomologique de France, ser. 6, 8*: ccviii-ccix.
- Simon E (1892) *Histoire naturelle des araignées*. Roret, Paris, 1: 1-256.
- Song DX, Zhu MS, Chen J (1999) *The spiders of China*. Hebei Science and Technology Publishing House, Shijiazhuang, 640 pp.
- Song DX, Zhu MS, Chen J (2001) *The fauna of Hebei, China: Araneae*. Hebei Science and Technology Publishing House, Shijiazhuang, 508 pp.