

INSECTA MUNDI

A Journal of World Insect Systematics

0008

A new species of *Tetraopes* Schoenherr (Coleoptera: Cerambycidae)

Frederick W. Skillman, Jr.

P. O. Box 375

Pearce, Arizona 85625

Date of Issue: 25 April 2007

Frederick W. Skillman, Jr.

A new species of *Tetraopes* Schoenherr (Coleoptera: Cerambycidae)

Insecta Mundi 0008: 1-3

Published in 2007 by

Center for Systematic Entomology, Inc.

P. O. Box 147100

Gainesville, FL 32604-7100 U. S. A.

<http://www.centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod taxon. Manuscripts considered for publication include, but are not limited to, systematic or taxonomic studies, revisions, nomenclatural changes, faunal studies, book reviews, phylogenetic analyses, biological or behavioral studies, etc. **Insecta Mundi** is widely distributed, and referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc.

As of 2007, **Insecta Mundi** is published irregularly throughout the year, not as a quarterly issues. As manuscripts are completed they are published and given an individual number. Manuscripts must be peer reviewed prior to submission, after which they are again reviewed by the editorial board to insure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Managing editor: Paul E. Skelley, e-mail: skellep@doacs.state.fl.us

Production editor: Michael C. Thomas, e-mail: thomasm@doacs.state.fl.us

Printed copies deposited in libraries of:

CSIRO, Canberra, ACT, Australia

Museu de Zoologia, São Paulo, Brazil

Agriculture and Agrifood Canada, Ottawa, Ontario, Canada

The Natural History Museum, London, England

Muzeum I Instytut Zoologii Pan, Warsaw, Poland

National Taiwan University, Taipei, Taiwan

California Academy of Sciences, San Francisco, CA, USA

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

Field Museum of Natural History, Chicago, IL, USA

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Electronic copies in PDF format:

Printed CD mailed to all members at end of year.

Florida Center for Library Automation: purl.fcla.edu/fcla/insectamundi

Author instructions available on the Insecta Mundi page at:

<http://www.centerforsystematicentomology.org/insectamundi/>

ISSN 0749-6737

A new species of *Tetraopes* Schoenherr (Coleoptera: Cerambycidae)

Frederick W. Skillman, Jr.

P. O. Box 375
Pearce, Arizona 85625
bycid@vtc.net

Abstract. A new species of the genus *Tetraopes* Schoenherr, *T. huetheri* Skillman, is described from South Dakota, USA. A modification to the existing key is provided. The new species is illustrated. The probable host plant is identified and illustrated.

Key Words. Insect, Coleoptera, Cerambycidae, *Tetraopes*, new species, key

Introduction

Recent collecting in western South Dakota has yielded a new species of the genus *Tetraopes* Schoenherr. The new species was discovered while sweeping near Mirror Lakes in Lawrence County.

Collection abbreviations listed in text are as follows: JPHC, Jeffrey P. Huether, Geneva, NY; RHTC, Robert H. Turnbow, Fort Rucker, AL; FSCA, Florida State Collection of Arthropods, Gainesville, FL; FWSC, Fred W. Skillman, Pearce, AZ; RMBC, R. Michael Brattain, Lafayette, IN; EMEC, Essig Museum of Entomology, Berkeley, CA.

***Tetraopes huetheri* Skillman, new species**
(Fig. 1, 2, 3)

Description: Male: Form (Fig. 1-2) small, slightly flattened; integument pale orange and reddish, underside dark chestnut brown; pubescence not obscuring punctuation, moderately dense, appressed, yellowish white, few, long, erect, pale, setae.

Head reddish, narrower than pronotum, front convex, moderately coarsely, densely, separately punctate, most punctures giving rise to a short, yellowish white, erect seta; pubescence fine, sparse, appressed, yellowish white, thinning towards vertex; genae slightly longer than lower eye lobes; maxillary palpi slightly enlarged, acute at apices; mandibles feebly arcuate, deeply excavated externally at base; labrum reddish, narrow, short, wedge shaped, broadly rounded at apex, clothed with a few, erect, white setae; eyes small, very finely faceted, upper lobes widely separated; area between antennal tuberculi shallowly impressed; antennae vary from dark reddish brown to dark brown, extending to fourth abdominal segment, first three segments shining, dorsal surface with numerous short black hairs, ventral surface sparsely clothed with white pubescence and a few, short, erect, black hairs, shining beneath; segments from fourth densely clothed with very fine, appressed, dark brown pubescence, narrowly yellowish white annulate at bases and apices with a few erect, black hairs at apices, more numerous beneath.

Pronotum broader than long, sides prominently obtusely tuberculate; apex shallowly impressed; base very shallowly impressed laterally; umbone reddish, broad, large, almost hexagonal, shallowly convex, slightly to moderately elevated, sides not delimited, disk with fine, scattered to moderate punctuation, some giving rise to very short, pale erect hairs, each side with two black spots; area surrounding umbone with sparse to moderately dense, short, appressed, yellowish white pubescence, especially dense on basal side of lateral tubercles; most tubercles having a few, long, erect, pale setae; base slightly elevated at scutellum, fringed with mostly short, yellowish white hair and having a transverse band of color ranging from slightly darker than surrounding area to a wide, dark brown band that can extend to the basal black spots. Prosternum narrow, feebly impressed at apex, shining, finely, transversely wrinkled, sparsely clothed with yellowish white, appressed pubescence, a few, longer, suberect, pale hairs present; meso- and metasterna micropunctate, moderately clothed with appressed yellowish white pubescence. Scutellum dark brown to black scantily clothed with black pubescence.

Elytra about 2 times as long as broad, sides parallel; punctures over basal ½ coarse, dense, subcontiguous becoming smaller and less dense toward apex; pubescence moderate, pale, suberect,



Figure 1-3. *Tetraopes huetheri* Skillman, n.sp. 1) holotype male; 2) *Tetraopes huetheri* Skillman, holotype male umbone; 3) *Tetraopes huetheri* Skillman, allotype female fifth abdominal segment showing notch and longitudinal impression.

underlying pubescence dense, short, appressed, yellowish-white, denser over apical 1/3; humeri black; two black, shining, sutural spots present at about basal 1/4, two larger black spots also present at midelytra at beginning of apical 1/3, apices rounded and slightly darkened. Legs long, micropunctate; femora moderately to densely clothed with yellowish-white, appressed pubescence, long, pale hairs sparse; tibiae clothed with yellowish-white, suberect pubescence; tarsi with claws appendiculate.

Abdomen densely micropunctate, densely clothed with fine, appressed, yellowish-white pubescence; fifth sternite subtruncate at apex. Length 7.0 - 8.5 mm.

Female: Form and color similar to male, but more robust. Front of head strongly convex. Antenna shorter. Fifth abdominal segment broadly truncate with small notch at apex (Fig. 3) and impressed longitudinally at middle extending from base half way to apex. Length 8.0 - 9.8 mm

Type material. HOLOTYPE male and allotype from "U.S.A., South Dakota: Lawrence Co., Upper Mirror Lake west end, 23 Aug. 2006, on *Asclepias verticillata*, J and C Huether" (deposited in FSCA). Paratypes (10 males, 4 females): same data as holotype (6); "U.S.A. South Dakota: Lawrence Co., Mirror Lake fish hatchery NW side, 24 Aug. 2005, J. Huether" (3); "S.D. Lawrence Co., Mirror Lakes, 24 Aug. 2005, R. Turnbow" (5). Paratypes deposited in JPHC, RHTC, FWSC, RMBC, EMEC.

A single rubbed and greased specimen not included in the type series is labeled: "WY: Cook Co., Vic. Entrance Devil's Tower Natl. Mnt., 22 Aug. 2006, J. and C. Huether".

Whorled milkweed, *Asclepias verticillata* L. (Fig. 4), is assumed to be the host as all specimens were swept from this plant. The presumed host ranges into 41 states, including WY. More collecting is needed in this area to help determine the range of *T. huetheri*.

Diagnosis. *Tetraopes huetheri* cannot be confused with any other described North American species. The combination of small size, prominent pronotal tubercles, low to moderately elevated umbone, appendiculate tarsal claws, and lack of elytral chevron serve to distinguish it.

Etymology. This species is dedicated to Jeffrey P. Huether who caught the first specimens and provided the picture of the presumed host.



Figure 4. Whorled milkweed, *Asclepias verticillata* L.

The following couplets should be inserted into Chemsak and Noguera's 2003 key to known species of *Tetraopes*:

- 2(1). Thoracic umbone narrow, disk convex; sides of pronotum not lobed; elytral chevron, if present, broad; form cylindrical 3
 — Thoracic umbone broad, disk flattened to shallowly convex; sides of pronotum lobed or obtusely tuberculate; elytral chevron, if present, narrow, transverse; form broad 2A
- 2A(2). Thoracic umbone almost quadrate, disk flattened; sides of pronotum moderately lobed; median elytral chevron, if present, narrow, transverse, located at basal one-third; form broad; length 12mm; Mexico (Jalisco) ***T. crassus* Chemsak and Noguera**
 — Thoracic umbone almost hexagonal, disk shallowly convex; sides of pronotum prominently, obtusely tuberculate; no elytral chevron present; form broad, slightly flattened; length 7.2 - 9.8 mm; U.S.A (South Dakota) ***T. huetheri* Skillman, n. sp.**

Acknowledgments

I thank the following: Jeffrey P. Huether, Geneva, NY and Robert H. Turnbow, Fort Rucker, AL, for loan of specimens; Paul E. Skelley, FSCA, Gainesville, FL, Trevor R. Smith, FSCA, Gainesville, FL, and Warren E. Boutchia, Tampa, FL, for technical assistance; Edward Cope, Bailey Hortorium Herbarium, Ithica, NY, for host plant identification; Michael C. Thomas and Dr. Skelley for manuscript review; Florida State Collection of Arthropods, Gainesville, FL, for facilities and equipment.

Literature cited

Chemsak, J. A., and Noguera, F. A. 2003. New Species of the Genus *Tetraopes* Schoenherr (Coleoptera: Cerambycidae). Pan-Pacific Entomologist 79: 237-244.

Accepted: April 20, 2007

