

INSECTA MUNDI

A Journal of World Insect Systematics

0039

First record of *Diaphorina citri* Kuwayama
(Hemiptera: Psyllidae) from the Sultanate of Oman

Talal S. Al-Zadjali
Directorate General of Agricultural Research (Rumais)
Ministry of Agriculture
P.O. Box 2512, P.C. 111, CPO
Muscat, Sultanate of Oman

Rohani Ibrahim
Department of Plant Protection
University Putra Malaysia (UPM)
Selangor, Malaysia

Ahmed K. Al-Rawahi
University of Nizwa
Barkat Al-MuZ
Nizwa, Sultanate of Oman

Date of Issue: June 6, 2008

Talal S. Al-Zadjali, Rohani Ibrahim, and Ahmed K. Al-Rawahi
First record of *Diaphorina citri* Kuwayama
(Hemiptera: Psyllidae) from the Sultanate of Oman
Insecta Mundi 0039: 1-3

Published in 2008 by

Center for Systematic Entomology, Inc.
P. O. Box 147100
Gainesville, FL 32614-7100 U. S. A.
<http://www.centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod taxon. Manuscripts considered for publication include, but are not limited to, systematic or taxonomic studies, revisions, nomenclatural changes, faunal studies, book reviews, phylogenetic analyses, biological or behavioral studies, etc. **Insecta Mundi** is widely distributed, and referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc.

As of 2007, **Insecta Mundi** is published irregularly throughout the year, not as quarterly issues. As manuscripts are completed they are published and given an individual number. Manuscripts must be peer reviewed prior to submission, after which they are again reviewed by the editorial board to insure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Managing editor: Paul E. Skelley, e-mail: insectamundi@gmail.com
Production editor: Michael C. Thomas, e-mail: thomasm@doacs.state.fl.us
Editorial board: J. H. Frank, M. J. Paulsen

Printed copies deposited in libraries of:

CSIRO, Canberra, ACT, Australia
Museu de Zoologia, São Paulo, Brazil
Agriculture and Agrifood Canada, Ottawa, Ontario, Canada
The Natural History Museum, London, England
Muzeum I Instytut Zoologii Pan, Warsaw, Poland
National Taiwan University, Taipei, Taiwan
California Academy of Sciences, San Francisco, CA, USA
Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA
Field Museum of Natural History, Chicago, IL, USA
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Electronic copies in PDF format:

Printed CD mailed to all members at end of year.
Florida Center for Library Automation: purl.fcla.edu/fcla/insectamundi
University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Author instructions available on the *Insecta Mundi* page at:
<http://www.centerforsystematicentomology.org/insectamundi/>

Printed Copy	ISSN 0749-6737
On-Line	ISSN 1942-1354
CD-ROM	ISSN 1942-1362

First record of *Diaphorina citri* Kuwayama
(Hemiptera: Psyllidae) from the Sultanate of Oman

Talal S. Al-Zadjali

Directorate General of Agricultural and Livestock Research (Rumais)

Ministry of Agriculture

P.O. Box 2512, P.C. 111, CPO

Muscat, Sultanate of Oman

Rohani Ibrahim

Department of Plant Protection

University Putra Malaysia (UPM)

Selangor, Malaysia

Ahmed K. Al-Rawahi

University of Nizwa

Barkat Al-MuZ

Nizwa, Sultanate of Oman

Mexican acid lime, *Citrus aurantifolia* (L.) Swingle (Rutaceae), is the most popular citrus species in Oman and is the second most important economic crop. The cultivated area of acid lime in the period of 2001-2004 ranged between 4460-2880 feddans (1 feddan = 1.038 acres = 0.420 hectares), with an annual production of about 8160-5875 metric tonnes (Anonymous 2004).

Previous surveys did not reveal any record of psyllids on citrus in Oman. However, the most important psyllid species attacking citrus, *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae), was recently observed on *C. aurantifolia*. It was observed first at Barka (Al-Batinah region) on 25 September 2005. Since then it was regularly observed in most citrus orchards of the Al-Batinah region. *Diaphorina citri* was found to be present in three locations in the Sultanate; Barka (23.709°N 57.887°E), Al-Rustaq (23.391°N 57.424°E) in the Al-Batinah region, and Masirat Al-Rawajeh (23.044°N 57.670°E) at Jabal Al-Akhdar near Nizwa in the A' Dakhliyah region (Figure 1).

In Barka, a D-vac suction trap was used to collect psyllids in the period from November 2007 to April 2008 in fortnightly visits to 10 acid lime trees. The total number of psyllids collected per monthly sample from the months of November 2007 through April 2008 was 560, 420, 1150, 103, 2454 and 2100 adults respectively. Sixty adults were collected by sweep net from Al-Rustaq on 19 June 2007 and one adult from Masirat Al-Rawajeh on 11 July 2007.

All stages of *D. citri* were observed in the citrus fields. Acid lime trees were found to be infested with clear symptoms of *D. citri* at Barka (Figure 2). Symptoms of damage include malformation and distortion of leaves (Figure 3). Nymphs and adults (Figure 4) were found feeding on acid lime leaves, and sooty mold had developed from the honeydew secretion of the psyllids (Figure 5).

Diaphorina citri was reared successfully on *C. aurantifolia* and orange jasmine (*Murraya paniculata* (L.) Jack), and all the above-mentioned symptoms were observed on *C. aurantifolia* in the laboratory at Rumais.

The importance of *D. citri* lies in its direct feeding that result in the distortion of leaves and even the death of new growth. But most importantly, it is a vector of citrus huanglongbing (HLB), or citrus greening disease (Halbert and Manjunath 2004) caused by three species of bacteria, *Candidatus Liberibacter* spp. From the Arabian Peninsula, *D. citri* has been reported from Saudi Arabia, Yemen, and very recently from United Arab Emirates (Wooler et al. 1974; Burckhardt 1981, 1985; Bové 1986; da Graca 1991; Burckhardt 2008). Also, it has been reported from eastern Iran (Bové et al. 2000). *Candidatus Liberibacter asiaticus* and *Candidatus Liberibacter africanus* have been reported in Saudi Arabia and Yemen (Toorawa 1998).

To our knowledge, this is the first report of *D. citri* in Oman. We believe its introduction occurred some time before its detection in 2005 because of its wide distribution in the Sultanate. Therefore, moni-

toring for citrus huanglongbing should start in the Sultanate as it has been detected in the neighboring countries such as Saudi Arabia and Yemen (Toorawa 1998, Halbert and Manjunath 2004). Monitoring can start by inspecting citrus trees and looking for disease symptoms, which include yellow shoots, asymmetrical mottling, and chlorosis of the leaves (Capoor et al. 1974), and by using molecular approaches such as PCR, strain-specific DNA probes, electron microscopy, and ELISA tests for detection of the bacteria (Roistacher 1991, Garnier and Bové 1993, Halbert and Manjunath 2004).

Acknowledgments

The authors are grateful to Dr. Daniel Burckhardt (Naturhistorisches Museum Basel, Switzerland) and Dr. Diana Percy (UBC Botanical Garden and Centre for Plant Research, University of British Columbia, Canada) for confirming *D. citri* identity. We are extremely grateful to Dr. Daniel Burckhardt (Naturhistorisches Museum Basel, Switzerland) and Prof. Yusof Ibrahim (Department of Plant Protection, University of Putra Malaysia) for their useful suggestions and critical comments on the manuscript. Our thanks are extended to Dr. Ahmed Al-Bakri, DG of Agricultural and Livestock Research (Rumais) for providing the financial support and Mr. Ahmed Al-Darmaki, assistant technician at the laboratory.

Literature Cited

- Anonymous. 2004.** Agriculture Census 2004/2005. Ministry of Agriculture and Fisheries; Muscat, Sultanate of Oman. [available at: <http://www.mafstat.gov.om/stats/home.asp> (in Arabic)].
- Bové, J. M. 1986.** Greening in the Arab Peninsula: Towards new techniques for detection and control. Food and Agriculture Organization of the United Nation (FAO), Plant Protection Bulletin 34(1): 7-14.
- Bové, J. M., J. L. Danet, K. Bananej, N. Hassanzadeh, M. Taghizadeh, M. Salehi, and M. Garnier. 2000.** Witches' broom disease of lime (WBDL) in Iran. p. 207-212. *In: Proceedings of 14th Conference of International Organization of Citrus Virologists. IOCV; Riverside, California, US. p. 152.*
- Burckhardt, D. 1981.** Insects of Saudi Arabia, Sternorrhyncha: Suborder Psylloidea. Fauna of Saudi Arabia 3: 213-226.
- Burckhardt, D. 1985.** Sternorrhyncha: Suborder Psylloidea of Saudi Arabia (Part 2). Fauna of Saudi Arabia 7: 141-159.
- Burckhardt, D. 2008.** Order Sternorrhyncha, superfamily Psylloidea. Arthropod fauna of the United Arab Emirates (UAE) 1: 159-169.
- Capoor, S. P., D. G. Rao, and S. M. Viswanath. 1974.** Greening disease of citrus in the Deccan Trap Country and its relationship with the vector, *Diaphorina citri* Kuwayama. p. 43-49. *In: Proceedings of the 6th Conference of International Organization of Citrus Virologists. IOCV; Berkeley, California, US. p. 232.*
- da-Graca, J. V. 1991.** Citrus greening disease. Annual Review of Phytopathology 29: 109-136.



Figure 1a,b. Distribution of *D. citri* in the Sultanate of Oman (map 1b provided by: www.worldatlas.com).



Figure 2-5. *Diaphorina citri* and tree damage. **2)** *D. citri* infested tree (Barka). **3)** Distorted acid lime leaves. **4)** Adult of *D. citri* (lateral view). **5)** Sooty mold on acid lime leaves.

- Garnier, M., and J. M. Bové. 1993.** Citrus greening disease and the greening bacterium. p. 212-219. *In: Proceedings of 12th Conference of International Organization of Citrus Virologists. IOCV; Riverside, California, US. p. 428.*
- Halbert, S. E., and K. L. Manjunath. 2004.** Asian citrus psyllids (Sternorrhyncha: Psyllidae) and greening disease of citrus: a literature review and assessment of risk in Florida. *Florida Entomologist* 87: 330-353.
- Roistacher, C. N. 1991.** Techniques for biological detection of specific citrus graft transmissible diseases (Greening). Food and Agriculture Organization of the United Nation (FAO); Rome. p. 35-45.
- Toorawa, P. 1998.** La maladie du huanglongbing (greening) des agrumes a L'Île Maurice. Detection de "*Candidatus Liberobacter asiaticum*" et "*Candidatus Liberobacter africanum*" dans les agrumes et les insectes vecteurs. Doctoral Thesis, L'University de Bordeaux. 186 p.
- Wooler A, D. Padgham, and A. Arafat. 1974.** Outbreaks and new records. Saudi Arabia. *Diaphorina citri* on citrus. Food and Agriculture Organization of the United Nation (FAO), Plant Protection Bulletin 22: 93-94.

Received April 14, 2008; accepted May 20, 2008.

