



Results in descriptive set theory on some represented spaces

Mathieu Hoyrup

► **To cite this version:**

| Mathieu Hoyrup. Results in descriptive set theory on some represented spaces. 2018. hal-01657883

HAL Id: hal-01657883

<https://hal.inria.fr/hal-01657883>

Preprint submitted on 7 Dec 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Results in descriptive set theory on some represented spaces

Mathieu Hoyrup

December 7, 2017

Descriptive set theory was originally developed on Polish spaces. It was later extended to ω -continuous domains [Sel04] and recently to quasi-Polish spaces [dB13]. All these spaces are countably-based. Extending descriptive set theory and its effective counterpart to general represented spaces, including non-countably-based spaces has been started in [PdB15].

We study the spaces $\mathcal{O}(\mathbb{N}^{\mathbb{N}})$, $\mathcal{C}(\mathbb{N}^{\mathbb{N}}, 2)$ and the Kleene-Kreisel spaces $\mathbb{N}\langle\alpha\rangle$. We show that there is a Σ_2^0 -subset of $\mathcal{O}(\mathbb{N}^{\mathbb{N}})$ which is not Borel. We show that the open subsets of $\mathbb{N}^{\mathbb{N}^{\mathbb{N}}}$ cannot be continuously indexed by elements of $\mathbb{N}^{\mathbb{N}}$ or even $\mathbb{N}^{\mathbb{N}^{\mathbb{N}}}$, and more generally that the open subsets of $\mathbb{N}\langle\alpha\rangle$ cannot be continuously indexed by elements of $\mathbb{N}\langle\alpha\rangle$. We also derive effective versions of these results.

These results give answers to recent open questions on the classification of spaces in terms of their base-complexity, introduced in [dBSS16]. In order to obtain these results, we develop general techniques which are refinements of Cantor's diagonal argument involving multi-valued fixed-point free functions and that are interesting on their own right.

1 Introduction

On Polish spaces, the definitions of pointclasses are extensional: they describe how the sets in the classes $\Sigma_0^1, \Sigma_1^0, \Sigma_2^0$, etc. are built starting from basic open sets in a countable basis. In particular these definitions imply that all these classes can be parametrized by the Baire space $\mathcal{N} = \mathbb{N}^{\mathbb{N}}$, which is usually expressed by saying that on a Polish space \mathbf{X} , each class $\Gamma(\mathbf{X})$ has a \mathcal{N} -universal set, i.e. a set U in $\Gamma(\mathcal{N} \times \mathbf{X})$ such that $\Gamma(\mathbf{X}) = \{U_f : f \in \mathcal{N}\}$ where $U_f = \{x \in \mathbf{X} : (f, x) \in U\}$.

On a general represented space, the definitions are intensional: they describe what it means for a subset to be $\Sigma_0^1, \Sigma_1^0, \Sigma_2^0$, etc. but they do not give information about what those sets look like and how to obtain them.

Given a represented space, we are interested in understanding more concretely what the elements of the various pointclasses look like, or how they can be described. We provide here essentially impossibility results, stating that

there is no simple way of describing them. A typical result is that for some particular represented spaces \mathbf{X}, \mathbf{Y} , there is no \mathbf{Y} -universal open set, i.e. there is no continuous surjection from \mathbf{Y} to $\mathcal{O}(\mathbf{X})$. The space \mathbf{Y} is usually \mathcal{N} or some Kleene-Kreisel space $\mathbb{N}\langle\alpha\rangle$. We also derive effective versions of such a result: there is no computable surjection from any Σ_k^1 -subset of \mathbb{N} to the computable open subsets of \mathbf{X} , or there is no Δ_k^1 -oracle enumerating those computable open sets.

We develop general tools that enable to prove these impossibility results in a number of interesting cases, in particular for the space of clopen subsets of \mathcal{N} or the Kleene-Kreisel spaces $\mathbb{N}\langle\alpha\rangle$. These tools treat the continuous and computable settings in the same way.

Our tools and techniques are based on Cantor's diagonal argument: the existence of a fixed-point free function is used to disprove the existence of an enumeration. However we usually work on spaces that are lattices of open sets, where every function has a fixed-point but where a *multi-valued* fixed-point free function may exist, and we need to develop techniques enabling us to deal with multi-valued functions. In particular we introduce the notion of a *canonizable space*.

The paper is organized as follows. In Section 2 we give all the necessary notions about represented spaces and their topologies. In Section 3 we prove that the space $\mathcal{O}(\mathcal{N})$ contains a Σ_2^0 -subset that is not Borel. In Section 4 we develop the techniques based on the diagonal argument to prove some impossibility results. In Section 5 we apply these techniques to prove that some particular spaces have no \mathcal{N} -universal open set. In Section 6 we refine our techniques and introduce the notion of a canonizable space to include other parameters spaces than \mathcal{N} and its subspaces. We apply this technique to classify the Kleene-Kreisel spaces and to prove a result about Markov computability.

2 Background

A *represented space* is a pair $\mathbf{X} = (X, \delta_X)$, where $\delta_X : \subseteq \mathcal{N} \rightarrow X$ is a partial onto map. We often identify \mathbf{X} with its carrier set X , for commodity. If $\delta_X(p) = x$ then we say that p is a *name* of x . A point $x \in \mathbf{X}$ is *computable* if it has a computable name. We denote by \mathbf{X}_c the set of computable points of \mathbf{X} . A function $f : \mathbf{X} \rightarrow \mathbf{Y}$ is *continuous* (resp. *computable*) if there exists a continuous (resp. computable) function $F : \text{dom}(\delta_X) \rightarrow \text{dom}(\delta_Y)$ mapping any name of any $x \in \mathbf{X}$ to a name of $f(x)$. We call F a *realizer* of f .

Two represented spaces \mathbf{X}, \mathbf{Y} are computably isomorphic, denoted $\mathbf{X} \cong \mathbf{Y}$, if there is a bijection between that is computable in both directions.

Cartesian product. The product of two represented spaces \mathbf{X}, \mathbf{Y} is denoted by $\mathbf{X} \times \mathbf{Y}$. A name of (x, y) is simply a pair of names of x and y (two elements of \mathcal{N} can be paired into another element of \mathcal{N}). One has $\mathbf{X} \times \mathbf{Y} \cong \mathbf{Y} \times \mathbf{X}$.

Exponentiation. The set of continuous functions from \mathbf{X} to \mathbf{Y} is itself a represented space denoted by $\mathcal{C}(\mathbf{X}, \mathbf{Y})$. A name for a continuous function $f : \mathbf{X} \rightarrow \mathbf{Y}$ is a name for any continuous realizer F of f (such a name gives the information needed to evaluate F , for instance it encodes a list of pairs (u, v) of finite sequences such that for every $p \in \text{dom}(\delta_X)$, (i) if u is a prefix of p and (u, v) appears in the list then $F(p)$ extends v and (ii) for every prefix v of $F(p)$ there exists a prefix u of p such that (u, v) appears in the list).

The evaluation map $\text{eval} : \mathbf{X} \times \mathcal{C}(\mathbf{X}, \mathbf{Y}) \rightarrow \mathbf{Y}$ is computable and $\mathcal{C}(\mathbf{X} \times \mathbf{Y}, \mathbf{Z}) \cong \mathcal{C}(\mathbf{X}, \mathcal{C}(\mathbf{Y}, \mathbf{Z}))$.

Topology. Every represented space $\mathbf{X} = (X, \delta_X)$ has a canonical topology τ_X , which is the final topology of δ_X : a set $U \subseteq X$ is open iff $\delta_X^{-1}(U)$ is the intersection of an open subset of \mathcal{N} with $\text{dom}(\delta_X)$. As $\text{dom}(\delta_X) \subseteq \mathcal{N}$ is countably-based, hence sequential, the final topology of δ_X is sequential. The set of open subsets of \mathbf{X} is itself a represented space obtained as follows. The Sierpiński space $\mathbb{S} = (\{\perp, \top\}, \delta_{\mathbb{S}})$ where $\delta_{\mathbb{S}}(0^{\mathbb{N}}) = \perp$ and $\delta_{\mathbb{S}}(p) = \top$ for $p \neq 0^{\mathbb{N}}$. A subset U of \mathbf{X} is then open exactly when its characteristic function $\mathbf{1}_U : \mathbf{X} \rightarrow \mathbb{S}$ is continuous. The space of open subsets of X is then $\mathcal{O}(\mathbf{X}) := \mathcal{C}(\mathbf{X}, \mathbb{S})$ with its canonical representation. In turn, the topology on $\mathcal{O}(\mathbf{X})$ is the Scott topology, which coincides with the ω -Scott topology (see [dBSS16] for more details).

Subspace. If $\mathbf{X} = (X, \delta_X)$ is a represented space then any subset $A \subseteq X$ has a canonical representation δ_A , which is simply the restriction of δ_X to $\delta_X^{-1}(A)$. Thus it immediately induces notions of continuous and computable functions from A to any other represented space. It is known that sequential topological spaces are not stable under taking subspace topologies. Similarly, the topology on A induced by δ_A is not in general the subspace topology, but is richer. In the same way, the topology on a product of represented spaces is not in general the product topology but is richer.

Admissibility. One has to be careful as there are two a priori distinct notions of continuity for functions from \mathbf{X} to \mathbf{Y} : the functions having a continuous realizer, and the functions that are continuous w.r.t. to the induced topologies. One easily checks that a continuous (i.e. continuously realizable) function is always continuous w.r.t. the induced topologies, however the converse implication fails without further assumption on the representation. The appropriate condition that makes this implication an equivalence is the admissibility of the representation. A representation is admissible if, intuitively, representing a point x is equivalent to being able to know which open sets contain x . The evaluation map $\text{eval} : \mathbf{X} \times \mathcal{O}(\mathbf{X}) \rightarrow \mathbb{S}$, equivalently $\text{eval} : \mathbf{X} \rightarrow \mathcal{O}(\mathcal{O}(\mathbf{X}))$ is computable: the space \mathbf{X} is *admissible* (resp. *computably admissible*) if the inverse of $\text{eval} : \mathbf{X} \rightarrow \mathcal{O}(\mathcal{O}(\mathbf{X}))$ is continuous (resp. computable). In this article, for the general results do not require the representations to be admissible, which is useful when one deals with Markov computability for instance (see Section 6.3).

Multifunctions. Often, a computation taking a name of $x \in \mathbf{X}$ as input and producing a name of some $y \in \mathbf{Y}$ as output does not induce a function from \mathbf{X} to \mathbf{Y} , as y might depend not only on x , but on the name of x given as input. For instance, given a real number $x \in [0, 1]$ it is possible to compute some $y \in [0, 1]$ such that $y \neq x$. However it is not possible to compute the same y for all name of any x , as it would induce a computable, hence continuous function from $[0, 1]$ to $[0, 1]$ without fixed-point, violating the Intermediate Value Theorem. A *multifunction* is a set-valued function $f : \mathbf{X} \rightarrow 2^{\mathbf{Y}}$, denoted by $f : \mathbf{X} \rightrightarrows \mathbf{Y}$. A multifunction $f : \mathbf{X} \rightrightarrows \mathbf{Y}$ is continuous (resp. computable) if there exists a continuous (resp. computable) function $F : \text{dom}(\delta_X) \rightarrow \mathcal{N}$ such that $\delta_Y(F(p)) \in f(\delta_X(p))$ for all $p \in \text{dom}(\delta_X)$.

Descriptive set theory. Notions from descriptive set theory originally developed on Polish spaces have a straightforward extension to any represented space. The question is then whether the results also extend, and how.

Any pointclass $\Gamma(\mathcal{N})$ on \mathcal{N} induces a pointclass on \mathbf{X} as follows: $A \subseteq \mathbf{X}$ belongs to $\Gamma(\mathbf{X})$ if there is a set $B \subseteq \mathcal{N}$ in $\Gamma(\mathcal{N})$ such that $\delta_X^{-1}(A) = B \cap \text{dom}(\delta_X)$.

In a topological space, the class of Borel sets is the smallest class containing the open sets and closed under taking complements and countable unions.

2.1 Retractions

As we have already mentioned, if A is a subset of a represented space \mathbf{X} , then the represented subspace $\mathbf{A} = (A, \delta)$ is not a topological subspace in general. Similarly, if there are continuous (i.e., continuously realizable) functions $f : \mathbf{Y} \rightarrow \mathbf{X}$ and $g : f(\mathbf{Y}) \rightarrow \mathbf{Y}$ such that $g \circ f = \text{id}_{\mathbf{Y}}$, then f is not necessarily a topological embedding because g is not necessarily continuous for the subspace topology on $f(\mathbf{Y})$. However if g has a continuous extension over \mathbf{X} then f is a topological embedding.

Definition 2.1. A *section-retraction* is a pair (s, r) of functions $s : \mathbf{Y} \rightarrow \mathbf{X}$, $r : \mathbf{X} \rightarrow \mathbf{Y}$ such that $r \circ s = \text{id}_{\mathbf{Y}}$. If both s and r are continuous, then \mathbf{Y} is called a *continuous retract* of \mathbf{X} . If both s and r are computable, then \mathbf{Y} is called a *computable retract* of \mathbf{X} .

Intuitively, elements of \mathbf{Y} can be encoded by elements of \mathbf{X} . If $r(x) = y$ then x is a code for y , and every y has a canonical code, given by $s(y)$. It implies in particular that \mathbf{Y} is a continuous (resp. computable) image of \mathbf{X} .

Lemma 2.1. *If \mathbf{Y} and \mathbf{Y}' are continuous (resp. computable) retracts of \mathbf{X} and \mathbf{X}' respectively, then $\mathcal{C}(\mathbf{Y}, \mathbf{Y}')$ is a continuous (resp. computable) retract of $\mathcal{C}(\mathbf{X}, \mathbf{X}')$. In particular $\mathcal{O}(\mathbf{Y})$ is a continuous (resp. computable) retract of $\mathcal{O}(\mathbf{X})$.*

Proof. Let (s, r) and (s', r') be sections-retractions for \mathbf{Y}, \mathbf{X} and \mathbf{Y}', \mathbf{X}' respectively. Let $S : \mathcal{C}(\mathbf{Y}, \mathbf{Y}') \rightarrow \mathcal{C}(\mathbf{X}, \mathbf{X}')$ map f to $s' \circ f \circ r$ and $R : \mathcal{C}(\mathbf{Y}, \mathbf{Y}') \rightarrow$

$\mathcal{C}(\mathbf{X}, \mathbf{X}')$ map g to $r' \circ g \circ s$. The pair (S, R) is easily a section-retraction. The case of $\mathcal{O}(\mathbf{Y}), \mathcal{O}(\mathbf{X})$ is obtained by taking $\mathbf{X}' = \mathbf{Y}' = \mathbb{S}$. \square

If A is a *closed* subset of X then the topology of the represented space \mathbf{A} coincides with the subspace topology. Moreover,

Lemma 2.2. *If A is a closed subset of \mathbf{X} then $\mathcal{O}(\mathbf{A})$ is a continuous retract of $\mathcal{O}(\mathbf{X})$. If A is a Π_1^0 -subset of \mathbf{X} then $\mathcal{O}(\mathbf{A})$ is a computable retract of $\mathcal{O}(\mathbf{X})$.*

Proof. Let $r : \mathcal{O}(\mathbf{X}) \rightarrow \mathcal{O}(\mathbf{A})$ map V to $V \cap A$ and $s : \mathcal{O}(\mathbf{A}) \rightarrow \mathcal{O}(\mathbf{X})$ map U to $U \cup A^c$. \square

We will need the following result in several places.

Lemma 2.3. *\mathcal{N} is a computable retract of $\mathcal{C}(\mathcal{N}, 2)$.*

Proof. Each clopen set $C \in \mathcal{C}(\mathcal{N}, 2)$ encodes a function $f : \mathbb{N} \rightarrow \mathbb{N}$, which is a kind of modulus of continuity of C at certain points, defined as follows. For each n , let $c_n : \mathbb{N} \rightarrow \mathbb{N}$ be the constant function with value n , and $c_{n,k}(i) = n$ for $i \neq k$, $c_{n,k}(k) = n + 1$. If C does not contain c_n , then let $f(n) = 0$. If C contains c_n , then let $f(n) = \max\{k \geq 1 : c_{n,k} \notin C\}$. We then define $r(C) := f$.

Every $f : \mathbb{N} \rightarrow \mathbb{N}$ is encoded in some $C \in \mathcal{C}(\mathcal{N}, 2)$, defined as follows. For each $n \in \mathbb{N}$, if $f(n) = 0$ then let $C \cap [n] = \emptyset$. If $f(n) = k + 1$ then let $C \cap [n] = \{g : g(k + 1) = n\}$. We then define $s(f) := C$.

The functions r and s are computable and one easily checks that $r \circ s(f) = f$ for all f . \square

3 A non-Borel set

On Polish spaces, the Borel hierarchy built from the open sets coincides with the hierarchy lifted from \mathcal{N} by the representation: for instance a set is a countable union of closed sets if and only if its pre-image is a countable union of closed sets.

On quasi-Polish spaces, the same holds if the definition of the Borel hierarchy is slightly amended: for instance Σ_2^0 -sets are not countable unions of closed sets, but countable unions of differences of open sets.

Here we show that on admissibly represented spaces this is no more true in general: in the space $\mathcal{O}(\mathcal{N})$ there exists a Σ_2^0 -subset of $\mathcal{O}(\mathcal{N})$ that is not even Borel.

To build this set, we first work on an intermediate space.

3.1 Product topology

We have seen that two represented spaces \mathbf{X} and \mathbf{Y} naturally induce a third represented space $\mathbf{X} \times \mathbf{Y}$. The topology induced by that representation is not in general the product topology, but its sequentialization.

A simple example is given by $\mathbf{X} = \mathcal{N}$ and $\mathbf{Y} = \mathcal{O}(\mathcal{N})$. The evaluation map $\mathcal{N} \times \mathcal{O}(\mathcal{N}) \rightarrow \mathbb{S}$ is continuous (and computable), however it is not continuous w.r.t. the product topology, because \mathcal{N} is not locally compact (see [EH02] for more details on this topic). In other words the set $\{(f, O) \in \mathcal{N} \times \mathcal{O}(\mathcal{N}) : f \in O\}$ is not open for the product topology (but it is sequentially open, or open for the topology induced by the representation). It is even worse.

Proposition 3.1. *$E = \{(f, O) \in \mathcal{N} \times \mathcal{O}(\mathcal{N}) : f \in O\}$ is not Borel for the product topology.*

Proof. We prove that for every Borel set A , there exists a dense G_δ -set $G \subseteq \mathcal{N}$ such that for every $f \in G$, $(f, \mathcal{N} \setminus \{f\}) \in A \iff (f, \mathcal{N}) \in A$. It implies the result as it is obviously false for the set E . To prove it, we show that the class of sets satisfying this condition contains the open sets in the product topology and is closed under taking complements and countable unions, which implies that this class contains the Borel sets.

First, consider a basic open set $A = [u] \times \mathcal{U}_K$ where u is a finite sequence of natural numbers, K is a compact subset of \mathcal{N} and $\mathcal{U}_K = \{O \in \mathcal{O}(\mathcal{N}) : K \subseteq O\}$. Define $G = [u]^c \cup [u] \setminus K$, which is a dense open set. For $f \in [u]^c$, no (f, O) belongs to A . For $f \in [u] \setminus K$, both $(f, \mathcal{N} \setminus \{f\})$ and (f, \mathcal{N}) belong to A .

If A satisfies the condition with a dense G_δ -set G , then A^c satisfies the condition with the same G . If A_i satisfy the condition with dense G_δ -sets G_i then $\bigcup_i A_i$ satisfies the condition with $G = \bigcap_i G_i$. \square

3.2 The space of open subsets of the Baire space

We can now prove the result.

Theorem 3.1. *There is a Σ_2^0 -subset of $\mathcal{O}(\mathcal{N})$ which is not Borel.*

Proof. We show that $\mathcal{N} \times_{\text{prod}} \mathcal{O}(\mathcal{N})$ is a retract of $\mathcal{O}(\mathcal{N})$ in some way. We build:

- A continuous function $s : \mathcal{N} \times_{\text{prod}} \mathcal{O}(\mathcal{N}) \rightarrow \mathcal{O}(\mathcal{N})$,
- A Σ_2^0 -measurable function $r : \mathcal{O}(\mathcal{N}) \rightarrow \mathcal{N} \times \mathcal{O}(\mathcal{N})$,
- Such that $r \circ s = \text{id}$.

First, these ingredients enable to derive the result. Indeed, let E from Proposition 3.1 and $F := r^{-1}(E) \subseteq \mathcal{O}(\mathcal{N})$. As E is open in $\mathcal{N} \times \mathcal{O}(\mathcal{N})$, F is Σ_2^0 . However F is not Borel, otherwise $E = s^{-1}(F)$ would be Borel in $\mathcal{N} \times_{\text{prod}} \mathcal{O}(\mathcal{N})$.

Let us now build s and r . First observe that $\mathcal{O}(\mathcal{N}) \cong \mathcal{O}(\mathcal{N}) \times \mathcal{O}(\mathcal{N})$ and that the topology on $\mathcal{O}(\mathcal{N}) \times \mathcal{O}(\mathcal{N})$ coincides with the product topology.

Lemma 2.3 provides two computable functions $r_0 : \mathcal{C}(\mathcal{N}, 2) \rightarrow \mathcal{N}$ and $s_0 : \mathcal{N} \rightarrow \mathcal{C}(\mathcal{N}, 2)$ such that $r_0 \circ s_0 = \text{id}$. Observing that \mathbb{S} and 2 are two represented spaces with the same underlying set and that $\mathcal{O}(\mathcal{N}) \cong \mathcal{C}(\mathcal{N}, \mathbb{S})$, those functions can be interpreted as $r_1 : \mathcal{O}(\mathcal{N}) \rightarrow \mathcal{N}$ and $s_1 : \mathcal{N} \rightarrow \mathcal{O}(\mathcal{N})$ with $r_1 \circ s_1 = \text{id}$. However, while s_1 is still computable, r_1 is not computable but is Σ_2^0 -measurable.

Let us simply pair s_1 and r_1 with the identity on $\mathcal{O}(\mathcal{N})$: let $s(f, O') = (s_1(f), O')$ and $r(O, O') = (r_1(O), O')$. \square

In particular, that Σ_2^0 -set is not a countable union of differences of open sets, as it should be on Polish or quasi-Polish spaces. More generally, it is not a countable boolean combination of open sets.

In order to overcome the mismatch between the hierarchy inherited from \mathcal{N} via the representation and the class of Borel sets, one may attempt to change the definition of Borel sets. In [NV97] the Borel sets are redefined as the smallest class containing the open sets and the saturated compact sets. We observe here that this class is too large in the space $\mathcal{O}(\mathcal{N})$. First, if $U \subseteq \mathcal{N}$ is open then the set $\{V \in \mathcal{O}(\mathcal{N}) : U \subseteq V\}$ is compact and saturated in $\mathcal{O}(\mathcal{N})$. From this it is easy to see that the set built above is Borel in this weaker sense. However this notion of Borel sets is too loose, because compact saturated sets do not usually have a Borel pre-image. For instance, the singleton $\{\mathcal{N}\}$ is compact saturated but its pre-image is a $\mathbf{\Pi}_1^1$ -complete set, hence is not Borel.

We leave the following questions for future work:

- What do the Σ_2^0 -subsets of $\mathcal{O}(\mathcal{N})$ look like?
- Is it possible to modify the definition of Borel sets on $\mathcal{O}(\mathcal{N})$ to match exactly the sets that have a Borel pre-image under the representation?

4 Parametrizations

In this section we investigate whether a collection of objects A can be parametrized by another collection B . Formally, A and B are represented spaces and A is parametrized by B if A is a continuous image of B . We will also be interested in the computable version.

Given A , one tries to find the “simplest” space B such that A is a continuous image of B . Simplicity can be measured by restricting our attention to spaces B that are subspaces of \mathcal{N} , and identifying their minimal descriptive complexity. The Borel hierarchy is too fine-grained: A is a continuous image of some Borel subset of \mathcal{N} iff it is a continuous image of a Σ_1^1 -subset of \mathcal{N} iff it is a continuous image of \mathcal{N} . Therefore, the complexity of parametrizing a space is better measured by the hyperprojective hierarchy $\{\Sigma_\alpha^1\}$ where α ranges over the countable ordinals. It happens that the levels of this hierarchy correspond to a hierarchy of spaces, the Kleene-Kreisel spaces $\mathbb{N}\langle\alpha\rangle$, as stated by Theorem 4.2 below: a represented space is a continuous image of some Σ_α^1 -subset of \mathcal{N} if and only if it is a continuous image of $\mathbb{N}\langle\alpha\rangle$.

4.1 Kleene-Kreisel functionals

If α is a countable ordinal then the space $\mathbb{N}\langle\alpha\rangle$ is defined inductively as follows.

Let $\mathbb{N}\langle 0 \rangle = \mathbb{N}$, $\mathbb{N}\langle \alpha + 1 \rangle = \mathcal{C}(\mathbb{N}\langle \alpha \rangle, \mathbb{N})$ and $\mathbb{N}\langle \lambda \rangle = \prod_{i \in \omega} \mathbb{N}\langle \beta(\lambda, i) \rangle$ for a limit ordinal λ and a fixed numbering $\beta(\lambda, i)$ of the ordinals smaller than λ .

We will need the following results.

Functionals of finite type. The following results are proved in [Nor80].

Lemma 4.1. *For each $k \in \mathbb{N}$, $\mathbb{N}\langle k \rangle$ contains a dense computable sequence.*

Theorem 4.1 (Theorem 5.22 in [Nor80]). *Let $k \in \mathbb{N}$ and $A \subseteq \mathcal{N}$. The following conditions are equivalent:*

- $A \in \Sigma_k^1(\mathcal{N})$,
- *There exists a computable predicate $R \subseteq \mathcal{N} \times \mathbb{N}\langle \alpha \rangle \times \mathbb{N}$ such that*

$$x \in A \iff \exists y \in \mathbb{N}\langle \alpha \rangle \forall n, R(x, y, n).$$

In particular, every Σ_k^1 -set is a computable image of a Π_1^0 -subset of $\mathbb{N}\langle k \rangle$, but Theorem 4.1 says more and we will need it later.

Functionals of countable type. In [SS15] an admissible representation δ_α of $\mathbb{N}\langle \alpha \rangle$ is built. Let $D_\alpha = \text{dom}(\delta_\alpha)$.

Theorem 4.2 ([SS15]). *Let α be a countable ordinal. One has $D_{\alpha+1} \in \mathbf{\Pi}_\alpha^1(\mathcal{N})$. A set $A \subseteq \mathcal{N}$ belongs to $\mathbf{\Sigma}_\alpha^1(\mathcal{N})$ iff it is a continuous image of $\mathbb{N}\langle \alpha \rangle$.*

4.2 Diagonal argument.

We now present the basic tool used to prove impossibility results about continuous parametrizations. It is a variation on Cantor's diagonal argument. It might be possible to make it an instance of Lawvere's categorical formulation of the diagonal argument [Law69].

Let \mathbf{Y} be a represented space. If $h : \mathbf{Y} \rightrightarrows \mathbf{Y}$ is a multifunction then a point $y \in Y$ is a **fixed-point** of h if $y \in h(y)$. We say that h is **fixed-point free** if h has no fixed-point. A computable fixed-point free multifunction can be thought as an algorithm taking some y as input (represented by a name) and producing some $y' \neq y$ as output (which may not be the same for all names of y).

Proposition 4.1. *Let \mathbf{Y} be a represented space. If \mathbf{Y} admits a continuous fixed-point free multifunction, then $\mathcal{C}(\mathcal{N}, \mathbf{Y})$ is not a continuous image of \mathcal{N} .*

Proof. Let $h : \mathbf{Y} \rightrightarrows \mathbf{Y}$ have no fixed-point and $H : \text{dom}(\delta_Y) \rightarrow \text{dom}(\delta_Y)$ be a continuous realizer of h . Let $\phi : \mathcal{N} \rightarrow \mathcal{C}(\mathcal{N}, \mathbf{Y})$ be continuous. The function ϕ can equivalently be typed as $\phi : \mathcal{N} \times \mathcal{N} \rightarrow \mathbf{Y}$, let $\Phi : \mathcal{N} \times \mathcal{N} \rightarrow \text{dom}(\delta_Y)$ be a continuous realizer of ϕ . We define $F : \mathcal{N} \rightarrow \mathbf{Y}$ by $F(f) := \delta_Y(H(\Phi(f)(f)))$. The function F is continuous and is not in the range of ϕ because for each f , F differs from $\phi(f)$ at f : $F(f) = \delta_Y(H(\Phi(f)(f))) \neq \delta_Y(\Phi(f)(f)) = \phi(f)(f)$. Therefore ϕ is not onto. \square

In other words, in that case $\mathcal{C}(\mathcal{N}, \mathbf{Y})$ is not a continuous image of any $\mathbf{\Sigma}_1^1$ -subset of \mathcal{N} .

Remark 4.1. Under the same assumption about \mathbf{Y} , the same argument shows that for any set $A \subseteq \mathcal{N}$, $\mathcal{C}(A, \mathbf{Y})$ is not a continuous image of A .

4.3 Complexity of computable enumerations

We will also investigate the computable counterpart of parametrizations.

Definition 4.1. Let \mathbf{X} be a represented space and $\Gamma(\mathbb{N})$ a pointclass. We say that \mathbf{X}_c is Γ -*enumerable* if \mathbf{X}_c is a computable image of some set in $\Gamma(\mathbb{N})$.

The proof of Proposition 4.1 is effective and immediately implies the following effective version.

Proposition 4.2. *If \mathbf{Y} admits a computable fixed-point free multifunction then there is no computable function from \mathcal{N} to $\mathcal{C}(\mathcal{N}, \mathbf{Y})$ containing all the computable elements in its range.*

Proof. In the proof of Proposition 4.1, if H and Φ are computable, then the function F is computable and is not in the range of ϕ . \square

Proposition 4.2 implies that the computable functions from \mathcal{N} to \mathbf{Y} are not Σ_1^0 -enumerable, as the subsets of \mathbb{N} that are computable images of \mathcal{N} are the Σ_1^0 -sets. However one would expect the much stronger result that they are not Σ_1^1 -enumerable. It is possible to obtain such a result under a mild assumption on $\delta_{\mathbf{Y}}$, usually satisfied in practice.

Definition 4.2. We say that $A \subseteq \mathcal{N}$ is *computably separable* if A contains a computable sequence that is dense in A .

If A is computably separable then the set of cylinders intersecting A is c.e., and there is a computable function mapping any cylinder intersecting A to an element of A in that cylinder. This property has the following interesting consequence.

Lemma 4.2. *Assume that $\text{dom}(\delta_{\mathbf{Y}})$ is computably separable. Every computable function from a Π_1^0 -subset of \mathcal{N} to \mathbf{Y} has a total computable extension.*

Proof. Let P be a Π_1^0 -subset of \mathcal{N} , $\phi : P \rightarrow \mathbf{Y}$ be computable and $\Phi : P \rightarrow \mathcal{N}$ be a computable realizer for ϕ . We define a total computable extension ϕ' of ϕ by defining a computable realizer of ϕ' as follows. Given f , run two algorithms in parallel. On the one hand, start computing $\Phi(f)$ and output the result as long as it is compatible with (i.e. has an extension in) $\text{dom}(\delta_{\mathbf{Y}})$. On the other hand, test whether $f \notin P$. If one eventually discovers that $f \notin P$, then extend the current output with some canonical element of $\text{dom}(\delta_{\mathbf{Y}})$. \square

We now get the effective version of Proposition 4.1.

Proposition 4.3. *Assume that $\text{dom}(\delta_{\mathbf{Y}})$ is computably separable and \mathbf{Y} admits a computable fixed-point free multifunction. The computable functions from \mathcal{N} to \mathbf{Y} are not Σ_1^1 -enumerable.*

Proof. Every Σ_1^1 -subset of \mathbb{N} is a computable image of a Π_1^0 -subset of \mathcal{N} , so it is sufficient to prove that the computable elements of $\mathcal{C}(\mathcal{N}, \mathbf{Y})$ is not a computable image of such a set.

Let P be a Π_1^0 -subset of \mathcal{N} and $\delta : P \rightarrow \mathcal{C}(\mathcal{N}, \mathbf{Y})$ be computable. The computable separability of the domain of δ_Y enables one to extend δ outside P . Indeed, δ is equivalently seen as a computable function from $P \times \mathcal{N}$ to \mathbf{Y} , and $P \times \mathcal{N}$ is a Π_1^0 -set, so Lemma 4.2 can be applied. We now apply Proposition 4.2. \square

Example 4.1. There is a strong contrast between the two following cases:

- The left-c.e. functions from \mathcal{N} to $[0, 1]$ can be effectively enumerated (from \mathbb{N}). Indeed, if q_n is a computable enumeration of the rational numbers in $[0, 1]$ then from any computable sequence $(U_n)_{n \in \mathbb{N}}$ of open subsets of \mathcal{N} (those sequences can be effectively enumerated), one can build the left-c.e. function $F(f) = \sup\{q_n : f \in U_n\}$ (with $\sup \emptyset = 0$).
- However the left-c.e. functions from \mathcal{N} to $[0, 1)$ are not Σ_1^1 -enumerable. Indeed, the computable function $h : [0, 1)_{<} \rightarrow [0, 1)_{<}$ mapping x to $(1 + x)/2 > x$ is fixed-point free, and the representation of $[0, 1)_{<}$ has a computably separable domain.

Oracles. So far we have measured the complexity of parametrizing points by measuring the descriptive complexity of the domain of a computable enumeration. A different but related approach is to measure the complexity of an oracle computing an enumeration.

Corollary 4.1. *Assume that $\text{dom}(\delta_Y)$ is computably separable and \mathbf{Y} admits a computable fixed-point free multifunction. There is no Δ_1^1 -oracle enumerating a list of indices of all the computable points of $\mathcal{C}(\mathcal{N}, \mathbf{Y})$.*

Proof. If there is such an oracle then the set of enumerated indices is also a Δ_1^1 -subset of \mathbb{N} , and the function mapping an index in that set to the corresponding point is a computable onto function with a Δ_1^1 domain, contradicting Proposition 4.3. \square

However it may happen that a Δ_1^1 -oracle enumerates all the computable points, producing names rather than indices. Again this is not possible under the same assumptions.

Proposition 4.4. *Assume that $\text{dom}(\delta_Y)$ is computably separable and \mathbf{Y} admits a computable fixed-point free multifunction. There is no Δ_1^1 -oracle enumerating all the computable points of $\mathcal{C}(\mathcal{N}, Y)$.*

Proof. Let $A \in \Delta_1^1(\mathbb{N})$ and $\phi : \mathbb{N} \rightarrow \mathcal{C}(\mathcal{N}, \mathbf{Y})$ be computable with oracle A . The set $\{A\}$ is a Δ_1^1 -subset (hence a Σ_1^1 -subset) of \mathcal{N} so it is a computable image of a Π_1^0 -subset P of \mathcal{N} . As a result, the function $\phi' : P \times \mathbb{N} \rightarrow \mathcal{C}(\mathcal{N}, \mathbf{Y})$ mapping (f, n) to $\phi(n)$ is a computable function defined on a Π_1^0 -subset of $\mathcal{N} \times \mathbb{N} \cong \mathcal{N}$. As in the proof of Proposition 4.3, ϕ' has a total computable extension, which cannot contain all the computable elements in its range. \square

5 Applications to open sets

Let \mathbf{X} be a represented space. $\mathbf{Y} := \mathcal{O}(\mathbf{X})$ is a represented space with a computably separable representation (every cylinder contains a representation of $X \in \mathcal{O}(\mathbf{X})$ consisting of a covering of \mathcal{N} , and that can be computed from the cylinder). In [dBSS16], a represented space \mathbf{X} is called \mathcal{N} -based if there is a continuous function from \mathcal{N} to $\mathcal{O}(\mathbf{X})$ whose image is a basis of the topology on \mathbf{X} . This is equivalent to saying that $\mathcal{O}(\mathbf{X})$ itself is a continuous image of \mathcal{N} .

Lemma 5.1. *A represented space \mathbf{X} is \mathcal{N} -based iff $\mathcal{O}(\mathbf{X})$ is a continuous image of \mathcal{N} .*

Proof. The space \mathbf{X} is hereditarily Lindelöf, i.e. every open cover contains a countable subcover (simply because \mathcal{N} and its subspaces satisfy this property and taking the final topology of the representation preserves this property). Now assume that \mathbf{X} is \mathcal{N} -based and let $\Phi : \mathcal{N} \rightarrow \mathcal{O}(\mathbf{X})$ be continuous, such that $\Phi[\mathcal{N}]$ is a basis. We can easily modify Φ so that the empty set is in its image. A function $f : \mathbb{N} \rightarrow \mathbb{N}$ can equivalently be seen as a sequence of functions $f_i : \mathbb{N} \rightarrow \mathbb{N}$. Let $\Psi(f) = \bigcup_i \Phi(f_i)$. The function $\Psi : \mathcal{N} \rightarrow \mathcal{O}(\mathbf{X})$ is continuous and onto, as every open subset of $\mathcal{O}(\mathbf{X})$ is a countable union of elements of the basis. \square

Observe that $\mathcal{C}(\mathcal{N}, \mathcal{O}(\mathbf{X})) \cong \mathcal{O}(\mathcal{N} \times \mathbf{X})$, hence Propositions 4.1, 4.3, and 4.4 immediately give the following results.

Theorem 5.1. *1. If $\mathcal{O}(\mathbf{X})$ admits a continuous fixed-point free multifunction then $\mathcal{N} \times \mathbf{X}$ is not \mathcal{N} -based.*

2. If $\mathcal{O}(\mathbf{X})$ admits a computable fixed-point free multifunction then the computable open subsets of $\mathcal{N} \times \mathbf{X}$:

- *Are not Σ_1^1 -enumerable,*
- *Are not enumerable relative to any Δ_1^1 oracle.*

Observe that by the Kleene fixed-point theorem, every continuous function from $\mathcal{O}(\mathbf{X})$ to $\mathcal{O}(\mathbf{X})$ has a fixed-point, so we really need to consider multifunctions rather than functions.

Also note that if \mathbf{X} is a countably-based space with its standard representation then $\mathcal{O}(\mathbf{X})$ has no continuous fixed-point free multifunction, because $\mathcal{N} \times \mathbf{X}$ is countably-based hence $\mathcal{O}(\mathcal{N} \times \mathbf{X})$ is a continuous image of \mathcal{N} .

5.1 Product space

Let us consider again the admissibly represented space $\mathbf{X} = \mathcal{N} \times \mathcal{O}(\mathcal{N})$. Its topology is not the product topology and contrary to the product topology it is not \mathcal{N} -based as shown below.

Lemma 5.2. *The space $\mathcal{O}(\mathcal{O}(\mathcal{N}))$ admits a computable fixed-point free multifunction.*

Proof. Given a name of $\mathcal{U} \in \mathcal{O}(\mathcal{O}(\mathcal{N}))$, we build a function $g : \mathbb{N} \rightarrow \mathbb{N}$ and define $\mathcal{V} = \{O \in \mathcal{O}(\mathcal{N}) : g \in O\}$. The function g is defined as follows. Start describing the function g mapping everything to 0. If eventually one finds that $\mathcal{U} \neq \emptyset$ then do as follows. So far, g belongs to some cylinder $[0^n]$. As $\mathcal{U} \neq \emptyset$, $\mathcal{N} \in \mathcal{U}$ so any enumeration of \mathcal{N} is accepted by the name of \mathcal{U} . Enumerate \mathcal{N} in such a way that $[0^n]$ is not covered in finite time. When this enumeration is accepted by \mathcal{U} , let g be an element of $[0^n]$ that is not yet covered by the enumeration of \mathcal{N} .

If $\mathcal{U} = \emptyset$ then $g = 0$ and \mathcal{V} is non-empty so $\mathcal{V} \neq \mathcal{U}$. If $\mathcal{U} \neq \emptyset$ then $\mathcal{N} \setminus \{g\}$ belongs to \mathcal{U} but not to \mathcal{V} , so $\mathcal{V} \neq \mathcal{U}$. \square

One can apply Theorem 5.1.

- $\mathcal{N} \times \mathcal{O}(\mathcal{N})$ is not \mathcal{N} -based,
- The computable open subsets of $\mathcal{N} \times \mathcal{O}(\mathcal{N})$ are neither Σ_1^1 -enumerable nor enumerable relative to a Δ_1^1 oracle.

However,

Proposition 5.1. $\mathcal{N} \times \mathcal{O}(\mathcal{N})$ is $\mathbb{N}\langle 2 \rangle$ -based.

Proof. $\mathcal{O}(\mathcal{O}(\mathcal{N}))$ admits a total representation $\delta : \mathcal{N} \rightarrow \mathcal{O}(\mathcal{O}(\mathcal{N}))$, equivalent to the canonical representation. Therefore the function $\mathcal{C}(\mathcal{N}, \mathcal{N}) \rightarrow \mathcal{C}(\mathcal{N}, \mathcal{O}(\mathcal{O}(\mathcal{N})))$ mapping f to $\delta \circ f$ is continuous and onto. Observe that $\mathcal{C}(\mathcal{N}, \mathcal{N}) \cong \mathbb{N}\langle 2 \rangle$ and $\mathcal{C}(\mathcal{N}, \mathcal{O}(\mathcal{O}(\mathcal{N}))) \cong \mathcal{O}(\mathcal{N} \times \mathcal{O}(\mathcal{N}))$. \square

Moreover $\mathcal{O}(\mathcal{N} \times \mathcal{O}(\mathcal{N}))$ is a computable image of a Π_1^1 -subset of \mathcal{N} , so the computable open subsets of $\mathcal{N} \times \mathcal{O}(\mathcal{N})$ are Π_1^1 -enumerable and enumerable relative to a Δ_2^1 -oracle.

5.2 Space of clopens

Let $\mathbf{X} = \mathcal{C}(\mathcal{N}, 2)$ be the space of clopen subsets of the Baire space.

Theorem 5.2. *The space $\mathcal{C}(\mathcal{N}, 2)$ is not \mathcal{N} -based. Its computable open sets are not Σ_1^1 -enumerable and are not enumerable relative to any Δ_1^1 oracle.*

To prove the theorem we need the following lemma.

Lemma 5.3. *The space $\mathcal{O}(\mathcal{C}(\mathcal{N}, 2))$ admits a computable fixed-point free multifunction.*

The proof is almost identical to the proof of Lemma 5.2.

Proof. Given a name of $\mathcal{U} \in \mathcal{O}(\mathcal{C}(\mathcal{N}, 2))$, start computing a function $g \in \mathcal{N}$ as explained below, and output $\mathcal{V} = \{C \in \mathcal{C}(\mathcal{N}, 2) : g \in C\} \in \mathcal{O}(\mathcal{C}(\mathcal{N}, 2))$.

Start defining $g = 0$ on the first inputs. At the same time, wait to see whether \mathcal{U} is non-empty. If eventually $\mathcal{U} \neq \emptyset$ then at that point g is defined up to input n . One can find some $C \in \mathcal{U}$ and $[0^n] \not\subseteq C$. Indeed, start from

some $A \in \mathcal{U}$ and generate a name of A which does not cover $[0^n]$ in finite time. This name is eventually accepted by the name of \mathcal{U} . Once C is found, let $g \in [0^n] \setminus C$.

We now check that $\mathcal{V} \neq \mathcal{U}$. If \mathcal{U} is empty then $\mathcal{V} \neq \mathcal{U}$ as \mathcal{V} is non-empty (and in that case, $g = 0$). If \mathcal{U} is non-empty then one finds $C \in \mathcal{U}$ and $g \notin C$, so $C \notin \mathcal{V}$ hence $\mathcal{V} \neq \mathcal{U}$. \square

Proof of Theorem 5.2. Let $\mathbf{X} = \mathcal{C}(\mathcal{N}, 2)$. Lemma 5.3 and Theorem 5.1 imply the result for the space $\mathcal{N} \times \mathbf{X}$.

We know from Lemma 2.3 that \mathcal{N} is a computable retract of \mathbf{X} , so $\mathcal{N} \times \mathbf{X}$ is a computable retract of $\mathbf{X} \times \mathbf{X} \cong \mathbf{X}$. As a result Lemma 2.1 implies that $\mathcal{O}(\mathcal{N} \times \mathbf{X})$ is a computable retract, hence a computable image of $\mathcal{O}(\mathbf{X})$, so the negative results about $\mathcal{O}(\mathcal{N} \times \mathbf{X})$ immediately apply to $\mathcal{O}(\mathbf{X})$. \square

Observe that Theorem 5.2 immediately applies to the space $\mathbb{N}\langle 2 \rangle = \mathcal{C}(\mathcal{N}, \mathbb{N})$, as $\mathcal{C}(\mathcal{N}, 2)$ is a computable retract of $\mathbb{N}\langle 2 \rangle$.

We will now prove the stronger result that $\mathcal{C}(\mathcal{N}, 2)$ and $\mathbb{N}\langle 2 \rangle$ are not even $\mathbb{N}\langle 2 \rangle$ -based (Theorem 6.3 below), and more generally $\mathbb{N}\langle k \rangle$ is not $\mathbb{N}\langle k \rangle$ -based for $k \geq 2$. For this, we need to refine our techniques, as we explain now.

6 Canonicity of names

The proof of Proposition 4.1 works because \mathcal{N} has a particular property: every element $f \in \mathcal{N}$ has a canonical name (which is f itself). This property has the consequence that every continuous (resp. computable) multifunction $F : \mathcal{N} \rightrightarrows \mathbf{X}$ has a continuous (resp. computable) choice function $F' : \mathcal{N} \rightarrow \mathbf{X}$, such that $F'(f) \in F(f)$ for all $f \in \mathcal{N}$. So the multifunction built using the diagonal argument and the fixed-point free multifunction is actually a single-valued function.

The spaces having this canonicity property are exactly the subspaces of \mathcal{N} . The proof of Proposition 4.1 does not carry over to spaces without this canonicity property, like \mathbb{R} or $\mathbb{N}\langle 2 \rangle$ for instance. However we provide a technique to overcome this problem.

Definition 6.1. A represented space \mathbf{X} is *canonizable* if there exists a subset $P \subseteq \mathbf{X}$ such that:

- P is closed,
- \mathbf{X} is a continuous image of P ,
- P is homeomorphic to some subspace of \mathcal{N} .

The first property implies that P as a represented subspace of \mathbf{X} has the subspace topology. The third property exactly says that P has the canonicity property, i.e. that there is a continuous function mapping every $x \in P$ to a name of x .

Example 6.1. \mathbb{R} is canonizable, because it is a continuous image of $\mathbb{N} \times 2^{\mathbb{N}}$ which embeds as a closed set in \mathbb{R} .

When a space is canonizable, the diagonal argument can be applied.

Theorem 6.1. *Let \mathbf{X} be a canonizable space. If $\mathcal{O}(\mathbf{Y})$ admits a continuous fixed-point free multifunction then $\mathbf{X} \times \mathbf{Y}$ is not \mathbf{X} -based.*

Proof. Let P make \mathbf{X} canonizable, and let $Q \subseteq \mathcal{N}$ be homeomorphic to P . Proposition 4.1 implies that $\mathcal{O}(Q \times \mathbf{Y}) \cong \mathcal{C}(Q, \mathcal{O}(\mathbf{Y}))$ is not a continuous image of Q , hence $\mathcal{O}(P \times \mathbf{Y})$ is not a continuous image of P . As P is a closed subset of \mathbf{X} , $\mathcal{O}(P \times \mathbf{Y})$ is a continuous retract, hence a continuous image of $\mathcal{O}(\mathbf{X} \times \mathbf{Y})$. As \mathbf{X} is a continuous image of P , one concludes by transitivity that $\mathcal{O}(\mathbf{X} \times \mathbf{Y})$ is not a continuous image of \mathbf{X} . \square

This result has a computable version. We say that \mathbf{X} is **computably canonizable** if the set P is a Π_1^0 -subset of \mathbf{X} , if \mathbf{X} is a computable image of P and P is computably homeomorphic to some subspace of \mathcal{N} . In that case, the same proof shows that there is no computable function from \mathbf{X} to $\mathcal{O}(\mathbf{X} \times \mathbf{Y})$ containing all the computable open sets in its range.

6.1 Kleene-Kreisel functionals as parameters

The notion of a canonizable space enables us to extend Theorem 5.1 from $\mathcal{N} = \mathbb{N}\langle 1 \rangle$ to the whole hierachies functionals of finite and countable type.

Theorem 6.2. *1. If $\mathcal{O}(\mathbf{X})$ admits a continuous fixed-point free multifunction then for each countable ordinal α , $\mathbb{N}\langle \alpha \rangle \times \mathbf{X}$ is not $\mathbb{N}\langle \alpha \rangle$ -based.*

2. If $\mathcal{O}(\mathbf{X})$ admits a computable fixed-point free multifunction then for each $k \in \mathbb{N}$ the computable open subsets of $\mathbb{N}\langle k \rangle \times \mathbf{X}$:

- *Are not Σ_k^1 -enumerable,*
- *Are not enumerable relative to any Δ_k^1 oracle.*

The proof essentially follows the same argument as the proof of Theorem 5.1, but we first need to show that the spaces $\mathbb{N}\langle \alpha \rangle$ are canonizable.

Lemma 6.1. *For every countable ordinal α , $\mathbb{N}\langle \alpha \rangle$ is canonizable.*

Proof. The proof is inspired by a construction appearing in the proof of Theorem 7.2 in [SS15].

We first prove the result for all successor ordinals. $D_{\alpha+1}$ is a $\mathbf{\Pi}_\alpha^1$ -set, so there exists a continuous surjective function $f : \mathbb{N}\langle \alpha \rangle \rightarrow \mathcal{N} \setminus D_{\alpha+1}$. We define $\psi : D_{\alpha+1} \rightarrow \mathbb{N}\langle \alpha + 1 \rangle$ as follows. For $x \in D_{\alpha+1}$ and $y \in \mathbb{N}\langle \alpha \rangle$, let $\psi(x)(y) = \min\{n \in \mathbb{N} : x_n \neq f(y)_n\}$. ψ is continuous, let $\Psi : D_{\alpha+1} \rightarrow \mathcal{N}$ be a continuous realizer of ψ . Let $P \subseteq \mathcal{N} \times \mathbb{N}\langle \alpha + 1 \rangle$ be the graph of ψ . P is homeomorphic to the graph of Ψ which is a subset of $\mathcal{N} \times \mathcal{N} \cong \mathcal{N}$. P is a closed subset of $\mathcal{N} \times \mathbb{N}\langle \alpha + 1 \rangle$. Indeed, a pair (x, F) belongs to P iff for all $y \in \mathbb{N}\langle \alpha \rangle$,

if $n = F(y)$ then $x_n \neq f(y)_n$ and $x_m = f(y)_m$ for all $m < n$, which can be checked for y in some countable dense subset of $\mathbb{N}\langle\alpha\rangle$. If this is not true then one can eventually see it. Finally, $\mathbb{N}\langle\alpha + 1\rangle$ is a continuous image of $D_{\alpha+1}$ which is the first projection of P .

For limit ordinals, we prove the result by induction. Let λ be a limit ordinal. For each $\beta < \lambda$, let P_β witness that $\mathbb{N}\langle\beta\rangle$ is canonizable. By definition, $\mathbb{N}\langle\lambda\rangle = \prod_{i \in \omega} \mathbb{N}\langle\beta(\lambda, i)\rangle$ for some numbering $\beta(\lambda, i)$ of the ordinals smaller than λ , and one can easily check that $\prod_{i \in \omega} P_{\beta(\lambda, i)}$ makes $\mathbb{N}\langle\lambda\rangle$ canonizable. \square

For the finite levels of the hierarchy we can prove an effective version.

Lemma 6.2. *For $k \geq 1$, $\mathbb{N}\langle k \rangle$ is computably canonizable.*

Proof. From Theorem 4.1, as $D_k \in \Pi_{k-1}^1$ there exists a computable predicate $R \subseteq \mathcal{N} \times \mathbb{N}\langle k-1 \rangle \times \mathbb{N}$ such that $x \in D_k \iff \forall y \in \mathbb{N}\langle k-1 \rangle, \exists n, R(x, y, n)$. We define a computable function $\psi : D_k \rightarrow \mathbb{N}\langle k \rangle$ by $\psi(x)(y) = \min\{n : R(x, y, n)\}$ for $x \in D_k, y \in \mathbb{N}\langle k-1 \rangle$. Let P be the graph of ψ . First, P is a Π_1^0 -subset of $\mathcal{N} \times \mathbb{N}\langle k \rangle$. Indeed, (x, H) belongs to the graph if and only if for all $y \in \mathbb{N}\langle k-1 \rangle$, if $n = H(y)$ then $R(x, y, n)$ and $\neg R(x, y, i)$ for all $i < n$, but this can be checked for all y in a computable dense sequence in $\mathbb{N}\langle k-1 \rangle$ (which exists by Lemma 4.1).

As ψ is computable, its graph is computably isomorphic to the graph of a computable realizer of ψ , which is a subset of $\mathcal{N} \times \mathcal{N} \cong \mathcal{N}$. Finally, $\mathbb{N}\langle k \rangle$ is a computable image of D_k which is the first projection of P . \square

It may be possible to extend the result to the constructive ordinals, however we do not address this question in this article.

Proof of Theorem 6.2. In the first case, as $\mathbb{N}\langle\alpha\rangle$ is canonizable we can apply Theorem 6.1.

In the second case, as $\mathbb{N}\langle k \rangle$ is computably canonizable there is no computable function from $\mathbb{N}\langle k \rangle$ to $\mathcal{O}(\mathbb{N}\langle k \rangle \times \mathbf{Y})$ containing all the computable open sets in its range, therefore there is no such computable function defined on some Π_1^0 -subset of $\mathbb{N}\langle\alpha\rangle$. We can conclude as every set in $\Sigma_k^1(\mathbb{N})$ is a computable image of some set in $\Pi_1^0(\mathbb{N}\langle\alpha\rangle)$, and for every set $A \in \Delta_2^1(\mathbb{N})$, $\{A\}$ is a computable image of some set in $\Pi_1^0(\mathbb{N}\langle\alpha\rangle)$. \square

6.2 Applications

We now apply Theorem 6.2 to particular spaces.

Back to the space of clopens. We saw in Theorem 5.2 that $\mathcal{C}(\mathcal{N}, 2)$ is not \mathcal{N} -based, i.e. that $\mathcal{O}(\mathcal{C}(\mathcal{N}, 2))$ is not a continuous image of \mathcal{N} . We can now apply Theorem 6.2 to improve that result.

Theorem 6.3. *The space $\mathcal{C}(\mathcal{N}, 2)$ is not $\mathbb{N}\langle 2 \rangle$ -based.*

The computable open subsets of $\mathcal{C}(\mathcal{N}, 2)$ are not Σ_2^1 -enumerable and not enumerable relative to any Δ_2^1 oracle.

The Kleene-Kreisel functionals.

Theorem 6.4. *For every countable ordinal $\alpha \geq 2$, $\mathbb{N}\langle\alpha\rangle$ is not $\mathbb{N}\langle\alpha\rangle$ -based.*

For every $k \geq 2$, the computable open subsets of $\mathbb{N}\langle k\rangle$ are not Σ_k^1 -enumerable and not enumerable relative to any Δ_k^1 oracle.

We just have to prove that $\mathcal{O}(\mathbb{N}\langle\alpha\rangle)$ has continuous fixed-point free multifunction and $\mathcal{O}(\mathbb{N}\langle k\rangle)$ has a computable one. They can be easily derived from the computable fixed-point free multifunction on $\mathcal{O}(\mathcal{C}(\mathcal{N}, 2))$ (Lemma 5.3) thanks to the following observation.

Lemma 6.3. *Let \mathbf{X} be a continuous (resp. computable) retract of \mathbf{Y} . If \mathbf{X} admits a continuous (resp. computable) fixed-point free multifunction then so does \mathbf{Y} .*

Proof. Let $r : \mathbf{Y} \rightarrow \mathbf{X}$ and $s : \mathbf{X} \rightarrow \mathbf{Y}$ be continuous such that $r \circ s = \text{id}$, and let R, S be continuous realizers. Let $h : \mathbf{X} \rightrightarrows \mathbf{X}$ be fixed-point free and H be a continuous realizer. The multifunction $k = s \circ h \circ r$ (defined by $k(y) = \{s(x) : x \in h(r(y))\}$) is continuous as it is realized by the continuous function $S \circ H \circ R$. Moreover it is fixed-point free, as if $y \in k(y)$ then $y = s(x)$ for some $x \in h(r(y))$, so $r(y)$ is a fixed-point for h , as $r(y) = r \circ s(x) = s \in h(r(y))$, contradicting the assumption about h . Everything works the same in the computable case. \square

Theorem 6.4 follows as $\mathcal{C}(\mathcal{N}, 2)$ is a computable retract of $\mathbb{N}\langle 2\rangle$ which is a computable retract of each $\mathbb{N}\langle k\rangle$ (Lemma 7.4 in [SS15]), $k \geq 2$, and a continuous retract of each $\mathbb{N}\langle\alpha\rangle$, $\alpha \geq 2$.

Implications on the hyperprojective base-hierarchy. In [dBSS16] the hierarchy $\text{Based}(\mathbb{N}\langle\alpha\rangle)$ is introduced: a space is in $\text{Based}(\mathbb{N}\langle\alpha\rangle)$ if it is $\mathbb{N}\langle\alpha\rangle$ -based. That hierarchy is related to another hierarchy called $\text{QCB}_0(\mathbf{\Pi}_\alpha^1)$. The following results are proved in [dBSS16]:

- Proposition 2.9: $\mathbb{N}\langle\alpha + 1\rangle \in \text{QCB}_0(\mathbf{\Pi}_\alpha^1)$,
- Proposition 6.11: $\text{QCB}_0(\mathbf{\Pi}_\alpha^1) \subseteq \text{Based}(\mathbb{N}\langle\alpha + 2\rangle)$,
- Corollary 6.12: The inclusion $\text{Based}(\mathbb{N}\langle\alpha\rangle) \subset \text{Based}(\mathbb{N}\langle\alpha + 3\rangle)$ is proper.

It is asked:

- Problem 6.2: for which α does the inclusions $\text{QCB}_0(\mathbf{\Pi}_\alpha^1) \subseteq \text{Based}(\mathbb{N}\langle\alpha + 1\rangle)$ and $\text{QCB}_0(\mathbf{\Pi}_\alpha^1) \subseteq \text{Based}(\mathbb{N}\langle\alpha\rangle)$ hold?
- Is the inclusion $\text{Based}(\mathbb{N}\langle\alpha\rangle) \subset \text{Based}(\mathbb{N}\langle\alpha + 1\rangle)$ proper?

Theorem 6.4 gives answers to these questions:

Corollary 6.1. 1. *For all countable ordinals $\alpha \geq 1$, $\text{QCB}_0(\mathbf{\Pi}_\alpha^1) \not\subseteq \text{Based}(\mathbb{N}\langle\alpha + 1\rangle)$.*

2. *For all countable ordinals α , the inclusion $\text{Based}(\mathbb{N}\langle\alpha\rangle) \subset \text{Based}(\mathbb{N}\langle\alpha + 1\rangle)$ is proper (witnessed by $\mathbb{N}\langle\alpha\rangle$),*

Proof. 1. The first assertion is witnessed by $\mathbb{N}\langle\alpha + 1\rangle$ (Theorem 6.4).

2. The second assertion is witnessed by $\mathbb{N}\langle\alpha\rangle$ for $\alpha \geq 2$ (Theorem 6.4), by $\mathcal{N} \times \mathcal{O}(\mathcal{N})$ for $\alpha = 1$ (see Section 5.1) and by $\mathcal{O}(\mathcal{N})$ for $\alpha = 0$. \square

6.3 Markov computability

Let \mathbf{X}_M be the set of computable elements of \mathbf{X} with the Markov representation mapping an index of a computable name of $x \in \mathbf{X}$ to x (an index $i \in \mathbb{N}$ can be identified with the constant function with value i , so that the domain of the representation is a subset of \mathcal{N}). Note that the identity from \mathbf{X}_M to \mathbf{X}_c is computable, but its inverse is not computable in general.

While the spaces \mathcal{N} and \mathcal{N}_c have the canonicity property, \mathcal{N}_M does not: it is not possible to associate to each computable function a canonical index that could be uniformly computed from any other index.

However,

Lemma 6.4. \mathcal{N}_M is computably canonizable.

Proof. Let $(\varphi_i)_{i \in \mathbb{N}}$ be a canonical enumeration of the partial computable functions from \mathbb{N} to \mathbb{N} . Let $\text{Tot} = \{i : \varphi_i \text{ is total}\}$. If $i \in \text{Tot}$, then let $t_i : \mathbb{N} \rightarrow \mathbb{N}$ map n to the halting time of $\varphi_i(n)$.

Let $P = \{(i, t) : \varphi_i \text{ is total and } t = t_i\}$, where (i, t) is understood as the function f mapping 0 to i and $n + 1$ to $t(n)$. P is a Π_1^0 -subset of \mathcal{N}_M (and even of \mathcal{N}_c). Indeed, given f , let $i = f(0)$: one has $f \notin P \iff \exists n, \varphi_i(n)$ does not halt in exactly $f(n)$ steps, which can be eventually discovered.

On P , the standard representation and the Markov representation are computably equivalent: given $f \in P$, one can compute an index of f , as one can compute an index of t_i where $i := f(0)$, so $P \subseteq \mathbf{X}_M$ is computably isomorphic to $P \subseteq \mathcal{N}$.

Finally, \mathcal{N}_M is a computable image of the first projection of P . \square

Lemma 6.5. $\mathcal{O}(\mathcal{N}_M)$ admits a computable fixed-point free multifunction.

Proof. One can think of a name of an element \mathcal{W} of $\mathcal{O}(\mathcal{N}_M)$ as an enumeration of a set $W \subseteq \mathbb{N}$ containing the indices of the computable functions in \mathcal{W} (and possibly indices of partial functions). Given an enumeration of a set of indices W , we enumerate another set W' coding some $\mathcal{W}' \in \mathcal{O}(\mathcal{N}_M)$ such that $\mathcal{W}' \neq \mathcal{W}$.

Let f_∞ be the null function, and $f_k(n) = 0$ if $n \neq k$, $f_k(k) = 1$.

Given an enumeration of W , we enumerate W' as follows. Start enumerating all the natural numbers. In parallel, test whether $f_\infty \in \mathcal{W}$. If it eventually happens, let n be larger than all the numbers enumerated so far in W' . Look for $n+1$ values of k such that $f_k \in \mathcal{W}$, and let K be larger than all of them. Now enumerate in W' all the numbers j such that φ_j coincides with some φ_l , $l < n$, on inputs $0, 1, \dots, K$.

Now we check that the construction fulfills the conditions.

- If $f_\infty \notin \mathcal{W}$ then $W' = \mathbb{N}$ and $\mathcal{W}' = \mathcal{N}_c$, so $f_\infty \in \mathcal{W}' \setminus \mathcal{W}$.

- If $f_\infty \in \mathcal{W}$ then there exist infinitely many k 's such that $f_k \in \mathcal{W}$, so the algorithm will eventually find K , and $\mathcal{W}' = \bigcup_{l < n} [\varphi_l \upharpoonright_K]$ (with the convention that $[\varphi_l \upharpoonright_K] = \emptyset$ if φ_l is not defined up to K). As there are at least $n + 2$ values of $k < K$ such that $f_k \in \mathcal{W}$, and all these functions differ on some input $< K$, one of them does not belong to \mathcal{W}' which a union of n cylinders of length K . As result, $f_k \in \mathcal{W} \setminus \mathcal{W}'$. \square

The computable elements of $\mathcal{O}(\mathcal{N}_M)$ are also called the Markov semidecidable properties. They correspond to the c.e. subsets of \mathbb{N} that are extensional w.r.t. the indices of total computable functions (for each total function, an extensional set contains all its indices or none of them). With the tools we have developed, we can resolve a question left open in [Hoy16].

Corollary 6.2. *The computable open subsets of \mathcal{N}_M are not Σ_3^0 -enumerable.*

Proof. One has $\mathcal{O}(\mathcal{N}_M \times \mathcal{N}_M) \cong \mathcal{O}(\mathcal{N}_M)$ so $\mathcal{O}(\mathcal{N}_M)$ is not a computable image of \mathcal{N}_M . The computable images of \mathcal{N}_M are exactly the Σ_3^0 -sets. \square

Finally, the computable open subsets of \mathcal{N}_M are Π_3^0 -enumerable, simply because the domain of the representation of $\mathcal{O}(\mathcal{N}_M)$ is Π_3^0 (said differently, for a c.e. set, being extensional is a Π_3^0 -property).

Acknowledgments

The author would like to thank Arno Pauly for suggesting the investigation of the Σ_2^0 -subsets of $\mathcal{O}(\mathcal{N})$ and the Σ_1^0 -subsets of $\mathcal{C}(\mathcal{N}, 2)$, as well as Matthias Schröder for discussions on this topic.

References

- [dB13] Matthew de Brecht. Quasi-polish spaces. *Ann. Pure Appl. Logic*, 164(3):356–381, 2013.
- [dBSS16] Matthew de Brecht, Matthias Schröder, and Victor Selivanov. Base-complexity classifications of qcb₀-spaces. *Computability*, 5(1):75–102, 2016.
- [EH02] Martin Escardó and Reinhold Heckmann. Topologies on spaces of continuous functions. *Topology Proceedings*, 26(2):545–564, 2001-2002.
- [Hoy16] Mathieu Hoyrup. The decidable properties of subrecursive functions. In *ICALP 2016, July 11-15, 2016, Rome, Italy*, volume 55 of *LIPICs*, pages 108:1–108:13. Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2016.
- [Law69] F. William Lawvere. *Diagonal arguments and cartesian closed categories*, pages 134–145. Springer Berlin Heidelberg, Berlin, Heidelberg, 1969.

- [Nor80] D. Normann. *Recursion on the Countable Functionals*. Lecture Notes in Mathematics. Springer Berlin Heidelberg, 1980.
- [NV97] Tommy Norberg and Wim Vervaat. Capacities on non-Hausdorff spaces. *Probability and Lattices*, 110:133–150, 1997.
- [PdB15] Arno Pauly and Matthew de Brecht. Descriptive set theory in the category of represented spaces. In *30th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2015, Kyoto, Japan, July 6-10, 2015*, pages 438–449. IEEE Computer Society, 2015.
- [Sel04] V. L. Selivanov. Difference hierarchy in φ -spaces. *Algebra and Logic*, 43(4):238–248, Jul 2004.
- [SS15] Matthias Schröder and Victor Selivanov. Some hierarchies of qcb_0 -spaces. *Mathematical Structures in Computer Science*, 25(8):1799–1823, 2015.