

1 Effect of CO₂ enrichment on phytoplankton photosynthesis in the
2 North Atlantic sub-tropical gyre

3
4 Gavin Tilstone*¹, Barbora Šedivá², Glen Tarran¹, Radek Kaňa^{2,3}, Ondřej Prášil^{2,3}.

5
6 ¹Plymouth Marine Laboratory

7 Prospect Place, The Hoe, Plymouth, PL1 3DH, UK.

8 ²Institute of Microbiology, Academy of Sciences of the Czech Republic, Opatovický mlyn,

9 CZ-379 81 Třeboň, the Czech Republic.

10 ³Faculty of Sciences, University of South Bohemia, Branišovská 31, 370 05 České

11 Budějovice, Czech Republic,

12
13 *Corresponding author: ghti@pml.ac.uk

14
15 Running head: Photosynthesis under high CO₂ in the Atlantic Ocean.

16
17 KEY WORDS: CO₂ enrichment, picoeukaryotes, dinoflagellates, phytoplankton,
18 photosynthesis, emission spectroscopy, sub-tropical Atlantic Ocean.

22 Abstract

23 The effects of changes in CO₂ concentration in seawater on phytoplankton community
24 structure and photosynthesis were studied in the North Atlantic sub-tropical gyre. Three
25 shipboard incubations were conducted for 48 h at ~760 ppm CO₂ and control (360 ppm CO₂)
26 from 49°N to 7°N during October and November 2010. Elevated CO₂ caused a decrease in pH
27 to ~7.94 compared to ~8.27 in the control. During one experiment, the biomass of nano- and
28 picoeukaryotes increased under CO₂ enrichment, but primary production decreased relative to
29 the control. In two of the experiments the biomass was dominated by dinoflagellates, and
30 there was a significant increase in the maximum photosynthetic rate (P_m^B) and light-limited
31 slope of photosynthesis (α^B) at CO₂ concentrations of 750 ppm relative to the controls. 77 K
32 emission spectroscopy showed that the higher photosynthetic rates measured under CO₂
33 enrichment increased the connection of reversible photosystem antennae, which resulted in an
34 increase in light harvesting efficiency and carbon fixation.

35
36

37 **1. Introduction**

38 There is global concern over increases in the CO₂ concentration of seawater and the effects of
39 ocean acidification on the functionality and productivity of marine ecosystems (Riebesell,
40 2008). Increases in atmospheric CO₂ from 280 to 370 ppm since the industrial revolution have
41 decreased surface ocean pH by 0.12 units (Riebesell, 2004). The “business as usual” scenario
42 predicts that CO₂ will rise to 700 ppm over the next 100 years (Houghton, 2001), which will
43 decrease seawater pH by a further 0.3 to 0.6 units (Riebesell, Schulz, Bellerby, Botros,
44 Fritsche et al., 2007), bicarbonate ion (HCO₃⁻) concentration by 50% (Riebesell, 2004) and
45 raise the sea surface temperature (SST) by 2 to 6°C (Alley, Berntsen, Bindoff, Chen & others,
46 2007; Bopp, Monfray, Aumont, Dufresne, Le Treut et al., 2001). The projected pH shift from
47 8.2 to 7.7 covers the entire range of variation in pH currently observed in open ocean surface
48 waters. The increase in SST is predicted to increase stratification and light availability of the
49 surface ocean due to shoaling of the upper mixed layer (Rost, Zondervan & Wolf-Gladrow,
50 2008; Sarmiento, Slater, Barber, Bopp, Doney et al., 2004), which could directly impact the
51 physiology of phytoplankton. These changes in pH, SST and light regime are predicted to
52 enhance primary production (PP) (Hein & SandJensen, 1997; Riebesell, Wolfgladrow &
53 Smetacek, 1993), especially in the North Atlantic (Bopp et al., 2001; Doney, Lindsay, Fung &
54 John, 2006; Feng, Hare, Leblanc, Rose, Zhang et al., 2009), though this will depend on the
55 associated dominant species and nutrient regime (Rost et al., 2008).

56 Information on the effects of CO₂ enrichment on phytoplankton photosynthesis in a
57 range of organisms and different trophic systems is essential to understand potential shifts in
58 the carbon cycle due to ocean acidification (Riebesell, 2004). Within the usual seawater pH
59 range (8.0-8.3), 90% of the total DIC is HCO₃⁻, and CO₂ is less than 1 % when the system is
60 in equilibrium with atmospheric CO₂ (Skirrow, 1975). The total inorganic carbon
61 concentration in seawater is 2 μmol L⁻¹ and the CO₂ content is 10 μmol L⁻¹. This is not

62 sufficient to saturate carbon fixation by the algal photosynthetic enzyme ribulose
63 biphosphate carboxylase oxygenase (RuBisCO), which has half-saturation constants of 20–
64 40 $\mu\text{mol L}^{-1}$ CO_2 for eukaryotic microalgae and up to 750 $\mu\text{mol L}^{-1}$ CO_2 for marine
65 cyanobacteria (Badger, Andrews, Whitney, Ludwig, Yellowlees et al., 1998; Hopkinson,
66 Dupont, Allen & Morel, 2011; Raven, 2011a; Raven & Johnston, 1991). Different
67 phytoplankton groups or species have therefore evolved a preference for different forms of
68 DIC, with some taking up CO_2 directly, whereas others draw on the pool of HCO_3^- present
69 (Elzenga, Prins & Stefels, 2000) and or mechanisms to concentrate CO_2 or HCO_3^- . Sensitivity
70 to CO_2 therefore varies in relation to the HCO_3^- : CO_2 preference and the affinity of
71 phytoplankton for carbon fixation. Some species can rapidly acclimate to changes in the
72 concentration of dissolved CO_2 , or total DIC (Nimer & Merrett, 1996). From an ecosystem
73 perspective, it has been suggested that the responses of phytoplankton to reduced pH in a high
74 CO_2 ocean are likely to be species-specific, with potential ‘winners’ and ‘losers’ (Hinga,
75 2002).

76 There is a growing body of literature on changes in photosynthesis due to increases in
77 CO_2 and related effects (Brading, Warner, Davey, Smith, Achterberg et al., 2011; Feng et al.,
78 2009; Feng, Warner, Zhang, Sun, Fu et al., 2008; Flynn, Clark, Mitra, Fabian, Hansen et al.,
79 2015). The majority of studies have been conducted on phytoplankton in laboratory culture
80 rather than on natural samples (Feng et al., 2008; Fu, Warner, Zhang, Feng & Hutchins, 2007;
81 King, Jenkins, Wallace, Liu, Wikfors et al., 2015; Shi, Li, Hopkinson, Hong, Li et al., 2015;
82 Wu, Gao & Riebesell, 2010). In diatoms, some cyanobacteria and coccolithophorids, elevated
83 CO_2 can lead to an increase in photosynthesis especially in large chain forming diatoms
84 (Tortell & Morel, 2002; Tortell, Payne, Li, Trimborn, Rost et al., 2008), *Synechococcus* spp.
85 (Fu et al., 2007) and *Emiliania huxleyi* (Leonardos & Geider, 2005). In the diatom
86 *Chaetoceros muelleri* for example, it has been shown that under saturating irradiance,

87 maximal photosynthetic rates are stimulated by increasing CO₂ availability (Ihnken, Roberts
88 & Beardall, 2011). For *Phaeodactylum tricornutum* grown at elevated CO₂ (1000 ppm)
89 corresponding to 7.8 pH, there was greater photoinhibition of the electron transport rate from
90 photosystem II (PSII) under high irradiance, whereas non-photochemical quenching was
91 reduced compared to low CO₂ grown cells (Wu et al., 2010).

92 Most studies on natural samples in the Atlantic Ocean have been conducted in
93 eutrophic and mesotrophic environments of the North Atlantic during natural or simulated
94 blooms. Some of these report an increase in the abundance of diatoms compared to
95 *Phaeocystis* spp. under CO₂ enrichment (Feng et al., 2008; Riebesell et al., 2007; Tortell et
96 al., 2008), others report that nano-phytoplankton or Prymnesiophytes replace diatoms under
97 elevated CO₂ and temperature in the Bering Sea (Hare, Leblanc, DiTullio, Kudela, Zhang et
98 al., 2007) and the North Atlantic (Feng et al., 2008), where there was a change from diatoms
99 to Prymnesiophytes under elevated CO₂ and a decrease in inorganic carbon production
100 (calcification). There have been few studies of CO₂ enrichment on changes in phytoplankton
101 community structure and photosynthesis in the oligotrophic gyres (Eggleston, Sabine & Morel,
102 2010), even though they occupy the largest areas of the ocean. To date there has only been
103 one study conducted in the sub-tropical North Atlantic at the Bermuda Atlantic Time Series
104 site (BATS), which showed no significant difference in carbon fixation rates at elevated pCO₂
105 (reduced pH) for phytoplankton assemblages dominated by *Prochlorococcus* sp. and
106 *Synechococcus* sp. (Lomas, Hopkinson, Losh, Ryan, Shi et al., 2012). By contrast, in the same
107 study nitrogen fixation rates in colonies of *Trichodesmium* increased by 54 % at pH 7.8 but
108 decreased by 21 % at pH 8.4.

109 The objective of this study was to assess the effect of CO₂ enrichment on
110 phytoplankton photosynthesis and community composition in the North Atlantic sub-tropical
111 gyre through a series of shipboard incubation experiments. Low temperature (LT) emission

112 spectra were used to assess changes in major pigment-protein complexes in the oxygenic
113 photosynthetic membranes. The LT spectra also provided information about the presence of
114 chlorophyll-containing light harvesting antenna complexes and their functional coupling to
115 photosystem reaction centres.

116

117 **2. Methods**

118 *2.1. Study area and experimental design*

119 Shipboard incubations were conducted aboard the *RRS James Cook* between 13 October and
120 21 November 2010 during the Atlantic Meridional Transect 20 (Cruise JC053). CO₂
121 enrichment experiments were carried out at three stations in the North Atlantic Gyre at 29°N,
122 34°W for experiment (Exp.) I; 18°N, 37°W for Exp. II and 7°N, 30°W for Exp. III (Fig. 1).
123 For each Exp., 80 L of near surface seawater was collected from the ship's underway supply
124 before dawn into a large Nalgene container. The water was sub-sampled into 18 acid cleaned
125 (IOC, 1994) 4 L clear polycarbonate bottles for incubation, leaving a head space of 0.5 L.
126 Nine of the bottles were bubbled with pre-mixed synthetic air with 360 ppm CO₂ (control)
127 and the other nine were bubbled with synthetic air with 760 ppm CO₂ (CO₂ treatment). The
128 gases were supplied to the experimental bottles from gas cylinders via nylon tubing fitted with
129 in-line 0.2 µm sterile Acrovent filters, to prevent contamination by particulates. The nylon
130 tubing entered the bottles through Nalgene caps with Kinesis Omni-Lok fittings and vent
131 tubes to prevent the build-up of pressure in the bottles. To minimise the effect of bubbling on
132 phytoplankton, each experimental bottle was bubbled for an initial 8 h period followed by a
133 further 4 h after 24 h to maintain the CO₂ over the 48 h period. The incubations were
134 conducted in an on-deck incubation system (78 x 60 x 68 cm) supplied with flow through sub-
135 surface seawater from the ship's underway supply to maintain the bottles at ambient
136 temperature. The incubation system had no light screen, but the sides of the incubator were

137 opaque which provided shading either side of zenith. pH, alkalinity, $p\text{CO}_2$, HCO_3^- , and Chl *a*
138 were measured in triplicate at 0 (T0), 6, (T6; except Chl *a*), 12 (T12), 24 (T24) and 48 (T48)
139 h. Samples for pico- and nanophytoplankton enumeration by flow cytometry, photosynthesis-
140 irradiance curves and low emission spectra were measured in triplicate at T0, T24 and T48.
141 Microscopy counts were made as single measurements at T0 and T48.

142

143 2.2. Carbonate chemistry: pH, $p\text{CO}_2$ and alkalinity

144 pH was determined spectrophotometrically onboard using $2 \mu\text{mol L}^{-1}$ of m-cresol from a 2
145 mmol L^{-1} stock solution which was prepared from pure sodium salt (Sigma-Aldrich, USA).
146 This method expresses pH on the total hydrogen concentration scale with 0.01 precision
147 (Dickson, Afghan & Anderson, 2003). Absorbance was recorded before and after addition of
148 m-cresol at 434, 578 and 730 nm on a PerkinElmer Lambda 35 spectrophotometer in a 100
149 mm cuvette. Seawater temperature was measured using an Amarell Ama-Digit 15
150 Temperature probe, and salinity was taken from a Seabird Electronics CTD calibrated against
151 salinity standards. For total alkalinity, samples were stored in 100 mL borosilicate dark bottles
152 with teflon cap liners spiked with 5% mercuric chloride (6.9 nmol L^{-1}). Alkalinity was
153 measured 2 months after the cruise using an automatic Apollo SciTech, Alkalinity Titrator
154 (Model AS-ALK2) in 0.8 M H_2SO_4 . $p\text{CO}_2$ was calculated from the pH, temperature, salinity
155 and alkalinity measurements using CO2SYS software (Pierrot & Wallace, 2006), using the
156 constants set according to Mehrbach et al. (1973) and Dickson and Millero (1987) and
157 corrected for differences between *in situ* and measured temperature.

158

159 2.3. Phytoplankton community structure

160 Nano and picoeukaryote phytoplankton cells from approx. 0.2 to 10 μm were enumerated
161 using a Becton Dickinson FACSortTM flow cytometer (Becton Dickinson, USA) equipped

162 with an air-cooled laser providing blue light at 488 nm, using the methods given in Tarran et
163 al. (Tarran, Heywood & Zubkov, 2006; Tarran, Zubkov, Sleight, Burkill & Yallop, 2001).
164 Picoeukaryotic phytoplankton and nanophytoplankton were analysed for between 2-4 minutes
165 to measure chlorophyll fluorescence (>650 nm), phycoerythrin fluorescence (585 ± 21 nm)
166 and side scatter (light scattered at 90° to the plane of the laser). Using a combination of
167 density plots as described in Tarran et al. (2001), it was possible to identify and quantify the
168 following planktonic groups in the samples: *Prochlorococcus* spp., *Synechococcus* spp.,
169 picoeukaryotic phytoplankton (approx. 0.6 to 3 μm , including prasinophyceae, chlorophyceae,
170 pelagophyceae) and nanoeukaryotic phytoplankton (approx. 3 to 10 μm , including
171 cryptophyceae and prymnesiophyceae). Instrument flow rate was calibrated daily using
172 Beckman Coulter Flowset fluorospheres of known concentration. Carbon biomass was
173 calculated using median cell diameters measured for each group using the methods described
174 in Tarran et al. (2006) and the carbon conversion factors given in (Bjørnsen, 1986) and
175 (Booth, 1988).

176 Microscopy was used to enumerate diatoms and dinoflagellates. For microscopic
177 counts, 100 mL of seawater samples were collected into brown borosilicate bottles and
178 preserved in Lugol's iodine (final concentration of 1 %) for stations II and III. Dinoflagellates
179 and diatoms were counted 2 months after the cruise using a Leica DM IRB microscope
180 (Leica, Germany). 100 mL of sample was settled in composite sedimentation chambers for 24
181 h. Samples were counted on 26 mm diameter sedimentation slides in replicate vertical and
182 horizontal transects using a x 20 objective. Cell volumes for these groups were calculated
183 using approximate geometric shapes and converted to biomass using the equations given in
184 Menden-Deuer and Lessard (2000).

185 For analysis of Chlorophyll-*a* (Chl *a*), 100 mL samples were filtered onto 25 mm
186 GF/F filters (pore size 0.7 μm , Whatman). The filter was then placed in a 15 mL FalconTM

187 tube (BD, UK) and 10 mL of 90 % acetone was added. The tubes were then stored in the dark,
188 in a -20°C freezer for 12 h. Chl *a* was then measured onboard fluorometrically following
189 [Holm-Hansen, Lorenzen, Holmes and Strickland \(1965\)](#) using a Turner Designs, Trilogy
190 Fluorometer.

191

192 2.4. *Phytoplankton photosynthesis*

193 2.4.1. *Photosynthesis-irradiance curves*

194 Photosynthetic-irradiance (P-E) experiments were conducted in linear photosynthetrons
195 illuminated with 50 W, 12 V tungsten halogen lamps following the methods described in
196 [Tilstone, Figueiras, Lorenzo and Arbones \(2003\)](#). Each incubator housed 16 sub-samples in
197 60 mL acid-cleaned polycarbonate bottles, which were inoculated with between 185 kBq (5
198 μCi) and 370 kBq (10 μCi) of ^{14}C labelled bicarbonate. The PE curves were conducted at the
199 same time each day to minimise the influence of diel light history. PAR in the incubator was
200 set to ambient levels measured over a 2 h period prior to incubation using a Skye Instruments
201 PAR Sensor (model SKP 200). For Exps I, II and III these were 1319, 2166 and 2215 $\mu\text{E m}^{-2}$
202 s^{-1} , respectively. The P-E curves were maintained at *in situ* temperature using the ship's non-
203 toxic seawater supply. After 1.5 h incubation, the suspended material was filtered through 25
204 mm GF/F filters which were then exposed to 37% hydrochloric acid fumes for 12 h and then
205 immersed in scintillation cocktail. ^{14}C disintegration time per minute (DPM) was measured
206 using the onboard 1414 liquid scintillation counter (PerkinElmer, USA) and the external
207 standard and the channel ratio methods to correct for quenching. Dark ^{14}C uptake was
208 subtracted from light uptake in the other 15 light incubation cells. Photosynthetic rates were
209 calculated from total dissolved inorganic carbon (DIC) and Chl *a*. DIC was calculated from
210 salinity and alkalinity. The broadband light-saturated Chl*a*-specific rate of photosynthesis, P_m^B
211 $[\text{mg C (mg Chl}a)^{-1} \text{h}^{-1}]$, and the light-limited slope, α^B $[\text{mg C (mg Chl}a)^{-1} \text{h}^{-1} (\mu\text{mol m}^{-2} \text{s}^{-1})^{-1}]$

212 ¹⁴C], were estimated by normalising ¹⁴C uptake to Chl *a* and fitting the data to the model of Platt
213 et al. (1980). Primary production (mg C m⁻³ d⁻¹) was subsequently calculated from
214 photosynthetically active radiation (E_{PAR}), Chl *a* and the photosynthetic parameters using the
215 brand model of Tilstone et al. (2003). E_{PAR} was modelled using the approach of Gregg &
216 Carder (1990) modified to include the effects of clouds (Reed, 1977) and using wind speed
217 and percentage cloud cover from the European Centre for Medium Range Weather
218 Forecasting (ECMWF) following Smyth et al. (2005).

219

220 2.4.2. Low temperature fluorescence emission spectra

221 Samples were filtered onto 25 mm GF/F filters from ~1 L of sea water and processed
222 following standard protocols (Suggett, Stambler, Prasil, Kolber, Quigg et al., 2009). A 4 x 10
223 mm piece of the filter was cut and placed onto a holder, flash frozen in liquid nitrogen and
224 measured immediately using a SM-9000 spectrometer (Photon Systems Instruments, Brno).
225 The instrument is based on a Zeiss MCS CCD spectrometer; the detector has 1044 pixels, the
226 grating images from 200-980 nm, the wavelength accuracy is 0.5 nm and the spectral
227 resolution is 3.5 nm (FWHM). Two gaussian shaped lines are separated at >3.5nm, with a
228 resolution of 3.2 nm from 0.8nm dispersion by 4 pixels. Whole fluorescence emission spectra
229 were determined using an integration time of 1000 ms. The fluorescence emission of blanks
230 (seawater pre-filtered through a 0.7 µm GF/F) was subtracted and the spectra were normalized
231 to 686 nm for de-convolution (Kaňa, Kotabová, Komárek, Papageorgiou, Govindjee et al.,
232 2012).

233

234 2.5. CTD and Nutrients.

235 A Seabird 911 plus CTD was deployed at each station at which experiments were conducted.

236 The data was processed using SeaBird software v7.21 and up and down casts were then

237 merged to 1 m binned resolution. Micro-molar nutrients were analysed using a 5 channel
238 nitrate (Brewer & Riley, 1965), nitrite (Grasshoff, 1976), phosphate, silicate (Kirkwood,
239 1989) and ammonia (Mantoura & Woodward, 1983) Bran & Luebbe AAIH segmented flow,
240 colourimetric, autoanalyser. Water samples were taken from a 24 x 20 litre bottle stainless
241 steel framed CTD / Rosette system (Seabird). These were sub-sampled into clean (acid-
242 washed) 60 mL HDPE (Nalgene) sample bottles. Subsequent nutrient analysis was complete
243 within 1-2 hours of sampling. Clean handling techniques were employed to avoid
244 contamination of the samples.

245

246 2.6. Statistical analysis

247 Paired T-tests samples were employed to test for significant differences between
248 treatments at T48 on individual Exp.'s and for all Exp.'s. Results of T-test analyses from all
249 Exp.'s (I-III) at 48 hrs are given in Table 2. Results of T-test analyses from individual Exp.'s
250 at 48 hrs are given on each sub-figure. The T-test results are given as $T_{1,19} = x$ and $P = y$
251 where T is the deviation of the sample mean from the normally distributed parametric mean to
252 parametric standard deviation ratio, the sub-script numbers (1, 23) denote the degrees of
253 freedom and P is the T-test critical significance value.

254

255 3. Results

256 3.1. Initial hydrographic conditions

257 Temperature varied from 26 °C to 29 °C in the surface waters of the sub-tropical Atlantic
258 (Figs. 1, 2). These stations were typically oligotrophic with a deep thermocline at between 50
259 and 75 m, low nitrate and nitrite (<0.02 µM), low surface Chl *a* (0.03-0.05 mg m⁻³) and deep
260 Chl *a* maxima between 70 and 130 m of between 0.1 and 0.3 mg m⁻³ (Fig. 2B, C, D). The

261 concentrations of nitrate+nitrite and silicate in the surface waters were beyond the detection
262 limit (Harris, 2011).

263

264 *3.2. Changes in carbonate chemistry*

265 For all experiments, by 48 h the CO₂ treatment resulted in an average concentration of 748
266 ppm and 7.97 pH units compared to 468 ppm and 8.14 pH units in the control (Fig. 3). For the
267 control, bubbling pre-mixed synthetic air with a CO₂ concentration of 360 ppm to natural
268 seawater, resulted in > 460 ppm (Fig. 3). At 48 h the difference in CO₂ content and pH
269 between the CO₂ treatment and control was ~280 ppm and 0.17 pH units, respectively (Fig.
270 3). Similarly, the mean HCO₃⁻ in the CO₂ treatment during these Exp.'s was 2013 ppm and
271 1860 ppm in the control (Fig. 4A, B, C) and the mean HCO₃⁻ : CO₂ ratio was 2.7 in the CO₂
272 treatment and 3.99 in the control (Fig. 4D, E, F).

273 The corresponding changes in total carbon (TC) and alkalinity (TA) during each experiment
274 are given in Table 1. For all Exp.'s, the TA in the control and CO₂ enrichment was similar and
275 consequently there was no significant difference in TA for Exp. II & III and all Exp.'s
276 together (Table 2). There was a significant difference in TA between CO₂ and control at T48
277 for Exp. I (T = 55.25, P = 0.012). By contrast, TC was significantly higher in the CO₂
278 treatment during Exp. II (T = 4.42, P = 0.048) & III (T = 7.08, P = 0.019) and all Exp.'s
279 together (Table 2), but this was not significant for Exp. I. Carbonate chemistry in the bottles
280 equilibrated at between 12 & 24 h. For all Exp.'s, there was a significant difference in CO₂,
281 pH, HCO₃⁻ and HCO₃⁻ : CO₂ TC, TA, (Table 2) between control and CO₂ treatments. At T48
282 in the CO₂ treatment, the CO₂ concentration was significantly higher than at T0 (Figure 3;
283 Table 2).

284

285 *3.3. Changes in phytoplankton community structure.*

286 During Exp.'s I & II there was no significant difference in Chl *a* between CO₂ and
287 control treatments (Fig. 5) and as a consequence, for all Exp.'s. there was no significant
288 difference between treatments and with T0 (Table 2). In Exp. III, Chl *a* was significantly
289 higher in the control (Fig. 5C). Of the phytoplankton groups enumerated by flow cytometry,
290 nanoeukaryotes had the highest biomass in Exp.'s I and II, whereas in Exp. III the biomass of
291 nano- and picoeukaryotes and *Synechococcus* spp. were similar (Fig. 6, 7). During all Exp.'s,
292 there was no difference in nanoeukaryotes and picoeukaryotes between control and CO₂
293 treatments (Table 2). During Exp. I, there was little change in the biomass of nanoeukaryotes
294 and picoeukaryotes over the duration of the Exp.'s and there was no significant difference
295 between control and CO₂ treatment (Fig. 6A, D). During Exp. II, the initial biomass of
296 nanoeukaryotes and picoeukaryotes decreased at 24 h followed by a slight increase at 48 h,
297 but there was no significant difference in biomass between control and CO₂ treatments at 48 h
298 (Fig. 6B, E). In Exp. III, the biomass of these groups initially decreased, but at 48 h both
299 nanoeukaryotes and picoeukaryotes were significantly higher in the control compared to the
300 CO₂ treatments and T0 (Fig. 6C, F). During all experiments there was no significant
301 difference between control and CO₂ treatments for *Synechococcus* spp. and *Prochlorococcus*
302 spp. (Fig. 7, Table 2). The biomass of *Prochlorococcus* spp. and *Synechococcus* spp.
303 decreased from T0 to T48 and T0 was significantly higher compared to the control and the
304 CO₂ treatment (Fig. 7, Table 2). For individual Exp.'s, there was no significant difference
305 between treatments for both *Synechococcus* spp. and *Prochlorococcus* spp. in Exp. I, II & III
306 (Fig. 7).

307 From the microscopy samples, there was no significant difference in diatoms biomass
308 between treatments during Exp.'s II and III (Table 2). The total biomass in Exp.'s II was low
309 3.5 and 4.5 µg C L⁻¹ at 48 h in CO₂ and control treatments, respectively when *Navicula* spp.
310 accounted for the highest biomass. In Exp. II, the diatom biomass was higher and 17 and 18

311 $\mu\text{g C L}^{-1}$ in CO_2 and control treatments, respectively when *Rhizosolenia* spp. and *Navicula*
312 spp. dominated (Fig. 8A, B). In this Exp. *Navicula* spp. dominated the biomass in the CO_2
313 treatment, whereas in the control *Rhizosolenia* spp. biomass was higher than any other group
314 (Fig. 8A, B). The biomass of diatoms was significantly higher in T0 compared to both the
315 control and CO_2 treatments (Fig. 8A, B, Table 2). The diatom biomass may have been
316 reduced through silicate limitation.

317 Dinoflagellates exhibited the highest biomass of all phytoplankton groups during
318 Exp.'s II and III, reaching on average $60 \mu\text{g C L}^{-1}$ under CO_2 treatment and $30 \mu\text{g C L}^{-1}$ in the
319 control, though the differences between treatments were not significant (Table 2).
320 *Gymnodinium* spp. accounted for the majority of the biomass, especially in Exp. III (Fig. 8D).
321 The dinoflagellate biomass was also significantly higher in the control at T0 compared to T48,
322 but this was not the case for the CO_2 treatment (Fig. 8C, D, Table 2). T-test was not
323 performed on individual Exp.'s, since only single samples were enumerated at T48.

324

325 3.4. Changes in photosynthesis and emission spectra

326 For all Exp.'s there was no significant difference in P_m^B between control and CO_2 treatment
327 (Fig. 9A, B, C, Table 2). For the individual Exp.'s, there was no difference in P_m^B between
328 control and CO_2 treatment in Exp. I (Fig. 9A), but P_m^B was significantly higher in the CO_2
329 treatment compared to the control in both Exp.'s II and III (Fig. 9B, C). In all Exp.'s, α^B was
330 significantly higher in the CO_2 treatment compared to the control (Fig. 9D, E, F, Table 2).
331 There was no significant difference in α^B in the control between T0 and T48, but α^B was
332 significantly higher at T48 in the CO_2 treatment compared to T0 (Table 2).

333 Using spectral de-convolution of LT emission spectroscopy measurements, the
334 fluorescence peak at 678 nm is associated with uncoupled chlorophyll-containing PS antenna,
335 while 686 nm represents a mixture of signals from PS II chlorophylls and terminal emitters of

336 phycobilisomes (PBS) (Rakhimberdieva, Vavilin, Vermaas, Elanskaya & Karapetyan, 2007).
337 Generally, the peak area at 678 nm decreased by more than three times from T0 to T48 in all
338 Exp.'s and in both treatments (Fig. 10, Table 3). In all of the CO₂ treatments however, there
339 was a significantly lower signal at 678 nm (Table 3), indicating reversible antenna connection
340 after CO₂ enrichment. In addition, the spectral emission of the PSII core antenna increased
341 from T0 to T48 (Fig. 10).

342

343 **4. Discussion**

344 *4.1. Changes in the phytoplankton community in relation to carbonate chemistry*

345 Experiments on CO₂ enrichment of natural samples in the Atlantic Ocean have mostly
346 focused on the interactions between diatoms, *Phaeocystis* spp. and prymnesiophytes. The sub-
347 tropical and tropical oligotrophic gyres are dominated by Cyanobacteria which make a
348 significant contribution to the carbon fixation in the global ocean (Bell & Kalff, 2001). In the
349 North Atlantic gyre, the phytoplankton assemblage at the deep Chl *a* maximum is typically
350 dominated by picoeukaryotes, *Prochlorococcus* spp. At the DCM and *Synechococcus* spp. in
351 the surface mixed layer (Tarran et al., 2006; Zubkov, Sleight, Tarran, Burkill & Leakey, 1998),
352 which exist at very low (beyond detection limit) nitrate and phosphate concentrations. Under
353 climate change scenarios of a warming ocean, increased stratification is likely to reduce
354 nutrient concentrations in the sub-tropical gyres (Sarmiento et al., 2004). We conducted
355 experiments with elevated CO₂ in the North Atlantic sub-tropical gyre during autumn to
356 evaluate the effects on phytoplankton community structure and photosynthetic rates. The
357 Exp.'s were conducted with no nutrient addition to mimic the oligotrophic conditions of the
358 gyre. The temperature at the experimental stations was 26 to 28°C and typical of stratified
359 sub-tropical waters where irradiance is high. During the Exp.'s, the phytoplankton community
360 was dominated by dinoflagellates, which was 55 to 80 times higher than the other groups, but

361 there was no significant difference between treatments and in some cases the biomass
362 decreased from T0 to T48 (Fig. 8).

363 Understanding the response of dinoflagellates to both increases in CO₂ and
364 temperature is key to detecting climate-driven perturbations in the ecosystem. In culture
365 experiments on autotrophic dinoflagellates, some strains of *Alexandrium fundyense*
366 (Hattenrath-Lehmann, Smith, Wallace, Merlo, Koch et al., 2015), *Karenia brevis* (Errera,
367 Yvon-Lewis, Kessler & Campbell, 2014), *Karlodinium veneficum* (Fu, Place, Garcia &
368 Hutchins, 2010), *Prorocentrum minimum* and *Heterosigma akashiwo* (Fu, Zhang, Warner,
369 Feng, Sun et al., 2008) grow significantly faster at high pCO₂. Additionally, *Alexandrium*
370 *fundyense* experiences a significant increase in cell toxicity under elevated CO₂ (Hattenrath-
371 Lehmann et al., 2015). There have been few studies on the response of *Gymnodinium* spp. and
372 *Gyrodinium* spp., which exhibited high biomass in this study. Most *Gymnodinium* spp. are
373 autotrophic, though some (e.g. *G. abbreviatum*, *G. heterostriatum*) are known heterotrophs
374 (Tomas, 1996). Similarly many *Gyrodinium* spp. are autotrophic, though some (e.g. *G.*
375 *lachryma*, *G. pingue*, *G. spirale*) are heterotrophic (Tomas, 1996), so the patterns shown in
376 Figures 8C, D may also include some heterotrophs as we were not able to identify all
377 *Gymnodinium* spp. and *Gyrodinium* spp. to species level. Calbet, Sazhin, Nejstgaard, Berger,
378 Tait et al. (2014) studied the response of natural phytoplankton communities in a Norwegian
379 Fjord using mesocosm experiments and found at elevated temperature and lower pH there was
380 no difference in *Gyrodinium* spp. abundance. Wynn-Edwards, King, Davidson, Wright,
381 Nichols et al. (2014) found that in continuous batch culture of Antarctic strains of
382 *Gymnodinium* spp. that there was no difference in growth rates and cell size between CO₂
383 enrichment to 993 ppm and ambient, but there was a significantly higher cell polyunsaturated
384 fatty acid content at the higher CO₂ concentrations.

385 In our Exp.'s Diatoms had the second highest biomass, but there was no difference
386 between the elevated CO₂ treatment and ambient (Fig. 8, Table 2). In other studies, the growth
387 rates of the diatoms *Skeletonema costatum* (Grev.) Cleve and *Chaetoceros* spp. increased at
388 elevated CO₂ by 100% and 24%, respectively (Beardall & Raven, 2004; Kim, Lee, Shin,
389 Kang, Lee et al., 2006). *Prochlorococcus* spp. exhibited the third highest biomass, however
390 there was also no significant difference in the biomass of *Prochlorococcus* spp., and that of
391 *Synechococcus* spp., picoeukaryotes, and nanoeukaryotes except in one Exp. III, in which
392 there was a higher biomass of both picoeukaryotes and nanoeukaryote in the control (Fig. 6).
393 The response of *Synechococcus* spp. in our experiments on natural samples contrasts the
394 trends observed in culture experiments. For example, Fu et al. (2007) found that 750 ppm CO₂
395 stimulated the growth rate of *Synechococcus* spp. The different responses between
396 *Synechococcus* spp. and *Prochlorococcus* spp. were thought to be due to differences in
397 inorganic carbon acquisition systems associated with carbon limitation at low pCO₂ in
398 *Synechococcus* spp. (Fu et al. 2007). In contrast, we observed no difference in biomass
399 between treatments under limiting nutrients. Similarly in natural phytoplankton communities
400 Lomas et al. (2012) found that under replete P and Fe there was no significant response in the
401 carbon-fixation of *Prochlorococcus* and *Synechococcus* spp. dominated communities to
402 elevated CO₂ and a decrease in pH. In their study, pH was adjusted by acid-base manipulation
403 (rather than direct enrichment with CO₂) and nutrient media were added to the incubations
404 even though nitrate and phosphate concentrations were below detection on first collection of
405 the samples. By contrast, in mesocosm exp.'s using natural phytoplankton communities with
406 replete nutrient concentrations, Paulino, Egge and Larsen (2008) reported a lower abundance
407 of *Synechococcus* spp. and a higher abundance of picoeukaryotes at high CO₂. By contrast,
408 Low-Decarie, Fussmann and Bell (2011) found in experimental mesocosms that increases in
409 CO₂ enhanced the competitive ability of chlorophytes relative to cyanobacteria. Similarly in

410 the North Atlantic, [Feng et al. \(2009\)](#) suggested that nanophytoplankton (coccolithophores)
411 may further increase in abundance relative to other phytoplankton groups in the later stages of
412 the spring bloom if future CO₂ concentrations and sea temperature continue to rise.

413 During our Exp.'s, dinoflagellates, diatoms and *Prochlorococcus* spp. decreased from
414 the initial T0 biomass, which may in part be due to the nitrate and silicate limitation or
415 grazing by heterotrophic and / or mixotrophic dinoflagellates, through confinement in bottles.
416 Nitrate and nitrite were low prior to confinement in all Exp.'s ([Fig. 2](#)) and silicate was beyond
417 the detection limit. The limited silicate availability meant that biomass of diatoms was low,
418 though it did increase in Exp. III. The biomass at T0 suggests that the cyanobacteria were
419 adversely affected by containment, which may have allowed the other groups to thrive due to
420 lack of competition and / or the recycled nutrients released during the decline of the
421 cyanobacteria biomass. Despite the initial decline, there was no clear response to elevated
422 CO₂ in the phytoplankton assemblage. Most studies on changes in the phytoplankton
423 community to elevated CO₂ report significant shifts or responses, however this may provide a
424 biased view since non-significant responses are rarely published ([Lomas et al., 2012](#)).

425

426 4.2. Effects of CO₂ enrichment on phytoplankton photosynthesis and primary production.

427 Aquatic photosynthesis is inherently under-saturated with respect to CO₂ ([Tortell, Rau](#)
428 [& Morel, 2000](#)), therefore increases in seawater CO₂ can enhance photosynthetic carbon
429 fixation ([Riebesell, 2004](#)). Some studies have observed a 40% increase in the uptake of DIC at
430 elevated CO₂ compared to present levels ([Riebesell et al., 2007](#)). It has been suggested that
431 the photosynthesis of dinoflagellates may be limited by current water column CO₂
432 concentrations and that they may benefit from rising atmospheric pCO₂ ([Eberlein, de Waal &](#)
433 [Rost, 2014](#)). There are a number of studies that contradict this hypothesis, for example, in
434 dilute batch cultures of the calcareous dinoflagellate *Scrippsiella trochoidea* and the toxic

435 dinoflagellate *Alexandrium tamarense* grown at a range of CO₂ concentrations (180–1200
436 ppm), there was no significant difference in photosynthetic and growth rates between
437 treatments, though *A. tamarense* did exhibit greater respiration rates at higher pCO₂ (Eberlein
438 et al., 2014). In our experiments, photosynthetic rates were significantly greater in the CO₂
439 enriched treatment (Fig. 9) and the dinoflagellates were probably able to maximise carbon
440 fixation due to lack of competition, since the biomass of the other groups was low (Fig. 6, 7,
441 8A, B). In Exp.'s II and III, when the dinoflagellates *Gymnodinium* and *Gyrodinium* spp.
442 dominated the biomass, significantly higher P_m^B and α^B at elevated CO₂ were associated with
443 a lower signal of unconnected PS antennae, suggesting that these previously unconnected PS
444 antennae become connected to facilitate the higher photosynthetic rates. In addition, there was
445 also a slight increase in fluorescence signal at 686 nm, due to a lower abundance of
446 *Prochlorococcus* spp. in the CO₂ treatment compared to the control, since this group do not
447 have PBS antennae (Partensky, Hess & Vault, 1999).

448 Photosynthetic carbon fixation is mediated by both pH and temperature dependent
449 enzymatic reactions via the Calvin-Benson Cycle (Giordano, Beardall & Raven, 2005; Portis
450 & Parry, 2007). Varying climate change scenarios of increased CO₂, temperature and a
451 decrease in pH may affect intra-cellular DIC uptake differently in different phytoplankton
452 groups or species (Badger et al., 1998). P_m^B and α^B and temperature are tightly coupled (Uitz,
453 Huot, Bruyant, Babin & Claustre, 2008), and different phytoplankton species are adapted to a
454 narrow temperature band with sharp decrease in photosynthetic rates beyond 20°C (Eppley,
455 1972; Raven & Geider, 1988) which can be modified through the synthesis of photoprotecting
456 rather than photosynthetic pigments (Kiefer & Mitchell, 1983). The enzyme that is primarily
457 responsible for photosynthetic carbon fixation is Ribulose Bisphosphate Carboxylase
458 (RuBisCO), though this does have some catalytic limitations (Giordano et al., 2005).
459 Phytoplankton have therefore adapted two strategies to maximise the performance of

460 RuBisCO; by either altering the affinity of RuBisCO or through CO₂-concentrating
461 mechanisms (CCMs). CCMs have a higher affinity for inorganic carbon (Giordano et al.,
462 2005; Raven, 2011a; Raven, 2011b; Reinfelder, 2011; Tortell et al., 2000), and through a
463 combination of RuBisCO activity or CCM, most species achieve photosynthetic saturation at
464 environmental CO₂ levels under light and nutrient replete conditions (Raven, 1997; Raven &
465 Johnston, 1991). The differences in photosynthetic efficiency and rates between different
466 phytoplankton groups can be partially explained by CCM and / or RuBisCO kinetics (Raven,
467 1997). The effect of an increase in CO₂ and decrease in pH may select phytoplankton species
468 or genotypes based on the efficiency of RuBisCO or CCM. Dinoflagellates possess Form II
469 RubisCO (Morse, Salois, Markovic & Hastings, 1995), which has low kinetic properties and
470 therefore a low photosynthetic efficiency (Tortell et al., 2000). They are abundant in low
471 nutrient oceanic waters (Margalef, 1979), have low growth rates attributed to their high basal
472 respiration rates (Tang, 1996), do not need to compete for their position in the ecosystem and
473 therefore rely on diffusive CO₂ uptake (Giordano et al., 2005). Dinoflagellates possess less
474 efficient CCMs and/or low affinities for CO₂, and may therefore benefit from living in a high
475 CO₂ world (Reinfelder 2011; Fu et al. 2012) and be highly sensitive to CO₂ (Rost et al.,
476 2008). Form II RUBISCO has a much lower selectivity for CO₂ over O₂ (Giordano et al.,
477 2005). To compensate for this, some marine dinoflagellates, such as *Prorocentrum minimum*,
478 *Heterocapsa triquetra*, *Ceratium lineatum*, *Protoceratium reticulatum* and *Peridinium*
479 *gatunense* have a CCM linked to carbonic anhydrases (CAs), the expression of which changes
480 in response to the external CO₂ concentration (Berman-Frank, Erez & Kaplan, 1998; Brading
481 et al., 2011; Ratti, Giordano & Morse, 2007; Rost, Riebesell & Sultemeyer, 2006). This
482 suggests that an increase in CO₂ stimulated efficiency in light absorption and utilization,
483 which resulted in the higher rates of carbon fixation shown in Figure 9. Using spectral de-
484 convolution of low temperature emission spectroscopy measurements, the peak at 691 nm is

485 emitted from active PS II chlorophylls and 711 nm from PS I chlorophylls (Kaňa et al., 2012).
486 The fluorescence peak at 678 nm was assigned to uncoupled chlorophyll-containing PS
487 antenna, while 686 nm represents a mixture of signals from PS II chlorophylls and terminal
488 emitters of phycobilisomes (PBS) (Rakhimberdieva et al., 2007). The peak at 686 nm
489 increased by a factor of two in all Exp.'s, which indicates either an increase in PBS content or
490 the presence of the iron-stress protein, IsiA (Chauhan, Folea, Jolley, Kouril, Lubner et al.,
491 2011). The lower signal of unconnected PS antennae in our Exp.'s maybe indicative of CCM
492 functionality as the dominant dinoflagellates connect more antennae to PS II to achieve the
493 higher P_m^B and α^B rates. The PS I chlorophyll peak at 711 nm remained constant and there
494 was no obvious difference between treatments (Fig. 10, Table 2). Although semi-quantitative,
495 the advantage of LT spectroscopy is that when it is applied to time series and bioassays,
496 gradual changes in phytoplankton photochemical processes can be monitored. This was the
497 case in the Exp.'s on this cruise, where more detailed biochemical or biophysical analyses
498 cannot usually be performed. Further work is required to ascertain whether the higher
499 photosynthetic rates we observed in *Gymnodinium*, and *Gyrodinium* spp. under CO₂
500 enrichment is a species or genotypic response. This response also needs to be further
501 investigated under different light conditions. For example, fluctuating light can affect
502 dinoflagellate growth and photosynthesis with constant light and high CO₂ having a negative
503 effect on the growth and quantum yield of *Prorocentrum micans* (Zheng, Giordano & Gao,
504 2015), which has implications for ocean warming and high stratification in a high CO₂ world.

505 Cyanobacteria and chlorophytes possess Form IB RuBisCO (Badger & Price, 2003).
506 Cyanobacteria utilize active HCO₃⁻ and CO₂ pumps as their CCM to facilitate CO₂ fixation
507 and maintain rapid growth at low external DIC concentrations (Badger & Price, 2003; Badger,
508 Price, Long & Woodger, 2006). In Exp. III, *Prochlorococcus* sp. biomass was significantly
509 higher in the control, possibly as a result of their high affinity for CO₂ uptake at low

510 concentrations or grazing of *Prochlorococcus* sp. biomass by heterotrophs or mixotrophs (Fig.
511 3D), but the photosynthetic rates in the control were significantly lower than the CO₂
512 treatment, principally because the dinoflagellate biomass was far higher in this Exp. By
513 comparison, Fu et al. (2007) also observed that the photosynthetic rate and light harvesting
514 efficiency of *Prochlorococcus* sp. *in vivo* was not affected by increasing CO₂. Similar to the
515 patterns we saw in Exp.'s II & III, Fu et al. (2007) also observed only slight differences in
516 growth and photosynthetic rates of *Synechococcus* spp. at 750 ppm CO₂ and control, though
517 this changed when the temperature was elevated by 4 °C. By comparison, Lomas et al. (2012)
518 observed that natural phytoplankton assemblages dominated by cyanobacteria were able to
519 adapt to changes in pH through modification of the assimilation number, which resulted in no
520 significant differences in photosynthetic rates. We found in two of our Exp.'s that the
521 assimilation number was significantly higher at elevated CO₂. The duration of our
522 experiments represent short term responses to CO₂ enrichment and do not account for longer
523 term adaptation and evolution, which can be complex (Collins & Bell, 2006). Our
524 experiments also do not account for interactive synergies between CO₂, temperature, nutrients
525 and light which may affect phytoplankton, as shown by other studies in the North Atlantic
526 (Feng et al., 2009; Hare et al., 2007). Further work is required to elucidate these dual effects
527 on phytoplankton community structure in the North Atlantic sub-tropical gyre.

528 What is the consequence of increases in CO₂ on water column integrated primary
529 production? Some studies have reported no significant changes in primary production under
530 elevated CO₂ (Goldman, 1999; Raven & Falkowski, 1999; Tortell & Morel, 2002; Tortell et
531 al., 2000). We found in the North Atlantic sub-tropical gyre when dinoflagellates dominated,
532 that there was a 25% increase in primary production over all Exp.'s at 760ppm CO₂ and that
533 this was significant in Exp.'s II & III (Fig. 11). This is similar to the findings of Hein and

534 [SandJensen \(1997\)](#) who suggested that increasing CO₂ concentrations in seawater would
535 increase primary production by 15 to 19%.

536

537 **5. Conclusions**

538 Three CO₂ enrichment experiments were conducted on natural phytoplankton
539 assemblages in the North Atlantic sub-tropical gyre. The dinoflagellates, *Gymnodinium*, and
540 *Gyrodinium* spp., dominated the biomass but there was no significant difference in the
541 biomass of these and other phytoplankton groups at elevated CO₂ concentrations of 760 ppm
542 and 7.94 pH compared to ambient (468 ppm CO₂ and 8.33 pH). There were however,
543 significantly higher photosynthetic rates in the elevated CO₂ treatment which was due to the
544 connection of reversible photosystem antennae at higher CO₂ concentrations, which resulted
545 in a 25% increase primary production.

546

547 **Acknowledgements.** We thank the Master and the Crew of the *RRS James Cook* for their
548 help during AMT20 (cruise JC053). The training on alkalinity analysis by Dr. Helen Findlay
549 is gratefully acknowledged. We also thank the Natural Environment Research Council
550 (NERC) Earth Observation Data Acquisition and Analysis Service (NEODAAS) at Plymouth
551 Marine Laboratory for supplying satellite imagery to the ship. We are grateful to Dr. Rob
552 Thomas for supplying Chl *a* data and to Carolyn Harris and Malcolm Woodward for the
553 nutrient data shown in Fig. 2. G.H.T. and G.A.T. were supported by the Natural Environment
554 Research Council's National Capability – The Atlantic Meridional Transect (AMT) and
555 Oceans 2025 Theme 10a. B.S. was supported by a fellowship from the Partnership for
556 Observations of the Global Oceans (POGO), for training on board *RRS James Cook* during
557 AMT20. B.S., O.P. and R.K were supported by GACR 206/08/1683 and the project Algatech

558 CZ1.05/2.1.00/03.01/0110. This study is a contribution to the International IMBER project
559 and is AMT contribution number 210.

560
561 **References**

- 562 Alley, R.B., Berntsen, T., Bindoff, N.L., Chen, Z., others, a., 2007. *Summary for*
563 *policymakers*. Cambridge and New York: Cambridge University Press.
- 564 Badger, M.R., Andrews, T.J., Whitney, S.M., Ludwig, M., Yellowlees, D.C., Leggat, W.,
565 Price, G.D., 1998. The diversity and coevolution of Rubisco, plastids, pyrenoids, and
566 chloroplast-based CO₂-concentrating mechanisms in algae. *Canadian Journal of Botany-*
567 *Revue Canadienne De Botanique*, 76, 1052-1071.
- 568 Badger, M.R., Price, G.D., 2003. CO₂ concentrating mechanisms in cyanobacteria: molecular
569 components, their diversity and evolution. *Journal of Experimental Botany*, 54, 609-622.
- 570 Badger, M.R., Price, G.D., Long, B.M., Woodger, F.J., 2006. The environmental plasticity
571 and ecological genomics of the cyanobacterial CO₂ concentrating mechanism. *Journal of*
572 *Experimental Botany*, 57, 249-265.
- 573 Beardall, J., Raven, J.A., 2004. The potential effects of global climate change on microalgal
574 photosynthesis, growth and ecology. *Phycologia*, 43, 26-40.
- 575 Bell, T., Kalff, J., 2001. The contribution of picophytoplankton in marine and freshwater
576 systems of different trophic status and depth. *Limnology and Oceanography*, 46, 1243-1248.
- 577 Berman-Frank, I., Erez, J., Kaplan, A., 1998. Changes in inorganic carbon uptake during the
578 progression of a dinoflagellate bloom in a lake ecosystem. *Canadian Journal of Botany-Revue*
579 *Canadienne De Botanique*, 76, 1043-1051.
- 580 Bjørnsen, P.K., 1986. Automatic determination of bacterioplankton biomass by means of
581 image analysis. *Applied and Environmental Microbiology*, 51, 1199-1204.
- 582 Booth, B.C., 1988. Size classes and major taxonomic groups of phytoplankton at two
583 locations in the subarctic Pacific Ocean in May and August, 1984. *Marine Biology*, 97, 275-
584 286.
- 585 Bopp, L., Monfray, P., Aumont, O., Dufresne, J.L., Le Treut, H., Madec, G., Terray, L., Orr,
586 J.C., 2001. Potential impact of climate change on marine export production. *Global*
587 *Biogeochemical Cycles*, 15, 81-99.
- 588 Brading, P., Warner, M.E., Davey, P., Smith, D.J., Achterberg, E.P., Suggett, D.J., 2011.
589 Differential effects of ocean acidification on growth and photosynthesis among phylotypes of
590 Symbiodinium (Dinophyceae). *Limnology and Oceanography*, 56, 927-938.
- 591 Brewer, P.G., Riley, J.P., 1965. The automatic determination of nitrate in seawater. *Deep-Sea*
592 *Research I*, 12, 765-772.
- 593 Calbet, A., Sazhin, A.F., Nejstgaard, J.C., Berger, S.A., Tait, Z.S., Olmos, L., Sousoni, D.,
594 Isari, S., Martinez, R.A., Bouquet, J.-M., Thompson, E.M., Bamstedt, U., Jakobsen, H.H.,
595 2014. Future Climate Scenarios for a Coastal Productive Planktonic Food Web Resulting in
596 Microplankton Phenology Changes and Decreased Trophic Transfer Efficiency. *Plos One*, 9.
- 597 Chauhan, D., Folea, I.M., Jolley, C.C., Kouril, R., Lubner, C.E., Lin, S., Kolber, D., Wolfe-
598 Simon, F., Golbeck, J.H., Boekema, E.J., Fromme, P., 2011. A Novel Photosynthetic Strategy
599 for Adaptation to Low-Iron Aquatic Environments. *Biochemistry*, 50, 686-692.
- 600 Collins, S., Bell, G., 2006. Evolution of natural algal populations at elevated CO₂. *Ecology*
601 *Letters*, 9, 129-135.
- 602 Dickson, A.G., Afghan, J.D., Anderson, G.C., 2003. Reference materials for oceanic CO₂
603 analysis: a method for the certification of total alkalinity. *Marine Chemistry*, 80, 185-197.

604 Dickson, A.G., Millero, F.J., 1987. A comparison of the equilibrium constants for the
605 dissociation of carbonic acid in seawater media. *Deep-Sea Research Part a-Oceanographic*
606 *Research Papers*, 34, 1733-1743.

607 Doney, S.C., Lindsay, K., Fung, I., John, J., 2006. Natural variability in a stable, 1000-yr
608 global coupled climate-carbon cycle simulation. *Journal of Climate*, 19, 3033-3054.

609 Eberlein, T., de Waal, D.B.V., Rost, B., 2014. Differential effects of ocean acidification on
610 carbon acquisition in two bloom-forming dinoflagellate species. *Physiologia Plantarum*, 151,
611 468-479.

612 Egleston, E.S., Sabine, C.L., Morel, F.M.M., 2010. Revelle revisited: Buffer factors that
613 quantify the response of ocean chemistry to changes in DIC and alkalinity. *Global*
614 *Biogeochemical Cycles*, 24.

615 Elzenga, J.T.M., Prins, H.B.A., Stefels, J., 2000. The role of extracellular carbonic anhydrase
616 activity in inorganic carbon utilization of *Phaeocystis globosa* (Prymnesiophyceae): A
617 comparison with other marine algae using the isotopic disequilibrium technique. *Limnology*
618 *and Oceanography*, 45, 372-380.

619 Eppley, R.W., 1972. Temperature and phytoplankton growth in the sea. *Fisheries Bulletin*, 70,
620 1063-1085.

621 Errera, R.M., Yvon-Lewis, S., Kessler, J.D., Campbell, L., 2014. Responses of the
622 dinoflagellate *Karenia brevis* to climate change: pCO₂ and sea surface temperatures.
623 *Harmful Algae*, 37, 110-116.

624 Feng, Y., Hare, C.E., Leblanc, K., Rose, J.M., Zhang, Y., DiTullio, G.R., Lee, P.A., Wilhelm,
625 S.W., Rowe, J.M., Sun, J., Nemcek, N., Gueguen, C., Passow, U., Benner, I., Brown, C.,
626 Hutchins, D.A., 2009. Effects of increased pCO₂ and temperature on the North Atlantic
627 spring bloom. I. The phytoplankton community and biogeochemical response. *Marine*
628 *Ecology-Progress Series*, 388, 13-25.

629 Feng, Y., Warner, M.E., Zhang, Y., Sun, J., Fu, F.-X., Rose, J.M., Hutchins, D.A., 2008.
630 Interactive effects of increased pCO₂, temperature and irradiance on the marine
631 coccolithophore *Emiliana huxleyi* (Prymnesiophyceae). *European Journal of Phycology*, 43,
632 87-98.

633 Flynn, K.J., Clark, D.R., Mitra, A., Fabian, H., Hansen, P.J., Glibert, P.M., Wheeler, G.L.,
634 Stoecker, D.K., Blackford, J.C., Brownlee, C., 2015. Ocean acidification with
635 (de)eutrophication will alter future phytoplankton growth and succession. *Proceedings of the*
636 *Royal Society B-Biological Sciences*, 282.

637 Fu, F.-X., Place, A.R., Garcia, N.S., Hutchins, D.A., 2010. CO₂ and phosphate availability
638 control the toxicity of the harmful bloom dinoflagellate *Karlodinium veneficum*. *Aquatic*
639 *Microbial Ecology*, 59, 55-65.

640 Fu, F.-X., Warner, M.E., Zhang, Y., Feng, Y., Hutchins, D.A., 2007. Effects of increased
641 temperature and CO₂ on photosynthesis, growth, and elemental ratios in marine
642 *Synechococcus* and *Prochlorococcus* (Cyanobacteria). *Journal of Phycology*, 43, 485-496.

643 Fu, F.-X., Zhang, Y., Warner, M.E., Feng, Y., Sun, J., Hutchins, D.A., 2008. A comparison of
644 future increased CO₂ and temperature effects on sympatric *Heterosigma akashiwo* and
645 *Prorocentrum minimum*. *Harmful Algae*, 7, 76-90.

646 Giordano, M., Beardall, J., Raven, J.A., 2005. CO₂ concentrating mechanisms in algae:
647 Mechanisms, environmental modulation, and evolution. *Annual Review of Plant Biology*, Vol.
648 56 (pp. 99-131).

649 Goldman, J.C., 1999. Inorganic carbon availability and the growth of large marine diatoms.
650 *Marine Ecology-Progress Series*, 180, 81-91.

651 Grasshoff, K., 1976. *Methods of Seawater analysis*. Weinheim: Verlag, Chemie.

652 Gregg, W.W., Carder, K.L., 1990. A Simple Spectral Solar Irradiance Model for Cloudless
653 Maritime Atmospheres. *Limnology and Oceanography*, 35, 1657-1675.

654 Hare, C.E., Leblanc, K., DiTullio, G.R., Kudela, R.M., Zhang, Y., Lee, P.A., Riseman, S.,
655 Hutchins, D.A., 2007. Consequences of increased temperature and CO₂ for phytoplankton
656 community structure in the Bering Sea. *Marine Ecology-Progress Series*, 352, 9-16.

657 Harris, C., 2011. AMT20 Nutrients. In A.P. Rees (Ed.), *AMT20 Cruise Report, RRS James*
658 *Cook JC053*. (p. 148). Plymouth, UK: Plymouth Marine Laboratory.

659 Hattenrath-Lehmann, T.K., Smith, J.L., Wallace, R.B., Merlo, L.R., Koch, F., Mittelsdorf, H.,
660 Goleski, J.A., Anderson, D.M., Gobler, C.J., 2015. The effects of elevated CO₂ on the growth
661 and toxicity of field populations and cultures of the saxitoxin-producing dinoflagellate,
662 *Alexandrium fundyense*. *Limnology and Oceanography*, 60, 198-214.

663 Hein, M., SandJensen, K., 1997. CO₂ increases oceanic primary production. *Nature*, 388,
664 526-527.

665 Hinga, K.R., 2002. Effects of pH on coastal marine phytoplankton. *Marine Ecology-Progress*
666 *Series*, 238, 281-300.

667 Holm-Hansen, O., Lorenzen, C.J., Holmes, R.W., Strickland, J.D.H., 1965. Fluorometric
668 determination of chlorophyll. *Journal du Conseil Permanent International pour l'Exploration*
669 *de la Mer.* , 30, 3-15.

670 Hopkinson, B.M., Dupont, C.L., Allen, A.E., Morel, F.M.M., 2011. Efficiency of the CO₂-
671 concentrating mechanism of diatoms. *Proceedings of the National Academy of Sciences of the*
672 *United States of America*, 108, 3830-3837.

673 Houghton, R.A., 2001. Counting terrestrial sources and sinks of carbon. *Climatic Change*, 48,
674 525-534.

675 Ihnken, S., Roberts, S., Beardall, J., 2011. Differential responses of growth and
676 photosynthesis in the marine diatom *Chaetoceros muelleri* to CO₂ and light availability.
677 *Phycologia*, 50, 182-193.

678 IOC, 1994. Protocols for the Joint Global Ocean Flux Study (JGOFS) core measurements.,
679 Vol. 29 (p. 126). IOC Manuals and Guides.

680 Kaňa, R., Kotabová, E., Komárek, O., Papageorgiou, G.C., Govindjee, Šedivá, B., Prášil, O.,
681 2012. *Biochimica et Biophysica Acta - Bioenergetics in press*.

682 Kiefer, D.A., Mitchell, B.G., 1983. A Simple, Steady-State Description of Phytoplankton
683 Growth Based on Absorption Cross-Section and Quantum Efficiency. *Limnology and*
684 *Oceanography*, 28, 770-776.

685 Kim, J.-M., Lee, K., Shin, K., Kang, J.-H., Lee, H.-W., Kim, M., Jang, P.-G., Jang, M.-C.,
686 2006. The effect of seawater CO₂ concentration on growth of a natural phytoplankton
687 assemblage in a controlled mesocosm experiment. *Limnology and Oceanography*, 51, 1629-
688 1636.

689 King, A.L., Jenkins, B.D., Wallace, J.R., Liu, Y., Wikfors, G.H., Milke, L.M., Meseck, S.L.,
690 2015. Effects of CO₂ on growth rate, C:N:P, and fatty acid composition of seven marine
691 phytoplankton species. *Marine Ecology Progress Series*, 537, 59-69.

692 Kirkwood, D.S., 1989. Simultaneous determination of selected nutrients in seawater. *ICES*,
693 *CM*, 29.

694 Leonardos, N., Geider, R.J., 2005. Elevated atmospheric carbon dioxide increases organic
695 carbon fixation by *Emiliania huxleyi* (Haptophyta), under nutrient-limited high-light
696 conditions. *Journal of Phycology*, 41, 1196-1203.

697 Lomas, M.W., Hopkinson, B.M., Losh, J.L., Ryan, D.E., Shi, D.L., Xu, Y., Morel, F.M.M.,
698 2012. Effect of ocean acidification on cyanobacteria in the subtropical North Atlantic. *Aquatic*
699 *Microbial Ecology*, 66, 211-222.

700 Low-Decarie, E., Fussmann, G.F., Bell, G., 2011. The effect of elevated CO₂ on growth and
701 competition in experimental phytoplankton communities. *Global Change Biology*, 17, 2525-
702 2535.

703 Mantoura, R.F.C., Woodward, E.M.S., 1983. Optimization of the indophenol blue method for
704 the automated-determination of ammonia in estuarine waters. *Estuarine Coastal and Shelf*
705 *Science*, 17, 219-224.

706 Margalef, R., 1979. Life Forms of Phyto Plankton as Survival Alternatives in an Unstable
707 Environment. *Oceanologica Acta*, 1, 493-509.

708 Mehrbach, C., Culberso, Ch, Hawley, J.E., Pytkowic, Rm, 1973. Measurement of Apparent
709 Dissociation-Constants of Carbonic-Acid in Seawater at Atmospheric-Pressure. *Limnology*
710 *and Oceanography*, 18, 897-907.

711 Menden-Deuer, S., Lessard, E.J., 2000. Carbon to volume relationships for dinoflagellates,
712 diatoms, and other protist plankton. *Limnology and Oceanography*, 45, 569-579.

713 Morse, D., Salois, P., Markovic, P., Hastings, J.W., 1995. A Nuclear-Encoded Form-Ii
714 Rubisco in Dinoflagellates. *Science*, 268, 1622-1624.

715 Nimer, N.A., Merrett, M.J., 1996. The development of a CO₂-concentrating mechanism in
716 *Emiliana huxleyi*. *New Phytologist*, 133, 383-389.

717 Partensky, F., Hess, E.R., Vaultot, D., 1999. Prochlorococcus, a marine photosynthetic
718 prokaryote of global significance. *Microbiol. Mol. Biol. Rev.*, 63, 106-127.

719 Paulino, A.I., Egge, J.K., Larsen, A., 2008. Effects of increased atmospheric CO₂ on small
720 and intermediate sized osmotrophs during a nutrient induced phytoplankton bloom.
721 *Biogeosciences*, 5, 739-748.

722 Pierrot, D.E., Wallace, D.W.R., 2006. MS Excel Program Developed for CO₂ System
723 Calculations. ORNL/CDIAC-105a., *Carbon Dioxide Information Analysis Center, Oak Ridge*
724 *National Laboratory*,. USA: Department of Energy, Oak Ridge, Tennessee.

725 Platt, T., Gallegos, C.L., Harrison, W.G., 1980. Photoinhibition of photosynthesis in natural
726 assemblage of marine phytoplankton. *J Mar Res*, 38, 687-701.

727 Portis, A.R., Jr., Parry, M.A.J., 2007. Discoveries in Rubisco (Ribulose 1,5-bisphosphate
728 carboxylase/oxygenase): a historical perspective. *Photosynthesis Research*, 94, 121-143.

729 Rakhimberdieva, M.G., Vavilin, D.V., Vermaas, W.F.J., Elanskaya, I.V., Karapetyan, N.V.,
730 2007. Phycobilin/chlorophyll excitation equilibration upon carotenoid-induced non-
731 photochemical fluorescence quenching in phycobilisomes of the *Synechocystis* sp PCC 6803.
732 *Biochimica Et Biophysica Acta-Bioenergetics*, 1767, 757-765.

733 Ratti, S., Giordano, M., Morse, D., 2007. CO₂-concentrating mechanisms of the potentially
734 toxic dinoflagellate *Protoceratium reticulatum* (Dinophyceae, Gonyaulacales). *Journal of*
735 *Phycology*, 43, 693-701.

736 Raven, J.A., 1997. Inorganic carbon acquisition by marine autotrophs. *Advances in Botanical*
737 *Research, Vol 27: Classic Papers, Vol. 27* (pp. 85-209).

738 Raven, J.A., 2011a. Effects on Marine Algae of Changed Seawater Chemistry with Increasing
739 Atmospheric CO₂. *Biology and Environment-Proceedings of the Royal Irish Academy*,
740 111B, 1-17.

741 Raven, J.A., 2011b. Inorganic carbon acquisition by eukaryotic algae: four current questions.
742 *Photosynthesis Research*, 106, 123-134.

743 Raven, J.A., Falkowski, P.G., 1999. Oceanic sinks for atmospheric CO₂. *Plant Cell and*
744 *Environment*, 22, 741-755.

745 Raven, J.A., Geider, R.J., 1988. Temperature and Algal Growth. *New Phytologist*, 110, 441-
746 461.

747 Raven, J.A., Johnston, A.M., 1991. Mechanisms of Inorganic-Carbon Acquisition in Marine-
748 Phytoplankton and Their Implications for the Use of Other Resources. *Limnology and*
749 *Oceanography*, 36, 1701-1714.

750 Reed, R., 1977. On estimating insolation over the ocean. *J. Phys. Oceanogr.*, 7, 482-485.

751 Reinfelder, J.R., 2011. Carbon Concentrating Mechanisms in Eukaryotic Marine
752 Phytoplankton. In C.A. Carlson, S.J. Giovannoni (Eds.), *Annual Review of Marine Science*,
753 *Vol 3*, Vol. 3 (pp. 291-315).

754 Riebesell, U., 2004. Effects of CO₂ enrichment on marine phytoplankton. *Journal of*
755 *Oceanography*, 60, 719-729.

756 Riebesell, U., 2008. Climate change - Acid test for marine biodiversity. *Nature*, 454, 46-47.

757 Riebesell, U., Schulz, K.G., Bellerby, R.G.J., Botros, M., Fritsche, P., Meyerhoefer, M., Neill,
758 C., Nondal, G., Oschlies, A., Wohlers, J., Zoellner, E., 2007. Enhanced biological carbon
759 consumption in a high CO₂ ocean. *Nature*, 450, 545-U510.

760 Riebesell, U., Wolfgladow, D.A., Smetacek, V., 1993. Carbon-Dioxide Limitation of
761 Marine-Phytoplankton Growth-Rates. *Nature*, 361, 249-251.

762 Rost, B., Riebesell, U., Sultemeyer, D., 2006. Carbon acquisition of marine phytoplankton:
763 Effect of photoperiod length. *Limnology and Oceanography*, 51, 12-20.

764 Rost, B., Zondervan, I., Wolf-Gladrow, D., 2008. Sensitivity of phytoplankton to future
765 changes in ocean carbonate chemistry: current knowledge, contradictions and research
766 directions. *Marine Ecology-Progress Series*, 373, 227-237.

767 Sarmiento, J.L., Slater, R., Barber, R., Bopp, L., Doney, S.C., Hirst, A.C., Kleypas, J.,
768 Matear, R., Mikolajewicz, U., Monfray, P., Soldatov, V., Spall, S.A., Stouffer, R., 2004.
769 Response of ocean ecosystems to climate warming. *Global Biogeochemical Cycles*, 18.

770 Shi, D., Li, W., Hopkinson, B.M., Hong, H., Li, D., Kao, S.-J., Lin, W., 2015. Interactive
771 effects of light, nitrogen source, and carbon dioxide on energy metabolism in the diatom
772 *Thalassiosira pseudonana*. *Limnology and Oceanography*, 60, 1805-1822.

773 Skirrow, G., 1975. *The dissolved gases - carbon dioxide*.

774 Smyth, T.J., Tilstone, G.H., Groom, S.B., 2005. Integration of radiative transfer into satellite
775 models of ocean primary production. *Journal of Geophysical Research-Oceans*, 110.

776 Suggett, D.J., Stambler, N., Prasil, O., Kolber, Z., Quigg, A., Vazquez-Dominguez, E.,
777 Zohary, T., Berman, T., Iluz, D., Levitan, O., Lawson, T., Meeder, E., Lazar, B., Bar-Zeev,
778 E., Medova, H., Berman-Frank, I., 2009. Nitrogen and phosphorus limitation of oceanic
779 microbial growth during spring in the Gulf of Aqaba. *Aquatic Microbial Ecology*, 56, 227-
780 239.

781 Tang, E.P.Y., 1996. Why do dinoflagellates have lower growth rates? *Journal of Phycology*,
782 32, 80-84.

783 Tarran, G.A., Heywood, J.L., Zubkov, M.V., 2006. Latitudinal changes in the standing stocks
784 of nano- and picoeukaryotic phytoplankton in the Atlantic Ocean. *Deep-Sea Research Part Ii-*
785 *Topical Studies in Oceanography*, 53, 1516-1529.

786 Tarran, G.A., Zubkov, M.V., Sleigh, M.A., Burkill, P.H., Yallop, M., 2001. Microbial
787 community structure and standing stocks in the NE Atlantic in June and July of 1996. *Deep-*
788 *Sea Research Part Ii-Topical Studies in Oceanography*, 48, 963-985.

789 Tilstone, G.H., Figueiras, F.G., Lorenzo, L.M., Arbones, B., 2003. Phytoplankton
790 composition, photosynthesis and primary production during different hydrographic conditions
791 at the Northwest Iberian upwelling system. *Marine Ecology-Progress Series*, 252, 89-104.

792 Tomas, C.R., 1996. *Identifying Marine Diatoms and Dinoflagellates*. San Diego, California:
793 Academic press Inc.

794 Tortell, P.D., Morel, F.M.M., 2002. Sources of inorganic carbon for phytoplankton in the
795 eastern Subtropical and Equatorial Pacific Ocean. *Limnology and Oceanography*, 47, 1012-
796 1022.

797 Tortell, P.D., Payne, C.D., Li, Y., Trimborn, S., Rost, B., Smith, W.O., Riesselman, C.,
798 Dunbar, R.B., Sedwick, P., DiTullio, G.R., 2008. CO₂ sensitivity of Southern Ocean
799 phytoplankton. *Geophysical Research Letters*, 35.

800 Tortell, P.D., Rau, G.H., Morel, F.M.M., 2000. Inorganic carbon acquisition in coastal Pacific
801 phytoplankton communities. *Limnology and Oceanography*, 45, 1485-1500.
802 Uitz, J., Huot, Y., Bruyant, F., Babin, M., Claustre, H., 2008. Relating phytoplankton
803 photophysiological properties to community structure on large scales. *Limnology and*
804 *Oceanography*, 53, 614-630.
805 Wu, Y., Gao, K., Riebesell, U., 2010. CO₂-induced seawater acidification affects
806 physiological performance of the marine diatom *Phaeodactylum tricornutum*. *Biogeosciences*,
807 7, 2915-2923.
808 Wynn-Edwards, C., King, R., Davidson, A., Wright, S., Nichols, P.D., Wotherspoon, S.,
809 Kawaguchi, S., Virtue, P., 2014. Species-Specific Variations in the Nutritional Quality of
810 Southern Ocean Phytoplankton in Response to Elevated pCO₂. *Water*, 6, 1840-1859.
811 Zheng, Y., Giordano, M., Gao, K., 2015. The impact of fluctuating light on the dinoflagellate
812 *Prorocentrum micans* depends on NO₃⁻ and CO₂ availability. *Journal of Plant Physiology*,
813 180, 18-26.
814 Zubkov, M.V., Sleigh, M.A., Tarran, G.A., Burkill, P.H., Leakey, R.J.G., 1998.
815 Picoplanktonic community structure on an Atlantic transect from 50 degrees N to 50 degrees
816 S. *Deep-Sea Research Part I-Oceanographic Research Papers*, 45, 1339-1355.

817
818 **Figure Legends.**

819 **Fig. 1.** Stations in the North Atlantic at which CO₂ enrichment experiments were conducted
820 overlaid on MODIS-aqua Sea Surface Temperature monthly composite image for October
821 2010.

822
823 **Fig. 2.** Initial profiles of temperature (blue line), nitrate (black line, open squares) and
824 chlorophyll a (green line, filled squares) at stations sampled for the CO₂ enrichment
825 experiments (A.) I, (B.) II, (C.) III. Dotted line is the euphotic depth.

826
827 **Fig. 3.** Changes in pCO₂ during experiments (A) I, (B) II, (C) III, and in pH during
828 experiments (D) I, (E) II, (F) III in control (open circles) and 760 ppm CO₂ (closed circles).
829 Broken line in (C) and (F) represents phases of (space) and dark (line) during the experiments.

830
831 **Fig. 4.** Changes in HCO₃⁻ during experiments (A) I, (B) II, (C) III, and in HCO₃⁻ : CO₂ during
832 experiments (D) I, (E) II, (F) III in control (open circles) and 760 ppm CO₂ (closed circles).
833 Broken line in (C) and (F) represents phases of light (space) and dark (line) during the
834 experiments.

835

836 **Fig. 5.** Changes in Chlorophyll *a* during experiments (A) I, (B) II, (C) III. Broken line in (C)
837 represents phases of light (space) and dark (line) during the experiments.

838

839 **Fig. 6.** Changes in the biomass of nanoeukaryotes in experiments (A.) I, (B.) II, (C.) III, and
840 changes in biomass of picoeukaryotes in experiments (D.) I, (E.) II, (F.) III in control (open
841 circles) and 760 ppm CO₂ (closed circles). Broken line in (C) and (F) represents phases of
842 light (space) and dark (line) during the experiments.

843

844 **Fig. 7.** Changes in the biomass of *Synechococcus* spp. in experiments (A.) I, (B.) II, (C.) III,
845 and changes in biomass of *Prochlorococcus* spp. in experiments (D.) I, (E.) II, (F.) III in
846 control (open circles) and 760 ppm CO₂ (closed circles). Broken line in (C) and (F) represents
847 phases of light (space) and dark (line) during the experiments.

848

849 **Fig. 8.** Changes in diatom biomass during experiments (A) II and (B) III, and dinoflagellate
850 biomass during experiments (C) II and (D) III at time(T) 0 and 48 h in control (AIR) and 760
851 ppm CO₂ enrichment. In (A) and (B) brown is *Navicula* spp., orange is pennate diatoms, light
852 green is *Nitzschia* spp., dark green is *Thalassiosira* spp., light blue is *Rhizosolenia* spp. and
853 dark blue is total diatom biomass. In (C) and (D) black is *Amphidinium* spp., dark blue is
854 *Gonyaulax* spp., light blue is *Gymnodinium* spp., green is *Gyrodinium* spp., yellow is total
855 dinoflagellate biomass (N=1; microscope transects = 4).

856

857 **Fig. 9.** Changes in P_m^B [mg C (mg Chl*a*)⁻¹ h⁻¹] during experiments (A.) I, (B.) II and (C.) III.

858 Changes in α^B [mg C (mg Chl*a*)⁻¹ h⁻¹ (μmol m⁻² s⁻¹)⁻¹] during experiments (D.) I, (E.) II and

859 (F.) III. Broken line in (C) and (F) represents phases of light (**space**) and dark (**line**) during
860 the experiments.

861

862 **Fig. 10.** Changes in low temperature emission during experiment (A) I CO₂ treatment, (B) I
863 control, (C) II CO₂ treatment, (D) II control, (E) III CO₂ treatment, and (F) III control (N=1).
864 Spectra are: black – phycoerythrin; red – phycocyanin; light green - unconnected antenna;
865 dark-blue – photosystem (PS) II or phycobilisomes anchoring pigments; yellow – PSII core
866 antenna; cyanin – PSI; light-blue – vibrational peak; grey – cumulative sum of all spectra.

867

868 **Fig. 11.** Changes in Primary production [$\text{mg C m}^{-3} \text{ d}^{-1}$] at 48h in control (grey bars) and 760
869 ppm CO₂ (black bars) for experiments I-II.