

University of Windsor

Scholarship at UWindsor

Biological Sciences Publications

Department of Biological Sciences

2000

Survival and growth of mussels subsequent to hemolymph sampling for DNA

J. F. Yanick

Daniel D. Heath
University of Windsor

Follow this and additional works at: <https://scholar.uwindsor.ca/biologypub>



Part of the [Biology Commons](#)

Recommended Citation

Yanick, J. F. and Heath, Daniel D., "Survival and growth of mussels subsequent to hemolymph sampling for DNA" (2000). *Journal of Shellfish Research*, 19, 2, 991-993.

<https://scholar.uwindsor.ca/biologypub/1139>

This Article is brought to you for free and open access by the Department of Biological Sciences at Scholarship at UWindsor. It has been accepted for inclusion in Biological Sciences Publications by an authorized administrator of Scholarship at UWindsor. For more information, please contact scholarship@uwindsor.ca.



<http://www.biodiversitylibrary.org/>

Journal of shellfish research.

[S.l. :National Shellfisheries Association,1981-

<http://www.biodiversitylibrary.org/bibliography/2179>

v. 19 (2000): <http://www.biodiversitylibrary.org/item/28581>

Article/Chapter Title: Survival and growth of mussels subsequent to hemolymph sampling for DNA

Author(s): Heath

Page(s): Page 991, Page 992, Page 993

Holding Institution: MBLWHOI Library

Sponsored by: MBLWHOI Library

Generated 5 October 2017 10:45 PM

<http://www.biodiversitylibrary.org/pdf4/070160600028581>

This page intentionally left blank.

SURVIVAL AND GROWTH OF MUSSELS SUBSEQUENT TO HEMOLYMPH SAMPLING FOR DNA

JENIA F. YANICK AND DANIEL D. HEATH*

Biology, College of Science and Management
University of Northern British Columbia
3333 University Way, Prince George
British Columbia, Canada, V2N 4Z9

ABSTRACT With the increasing use of molecular genetic techniques in ecology and evolution, it has become apparent that methods of non-destructive DNA sampling must be developed. In this study we collected 50 blue mussels (*Mytilus* spp.) in each of three size categories: small (10–20 mm), medium (20–30 mm), and large (30+ mm). Hemolymph was extracted from 25 mussels in each size category and the remaining 25 mussels served as controls. The hemolymph was extracted and control mussels were monitored for 384 days, during which time no significant differences in survival or growth were found. We extracted DNA from the hemolymph and successfully polymerase chain reaction-amplified the ITS and Glu-5' species-specific markers from 81% and 92% of the samples, respectively, and determined that all mussels were *Mytilus trossulus* (Lamarck). The extraction of hemolymph for DNA analysis allows for molecular investigations of populations or species which are either rare or in limited numbers, and for life history investigations where survival of the organism is necessary.

KEY WORDS: non-destructive, *Mytilus*, PCR, DNA, survival, growth, hemolymph

INTRODUCTION

Molecular genetic techniques and the genetic characterization of individuals have become common in the study of the ecology and evolution of marine invertebrates, particularly bivalves (Mitton 1994). The benefit of the polymerase chain reaction (PCR) is that very little DNA is required, thus making analysis possible when the quantity and/or quality of DNA is limited. PCR-based species markers have been used within the *Mytilus* species complex for conservation, ecological, and evolutionary applications (Heath et al. 1995, 1996, Rawson et al. 1996). Molecular genetic characterization has also clarified population genetic structure in a variety of other bivalves (Sarver and Foltz 1993, Manuel et al. 1996, David et al. 1997, Suchanek et al. 1997, Herbinger et al. 1998), as well as aiding in investigations into the ecology and life history of bivalves with planktonic phases (Toro 1998).

Typically, shellfish are destructively sampled in order to sample tissue for DNA extraction. Destructive sampling involves killing the animal to obtain the necessary tissue for genetic analysis (Taberlet et al. 1999). Although this is acceptable for some studies (Sarver and Foltz 1993, Heath et al. 1995, Hare et al. 1996, Heath et al. 1996, Suchanek et al. 1997, Herbinger et al. 1998, Toro 1998), destructive sampling is clearly not acceptable for studies involving growth or survival measurements, or for investigations of small or rare populations. For such studies there is a need for a technique that would allow the collection of DNA without harming the organism. Non-destructive sampling generally involves capturing the target organism, taking an invasive sample without killing it, and then releasing it (Taberlet et al. 1999). It is, therefore, important that any potential technique be tested for even minor adverse effects on the survival or growth of the target organism.

Here we describe sampling hemolymph for DNA extraction from shellfish. Specifically, we sampled hemolymph from three size categories of blue mussels (*Mytilus* spp.), extracted DNA, and

amplified fragments using PCR with species-specific markers. Hemolymph is made up of mostly water, but does contain cells, including nucleated hemocytes (Morse and Zardus 1997) and is responsible for the transportation of digestion products throughout the body (Brusca and Brusca 1990), among other functions. We followed the survival and growth of hemolymph-extracted and control mussel groups for over 1 y to ascertain whether this technique resulted in decreased survival and/or growth.

MATERIALS AND METHODS

Fifty mussels in each of three size categories were collected from the western coast of Quadra Island, located near Campbell River, British Columbia, Canada. The mussels were measured with calipers to the nearest 0.01 mm and sorted into small (10–20 mm), medium (20–30 mm), and large (30+ mm) size categories. In each group, hemolymph was extracted (50–200 μ L) from 25, while the other 25 (control) mussels were handled, but not sampled. A 1-cc syringe (22-gauge, 1.5-inch needle) was inserted through the rear hinge joint and hemolymph was extracted until no more fluid could be removed. The extracted hemolymph was expelled into 1.0 mL of 95% ethanol and stored at room temperature. After sampling, mussels were placed in six cages (3 hemolymph-extracted and 3 control) and hung approximately 1 m below the surface at the original collection site. The cages were 5 \times 5 \times 10 cm and were slotted to enable free water flow through the cages, but excluded potential predators.

The hemolymph and alcohol were transported to the laboratory where they were centrifuged (13,000 rpm, 15 min), the liquid was removed, and the pellet dried (LABCONCO Centrivap Concentrator) at 60 $^{\circ}$ C for 8 min. The dried cells were digested overnight in 200 μ L of lysis buffer (10 mM Tris-HCl, pH 8.0, 15 mM ethylenediamine tetra acetate, and 0.5% sodium dodecyl sulphate) and 125 μ g of proteinase K at 37 $^{\circ}$ C. The solution was then extracted once with an equal volume of phenol:chloroform:isoamyl alcohol (24:24:1), followed by isopropanol precipitation (Heath et al. 1995). The extracted DNA was resuspended in 100 μ L of double-distilled water and was then PCR-amplified following the ITS protocol described in Heath et al. (1995) and the

*Corresponding author, current address: Great Lakes Institute for Environmental Research, University of Windsor, Windsor, ON, Canada N9B 3P4.

Glu-5' protocol described in Rawson et al. (1996). Both markers give species-specific results (Heath et al. 1995, Rawson et al. 1996) and thus we are reasonably certain we amplified target DNA, and not contaminating DNA. The PCR products were visualized on a 1.8% agarose gel stained with ethidium bromide (Fig. 1). Individual mussels were scored for genotype at each marker locus on the basis of a diagnostic restriction fragment length polymorphism (ITS; Heath et al. 1995) or using an automated DNA sequencer to determine amplified fragment length (GLU-5'; Rawson et al. 1996).

Between April 3, 1998 and April 22, 1999, the mussels in this experiment were measured and the survivors were counted at three sampling times after transfer to the experimental cages (58, 140, and 384 days). Student's *t* test (shell length) and chi-square (survival) were used for statistical analysis to determine whether differences existed between the two groups at day 58 and day 384. Day 58 comparisons were made to test for short-term effects, while day 384 comparisons were made for long-term effects.

RESULTS

DNA was successfully PCR-amplified from 61 of the 75 samples (81%) for ITS, and from 69 of the 75 samples (92%) for GLU (Fig. 1). There was no consistent effect of mussel size (and hence hemolymph volume) on the success of the PCR amplification. All mussels were determined to be *Mytilus trossulus*. The hemolymph technique was found to have little effect on either survival or growth (Fig. 2). At day 58 and day 384, the survivorship of the hemolymph-extracted mussels was not found to be significantly different than the survivorship of the control mussels in any of the size categories ($P > 0.10$). At day 58, the control mussels were slightly larger than the hemolymph-extracted mussels in the small size category ($P = 0.018$), but there was no significant difference in either the large or the medium size categories ($P > 0.50$). At day 384, the hemolymph-extracted mussels

were larger than the control mussels in the large size category ($P = 0.033$), but there was no significant difference in the medium or small size categories ($P > 0.10$).

There were also no consistent differences found in survival or growth among the mussel size categories. The smallest mussels we sampled were between 10 and 20 mm and had mortality and growth similar to the larger size categories.

DISCUSSION

We sampled a wide range of sizes of mussels using a non-destructive method of DNA sampling and successfully extracted DNA for PCR purposes from most of the mussels, including those in the small category. Our PCR success rates were comparable to those of Heath et al. (1995) who used destructive tissue sampling methods. The technique described here is a useful tool for field-work, as it does not require the killing of the organism under study. Furthermore, we found no consistent effect of hemolymph sampling on either survival or growth of the mussels. Although this is not surprising for the larger mussels, it is unexpected for the small animals, as the extraction of a large portion of the organism's body fluid would be expected to negatively effect the organism's growth and/or survival.

Although we used mussels, our technique is applicable to other bivalve species. For example, Manuel et al. (1996) used a similar technique on scallops (*Placopecten magellanicus*); however, they extracted approximately 5 to 10 times the volume of hemolymph and did not test for potential growth or survival effects of their sampling method. Other researchers have reported hemolymph sampling in bivalves for various purposes, including DNA extraction (Marsh et al. 1995), hemocyte pathology and function (Moore et al. 1991; Oliver & Fisher 1995), and ploidy analysis (Komaru et al. 1988). However, no attempt was made to determine the effect of that sampling on the viability of the animals. Our study also




Figure 1. Agarose gel electrophoresis of PCR-amplified DNA fragments using the ITS and Glu-5' species-specific primer sets. The various lanes are PCR results using DNA from a regular extraction method (R) and DNA extracted from hemolymph taken from mussels in three size classes (L, large; M, medium; S, small).




Figure 2. Comparison of mean shell length (± 1 SE) and survival for hemolymph-extracted and control mussels from three size classes over a 385-day period, post-treatment. The circles represent the large size class, the squares represent the medium size class, and the triangles represent the small size class. The filled symbols with the solid lines are the hemolymph-extracted mussels, while the open symbols with the dashed lines are the control mussels.

showed the syringe-extracted hemolymph consistently provides PCR-quality DNA from large numbers of animals sampled under field conditions. Hemolymph extraction clearly has considerable potential for studies requiring the non-destructive sampling of DNA from bivalves, and thus has applications for growth and survival studies. This technique will also be useful for studies of the ecology and population dynamics of bivalves where destructive sampling of the organism is either not permitted or not desirable for the experimental design.

ACKNOWLEDGMENTS

We would like to thank C. Bryden, R. Hepburn, and S. Henry for their field assistance. This study was supported by a Natural Science and Engineering Research Council of Canada grant (to D.D.H.). Fieldwork was funded by Yellow Island Aquaculture, Ltd., the Natural Science and Engineering Research Council of Canada, and the Science Council of British Columbia supplied post-graduate funding to J.F.Y.

LITERATURE CITED

- Brusca, R. C. & G. J. Brusca. 1990. Invertebrates. Sinauer Associates Inc., Sunderland, MA.
- David, P., M.-A. Perdieu, A-F. Pernot & P. Jarne. 1997. Fine-grained spatial and temporal population genetic structure in the marine bivalve *Spisula ovalis*. *Evolution* 51:1318-1322.
- Hare, M. P., S. A. Karl & J. C. Avise. 1996. Anonymous nuclear DNA markers in the American oyster and their implications for the heterozygote deficiency problem in marine bivalves. *Mol. Biol. Evol.* 13:334-345.
- Heath, D. D., P. D. Rawson & T. J. Hilbish. 1995. PCR-based nuclear markers identify alien blue mussel (*Mytilus* spp.) genotypes on the west coast of Canada. *Can. J. Fish. Aquat. Sci.* 52:2621-2627.
- Heath, D. D., D. R. Hatcher & T. J. Hilbish. 1996. Ecological interaction between sympatric *Mytilus* species on the west coast of Canada investigated using PCR markers. *Mol. Ecol.* 5:443-447.
- Herbinger, C. M., B. M. Vercaemer, B. Gjetvaj & R. K. O'Dor. 1998. Absence of genetic differentiation among geographically close sea scallop (*Placopecten magellanicus* G.) beds with cDNA and microsatellite markers. *J. Shellfish Res.* 17:117-122.
- Komaru, A., Y. Uchimura, H. Ieyama & K.T. Wada. 1988. Detection of induced triploid scallop, *Chlamys nobilis*, by DNA microfluorimetry with DAPI staining. *Aquaculture* 69:201-209.
- Manuel, J. L., S. Burbridge, E. L. Kenchington, M. Ball & R. K. O'Dor. 1996. Veligers from two populations of scallop *Placopecten magellanicus* exhibit different vertical distributions in the same mesocosm. *J. Shellfish Res.* 15:251-257.
- Marsh, A.G., J.D. Gauthier & G.R. Vasta. 1995. A semiquantitative PCR assay for assessing *Perkinsus marinus* infections in the eastern oyster, *Crassostrea virginica*. *J. Parasitology* 81:577-583.
- Mitton, J. B. 1994. Molecular approaches to population biology. *Annu. Rev. Ecol. Syst.* 25:45-69.
- Moore, J.D., R.A. Elston, A.S. Drum & M.T. Wilkinson. 1991. Alternate pathogenesis of systemic neoplasia in the bivalve mollusc *Mytilus*. *J. Invert. Pathol.* 58:231-243.
- Morse, M. P. & J. D. Zardus. 1997. Bivalvia. pp. 7-118. In: F.W. Harrison (ed.). *Microscopic Anatomy of Invertebrates*, Vol. 6A, Mollusca II. Wiley-Liss, New York.
- Oliver, L.M. & W.S. Fisher. 1995. Comparative form and function of oyster *Crassostrea virginica* hemocytes from Chesapeake Bay (Virginia) and Apalachicola Bay (Florida). *Dis. Aquat. Org.* 22:217-225.
- Rawson, P. D., K. L. Joyner, K. Meetze & T. J. Hilbish. 1996. Evidence for intragenic recombination within a novel genetic marker that distinguishes mussels in the *Mytilus edulis* species complex. *Heredity* 77: 599-607.
- Sarver, S. K. & D. W. Foltz. 1993. Genetic population structure of a species' complex of blue mussels (*Mytilus* spp.). *Mar. Biol.* 117:105-112.
- Suchanek, T. H., J. B. Geller, B. R. Kreiser & J. B. Mitton. 1997. Zoo-geographic distributions of the sibling species *Mytilus galloprovincialis* and *M. trossulus* (Bivalvia: Mytilidae) and their hybrids in the North Pacific. *Biol. Bull.* 193:187-194.
- Taberlet, P., L. P. Waits & G. Luikart. 1999. Noninvasive genetic sampling: look before you leap. *Tree* 14:323-327.
- Toro, J.E. 1998. Molecular identification of four species of mussels from Southern Chile by PCR-based nuclear markers: the potential use in studies involving planktonic surveys. *J. Shellfish Res.* 17:1203-1205.

The following text is generated from uncorrected OCR.

[Begin Page: Page 991]

Journal of Shellfish Research, Vol. 14, No. 2, W1-W3, 2000.

SURVIVAL AND GROWTH OF MUSSELS SUBSEQUENT TO HEMOLYMPH

SAMPLING FOR DNA

JENIA F. YANICK AND DANIEL D. HFIATH*

Biology, College of Science and Management

University of Northern British Columbia

3333 University Way, Prince George

British Columbia, Canada. V2N 4Z9

ABSTRACT With the increasing use of molecular genetic techniques in ecology and evolution, it has become apparent that methods

of non-destructive DNA sampling must be developed. In this study we collected 70 blue mussels (*Mytilus* spp.) in each of three size

categories: small (10-20 mm), medium (20-30 mm), and large (30+ mm). Hemolymph was extracted from 25 mussels in each size

category and the remaining 25 mussels served as controls. The hemolymph was extracted and control mussels were monitored for 384

days, during which time no significant differences in survival or growth were found. We extracted DNA from the hemolymph and

successfully polymerase chain reaction-amplified the ITS and Glu-5' species-specific markers from 81% and 92% of the samples,

respectively, and determined that all mussels were *Mytilus trossulus* (Lamarck). The extraction of hemolymph for DNA analysis allows

for molecular investigations of populations or species which are either rare or in limited numbers, and for life history investigations

where survival of the organism is necessary.

KEY WORDS: non-destructive. *Mytilus*. PCR. DNA. survival, growth, hemolymph

INTRODUCTION

Molecular genetic techniques and the genetic characterization of individuals have become common in the study of the ecology and evolution of marine invertebrates, particularly bivalves (Mitten 1994). The benefit of the polymerase chain reaction (PCR) is that very little DNA is required, thus making analysis possible when the quantity and/or quality of DNA is limited. PCR-based species markers have been used within the *Mytilus* species complex for conservation, ecological, and evolutionary applications (Heath et al. 1995, 1996, Rawson et al. 1996). Molecular genetic characterization has also clarified population genetic structure in a variety of other bivalves (Sarver and Foltz 1993, Manuel et al. 1996, David et al. 1997, Suchanek et al. 1997, Herbinger et al. 1998), as well as aiding in investigations into the ecology and life history of bivalves with planktonic phases (Toro 1998).

Typically, shellfish are destructively sampled in order to sample tissue for DNA extraction. Destructive sampling involves killing the animal to obtain the necessary tissue for genetic analysis (Taberlet et al. 1999). Although this is acceptable for some studies (Sarver and Foltz 1993, Heath et al. 1995, Hare et al. 1996, Heath et al. 1996, Suchanek et al. 1997, Herbinger et al. 1998, Toro 1998), destructive sampling is clearly not acceptable for studies involving growth or survival measurements, or for investiga-

tions of small or rare populations. For such studies there is a need for a technique that would allow the collection of DNA without harming the organism. Non-destructive sampling generally involves capturing the target organism, taking an invasive sample without killing it, and then releasing it (Taberlet et al. 1999). It is, therefore, important that any potential technique be tested for even minor adverse effects on the survival or growth of the target organism.

Here we describe sampling hemolymph for DNA extraction from shellfish. Specifically, we sampled hemolymph from three size categories of blue mussels (*Mytilus* spp.), extracted DNA, and

*Corresponding author, current address: Great Lakes Institute for Environmental Research, University of Windsor, Windsor, ON, Canada N9B 3P4.

amplified fragments using PCR with species-specific markers. Hemolymph is made up of mostly water, but does contain cells, including nucleated hemocytes (Morse and Zardus 1997) and is responsible for the transportation of digestion products throughout the body (Brusca and Brusca 1990), among other functions. We followed the survival and growth of hemolymph-extracted and control mussel groups for over 1 y to ascertain whether this technique resulted in decreased survival and/or growth.

MATERIALS AND METHODS

Fifty mussels in each of three size categories were collected from the western coast of Quadra Island, located near Campbell

River, British Columbia, Canada. The mussels were measured with calipers to the nearest 0.01 mm and sorted into small (10-20 mm), medium (20-30 mm), and large (30-40 mm) size categories. In each group, hemolymph was extracted (50-200 µL) from 25, while the other 25 (control) mussels were handled, but not sampled. A 1-cc syringe (22-gauge, 1.5-inch needle) was inserted through the rear hinge joint and hemolymph was extracted until no more fluid could be removed. The extracted hemolymph was expelled into 1.0 mL of 95% ethanol and stored at room temperature. After sampling, mussels were placed in six cages (3 hemolymph-extracted and 3 control) and hung approximately 1 m below the surface at the original collection site. The cages were 5 x 5 x 10 cm and were slotted to enable free water flow through the cages, but excluded potential predators.

The hemolymph and alcohol were transported to the laboratory where they were centrifuged (13,000 rpm, 15 min), the liquid was removed, and the pellet dried (LABCONCO Centrivap Concentrator) at 60 °C for 8 min. The dried cells were digested overnight in 200 µL of lysis buffer (10 mM Tris-HCl, pH 8.0, 15 mM ethylenediamine tetra acetate, and 0.5% sodium dodecyl sulphate) and 125 µg of proteinase K at 37 °C. The solution was then extracted once with an equal volume of phenol:chloroform:isoamyl alcohol (24:24:1), followed by isopropanol precipitation (Heath et al. 1995). The extracted DNA was resuspended in 100 µL of double-distilled water and was then PCR-amplified following the ITS protocol described in Heath et al. (1995) and the

[Begin Page: Page 992]

Yanick and Heath

Glu-5' protocol described in Rawson et al. (1996). Both markers give species-specific results (Heath et al. 1995. Rawson et al. 1996) and thus we are reasonably certain we amplified target DNA, and not contaminating DNA. The PCR products were visualized on a 1.8% agarose gel stained with ethidium bromide (Fig. 1). Individual mussels were scored for genotype at each marker locus on the basis of a diagnostic restriction fragment length polymorphism (ITS: Heath et al. 1995) or using an automated DNA sequencer to determine amplified fragment length (GLU-5'; Rawson et al. 1996).

Between April 3, 1998 and April 22, 1999, the mussels in this experiment were measured and the survivors were counted at three sampling times after transfer to the experimental cages (58, 140, and 384 days). Student's *t* test (shell length) and chi-square (survival) were used for statistical analysis to determine whether differences existed between the two groups at day 58 and day 384. Day 58 comparisons were made to test for short-term effects, while day 384 comparisons were made for long-term effects.

RESULTS

DNA was successfully PCR-amplified from 61 of the 75 samples (81%) for ITS, and from 69 of the 75 samples (92%) for GLU (Fig. 1). There was no consistent effect of mussel size (and hence hemolymph volume) on the success of the PCR amplification. All mussels were determined to be *Mxtihis trosstilus*. The hemolymph technique was found to have little effect on either survival or growth (Fig. 2). At day 58 and day 384, the survivorship of the hemolymph-extracted mussels was not found to be significantly different than the survivorship of the control mussels in any of the size categories ($P > 0.10$). At day 58, the control mussels were slightly larger than the hemolymph-extracted mussels in the small size category ($P = 0.018$), but there was no significant difference in either the large or the medium size categories ($P > 0.50$). At day 384, the hemolymph-extracted mussels were larger than the control mussels in the large size category ($P = 0.033$), but there was no significant difference in the medium or small size categories ($P > 0.10$).

There were also no consistent differences found in survival or growth among the mussel size categories. The smallest mussels we sampled were between 10 and 20 mm and had mortality and growth similar to the larger size categories.

DISCUSSION

We sampled a wide range of sizes of mussels using a non-

destructive method of DNA sampling and successfully extracted DNA for PCR purposes from most of the mussels, including those in the small category. Our PCR success rates were comparable to those of Heath et al. (1995) who used destructive tissue sampling methods. The technique described here is a useful tool for field-work, as it does not require the killing of the organism under study. Furthermore, we found no consistent effect of hemolymph sampling on either survival or growth of the mussels. Although this is not surprising for the larger mussels, it is unexpected for the small animals, as the extraction of a large portion of the organism's body fluid would be expected to negatively effect the organism's growth and/or survival.

Although we used mussels, our technique is applicable to other bivalve species. For example, Manuel et al. (1996) used a similar technique on scallops (*Placopecten magetlanicus*): however, they extracted approximately 5 to 10 times the volume of hemolymph and did not test for potential growth or survival effects of their sampling method. Other researchers have reported hemolymph sampling in bivalves for various purposes, including DNA extraction (Marsh et al. 1995), hemocyte pathology and function (Moore et al. 1991; Oliver* Fisher 1995), and ploidy analysis (Komaru et al. 1988). However, no attempt was made to determine the effect of that sampling on the viability of the animals. Our study also

ITS

Glu-5'

RLMS RLMS

1000 bp-

500 bp -

300 bp -

Kijuruf I. Aj-arosc gel clec-tri>ph<iri'sis n[I'CK-ampiilu'd DNA Ir:i(;nuiils usiny llic I IS and (ilii-S' ^)^^.il•^~^(H•li(k
prinur mls. lhc various lanes

are l't'U resiills iisinu 1)N.\ from a regular exlruction iik'thocl (UI and DNA extracted from hemolymph taken from
mussels in three size classes

(L, large; M, medium; S, small).

[Begin Page: Page 993]

Survival and Growth ok Hkmo'i mph-Samples Mussels

993

Control

Hemolymph Extracted

400

Time (days)

Figure 2. Comparison of mean shell length (± 1 SE) and survival for hemolymph-extracted and control mussels from three size classes over a

385-day period, post-treatment. The circles represent the large size class, the squares represent the medium size class, and the triangles represent

the small size class. The Tilled symbols with the solid lines are the hemolymph-extracted mussels, while the open symbols with the dashed lines

are the control mussels.

showed the syringe-extracted hemolymph consistently provides PCR-quality DNA from large numbers of animals sampled under field conditions. Hemolymph extraction clearly has considerable potential for studies requiring the non-destructive sampling of DNA from bivalves, and thus has applications for growth and survival studies. This technique will also be useful for studies of the ecology and population dynamics of bivalves where destructive sampling of the organism is either not permitted or not desirable for the experimental design.

ACKNOWLEDGMENTS

We would like to thank C. Bryden, R. Hepburn, and S. Henry for their field assistance. This study was supported by a Natural Science and Engineering Research Council of Canada grant (to D.D.H.). Fieldwork was funded by Yellow Island Aquaculture, Ltd., the Natural Science and Engineering Research Council of Canada, and the Science Council of British Columbia supplied post-graduate funding to J.F.Y.

LITERATURE CITED

Brusca. R. C. & G. J. Brusca. IWO. Invertebrates. Sinauer Associates Inc..

Sunderland. MA.

David. P., M.-A. Perdieu, A-F. Pemot & P. Jarne. 1997. Fine-grained spatial and temporal population genetic structure in the marine bivalve *Spisulit ovalis*. *Evolution* 51:1318-1322.

Hare. M. P., S. A. Karl & J. C. Avise. 1996. Anonymous nuclear DNA markers in the American oyster and their implications for the heterozygote deficiency problem in marine bivalves. *Mol. Biol. Evol.* 13:334-345.

Heath. D. D., P. D. Rawson & T. J. Hilbish. 1995. PCR-based nuclear markers identify alien blue mussel (*Mytilus* spp.) genotypes on the west coast of Canada. *Can. J. Fish. Aquat. Sci.* 52:2621-2627.

Heath. D. D., D. R. Hatcher & T. J. Hilbish. 1996. Ecological interaction between sympatric *Mytilus* species on the west coast of Canada investigated using PCR markers. *Mol. Ecol.* 5:443-447.

Herbinger. C. M., B. M. Vercaemer, B. Gjetvaj & R. K. O'Dor. 1998. Absence of genetic differentiation among geographically close sea scallop (*Placofyeclen magelkinicus* G.) beds with cDNA and microsatellite markers. *Mar. Biol.* 131:117-122.

Komaru, A., Y. Uchimura, H. Ieyama & K.T. Wada. 1988. Detection of induced triploid scallop, *Chlamys nobilis*, by DNA microfluorimetry with DAPI staining. *Aquaculture* 69:201-209.

Manuel. J. L., S. Burbridge, E. L. Kenchington, M. Ball & R. K. O'Dor.

1996. Veligers from two populations of scallop *Ptarmacolen maile-*
lunicus exhibit different vertical distributions in the same mesocosm. J.
Shellfish Res. 15:251-257.

Marsh, A.G., J.D. Gauthier & G.R. Vasta. 1995. A semiquantitative PCR
assay for assessing *Perkinsus marinus* infections in the eastern oyster,
Crassostrea virginica. J. Parasitology 81:577-583

Mitton, J. B. 1994. Molecular approaches to population biology. Annu.
Rev. Ecol. Syst. 25:45-69.

Moore, J.D., R.A. Elston, A.S. Drum & M.T. Wilkinson. 1991. Alternate
pathogenesis of systemic neoplasia in the bivalve mollusc *Mytilus*. J.
Invert. Pathol. 58:231-243.

Morse, M. P. & J. D. Zardus. 1997. Bivalvia. pp. 7-18. In: F.W. Harrison
(ed.). Microscopic Anatomy of Invertebrates, Vol. 6A. Mollusca II.
Wiley-Liss. New York.

Oliver, L.M. & W.S. Fisher. 1995. Comparative form and function of
oyster *Crassostrea virginica* hemocytes from Chesapeake Bay (Vir-
ginia) and Apalachicola Bay (Florida). Dis. Aquat. Org. 22:217-225.

Rawson, P. D., K. L. Joyner, K. Meetze & T. J. Hilbish. 1996. Evidence for
intra-genic recombination within a novel genetic marker that distin-
guishes mussels in the *Mytilus edulis* species complex. Heredity 11:
599-607.

Sarver, S. K. & D. W. Foltz. 1993. Genetic population structure of a species' complex of blue mussels (*Mytilus* spp.). *Mar. Biol.* 117:105-112.

Suchanek, T. H., J. B. Geller, B. R. Kreiser & J. B. Minon. 1997. Zoogeographic distributions of the sibling species *Mytilus galloprovincialis* and *M. trossulus* (Bivalvia: Mytilidae) and their hybrids in the North Pacific. *Biol. Bull.* 193:187-194.

Taherlet, P., L. P. Waits & G. Luikart. 1999. Noninvasive genetic sampling: look before you leap. *Tree* 14:323-327.

Toro, J.E. 1998. Molecular identification of four species of mussels from Southern Chile by PCR-based nuclear markers: the potential use in studies involving planktonic surveys. *J. Shellfish Res.* 17:1203-1205.