

Nomenclatural and taxonomic changes, new distribution and biological records for jewel beetles (Coleoptera: Buprestidae)

C. L. Bellamy

Plant Pest Diagnostic Lab, California Department of Food & Agriculture,
3294 Meadowview Road, Sacramento, California, 95832, U.S.A.
email: cbellamy@cdfa.ca.gov

Abstract: Replacement names, nomenclatural, distributional and biological notes are presented for 21 species of Buprestidae (Coleoptera). *Agrilodes strandi* ssp. *meranus* Obenberger, 1942, and *Polybothris (Amphisbeta) vitalisi* var. *stygia*, Obenberger, 1942, are proposed to allow the subspecies or variety names define the species, respectively, with *strandii* and *vitalisi* remaining in synonymy as nomina nuda. *Acmaeodera ruficaudis macfadyeni* is proposed as a new replacement name for *Acmaeodera ruficaudis pinguis* Holm, 1985, preoccupied by *A. pinguis* Fairmaire, 1902; *Buprestis aenescens* Wiedemann, 1823 is synonymized under *Buprestis albomarginata* Herbst, 1801; *Buprestis planus* Fabricius, 1798 is transferred to *Dismorpha* Gistel, 1848; *Damarsila conturbata* Thomson, 1879 is removed from synonymy under *Buprestis amaurotica* Klug, 1855 and reinstated as a valid species; *Damarsila obsti* Cobos, 1957 is rejected as an unnecessary replacement name for *Psiloptera ornata* Obst, 1903; *Dicerca reticulatoides* is proposed as a new replacement name for *D. reticulata* Assmann, 1870, a fossil taxon, preoccupied by *Buprestis reticulata* Fabricius, 1794, a junior subjective synonym of *Dicerca aenea* (Linné, 1758); *Kamosia luciae* Obenberger, 1935, is synonymized under *Kamosiella dermestoides* (Thomson, 1878); *Melobasis novaeguinae* is proposed as a new replacement name for *Melobasis papuana* Obenberger, 1938, preoccupied by *M. (Briseis) papuana* Obenberger, 1924; *Sjoestedtius divinus* Obenberger, 1935, is transferred to *Malawiella* Bellamy, 1990, and *Sjoestedtius atakorensis* Descarpentries, 1952 is proposed as its new subjective synonym. *Buprestis albomarginata* Herbst, *Damarsila contubata* Thomson, and *Psiloptera ornata* Obst are transferred to *Lampetis* Dejean, 1833. Biological or distributional notes are given for species of *Agrilaxia* Kerremans, 1903, *Agrilus* Curtis, 1825, *Conognatha* Eschscholtz, 1829, *Euplectalecia* Obenberger, 1924, *Halecia* Laporte and Gory, 1837, *Ovalisia* Kerremans, 1900, *Sambomorpha* Obenberger, 1924, and *Spectralia* Casey, 1909.

Introduction

With several large catalogue projects under way (e.g. world buprestids, Bellamy; North American (north of Mexico) buprestids, Nelson, both *in prep.*), the need to extend distribution, add biological associations and propose new synonyms and replacement names for numerous species continues. This paper continues in the style of data presented earlier by Bellamy (1998, 1999).

Materials and Methods

The species discussed below are presented in alphabetical sequence and the order should not imply any classification detail nor opinion.

The annotations used in the text are: (h) for handwritten label data; (p) for printed label data; the forward slash '/' to separate data from consecutive labels and subsequently added data are added in square brackets '[]' and the following collection codens:

CASC	California Academy of Sciences, San Francisco, California, U.S.A.
CLBC	C. L. Bellamy collection, Sacramento, California, U.S.A.
CMNC	Canadian Museum of Nature, Ottawa, Canada
HNHM	Hungarian Natural History Museum, Budapest
INBC	Instituto Nacional de Biodiversidad, Santo Domingo de Heredia, Costa Rica
LACM	Natural History Museum of Los Angeles County, California, U.S.A.
MNHN	Museum National d'histoire Naturelle, Paris
NMPC	National Museum, Prague, Czech Republic
RLWE	R. L. Westcott collection, Salem, Oregon, U.S.A.
UCDC	R. M. Bohart Museum of Entomology, University of California, Davis, California, U.S.A.
ZMUC	Zoological Museum, University of Copenhagen, Denmark

Nomenclatural and Taxonomic Changes

Acmaeodera ruficaudis macfadyeni, new replacement name

Acmaeodera ruficaudis pinguis Holm, 1985: 156
(name preoccupied).

Remarks. The subspecies epithet *Acmaeodera ruficaudis pinguis* Holm, 1985, is preoccupied by *Acmaeodera pinguis* Fairmaire, 1902, now placed in *Sponsor* Gory and Laporte, 1839, under the replacement name of *S. cobosi* Bellamy, 1998, itself preoccupied in that combination by *S. pinguis* Guérin-Méneville 1840: 357. The replacement name is chosen to honor Duncan MacFadyen, formerly of the Transvaal Museum, for his friendship, collecting prowess and many favors during the time we worked together.

Agriloides meranus Obenberger, 1942

Remarks. Obenberger (1942a) described a new subspecies of *Agriloides strandi* ssp. *meranus* from Ecuador, but as later noted by Cobos (1959), *A. strandi* itself was never described. There is no labelled type specimen of the nominate subspecies in the Obenberger collection (NMPC), so I propose the type of *A. meranus* should define the species, with the name *strandi* remaining in synonymy as a nomen nudum.

Dismorpha plana (Fabricius, 1798), new combination

Remarks. This species' name was noted as preoccupied (by *B. plana* Olivier, 1790, now *Latipalpis plana*), replaced by the epithet *planula* and transferred to *Stenogaster* Solier, 1833, by Saunders (1871). Obenberger (1934) transferred this species, with a question mark, to *Mychommatus* Murray, 1868. Théry (1947) noted the similarity between two species of *Mychommatus* and *S. planula*. The generic name *Dismorpha* was proposed by Gistel (1848), since *Stenogaster* was preoccupied by the hymenopteran *Stenogaster* Guérin-Méneville 1831, and this was first noted by Hespenheide (1979). Neither Hespenheide (1979) nor Cobos (1990), in his revision of *Dismorpha*, listed *S. planula*. In Cobos' revision, *Dismorpha* was presumed to have a Neotropical distribution. I have examined the type of *B. plana* F. (ZMUC) and note that it agrees well, in general facies and character states, to

species placed in *Dismorpha*. It comes close to specimens in my possession identified as *D. irrorata* (Gory and Laporte, 1839) from Brazil. Since *B. plana* was originally described from Senegal, several conclusions are possible, including: 1) the original label data was transposed and 'Senegal' is an error; or 2) the range of *Dismorpha* actually includes West Africa. Since I am not aware of additional specimens of *B. plana* or any specimens of *Dismorpha* from West Africa, it seems more certain that this specimen must have been mislabelled or otherwise confused. Without the opportunity now to compare the Fabrician type to the types of other *Dismorpha* spp., I will simply note the change of combination.

Dicerca reticulatoides, new replacement name

Dicerca reticulata Assmann, 1870: 60 (name preoccupied).

Remarks. The fossil taxon *Dicerca reticulata* Assmann (1870) is preoccupied by *Buprestis reticulata* Fabricius, 1794, currently a junior subjective synonym of *Dicerca aenea* (Linné, 1758) and this new name is proposed to replace it.

Kamosiella dermestoides (Thomson, 1878)

Meliboeus dermestoides Thomson, 1878: 81.
Amorphosoma originaria Péringuey, 1908: 303.
Kamosia dermestoides: Théry, 1926: 59; Obenberger 1934: 877.
Kamosia originaria: Obenberger 1934: 877; Bellamy 1988: 187.
Kamosiella dermestoides: Bellamy 1988: 187.
Kamosia luciae Obenberger, 1935: 53. **new synonymy**

Specimens examined. The single type specimen (NMPC 23045) is labelled: "St. Lucia Lake, H W B-M Oct. 1927", although the description lists "Transvaal (sic): Sta Lucia".

Remarks. This South African species has been described three times. At the time Obenberger (1935) described *Kamosia luciae*, he related it to *K. originaria*.

Lampetis albomarginata (Herbst, 1801), new combination

Buprestis albomarginata Herbst, 1801: 132.

Buprestis aenescens Wiedemann, 1823: 100. new synonymy

Type specimen: sex undetermined (ZMUC), Mus. Westerm. (p)/Type (red, p)/Cap. b. sp., Aenescens Wiedem. (h).

Remarks. Following the original description, *Buprestis aenescens* Wiedemann has been an anomaly since it has only been listed in one subsequent work, that by Saunders (1871: 137) under "Species, the genera of which are doubtful". I have not found it listed in the catalogues of Gemminger and Harold (1869), Kerremans (1903) nor Obenberger (1926, 1930) under *Buprestis* nor under any other combination. Since it is conspecific with the widespread African species *B. albomarginata* Herbst, 1801, the new synonymy is proposed. The placement of these taxa in the genus *Lampetis* Dejean, 1833, follows the proposals of Kurosawa (1993).

***Lampetis conturbata* Thomson, 1879,
name resurrected and new combination**

Damarsila conturbata Thomson, 1879: 171.

Psiloptera (Damarsila) conturbata: Théry, 1955: 392; Descarpentries, 1970: 199.

Damarsila amaurotica (Klug, 1855) var. *conturbata*: Obenberger, 1926: 174; da Veiga-Ferreira, 1959: 445.

Remarks. At time of description, Thomson (1879) remarked that his new species was close to *Buprestis amaurotica* Klug, 1855. All subsequent authors regarded it as distinct until Obenberger (1926) listed it as a variety of *B. amaurotica* and he was followed in this by da Veiga-Ferreira (1959). However, both Théry (1955) and Descarpentries (1970) considered it a valid species. It is transferred to *Lampetis* following the concepts of Kurosawa (1993). This species is common in southern Africa and is collected from the foliage of *Terminalia sericea* Burch. ex DC (Combretaceae).

***Lampetis ornata* (Obst, 1903),
new combination**

Psiloptera ornata Obst, 1903: 144.

Damarsila obsti Cobos, 1957: 192 (superfluous replacement name for *P. ornata* Obst). new synonymy

Remarks. Cobos (1957) proposed this combination as a replacement name for *Psiloptera ornata* Obst (1903), erroneously concluding that the taxon currently known as *Chalcopoecila ornata* (Gory, 1840) had been originally described in the genus *Psiloptera* Dejean, 1833, when the original combination was in *Buprestis* L., 1758. The transfer to *Lampetis* follows Kurosawa (1993) as with the preceding two species.

***Malawiella divina* Obenberger, 1935,
new combination**

Sjoestedtius divinus Obenberger, 1935: 53.

Sjoestedtius atakorensis Descarpentries, 1952: 1155. new synonymy

Specimens examined. Holotype of *divinus* (NMPC 24054): Kameroun; holotype, paratype of *atakorensis* (MNHN): [BENIN] Dahomey, Koussokoingou Atakore, 600-700m; 6 ex. (HNHM, CLBC): Ghana, Northern Region, Nyankpala 200m, N09.25 W01.00, 5.viii.1965, S. Endrödy-Younga.

Remarks. The types of these two taxa were compared directly. The holotype and paratype of *S. atakorensis* are slightly smaller and more golden green but otherwise they are identical to the holotype of *S. divinus*. The collective locality data indicates that this species is distributed, at least, from Ghana to Cameroun. This species is transferred to *Malawiella* Bellamy, 1990, as it much better agrees with character states from the original generic diagnosis and in the key to African agrilid genera by Bellamy (1990).

***Melobasis novaeguineae*,
new replacement name**

Melobasis papuana Obenberger, 1938: 80 (name preoccupied by *M. papuana* (Obenberger, 1924)).

Remarks. The Australasian genus *Melobasis* Laporte and Gory, 1837, originally a subgenus of *Buprestis*, is currently defined to contain four subgenera (sensu Obenberger 1930:427): s. str., *Briseis* Saunders, 1871, *Diceropygus* Deyrolle, 1864 and *Paramelobasis* Théry, 1923. Since this subgeneric definition has become accepted, the combination *M. papuana* Obenberger, 1938 is preoccupied by *Briseis papuana* Obenberger, 1924 and a new name is proposed.

Polybothris (Amphisbeta) stygia
Obenberger, 1942

Remarks. In a similar situation to *Agriloides meranus* (see above), in another work of the same year, Obenberger (1942b) described a new variety of *Polybothris (Amphisbeta) vitalisi*, var. *stygia*. According to Dr. S. Bílý, NMPC, there is no record that *P. vitalisi* was ever described and no types are present that would validate this name. The description of var. *stygia* is two lines in Latin, but would be sufficient to define the name and would require *vitalisi* be left in synonymy as a nomen nudum.

**Biological, Distributional
and Predation Notes**

***Agrilaxia hespenheidei* (Bílý, 1984)**

This species is known from the Huachuca Mountains of southeastern Arizona (Bílý, 1984) and from Chiapas, Mexico (Nelson, 1987), but nothing has been recorded about its biology. One specimen was collected at the type locality, Copper Canyon, 11.viii.2001, J. Rifkind, on flowers of thistle, *Cirsium* sp. (**new adult host record**).

***Agrius restrictus* Waterhouse, 1889**

This beautifully dichromatic species is known from Chihuahua, Mexico (Fisher, 1928) and the Huachuca Mountains, Arizona (Nelson, 1965) from foliage of *Rhus choriophylla* Woot. & Standl. (Anacardiaceae). Additional data are: Arizona, Santa Cruz Co., Madera Canyon, Santa Rita Mountains, 30.vii.1989, A. V. Evans, W. B. Warner, one specimen at rest on leaf of unidentified plant (CLBC); Sycamore Canyon, N31° 25' W111° 11', 12.viii.2001, C. L. Bellamy, one male specimen beaten from *Quercus emoryi* Torr. (Fagaceae) (CLBC) (**new adult host record**).

***Conognatha (Pithiscus) vulnerata* (Perty,
1830)**

Two specimens of this Brazilian species were collected recently with the following data: Brazil: Minas Gerais: Municipio of Santana do Riacho, Serra do Cipó, at km 118 along the road from Lagos Santa to Conceição do Mato Dentro, Campo rupestre and Camp graminoso at 1227m./ 19° 18.802'S 43°33.479'W; feeding on anthers of *Lavoisiera macrocarpa* Naud. (Melastomataceae); 24 October 2001,

F. Almeda collector (CASC) (**new adult host record**).

***Euplectacea beltii* (Saunders, 1874)**

A single specimen purported to be this species, previously known only from Nicaragua, was erroneously recorded from Panamá by Bellamy and Westcott (1995). I have recently re-examined the types, or photographs of types, of all *Euplectalecia* spp. from Central America and must retract the record of *E. beltii* from Panamá. This specimen is actually *E. sordidenotata* (Obenberger, 1924), discussed below.

***Euplectalecia sordidenotata* (Obenberger,
1924)**

This beautiful species was described from Costa Rica and additional specimens (CLBC) are from: Panamá: Panamá Prov., Cerro Campana, 850m, 8°40'N, 79°56'W, 27.viii.1972, Stockwell (1 ex.); 7.5-13 km N El Llano, 13/14.v.1994, F. T. Hovore, on *Cecropia* leaves (2 ex.); 12 km N El Llano, 4.vi.1986, E. Giesbert (1 ex.); 8-10 km N El Llano, 26.iv.-4.v.1992, E. Giesbert (1 ex) (**new country and adult host records**). The last specimen is greenish, rather than the more typical red-cupreous of the other specimens and the type (NMPC), and was erroneously recorded earlier as *E. beltii* (Saunders, 1874) (see above).

***Euplectalecia suffusa* (Waterhouse, 1889)**

This species was originally described from Chiriquí in Panamá, and a number of specimens have been examined from various localities in that country, including: Canal Zone, Barro Colorado Is., 3 km SSW Balboa, Farfan Beach, 7km SW Gatun Lock; Panamá Prov., Cerro Campana (CLBC, RLWE). One additional specimen (RLWE), compared to the type (BMNH), extends the distribution significantly north: México, Oaxaca, 8 mi. SW Tuxtla, 200', 26.vi.1983, R. Anderson (**new country record**).

***Halecia chrysodemooides* Saunders, 1874**

This species is recorded from Belize, Costa Rica, Honduras, Nicaragua and Panamá and was last discussed by Bellamy and Westcott (1995). It is not surprising to find that it occurs in Guatemala: Dept. Izabal, Finca Firmez, Sierra del Caral, N15°

23.79 W88° 44.18, 935 ft., 23.v.2001, J. F. Limón, flying around foliage of *Cecropica* (CLBC, LACM) (**new country and adult host records**).

Ovalisia plasoni (Théry, 1934)

This beautiful little species is apparently known only from the unique type from "Warco, New Guinea" (see Bílý, 1993). A short series was collected as follows: PAPUA NEW GUINEA GULF: Ivimka Res. Station, Lakekumu Basin, 120m, 7° 44'S, 146° 30'E, 21.iii.2000, T.A. Sears, malaise trap (UCDC, NMPC, CLBC, RLWE).

Sambomorpha chiapas Bellamy, 1997

This species was described from a single specimen from Chiapas, Mexico (Bellamy, 1997). Additional specimens are from: GUATEMALA: Baja Verapaz, 14.5 kn N, Salama 1620 m, 23.v.1991, R. Anderson, dry oak/pine woodland, 91-13; same data except 16 mi N. Salama, 1550 m, 1.vi.1991, oak pastureland, 91-40 (**new country record**) (CMNC, CLBC).

Spectralia purpurascens (Schaeffer, 1905)

On June 19, 2001, approximately 2 mi W Ocotillo, along Interstate 8 in western Imperial County, California, U.S.A., while searching the foliage of *Justicia californica* (Benth.) D. Gibson (Acanthaceae) for this uncommon beetle, a male robber fly, *Saropogon mohawki* Wilcox, was observed with prey in its grasp. Since the prey appeared to be a buprestid, both were collected and I was surprised to find that the fly had collected the beetle I had not been able to locate on the host plant (**new predator record**).

Spectralia sulcifera (Laporte and Gory, 1837)

This species was originally described from Cayenne with subsequent records only for Para in Brazil. Material sent for identification included a number of specimens (INBC, CLBC): Costa Rica: Guanacaste Prov., Estacion Maritza, 600m, W side Volcan Orosi, malaise trap, 1988; same data except, R. Vargas, 27 feb. - 10 mar, 1992; 6 km W Santa Elena de Monte Verde, 4.vii.1983, W. Colby (CLBC) (**new country record**).

Acknowledgments

I thank Svatopluk Bílý, Narodni Museum, Prague, Czech Republic for his continuing assistance over the years and for his and his wife Vlasta's hospitality during my visits to Prague; J. J. Menier, MNHN, Paris, for his warm greeting and hospitality during visits; Ole Martin, Zoology Museum, University of Copenhagen, Denmark for the loan of the Fabrician and Weidemann types; Eric Fisher, this address, for the identification of the robber fly; and Bob Anderson, Canadian Museum of Nature, Ottawa; Roberta Brett, California Academy of Sciences, San Francisco; and Rick Westcott, Oregon Department of Agriculture, Salem, for forwarding specimens or records that have been included in this paper; J. M. Davidson, J. F. Limón, and J. R. Rifkind, for specimens or collaborating on collecting trips. S. Bílý and M. G. Volkovitsh, Zoological Institute, Russian Academy of Sciences, St. Petersburg are thanked for reviewing the manuscript.

Literature Cited

- Assmann, A. 1870. Paleontologie. II. Beitrag. Fossile Insekten aus der tertiären (oligocenischen) Braunkohle von Naumburg am Bober. Zeitschrift für Entomologie, Breslau (N.F.) 1: 1-62.
- Bellamy, C. L. 1988. New genera, species and synonymy in African Coroebini (Coleoptera, Buprestidae, Agrilinae). Coleopterists Bulletin 42(2): 173-192.
- Bellamy, C. L. 1990. Studies in the African Agrilinae, Agrilini II (Coleoptera: Buprestidae). Revue de Zoologique Africaine 104: 573-579.
- Bellamy, C. L. 1997. A revision of the Neotropical coroebine genus *Sambomorpha* Obenberger 1924 (Coleoptera: Buprestidae Agrilinae). Tropical Zoology 10: 143-156.
- Bellamy, C. L. 1998. Further replacement names in the family Buprestidae (Coleoptera). African Entomology 6(1): 91-99.
- Bellamy, C. L. 1999. Nomenclatural notes in Buprestidae (Coleoptera). Folia Heyrovskyana 7(1): 1-11.
- Bellamy, C. L. 2001. Proposed conservation of 31 species-group names originally published as junior primary homonyms in *Buprestis* Linnaeus, 1758 (Insecta, Coleoptera). Bulletin of Zoological Nomenclature 58(1): 24-31.
- Bellamy, C. L. and R. L. Westcott. 1995. New records for the *Halecia* Laporte and Gory ge-

- nus-group from Central America (Coleoptera: Buprestidae). Coleopterists Bulletin 49(2): 195-199.
- Bilý, S.** 1984. Taxonomic notes on *Anthaxia*, with descriptions of new taxa (Coleoptera, Buprestidae). Acta Entomologica Bohemoslovaca 81: 212-222.
- Bilý, S.** 1993. A revision of the genera *Philanthaxia* Deyrolle and *Pagdeniella* Théry (Coleoptera, Buprestidae). Folia Heyrovskyana 1(9-10): 84-115.
- Bilý, S.** 1997. Supplement to the revisions of *Philanthaxia* and *Pagdeniella* and notes on the related genera (Coleoptera: Buprestidae). Folia Heyrovskyana 5(1): 1-13.
- Cobos, A.** 1957. Séptima nota sobre Bupréstidos neotropicales. Sobre los Chalcophorini Kerremans, de América meridional, con descripción de un nuevo género y dos nuevas especies. Archivos de Instituto de Aclimatación 6: 191-198.
- Cobos, A.** 1959. Octava nota sobre Bupréstidos neotropicales. Rectificaciones y descripciones diversas (Coleoptera, Buprestidae). Bulletin de l'Institut Royal Sciences Naturelles Belgique 35(2): 1-47.
- Cobos, A.** 1990 Revisión del género *Dismorpha* Gistel (Coleoptera, Buprestidae). Revista Brasileira de Entomologia 34(3): 539-559.
- da Veiga-Ferreira, G.** 1959. Revisão das espécies Sul-Africanas da subtribo Psilopterina. Revista de Entomologia de Moçambique 2(2): 423-546.
- Descarpentries, A.** 1952. Mission A. Villiers au Togo et au Dahomey (1950). XIX. Coléoptères Buprestides. Bulletin de l'Institut Fondamental d'Afrique Noire 14(4): 1152-1158.
- Descarpentries, A.** 1970. Results of the Swedish Museum Lund expedition to Southern Africa 1950-1951. Chapter 6. Coleoptera: Buprestidae. In: Hanström, B., P. Brinck and G. Rudebeck (Eds.), South African Animal Life, Lund, Sweden, 14(6):190-215.
- Fabricius, I. C.** 1794. Entomologia systematica. Appendix. Volume 4, Hafniae, 472 pp.
- Fabricius, I. C.** 1798. Supplementum entomologiae systematicae, Hafniae 572 pp.
- Fairmaire, L. M. H.** 1902. Buprestides nouveaux recueillis par M. Ch. Alluaud dans le sud de Madagascar [Col.]. Bulletin de la Société Entomologique de France 1902: 77-81.
- Fisher, W. S.** 1928. A revision of the North American species of buprestid beetles belonging to the genus *Agrius*. United States National Museum Bulletin 145: 1-347.
- Gemminger, M. and E. von Harold.** 1869. Catalogus coleopterorum hucusque descriptorum synonymicus et systematicus. Volume 5, pp. 1347-1608, Monachii, London. (Buprestidae, Trixagidae, Monommidae, Eucnemidae, Elateridae, Cebrionidae).
- Gistel, J. N. F. X.** 1848. Naturgeschichte des Thierreichs für höhere Schulen. 216 pp., 32 plates. Stuttgart.
- Gory, H. L.** 1840. Histoire naturelle et iconographie des insectes Coléoptères. Supplément aux Buprestides. P. Duménil, Paris. Volume 4, livraisons 36-42, genus *Buprestis* (pp. 81-106; 112-125).
- Gory, H. L. and F. L. Laporte de Castelnau.** 1839. Histoire naturelle et iconographie des insectes Coléoptères. Monographie des buprestides. P. Duménil, Paris. Volume 2, Volume 2, livraisons 25-35 (genera paged separately).
- Guérin-Méneville, F. E.** 1840. Note monographique sur le genre *Sponsor* de M. Gory, démembré du grand genre des Buprestides. Revue Zoologique 1840: 356-358.
- Herbst, J. F. W.** 1801. Natursystem aller bekannten in- und auslandischen Insecten, als eine Fortsetzung der Büssonschen Naturgeschichte. Der Käfer neunter Theil, Volume 9, pp. 1-344, illustrated.
- Hespenheide, H. A.** 1979. Nomenclatural notes on the Agrilinae (Buprestidae). IV. The Coleopterists Bulletin 33(1): 105-120.
- Holm, E.** 1985. Supplementary notes and descriptions of Ethiopian Acmaeoderinae, Julodinae and Polycestinae (Coleoptera: Buprestidae). Journal of the Entomological Society of Southern Africa 48(1): 135-161.
- ICZN.** 2002. Opinion 2008. 30 species-group names originally published as junior primary homonyms in *Buprestis* Linnaeus, 1758 (Insecta, Coleoptera): conserved. Bulletin of Zoological Nomenclature 59(3): 211-216.
- Kerremans, C.** 1903. Coleoptera Serricornia, Fam. Buprestidae. In: P. Wytsman (ed.). *Genera Insectorum*, Fasc. 12b; 12c; 12d. Verteneuil and Desmet, Bruxelles, pp. 49-338.
- Kurosawa, Y.** 1993. Reorganization of the genus *Psiloptera* (Coleoptera, Buprestidae). Japanese Journal of Entomology 61(3): 577-583.
- Laporte de Castelnau, F. L. and H. L. Gory.** 1837. Histoire naturelle de iconographie des insectes coléoptères. Monographie des bupres-

- tides. P. Duménil, Paris. Volume 1, livraisons 12-16, genera: *Buprestis* (pp. 65-160).
- Nelson, G. H.** 1965. Notes on the Buprestidae: part IV, with a new synonym in *Chrysobothris*. Bulletin of the Brooklyn Entomological Society 59/60: 37-41.
- Nelson, G. H.** 1987. Additional notes on the biology and distribution of Buprestidae in North America, II. Coleopterists Bulletin 41(1): 57-65.
- Obenberger, J.** 1924. Kritische studien über die Buprestiden (Col.). Archiv für Naturgeschichte 90 (A) 3: 1-171.
- Obenberger, J.** 1926. Buprestidae 1. In: W. Junk, S. Schlenkling (eds.). Coleopterorum Catalogus, W. Junk, Berlin, Part 84: 1-212.
- Obenberger, J.** 1930. Buprestidae 2. In: W. Junk, S. Schenkling (eds.). Coleopterorum Catalogus, W. Junk, Berlin, Part 111: 213-568.
- Obenberger, J.** 1934. Buprestidae 4. In: W. Junk, S. Schenkling (eds.). Coleopterorum Catalogus, W. Junk, 's-Gravenhage, Part 143: 782-934.
- Obenberger, J.** 1935. Agrilinorum diagnoses praeliminares (Col. Bupr.). Acta Societatis Entomologicae Cechosloveniae 32: 49-56.
- Obenberger, J.** 1938. Nový krasec z N. Guiney (Col. Bupr.). [New Buprestid beetle from New Guinea]. Acta Societatis Entomologicae Cechosloveniae 35: 80.
- Obenberger, J.** 1942a. *Agriloides strandi* ssp. *meranus* n. (Col. Bupr.). Acta Societatis Entomologicae Bohemiae 39: 120.
- Obenberger, J.** 1942b. De novis generis *Polybothris* speciebus et formis. (Col. Bupr.). Nové formy rodu *Polybothris* (Col. Bupr.). Acta Entomologica Musaei Nationalis Pragae 20: 30-49.
- Obst, P.** 1903. Diagnosen neuer afrikanischer Coleopteren, meistens Buprestiden. Annales de la Société Entomologique de Belgique 47: 141-146.
- Péringuey, L.** 1908. Seventh contribution to the South African Coleopterous fauna. Annals of the South African Museum 5: 271-344.
- Saunders, E.** 1871. Catalogus Buprestidarum Synonymicus et Systematicus. J. Janson, London, 171 pp.
- Théry, A.** 1926. Recherches synonymiques sur les Buprestides et descriptions d'espèces nouvelles. Bulletin et Annales de la Société Entomologique de Belgique 66(1-2): 33-74.
- Théry, A.** 1947. Buprestides d'Angola. Arquivos do Museo Boçage 18: 1-96.
- Théry, A.** 1955. Buprestides des provinces de Rhodésie du Sud, du Transvaal, du Natal et du Cap, au Transvaal Museum. Annals of the Transvaal Museum 22: 391-400.
- Thomson, J.** 1878. Typi Buprestidarum Musaei Thomsoniani. E. Deyrolle, Paris, 103 pp.
- Thomson, J.** 1879. Revue du groupe des Psiloptérites; Description d'espèces nouvelles de Psiloptérites. Revue et Magasin de Zoologie (3) 7(42): 161-177.
- Waterhouse, C. O.** 1889. Biologia Centrali-Americanana, Insecta, Coleoptera, Buprestidae, Volume 3, part 1, pp. 49-166, plates iv-viii and Supplement, pp. 167-193.
- Wiedemann, C. R. W.** 1823. Zweihundert neue Käfer von Java, Bengalen und dem Vorgebirge der guten Hoffnung. Zoologisches Magazin 2(1): 1-135.