


Staff training in PBS for adults with ID and challenging behaviour

Supplementary Fig. 1 ABC-C_t score over 12 months.

Supplementary Table 1 Sensitivity and exploratory multivariate analyses

| <i>Model</i> | <i>Difference</i> | <i>95% CI</i> | <i>P-value</i> |
|--|-------------------|---------------|----------------|
| Sensitivity analyses | | | |
| Primary model | -2.14 | (-8.79, 4.51) | 0.528 |
| Area deprivation | -2.39 | (-9.19, 4.41) | 0.491 |
| Completed by family/paid carer | -1.21 | (-8.20, 5.79) | 0.735 |
| Missing data (BOCF) | -1.83 | (-8.42, 4.76) | 0.586 |
| Heteroscedastic model | -2.35 | (-9.24, 4.55) | 0.505 |
| Imbalance in baseline characteristics | -0.81 | (-7.95, 6.32) | 0.824 |
| % of participants who had at least one intervention component (e.g. plan, observations, goodness of fit) | 1.41 | (-15.5, 18.3) | 0.870 |
| Multivariate analysis (ABC-C subdomains) | | | |
| Irritability, Agitation, Crying | -0.041 | (-0.22, 0.14) | |
| Lethargy, Social Withdrawal | -0.016 | (-0.22, 0.19) | |
| Stereotypic Behaviour | -0.050 | (-0.25, 0.14) | |
| Hyperactivity, Non-compliance | -0.049 | (-0.23, 0.13) | |

Supplementary Table 2 Secondary outcomes over 12 months

| | Descriptive: N (%) | | | Analysis over 12 months | | |
|------------------------------------|---------------------|---------------------|---------------------|-------------------------|---------------------------|---------------|
| | Baseline | 6 Months | 12 Months | N _{SU} | Odds Ratio/ Difference | 95% CI |
| Service users | | | | | | |
| Mini-PASADD | N (%) | N (%) | N (%) | | | |
| Common mental disorder | | | | | | |
| Control | 61 (46) | 43 (37) | 54 (44) | 230 | 1.07 | (0.61, 1.87) |
| Intervention | 56 (52) | 45 (46) | 42 (42) | | | |
| Severe mental illness (SMI) | | | | | | |
| Control | 27 (20) | 13 (11) | 21 (17) | 229 | 1.24 | (0.32, 4.81) |
| Intervention | 20 (19) | 17 (18) | 15 (15) | | | |
| Autistic spectrum | | | | | | |
| Control | 31 (23) | 31 (27) | 40 (33) | 230 | 0.70 | (0.26, 1.88) |
| Intervention | 19 (18) | 24 (24) | 22 (22) | | | |
| GCPLA | Median (IQR) | Median (IQR) | Median (IQR) | | | |
| Range | | | | | | |
| Control | 17 (12, 22) | 16.5 (13, 21) | 17 (13, 21) | 232 | 0.587 | (-0.57, 1.74) |
| Intervention | 19 (13, 23.5) | 19 (14, 23) | 17 (13.5, 22) | | | |
| Busy | | | | | | |
| Control | 10 (7, 13) | 11 (7, 13) | 11 (8, 13) | 232 | 0.377 | (-0.59, 1.34) |
| Intervention | 11 (8, 15) | 11 (8, 14) | 12 (8, 14) | | | |
| Family carers | | | | | | |
| Uplift | | | | | | |
| Control | 15 (13, 17) | 15 (12, 17) | 15 (13, 17) | | | |
| Intervention | 14 (13, 16) | 15 (14, 17) | 15 (13, 16) | | | |
| Burden | | | | | | |
| Control | 33 (28, 39) | 31 (25, 36) | 30 (25, 39) | | | |
| Intervention | 28 (26, 31) | 29 (25.5, 32.5) | 30 (28, 32.5) | | | |
| GHQ score | | | | | | |
| Control | 4 (1, 8) | 4 (2, 7) | 3 (1, 6) | | | |
| Intervention | 3 (0, 4) | 2.5 (1, 6.5) | 2 (0, 3) | | | |
| Paid carers | | | | | | |
| CDS* | | | | | | |
| Control | 24 (15, 37) | | | | | |
| Intervention | 21 (13, 31) | | | | | |

*Respondents changed over the 12 months, consequently the trends are not provided