

INSECTA MUNDI

A Journal of World Insect Systematics

0300

‘Retournement’ of the aedeagus in Curculionidae
(Coleoptera, Curculionoidea)

Pierre Jolivet
67 Boulevard Soult, 75012
PARIS, France

K. K. Verma
HIG 1/327
Housing Board Colony
Borsi, DURG (C.G.) – 491001, India

Rashmi Saxena
Department of Zoology
Govt. College, Kherwada
Resi.: ‘SOPAN’, 71, Madhav
Vihar, Shobhagpura, UDAIPUR – 313001, India

Date of Issue: May 10, 2013

Pierre Jolivet, K. K. Verma, and Rashmi Saxena
'Retournement' of the aedeagus in Curculionidae (Coleoptera, Curculionoidea)
Insecta Mundi 0300: 1-5

ZooBank Registered: urn:lsid:zoobank.org:pub:636D2AA9-8BA1-48F4-8D93-C7999711DE68

Published in 2013 by

Center for Systematic Entomology, Inc.
P. O. Box 141874
Gainesville, FL 32614-1874 USA
<http://www.centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology. Manuscript preparation guidelines are available at the CSE website.

Managing editor: Paul E. Skelley, e-mail: insectamundi@gmail.com

Production editor: Michael C. Thomas, Brian Armitage, Ian Stocks

Editorial board: J. H. Frank, M. J. Paulsen

Subject editors: G.B. Edwards, J. Eger, A. Rasmussen, F. Shockley, G. Steck, Ian Stocks, A. Van Pelt, J. Zaspel

Spanish editors: Julieta Brambila, Angélico Asenjo

Printed copies (ISSN 0749-6737) annually deposited in libraries:

CSIRO, Canberra, ACT, Australia
Museu de Zoologia, São Paulo, Brazil
Agriculture and Agrifood Canada, Ottawa, ON, Canada
The Natural History Museum, London, Great Britain
Muzeum i Instytut Zoologiczny PAN, Warsaw, Poland
National Taiwan University, Taipei, Taiwan
California Academy of Sciences, San Francisco, CA, USA
Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA
Field Museum of Natural History, Chicago, IL, USA
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA
Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (On-Line ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.
Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>
University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>
Goethe-Universität, Frankfurt am Main: <http://edocs.ub.uni-frankfurt.de/volltexte/2010/14363/>

Author instructions available on the Insecta Mundi page at:

<http://www.centerforsystematicentomology.org/insectamundi/>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

'Retournement' of the aedeagus in Curculionidae
(Coleoptera, Curculionoidea)

Pierre Jolivet
67 Boulevard Sout, 75012
PARIS, France
timarcha@sfr.fr

K. K. Verma
HIG 1/327
Housing Board Colony
Borsi, DURG (C.G.) – 491001, India
kkverma.sheel@gmail.com

Rashmi Saxena
Department of Zoology
Govt. College, Kherwada
Resi.: 'SOPAN', 71, Madhav
Vihar, Shobhagpura, UDAIPUR – 313001, India
karuneshsaxena@gmail.com

Abstract. Retournement or turning of the aedeagus about its longitudinal axis through about 180° during development is known in Chrysomeloidea (Coleoptera). This change in the orientation of the organ may be observed during the postembryonic development. This change produces certain morphological effects. By observing these morphological features in the imago the retournement may be inferred. Such morphological features in Curculionidae (Coleoptera) are here recorded. From this it has been inferred not only that retournement of the aedeagus is included in the ontogeny of curculionids, but also that the change of orientation of the organ occurs by the same mechanism as in Chrysomeloidea. These inferences attest the notion of a close phyletic relationship between the superfamilies Curculionoidea and Chrysomeloidea.

Introduction


'Retournement' of the aedeagus is the rotation of the developing aedeagus about its longitudinal axis through about 180° during the post-embryonic development. This change of orientation of the aedeagus is irreversible and generally clockwise when viewed from behind, though in some cases it may be counter-clockwise, and is prevalent, if not universal, among Chrysomeloidea (Verma 1994).

Like Chrysomeloidea, Curculionoidea are pseudotetramerous in their tarsi. The two superfamilies are phylogenetically close, which will be covered at some length under the section "Discussion" in this communication. This phyletic closeness suggests the possibility of aedeagal 'retournement' among Curculionidae; hence this study.

Progress of the 'retournement' during development has been observed and recorded in two chrysomelids, *Galerucella birmanica* (Jacoby, 1859) (Verma 1969), and *Aspidomorpha miliaris* (Fabricius, 1775) (Verma and Kumar 1972). Occurrence of the aedeagal rotation during development has been inferred in a number of chrysomeloid species from the observation of certain morphological changes, as seen in the adult, from the developmental turning of the aedeagus (Kumar and Verma 1971, 1980; Pawar and Verma 1977). For the present study we searched for similar morphological changes in adult curculionids.

Material and methods

Fresh specimens of a fairly large curculionid species, *Xanthochelus superciliosus* Gyllenhal, 1834 were dissected under magnification, and examined. A few dissections of a smaller species, *Myloccerus*



Figures 1-3. *Xanthochelus superciliosus* aedeagus. **1)** Aedeagal apparatus of adult in ventral view. The muscles GM1 and GM2 have been labeled following Burke (1959). **2)** Basal part of the aedeagal apparatus in lateral view, from the right side. **3)** Tracheal supply to the aedeagus base in ventral view.

fabricii Guerin, 1843 have also been done. Figures were drawn using a camera lucida, and extent of magnification in the figures was recorded with the help of a micrometer scale.

Observations

In both species of curculionids examined, a long spiculum gastrale, which arises posteriorly from the terminal abdominal sternum and extends forward beneath the aedeagus, is tilted upward and to the right side of the aedeagus in its anterior part (Fig. 1-2). Arising from its anterior tip is a unilateral muscle, the retractor of the spiculum. This muscle attaches to a tegminal ring at the base of the aedeagus. The tegminal ring is provided with a short apodeme.


The tracheal supply, reaching the base of the aedeagus, is twisted, i.e. the tracheae, from the right side, enter the muscular mass at the base of the aedeagus on the left side, and those from the left side reach the aedeagal base to enter the muscular mass on the right side (Fig. 3). The nature of the twisted tracheal supply to the aedeagus indicates a clockwise 'retournement' of the organ, when looked at from behind. The relative orientation of the right and left aedeagal tracheae, on completion of the rotation of the aedeagus in the clockwise direction, is well explained and illustrated in Kumar and Verma (1971).

The median ejaculatory duct presents a spiral coiling, when the aedeagus is in repose (Fig. 2-3). In all dissections the aedeagus appeared deversed to the right side. ('deversed' means having an inclined orientation of the aedeagus in repose.)

Discussion

The developmental phenomenon of 'retournement' of the aedeagus has been inferred from the following morphological features in the adult male curculionids:

- (i) Twisted nature of the tracheal supply for the aedeagus.
- (ii) Spiral coiling of the median ejaculatory duct.



Figures 4-6. 4) Diagrammatic cross section through a basal part of the aedeagal apparatus in a young pupa of *Galerucella birmanica* (after Verma 1969). 5) Cross section through a basal part of the developing genital tube in a pupa, within 24 hours of pupation, of *Galerucella birmanica* (after Verma 1969). 6) Spiculum 1 with parameral cap in *Aeolesthes induta*, dorsal view (after Kumar and Verma 1980).

- (iii) The asymmetrical tilting, upward and towards the right side, of the spiculum gastrale.
- (iv) The unpaired muscle arising from the anterior tip of the spiculum attaches to the tegminal ring.

From similar adult morphological features the 'retournement' has been inferred in several Chrysomeloidea (Kumar and Verma 1971, 1980; Pawar and Verma 1977).

It may be noted that there is a close similarity between the mechanism of the 'retournement' in Chrysomeloidea and that in Curculionidae. In the former the 'retournement' has been attributed to a unilateral pull on the dorsal face of the genital tube (= aedeagus + genital atrium) by the right protractor of the spiculum 1, while its counterpart of the left side undergoes degenerative changes (Fig. 4) (Verma 1969; Verma and Kumar 1972). This leads to a rotation of the developing aedeagus in the clockwise direction, as viewed from the caudal end. In some cases the right protractor degenerates, resulting in a counterclockwise 'retournement'.

In many Chrysomeloidea there are two spicula in the aedeagal apparatus, one behind the other. They have been referred to as spiculum 1 (the more anterior one), and spiculum 2 (Kumar and Verma 1980). It may be clarified here that the structure, which has been referred to as spiculum 2, is the homologue of the spiculum gastrale in curculionids.

In the chrysomelid *Galerucella birmanica* the spiculum 2 is represented by a soft diverticulum of the genital atrium, arising close in front of the last visible abdominal sternum (Verma 1969) (Fig. 4). Spiculum 1 (Kumar and Verma 1980), and simply named as spiculum in Verma (1969) and Verma and Kumar (1972) is actually a tegminal apodeme, which will become evident if the development of this structure in *Galerucella birmanica* (Verma 1969) is kept in view. In this chrysomelid the rudiment of the spiculum

appears as an apodemal invagination at the base of the aedeagal rudiment on the dorsal side (Fig. 5). The basal dorsal part of the aedeagal rudiment develops a parameral cap; hence it should be regarded as representing the tegmen. (For further development and the 'fate' of the parameral cap Verma, 1969 may be referred to). In support of this notion is the spiculum 1 in the cerambycid *Aeolesthes induta* Newman, 1842 carrying a bifid parameral cap dorsally (Fig. 6). It may be recalled that the tegmen in a curculionid includes a short tegminal apodeme.

In the curculionids the unilaterally developed muscle, arising from the anterior tip of the spiculum gastrale and attaches to the tegmen, is comparable to the asymmetrically developed right protractor of the spiculum 1 in Chrysomeloidea (Kumar and Verma 1980), as it arises from the spiculum gastrale/spiculum 2 and attaches to the spiculum 1/tegminal apodeme. If this homology is accepted, it becomes obvious that in a curculionid the muscle, which is responsible in bringing about the 'retournement', not only rotates the genital tube, but also pulls the anterior part of the spiculum gastrale sideways on the right side and upward; hence the tilt in the anterior part of the spiculum.

The morphological features, suggesting the developmental phenomenon of 'retournement' of the aedeagus, seem common, if not universal, among Curculionoidea. The asymmetrical upward and rightward tilting in an anterior part of the spiculum gastrale has been recorded also by Burke (1959) in the cotton boll weevil (*Anthonomus grandis* Boheman, 1843), by Hieke (1966) in the curculionid *Liparus Olivier*, by Francke-Grosmann (1948) in a scolytid (*Dendroctonus micans* Kugel, 1794), by Bissell (1937) in the pecan weevil (*Curculio caryae* Horn, 1873), and by Schoof (1942) in the curculionid *Conotrachelus Schönherr*. These authors also observed a strong muscle band arising from the anterior tip of the spiculum gastrale and attaching to the tegmen on the right side, in contrast to the similar left muscle being weakly developed or absent.

Our communication is not the first to report the aedeagal 'retournement' in Curculionidae. Wanat (2007) has comparatively studied male terminalia in Curculionoidea and other Coleoptera, and has inferred that basal weevil groups, Nemonychidae, Oxycoryninae, and Aglycyderinae exhibit genitalia rotation around the long body axis.

Phyletic closeness between Chrysomeloidea and Curculionoidea has been generally recognized. Farrell (1998) worked out phylogeny of Phytophaga, taking into account 212 morphological features and 115 complete DNA sequences. In Farrell's study it was inferred that Chrysomeloidea and Curculionoidea are sister groups. This opinion was expressed also by Marvaldi et al. (2002). Hunt et al. (2007) studied 1900 beetle species through phylogenetic analysis, based on sequences of three genes, and inferred that Chrysomeloidea and Curculionoidea, along with Erotylid series, Cucujid series, and Nitidulidae, appear to belong to the same clade.

The present study, pointing to 'retournement' of the aedeagus in Curculionidae taking place through essentially the same developmental mechanism, is additional evidence supporting the phyletic closeness between the two superfamilies, Chrysomeloidea and Curculionoidea. As Beutel et al. (2010) pointed out, even in the present age of phylogenomics, comparative morphology may still play a vital role in working out phyletic relations.

Acknowledgments

The authors are grateful to Dr. V. V. Ramamurthy, Principal Scientist, Division of Entomology, Indian Agricultural Research Institute, New Delhi for identifying the curculionids, studied. They are also thankful to the Museum Library of Paris for generously helping us in getting access to the required literature. We thank Ron N. Beenen, Naturalis Biodiversity Centre, Leiden, and Jean-François Voisin, Museum National d'Histoire Naturelle, Paris for reviewing the manuscript.

Literature Cited

- Beutel, R. G., F. Friedrich, T. Hornschemeyer, H. Pohl, F. Hunefeld, Beckmann, F. R. Meier, B. Misof, M. F. Whiting, and L. Vilhelmsen. 2010. Morphological and molecular evidence converge upon a robust phylogeny of the megadiverse Holometabola. *Cladistics* 26(2010): 1 – 15.
- Bissell, T. L. 1937. Structure of the reproductive system of the pecan weevil (Curculionidae). *Annals of the Entomological Society of America* 30: 242 – 249, 2 plates.

- Burke, H. R. 1959.** Morphology of the reproductive systems of the cotton boll weevil (Coleoptera, Curculionidae). *Annals of the Entomological Society of America* 52: 287-294.
- Farrell, B. D. 1998.** "Inordinate Fondness" explained: Why are there so many beetles. *Science* 281: 555 – 559.
- Francke-Grosmann, H. 1948.** Über den kopulationsapparat des Reisenbastkafers *Dendroctonus micans* Kug. *Verhandlungen der Deutschen Zoologischen Gesellschaft, Kiel* 1948: 219 -225.
- Hieke, F. 1966.** Vergleichende funktionelle Anatomie der abdominalmuskulatur einiger männlicher Coleopteren unter besonderer Berücksichtigung des Genitoanalkomplexes. *Deutsche Entomologische Zeitschrift, N.F. Heft (I – III):* 1 – 168.
- Hunt, T., J. Bergsten, Z. Levkanicova, A. Papadopoulou, O. St. John, R. Wild, P. M. Hammond, D. Ahrens, M. Balke, M. S. Caterino, J. Gomez-Zurita, I. Ribera, T. G. Barraclough, M. Bocakova, L. Bocak, and A. P. Vogler. 2007.** A comprehensive phylogeny of beetles reveals the evolutionary origins of a superradiation. *Science* 318: 1913 – 1916.
- Kumar, D., and K. K. Verma. 1971.** 'Retournement' of the aedeagus in Chrysomelidae. (Coleoptera, Phytophaga). *Journal of Natural History* 5: 635 – 642.
- Kumar, D. and K. K. Verma. 1980.** Aedeagal musculature in Phytophaga (Coleoptera). *Journal of Natural History* 14: 237 – 270.
- Marvaldi, A. E., A. S. Sequeira, C. W. O'Brien, and B. D. Farrell. 2002.** Molecular and morphological phylogenetics of weevils (Coleoptera, Curculionidae): do niche shifts accompany diversification? *Systematic Biology* 51: 761 – 785.
- Pawar, B. L., and K. K. Verma. 1977.** 'Retournement' of the aedeagus in Bruchidae (Coleoptera, Phytophaga). *Entomon* 2(2): 171 – 174.
- Schoof, H. F. 1942.** The genus *Conotrachelus* Dejean (Coleoptera, Curculionidae) in the North Central United States. *Illinois Biological Monographs* XIX(1-2): 1 – 170.
- Verma, K. K. 1969.** Functional and developmental anatomy of the reproductive organs in the male of *Galerucella birmanica* Jac. (Coleoptera, Chrysomelidae). *Annales des Sciences Naturelles (Zoologie)* 12XI(2): 139 – 234.
- Verma, K. K. 1994.** 'Retournement' of the aedeagus in Chrysomelidae (Coleoptera) p. 355-362. *In:* P.H. Jolivet, M.L. Cox, and E. Petitpierre, (eds.). *Novel aspects of the biology of Chrysomelidae*. Kluwer Academic Publishers; the Netherlands. xxiii + 582 p.
- Verma, K. K., and D. Kumar. 1972.** The aedeagus, its musculature and 'retournement' in *Aspidomorpha miliaris* (Coleoptera, Chrysomelidae). *Journal of Natural History* 6(6): 699 – 719.
- Wanat, M. 2007.** Alignment and homology of male terminalia in Curculionoidea and other Coleoptera. *Invertebrate Systematics* 21: 147 – 171.

Received December 9, 2012; Accepted March 29, 2013.

