

Original citation:

Ketterer, Christian and Mondino, Andrea (2018) *Sectional and intermediate Ricci curvature lower bounds via optimal transport*. *Advances in Mathematics*, 329 . pp. 781-818. doi:[10.1016/j.aim.2018.01.024](https://doi.org/10.1016/j.aim.2018.01.024)

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/97615>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

© 2018, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

SECTIONAL AND INTERMEDIATE RICCI CURVATURE LOWER BOUNDS VIA OPTIMAL TRANSPORT

CHRISTIAN KETTERER AND ANDREA MONDINO

ABSTRACT. The goal of the paper is to give an optimal transport characterization of sectional curvature lower (and upper) bounds for smooth n -dimensional Riemannian manifolds. More generally we characterize, via optimal transport, lower bounds on the so called p -Ricci curvature which corresponds to taking the trace of the Riemann curvature tensor on p -dimensional planes, $1 \leq p \leq n$. Such characterization roughly consists on a convexity condition of the p -Renyi entropy along L^2 -Wasserstein geodesics, where the role of reference measure is played by the p -dimensional Hausdorff measure. As application we establish a new Brunn-Minkowski type inequality involving p -dimensional submanifolds and the p -dimensional Hausdorff measure.

CONTENTS

1. Introduction	1
2. Preliminaries	5
3. Countably \mathcal{H}^p -rectifiable geodesics in Wasserstein space	7
4. Jacobi fields computations	11
5. OT characterization of sectional curvature upper bounds	15
6. OT characterization of sectional, and more generally p -Ricci, curvature lower bounds	18
References	24

1. INTRODUCTION

The interplay between Ricci curvature and optimal transport is well known and it has been a topic of tremendous interest in the last years. On the other hand it seems to be still an open problem to find the link between *sectional* curvature bounds (and more generally intermediate Ricci curvature bounds) and optimal transportation. The goal of the paper is to address such a question.

Inspired by the pioneering work on Ricci curvature lower bounds via optimal transport by Sturm and von Renesse [18], later extended to non-smooth spaces in the foundational works of Lott-Villani [8] and Sturm [16, 17], we analyze convexity properties of the p -Renyi entropy along L^2 -Wasserstein geodesics, where the role of the reference measure is played here by the p -dimensional Hausdorff measure. In a first approximation, one can think of studying the convexity of the p -Renyi entropy along an L^2 -Wasserstein geodesics made of probability measures concentrated on p -dimensional submanifolds of M .

The study of optimal transportation between measures supported on arbitrary submanifolds in an arbitrary Riemannian manifold seems to be quite a new topic in the literature. Nevertheless several authors treated remarkable particular cases and related questions:

- Gangbo-McCann [5] proved results about optimal transport between measures supported on hyper-surfaces in Euclidean space;

Key words and phrases. Sectional curvature, intermediate Ricci curvature, Optimal Transport, Brunn-Minkowski inequality.

C. Ketterer: Freiburg University, Mathematics Department, email: ckettere@math.toronto.edu.

A. Mondino: University of Warwick, Mathematics Institute, email: A.Mondino@warwick.ac.uk.

- McCann-Sosio [11] and Kitagawa-Warren [6] gave more refined results about optimal transport between two measures supported on a codimension one sphere in Euclidean space;
- Castillon [2] considered optimal transport between a measure supported on a submanifold of Euclidean space and a measure supported on a linear subspace;
- Lott [7] characterized the tangent cone (in the W_2 -metric) to a probability measure supported on a smooth submanifold of a Riemannian manifold.

In order to state the results, let us introduce some notation (for more details see Section 2). Let (M^n, g) be a smooth, complete, n -dimensional Riemannian manifold without boundary. For $p = \{1, \dots, n\}$, denote by \mathcal{H}^p the p -dimensional Hausdorff measure and consider the space $\mathcal{P}_c(M, \mathcal{H}^p)$ of probability measures with compact support which are absolutely continuous with respect to \mathcal{H}^p . Given $1 \leq p \leq p' < \infty$, the p' -Renyi entropy with respect to \mathcal{H}^p is defined as

$$S_{p'}(\cdot|\mathcal{H}^p) : \mathcal{P}_c(M, \mathcal{H}^p) \rightarrow [-\infty, 0], \quad S_{p'}(\mu|\mathcal{H}^p) = - \int \rho^{1-\frac{1}{p'}} d\mathcal{H}^p,$$

where ρ is the density of μ with respect to \mathcal{H}^p , i.e. $\mu = \rho\mathcal{H}^p$. Note that in the borderline case $p' = p = 1$, one gets

$$S_1(\mu|\mathcal{H}^1) = -\mathcal{H}^1(\text{supp}(\mu)).$$

The (relative) Shannon entropy is defined by

$$\text{Ent}(\cdot|\mathcal{H}^p) : \mathcal{P}_c(M, \mathcal{H}^p) \rightarrow [-\infty, \infty], \quad \text{Ent}(\mu|\mathcal{H}^p) = \lim_{\varepsilon \downarrow 0} \int_{\{\rho > \varepsilon\}} \rho \log \rho d\mathcal{H}^p.$$

This coincides with $\int_{\{\rho > 0\}} \rho \log \rho d\mathcal{H}^p$, provided that $\int_{\{\rho \geq 1\}} \rho \log \rho d\mathcal{H}^p < \infty$, and $\text{Ent}(\mu|\mathcal{H}^p) := \infty$ otherwise. Recall also the definition of the distortion coefficients. Given $K \in \mathbb{R}$, we set for $(t, \theta) \in [0, 1] \times \mathbb{R}^+$,

$$\sigma_{K,1}^{(t)}(\theta) := \begin{cases} \infty & \text{if } K\theta^2 \geq \pi^2, \\ \frac{\sin(t\theta\sqrt{K})}{\sin(\theta\sqrt{K})} & \text{if } 0 < K\theta^2 < \pi^2, \\ t & \text{if } K\theta^2 = 0, \\ \frac{\sinh(t\theta\sqrt{-K})}{\sinh(\theta\sqrt{-K})} & \text{if } K\theta^2 \leq 0. \end{cases}$$

A subset $\Sigma \subset M$ is said to be *countably \mathcal{H}^p -rectifiable* if, up to a \mathcal{H}^p -negligible subset, it can be covered by countably many p -dimensional Lipschitz submanifolds. We say that a W_2 -geodesic $\{\mu_t\}_{t \in [0,1]}$ is *countably \mathcal{H}^p -rectifiable* if for every $t \in [0, 1]$ the measure $\mu_t \in \mathcal{P}_c(M, \mathcal{H}^p)$ is concentrated on a countably \mathcal{H}^p -rectifiable set $\Sigma_t \subset M$ (see Section 3 for a through discussion of rectifiable W_2 -geodesics and in particular Remark 3.7 for a sufficient generic condition of rectifiability).

Our first main result is an optimal transport characterization of sectional curvature upper bounds.

Theorem 1.1 (OT characterization of sectional curvature upper bounds, Theorem 5.2). *Let (M, g) be a complete Riemannian manifold without boundary and let $K \geq 0$. Then the following statements (i) and (ii) are equivalent:*

- (i) *The sectional curvature of (M, g) is bounded above by K .*
- (ii) *Let $\{\mu_t\}_{t \in [0,1]}$ be a countably \mathcal{H}^1 -rectifiable W_2 -geodesic, and let Π be the corresponding dynamical optimal plan. Then, if $t_0, t_1 \in (0, 1)$ and $\tau(s) = (1-s)t_0 + st_1$, it holds*

$$\mathcal{H}^1(\text{supp } \mu_{\tau(s)}) \leq \int \left[\sigma_{K,1}^{(1-s)}(|\gamma \circ \tau|) \rho_{t_0}(\gamma(t_0))^{-1} + \sigma_{K,1}^{(s)}(|\gamma \circ \tau|) \rho_{t_1}(\gamma(t_1))^{-1} \right] d\Pi(\gamma), \quad \forall s \in [0, 1],$$

where ρ_t is the density of μ_t with respect to \mathcal{H}^1 .

In the case of $K = 0$ the inequality in (ii) becomes

$$\mathcal{H}^1(\text{supp } \mu_{\tau(s)}) \leq (1-s)\mathcal{H}^1(\text{supp } \mu_{t_0}) + s\mathcal{H}^1(\text{supp } \mu_{t_1}), \quad \forall s \in [0, 1].$$

See Remark 5.3 for the motivation why the upper bound K must be non-negative. Let us also stress that in the assertion (ii) one cannot relax the assumption to $t_0, t_1 \in [0, 1]$, see Remark 5.4 for a counterexample.

The second main result is an optimal transport characterization of sectional curvature lower bounds. In order to state it, some more notation must be introduced. First of all, given a \mathcal{H}^p -rectifiable W_2 -geodesic $\{\mu_t\}_{t \in [0,1]}$, thanks to the Monge-Mather shortening principle [19, Theorem 8.5] we know that, for every $t \in [0, 1]$, $\mu_t = (T_{1/2}^t)_\# \mu_{1/2}$ with $T_{1/2}^t : \Sigma_{1/2} \rightarrow \Sigma_t$ Lipschitz. For $\mu_{1/2}$ -a.e. x we can set (see Lemma 3.10 and Remark 3.11 for the details)

$$B_x(t) : T_x \Sigma_{1/2} \rightarrow T_{\gamma_x(t)} \Sigma_t, \quad B_x(t) := DT_{1/2}^t(x) \quad \forall t \in [0, 1].$$

In Lemma 3.9 we will prove a Monge-Ampère inequality implying that $B_x(t)$ is invertible. Let $\gamma_x(t) := T_{1/2}^t(x)$ be a geodesic performing the transport and consider

$$\begin{aligned} \mathcal{U}_x(t) &:= (\nabla_t B_x(t)) B_x(t)^{-1} : T_{\gamma_x(t)} \Sigma_t \rightarrow T_{\gamma_x(t)} M, \\ \mathcal{U}_x^\perp(t) &:= [\mathcal{U}_x(t)]^\perp : T_{\gamma_x(t)} \Sigma_t \rightarrow (T_{\gamma_x(t)} \Sigma_t)^\perp, \end{aligned}$$

where ∇_t denotes the covariant derivative along $\gamma_x(t)$ in M and \perp is the orthogonal projection on the orthogonal complement $(T_{\gamma(t)} \Sigma_t)^\perp$ of $T_{\gamma(t)} \Sigma_t$. If $|\dot{\gamma}_x| \neq 0$, we set

$$\kappa_{\gamma_x} : [0, |\dot{\gamma}_x|] \rightarrow \mathbb{R}, \quad \kappa_{\gamma_x}(|\dot{\gamma}_x| t) |\dot{\gamma}_x|^2 := \|\mathcal{U}_x^\perp(t)\|^2, \quad \forall t \in [0, 1],$$

if $|\dot{\gamma}_x| = 0$, we set $\kappa_{\gamma_x}(0) = 0$. We now introduce the generalized distortion coefficients σ_κ associated to a continuous function $\kappa : [0, \theta] \rightarrow \mathbb{R}$ (cf. [9]). First of all, the generalized sin-function associated to κ , denoted by \sin_κ , is defined as the unique solution $v : [0, \theta] \rightarrow \mathbb{R}$ of the equation

$$v'' + \kappa v = 0 \quad \& \quad v(0) = 0, \quad v'(\theta) = 1.$$

The generalized distortion coefficients $\sigma_\kappa^{(t)}(\theta)$, for $t \in [0, 1]$ and $\theta > 0$, are defined as

$$\sigma_\kappa^{(t)}(\theta) := \begin{cases} \frac{\sin_\kappa(t\theta)}{\sin_\kappa(\theta)} & \text{if } \sin_\kappa(t\theta) > 0 \text{ for all } t \in [0, 1], \\ \infty & \text{otherwise.} \end{cases}$$

In the case $\kappa = K = \text{const}$ one has $\sigma_\kappa^{(t)}(\theta) = \sigma_{K,1}^{(t)}(\theta)$. It is convenient to also set $\sigma_\kappa^{(\cdot)}(0) \equiv 1$, $\kappa^-(t) = \kappa(\theta - t)$ and $\kappa^+(t) := \kappa(t)$. Finally, consider the Green function $g : [0, 1] \times [0, 1] \rightarrow [0, 1]$ given by

$$g(s, t) := \begin{cases} (1-s)t & \text{if } t \in [0, s], \\ s(1-t) & \text{if } t \in [s, 1]. \end{cases}$$

We can now state the optimal transport characterization of sectional curvature lower bounds.

Theorem 1.2 (OT characterization of sectional curvature lower bounds). *Let (M, g) be a complete n -dimensional Riemannian manifold without boundary and fix $K \in \mathbb{R}$.*

- If $K \geq 0$ the next conditions are equivalent:

- M has sectional curvature bounded from below by K .
- Let $p \in \{2, \dots, n\}$ be arbitrary, let $\{\mu_t\}_{t \in [0,1]}$ be a \mathcal{H}^p -rectifiable W_2 -geodesic and Π be the corresponding dynamical optimal plan. Then, for any $p' \geq p$, for all $t \in [0, 1]$ it holds

$$S_{p'}(\mu_t | \mathcal{H}^p) \leq - \int \left[\sigma_{((p-1)K - \kappa_\gamma^-) / p'}^{(1-t)}(|\dot{\gamma}|) \rho_0^{-\frac{1}{p'}}(\gamma(0)) + \sigma_{((p-1)K - \kappa_\gamma^+) / p'}^{(t)}(|\dot{\gamma}|) \rho_1^{-\frac{1}{p'}}(\gamma(1)) \right] d\Pi(\gamma).$$

- The condition (ii) holds for $p = 2$.

- Let $p \in \{2, \dots, n\}$ be arbitrary, $\{\mu_t\}_{t \in [0,1]}$ and Π be as in (ii). Then for all $t \in [0, 1]$ it holds

$$\text{Ent}(\mu_t | \mathcal{H}^p) \leq (1-t) \text{Ent}(\mu_0 | \mathcal{H}^p) + t \text{Ent}(\mu_1 | \mathcal{H}^p) - \int \int_0^1 g(s, t) |\dot{\gamma}|^2 ((p-1)K - \kappa_\gamma(s|\dot{\gamma}|)) ds d\Pi(\gamma).$$

- The condition (iii) holds for $p = 2$.

- If $K \leq 0$ the next conditions are equivalent:
 - (i) M has sectional curvature bounded from below by K .
 - (ii) Let $p \in \{1, \dots, n\}$ be arbitrary, let $\{\mu_t\}_{t \in [0,1]}$ be a \mathcal{H}^p -rectifiable W_2 -geodesic and Π be the corresponding dynamical optimal plan. Then, for any $p' \geq p$, for all $t \in [0, 1]$ it holds

$$S_{p'}(\mu_t | \mathcal{H}^p) \leq - \int \left[\sigma_{(\bar{K} - \kappa_\gamma^-)/p'}^{(1-t)}(|\dot{\gamma}|) \rho_0^{-\frac{1}{p'}}(\gamma(0)) + \sigma_{(\bar{K} - \kappa_\gamma^+)/p'}^{(t)}(|\dot{\gamma}|) \rho_1^{-\frac{1}{p'}}(\gamma(1)) \right] d\Pi(\gamma),$$

where $\bar{K} := \min\{p, n-1\}K$.

- (ii)' The condition (ii) holds for $p = 1$.
- (iii) Let $p \in \{1, \dots, n\}$ be arbitrary, \bar{K} , $\{\mu_t\}_{t \in [0,1]}$ and Π be as in (ii). Then for all $t \in [0, 1]$ it holds

$$\text{Ent}(\mu_t | \mathcal{H}^p) \leq (1-t) \text{Ent}(\mu_0 | \mathcal{H}^p) + t \text{Ent}(\mu_1 | \mathcal{H}^p) - \int_0^1 \int_0^1 g(s, t) |\dot{\gamma}|^2 (\bar{K} - \kappa_\gamma(s|\dot{\gamma}|)) ds d\Pi(\gamma).$$

- (iii)' The condition (iii) holds for $p = 1$.

Note that, in case $p = n$, the correction term κ_γ vanishes (indeed it does not appear in the OT characterization of Ricci curvature lower bounds), but for $p < n$ Theorem 1.2 is sharp in the sense that one can not suppress κ_γ (see Remark 6.2); more strongly, for the very same example of Remark 6.2, all the inequalities involved in the proof Theorem 1.2 become identities (see Remark 6.4), showing the sharpness of the arguments.

Theorem 1.2 is actually a particular case of Theorem 6.1 (see also Remark 2.3, for the link between p -Ricci and sectional curvatures) where we characterize lower bounds on the p -Ricci curvature in terms of optimal transport, for any $p \in \{1, \dots, n\}$. For the rigorous definition and basic properties of the p -Ricci curvature we refer to Section 2, here let us just mention the intuitive idea behind: in the standard Ricci curvature (corresponding in this notation to the n -Ricci curvature), one considers the trace of the Riemann curvature tensor along *all the tangent space* to M at some point $x \in M$, while in the p -Ricci curvature one considers the trace of the Riemann curvature tensor *just along p -dimensional subspaces*. The notion of p -Ricci curvature has already been considered in the literature, in particular in connection with topological results (see for instance the works of Wu [22], Shen [15], Wilhelm [21], Petersen-Wilhelm [12] and Xu-Ye [23]). Just to fix the ideas, let us recall that if the sectional curvature is bounded below by $K \geq 0$, then the p -Ricci curvature is bounded below by $(p-1)K$; if instead the sectional curvature is bounded below by $K \leq 0$, then the p -Ricci curvature is bounded below by $\min\{p, n-1\}K$.

The paper is organized as follows: Section 2 settles the notation and the preliminaries. In Section 3 we analyze \mathcal{H}^p -rectifiable W_2 -geodesics and in Section 4 we perform the Jacobi fields computations/estimates that will be used to prove the main results. Section 5 is devoted to the proof of Theorem 1.1, namely the optimal transport characterization of sectional curvature upper bounds. Finally, in Section 6 we state and prove our main results characterizing sectional and p -Ricci lower bounds in terms of optimal transportation; as a consequence, we also obtain a new Brunn-Minkowski type inequality involving p -dimensional submanifolds and the p -Ricci curvature (see Corollary 6.5).

Acknowledgement: Most of the work was done while both authors were in residence at the Mathematical Sciences Research Institute in Berkeley, California during the Spring 2016 semester, supported by the National Science Foundation under Grant No. DMS-1440140. We thank the organizers of the Differential Geometry Program and MSRI for providing great environment for research and collaboration. In the final steps of the project, A. M. has been supported by the EPSRC First Grant EP/R004730/1. We also wish to express our gratitude to Robert McCann for suggesting the Remark 3.7, and to Martin Kell and Gerardo Sosa for their careful reading of the manuscript.

2. PRELIMINARIES

Optimal transport and Wasserstein geometry. It is out of the scopes of this short section to give a comprehensive introduction to optimal transport, for this purpose we refer to [19]. Instead, we will be satisfied by recalling those notions and results that we will use throughout the paper.

Let (X, d) be a complete, separable and proper metric space. A curve $\gamma : [0, 1] \rightarrow X$ is said to be a (length-minimizing, constant speed) geodesic if

$$d(\gamma(s), \gamma(t)) = |s - t| d(\gamma(0), \gamma(1)), \quad \forall s, t \in [0, 1].$$

We denote by $\text{Geo}(X) := \{\gamma : [0, 1] \rightarrow X \text{ s.t. } \gamma \text{ is a geodesic}\}$ the family of geodesics equipped with the L^∞ -topology. The evaluation map $e_t : \text{Geo}(X) \rightarrow X$ is given by $e_t(\gamma) = \gamma(t)$, and it is clearly continuous with respect to the sup-distance $d_\infty(\gamma, \tilde{\gamma}) = \sup_{t \in [0, 1]} d(\gamma(t), \tilde{\gamma}(t))$.

$\mathcal{P}_c(X)$ denotes the space of Borel probability measures with compact support and $\mathcal{P}_2(X)$ denotes the space of Borel probability measures μ with finite second moment, i.e. satisfying $\int_X d^2(x, x_0) d\mu(x) < \infty$ for some (and thus for any) $x_0 \in X$.

The space $\mathcal{P}_2(X)$ is naturally endowed with the L^2 -Wasserstein distance W_2 defined by

$$W_2(\mu_1, \mu_2)^2 := \inf \left\{ \int_{X \times X} d^2(x, y) d\pi(x, y) \text{ s.t. } \pi \in \text{Cpl}(\mu_1, \mu_2) \right\},$$

where $\text{Cpl}(\mu_1, \mu_2)$ is the family of all couplings between μ_1 and μ_2 , i.e. of all the probability measures $\pi \in \mathcal{P}(X^2)$ such that $(P_i)_\# \pi = \mu_i$, $i = 1, 2$, P_1, P_2 being the projection maps. $(\mathcal{P}_2(X), W_2)$ becomes a separable metric space that is a geodesic metric space provided X is a geodesic metric space.

A coupling $\pi \in \text{Cpl}(\mu_1, \mu_2)$ is optimal if

$$\int_{X^2} d(x, y)^2 d\pi(x, y) = W_2(\mu_1, \mu_2)^2.$$

Optimal couplings always exist, and if an optimal coupling π is induced by a map $T : Z \rightarrow X$ via $(T, \text{Id}_X)_\# \mu_1 = \pi$, where Z is a measurable subset of X , we say that T is an optimal map. A probability measure $\Pi \in \mathcal{P}(\text{Geo}(X))$ is called an optimal dynamical coupling or plan if $(e_0, e_1)_\# \Pi$ is an optimal coupling between the initial and final marginal distribution. For every W_2 -geodesic $\{\mu_t\}_{t \in [0, 1]}$ there exists an optimal dynamical plan $\Pi \in \mathcal{P}(\text{Geo}(X))$ such that $\mu_t = (e_t)_\# \Pi$ for all $t \in [0, 1]$.

In the present paper, a key role is played by the subspace $\mathcal{P}_2(X, \mathcal{H}^p) \subset \mathcal{P}_2(X)$ made of probability measures that are absolutely continuous with respect to the p -dimensional Hausdorff measure \mathcal{H}^p . We also denote with $\mathcal{P}_c(X, \mathcal{H}^p) \subset \mathcal{P}_2(X, \mathcal{H}^p)$ the subspace of absolutely continuous probability measures with compact support.

In the introduction, for simplicity, we defined the entropy functionals for compactly supported probability measures; the definitions carry over to probability measures with finite second moment, let us briefly recall them. Given $1 \leq p \leq p' < \infty$, the p' -Renyi entropy with respect to the p -dimensional Hausdorff measure \mathcal{H}^p is defined as

$$S_{p'}(\cdot | \mathcal{H}^p) : \mathcal{P}_2(X, \mathcal{H}^p) \rightarrow [-\infty, 0], \quad S_{p'}(\mu | \mathcal{H}^p) = - \int \rho^{1 - \frac{1}{p'}} d\mathcal{H}^p,$$

where ρ is the density of μ with respect to \mathcal{H}^p , i.e. $\mu = \rho \mathcal{H}^p$. Notice that, by Jensen's inequality, we have

$$[-\infty, 0] \ni -(\mathcal{H}^p(\text{supp } \mu))^{1/p'} \leq S_{p'}(\mu | \mathcal{H}^p).$$

In particular if ρ is concentrated on a set of finite \mathcal{H}^p -measure then $S_{p'}(\mu | \mathcal{H}^p) > -\infty$. Note that in the borderline case $p' = p = 1$, one gets

$$S_1(\mu | \mathcal{H}^1) = -\mathcal{H}^1(\text{supp}(\mu)).$$

Finally, the (relative) Shannon entropy is defined by

$$\text{Ent}(\cdot | \mathcal{H}^p) : \mathcal{P}_2(X, \mathcal{H}^p) \rightarrow [-\infty, \infty], \quad \text{Ent}(\mu | \mathcal{H}^p) = \lim_{\varepsilon \downarrow 0} \int_{\{\rho > \varepsilon\}} \rho \log \rho d\mathcal{H}^p.$$

This coincides with $\int_{\{\rho>0\}} \rho \log \rho d\mathcal{H}^p$, provided that $\int_{\{\rho \geq 1\}} \rho \log \rho d\mathcal{H}^p < \infty$, and $\text{Ent}(\mu|\mathcal{H}^p) := \infty$ otherwise.

Rectifiable sets. Let $\Sigma \subset \mathbb{R}^n$ and $m \in \mathbb{N}, m \leq n$. We say that Σ is *countably m -rectifiable* if there is a countable family of Lipschitz maps $f_i : \mathbb{R}^m \rightarrow \mathbb{R}^n$, such that $\Sigma \subset \bigcup_i f_i(\mathbb{R}^m)$. The set Σ is *countably \mathcal{H}^m -rectifiable* if there is a countably m -rectifiable set $\Sigma' \subset \mathbb{R}^n$ such that $\mathcal{H}^m(\Sigma \setminus \Sigma') = 0$.

As it is well known, using Whitney extension Theorem, it is possible to show that a subset $\Sigma \subset \mathbb{R}^n$ is countably \mathcal{H}^m -rectifiable if and only if there exists a sequence of m -dimensional C^1 -submanifolds $\{S_i\}_{i \in \mathbb{N}}$ such that

$$\mathcal{H}^m\left(\Sigma \setminus \bigcup_{i \in \mathbb{N}} S_i\right) = 0.$$

Clearly, by considering local coordinates (or by Nash isometric embedding Theorem), one can define the same notions for subsets of an n -dimensional Riemannian manifold.

Intermediate Ricci curvature. Let (M, g) be an n -dimensional Riemannian manifold and let

$$R : TM \times TM \times TM \rightarrow TM, \quad R(X, Y)Z := \nabla_Y \nabla_X Z - \nabla_X \nabla_Y Z + \nabla_{[X, Y]} Z$$

the Riemannian curvature tensor (of course ∇ denotes the Levi-Civita connection of (M, g) and $[\cdot, \cdot]$ denotes the Lie bracket). Sometimes we will use the notation $|v| := \sqrt{g(v, v)}$ and $\langle v, w \rangle := g(v, w)$. Using the standard notation, $T_x M$ is the tangent space of M at the point $x \in M$. For a 2-plane $P \subset T_x M$ spanned by $v, w \in T_x M$, let

$$\text{Sec}(P) = \text{Sec}(v, w) := \frac{\langle R(v, w)v, w \rangle}{|v|^2 |w|^2 - \langle v, w \rangle^2}$$

be the sectional curvature. Recall that, given $w \in T_x M$, the Ricci curvature $\text{Ric}(w, w)$ is defined by

$$\text{Ric}(w, w) := \text{tr} [R(w, \cdot)w].$$

Definition 2.1 (p -Ricci Curvature). Let $p \in \{1, \dots, n\}$. For a p -dimensional plane P in $T_x M$ and a vector $w \in T_x M$, we define the p -Ricci curvature of P in the direction of w as

$$(1) \quad \text{Ric}_p(P, w) := \text{tr} [\top_P \circ (R(w, \cdot)w)|_P] = \sum_{i=1}^p \text{Sec}(e_i, w)(|w|^2 - \langle e_i, w \rangle^2),$$

where e_1, \dots, e_p is an orthonormal basis of P , and $\top_P : T_x M \rightarrow P$ is the orthogonal projection of $T_x M$ onto P .

Note that, in particular, if $|w| = 1$ and w is orthogonal to P then

$$\text{Ric}_p(P, w) = \sum_{i=1}^p \text{Sec}(e_i, w).$$

It is standard to check that Ric_p is well-defined and independent of the choice of a basis for P . Notice also that, if $w \notin P$, then

$$(2) \quad \text{Ric}_p(P, w) = \text{Ric}_{p+1}(\text{span}(P, w), w) = \text{Ric}_p(\text{span}(P, w) \cap w^\perp, w) = \sum_{i=1}^p \text{Sec}(e_i, w)|w|^2,$$

where $\{e_i\}_{i=1, \dots, p}$ is an orthonormal basis of $\text{span}(P, w) \cap w^\perp$, $w^\perp \subset T_x M$ being the orthogonal subspace to w .

Definition 2.2 (p -Ricci upper and lower bounds). We say that (M, g) has p -Ricci curvature bounded from below (resp. from above) by K if, for any $x \in M$ and any p -dimensional plane $P \subset T_x M$, we have $\text{Ric}_p(P, w) \geq K|w|^2$ (resp. $\text{Ric}_p(P, w) \leq K|w|^2$); in this case we write $\text{Ric}_p \geq K$ (resp. $\text{Ric}_p \leq K$).

Remark 2.3 (Some notable cases). The cases $p = 1, 2$ are strictly linked with the sectional curvature while $p = n - 1, n$ are related to the standard Ricci curvature. More precisely

- $p = 1$: if P is the real line spanned by v , $\langle v, w \rangle = 0$, $|v| = |w| = 1$, then

$$\text{Ric}_1(P, w) = \text{Sec}(v, w);$$

on the other hand $\text{Ric}_1(P, v) = 0$, i.e. the 1-Ricci curvature always vanishes in the direction of P itself. In particular no Riemannian manifold has 1-Ricci curvature bounded from below (resp. above) by a strictly positive (resp. negative) constant. Nevertheless M has non-negative (resp. non-positive) 1-Ricci curvature if and only if the sectional curvature is non-negative (resp. non-positive).

- $p = 2$: if P is the 2-plane spanned by the orthonormal vectors e_1, e_2 then

$$(3) \quad \text{Ric}_2(P, e_1) = \text{Ric}_2(P, e_2) = \text{Sec}(e_1, e_2).$$

Moreover, if w is orthogonal to P with $|w| = 1$ then

$$\text{Ric}_2(P, w) = \text{Sec}(e_1, w) + \text{Sec}(e_2, w).$$

In particular for every $K \geq 0$ (resp. $K \leq 0$), it holds $\text{Ric}_2 \geq K$ (resp. $\text{Ric}_2 \leq K$) if and only if $\text{Sec} \geq K$ (resp. $\text{Sec} \leq K$). Note also that if $\text{Sec} \geq K \geq 0$ then for every $p \in \{2, \dots, n\}$ it holds $\text{Ric}_p \geq (p-1)K$.

- $p = n-1$: if P is an $n-1$ -plane and w is orthogonal to P , then

$$\text{Ric}_{n-1}(P, w) = \text{Ric}(w, w).$$

- $p = n$: in this case one has $P = T_x M$, and for every $w \in T_x M$ it holds

$$\text{Ric}_n(T_x M, w) = \text{Ric}(w, w).$$

- If $\text{Sec} \geq K$, depending on the sign of $K \in \mathbb{R}$ we have:
 - $\text{Sec} \geq K \geq 0$ implies that $\text{Ric}_p \geq (p-1)K$, for all $p \in \{1, \dots, n\}$
 - $\text{Sec} \geq K$ with $K \leq 0$ implies that $\text{Ric}_n \geq (n-1)K$ and $\text{Ric}_p \geq pK$ for all $p \in \{1, \dots, n-1\}$.

3. COUNTABLY \mathcal{H}^p -RECTIFIABLE GEODESICS IN WASSERSTEIN SPACE

The next result is a well known consequence of the Monge-Mather shortening principle [19, Theorem 8.5].

Theorem 3.1. *Consider a Riemannian manifold (M, g) , fix a compact subset $E \subset\subset M$ and let Π be a dynamical optimal plan such that $(e_t)_\# \Pi$ is supported in E for every $t \in [0, 1]$.*

Then Π is supported on a set of geodesics $S \subset \text{Geo}(M)$ satisfying the following: for every $t_0 \in (0, 1)$ there exists $C_E(t_0) > 0$ such that for any two geodesics $\gamma, \eta \in S$ it holds

$$\sup_{t \in [0, 1]} d(\gamma(t), \eta(t)) \leq C_E(t_0) d(\gamma(t_0), \eta(t_0)),$$

where d is the Riemannian distance on (M, g) .

Remark 3.2. As a consequence of Theorem 3.1, if $\{\mu_t\}_{t \in [0, 1]}$ is a W_2 -geodesic such that μ_0, μ_1 are compactly supported probability measures on M , and $t_0 \in (0, 1)$ is given, then for any $t \in [0, 1]$ the map $T_{t_0}^t : \gamma(t_0) \mapsto \gamma(t)$ is well-defined μ_{t_0} -almost everywhere and Lipschitz continuous on its domain; moreover it is the unique optimal transport map between μ_{t_0} and μ_t . In other words, the optimal coupling $(e_{t_0}, e_t)_\# \Pi$ is induced by $T_{t_0}^t$, i.e. $(e_{t_0}, e_t)_\# \Pi = (\text{Id}, T_{t_0}^t)_\# \mu_{t_0}$.

Lemma 3.3. *Let (M, g) be a complete n -dimensional Riemannian manifold without boundary, and let $p \in \{1, \dots, n\}$. Let $\mu_0, \mu_1 \in \mathcal{P}_c(M, \mathcal{H}^p)$ and assume $\{\mu_t\}_{t \in [0, 1]}$ is a W_2 -geodesic between μ_0, μ_1 such that for some $t_0 \in (0, 1)$ the measure μ_{t_0} is concentrated on a countably \mathcal{H}^p -rectifiable set $\Sigma_{t_0} \subset M$.*

Then for every $t \in [0, 1]$ there exists a countably \mathcal{H}^p -rectifiable set $\Sigma_t \subset M$ such that μ_t is concentrated on Σ_t ; moreover $\mu_t = \rho_t \mathcal{H}^p \llcorner \Sigma_t \in \mathcal{P}_c(M, \mathcal{H}^p)$ for a suitable probability density $\rho_t \in L^1(M, \mathcal{H}^p)$.

Proof. Step 1. By Theorem 3.1 and Remark 3.2 we know that for every $t \in [0, 1]$ there exists a Lipschitz map $T_{t_0}^t : \text{supp } \mu_{t_0} \rightarrow \text{supp } \mu_t$ such that $\mu_t = (T_{t_0}^t)_\# \mu_{t_0}$. Since by assumption μ_{t_0} is concentrated on the countably \mathcal{H}^p -rectifiable set Σ_{t_0} , it is then clear that μ_t is concentrated on $\Sigma_t := T_{t_0}^t(\Sigma_{t_0})$ which is countably \mathcal{H}^p -rectifiable set too, as Lipschitz image of a countably \mathcal{H}^p -rectifiable set. In order to conclude that

$\mu_t = \rho_t \mathcal{H}^p \llcorner \Sigma_t \in \mathcal{P}_c(M, \mathcal{H}^p)$ it is then enough to show that $\mu_t(A) = 0$ for every $A \subset \text{supp } \mu_t$ satisfying $\mathcal{H}^p(A) = 0$. This will be proved in Step 3, using the discussion of Step 2.

Step 2. Consider $\mu_0, \mu_1 \in \mathcal{P}_c(M, \mathcal{H}^p)$ and write $\mu_i = \rho_i \mathcal{H}^p$ for $i = 0, 1$. Let $\Pi \in \mathcal{P}(\text{Geo}(M))$ a dynamical optimal plan between μ_0 and μ_1 , and let $\{\mu_t := (e_t)_\# \Pi\}_{t \in [0,1]}$ be the induced L^2 -Wasserstein geodesic. We denote with $\pi_{t,s} = (e_t, e_s)_\# \Pi$ the corresponding optimal coupling between μ_t and μ_s for any $t, s \in [0, 1]$. Since μ_0 and μ_1 have compact support, then there exists a compact subset $E \subset\subset M$ such that $\text{supp } \mu_t \subset E$ for every $t \in [0, 1]$.

By Theorem 3.1, the dynamical optimal plan Π is supported on set $S \subset \text{Geo}(M)$ satisfying the following: for any $t \in (0, 1)$ there exists $C_E(t) > 0$ such that for any $s \in [0, 1]$ it holds

$$(4) \quad d(\gamma(s), \eta(s)) \leq C_E(t) d(\gamma(t), \eta(t)) \quad \text{for any pair } \gamma, \eta \in S.$$

As observed in Remark 3.2, the optimal plan $\pi_{t,s}$ is then induced by a Lipschitz-continuous optimal transport map $T_t^s : \text{supp } \mu_t \rightarrow \text{supp } \mu_s$ with Lipschitz constant bounded above by $C_E(t)$. In particular $(T_t^s)_\# \mu_t = \mu_s$.

Step 3. Let $t \in (0, 1)$, and consider $\pi_{t,0} := (e_t, e_0)_\# \Pi$. Our goal is to show that if $A \subset \text{supp } \mu_t$ satisfies $\mathcal{H}^p(A) = 0$, then $\mu_t(A) = 0$ as well. Since by Step 2 the plan $\pi_{t,0}$ is induced by the map T_t^0 , we have

$$(5) \quad \mu_t(A) = \pi_{t,0}(A, M) = \pi_{t,0}(A, T_t^0(A)).$$

On the other hand

$$(6) \quad \pi_{t,0}(A, T_t^0(A)) \leq \pi_{t,0}(M, T_t^0(A)) = \mu_0(T_t^0(A)).$$

Since $T_t^0 : \text{supp } \mu_t \rightarrow \text{supp } \mu_0$ is Lipschitz and $\mathcal{H}^p(A) = 0$, then it also holds $\mathcal{H}^p(T_t^0(A)) = 0$. Recalling that by assumption $\mu_0 \ll \mathcal{H}^p$, we then get that $\mu_0(T_t^0(A)) = 0$. The claim follows then by the combination of (6) and (5). \square

Definition 3.4. We say that a W_2 -geodesic $\{\mu_t\}_{t \in [0,1]}$ is *countably \mathcal{H}^p -rectifiable* if for every $t \in [0, 1]$ the measure $\mu_t \in \mathcal{P}_c(M, \mathcal{H}^p)$ is concentrated on a countably \mathcal{H}^p -rectifiable set $\Sigma_t \subset M$.

Remark 3.5. By Lemma 3.3, a W_2 -geodesic $\{\mu_t\}_{t \in [0,1]}$ is countably \mathcal{H}^p -rectifiable if and only if $\mu_0, \mu_1 \in \mathcal{P}_c^2(M, \mathcal{H}^p)$ and there exists $t_0 \in (0, 1)$ such that the measure μ_{t_0} is concentrated on a countably \mathcal{H}^p -rectifiable set $\Sigma_{t_0} \subset M$.

Remark 3.6. Note that, in Definition 3.4, one can replace Σ_t by $\Sigma_t \cap \text{supp } \mu_t$; thus from now on we will always tacitly assume that $\Sigma_t = \Sigma_t \cap \text{supp } \mu_t$, for all $t \in [0, 1]$. Also, since for $s \in (0, 1)$ and $t \in [0, 1]$ the optimal transport map T_s^t given in Remark 3.2 is well defined μ_s -a.e., from now on we will just consider the restriction $T_s^t \llcorner \Sigma_s$ and, for simplicity of notation, write T_s^t to indicate the map $T_s^t \llcorner \Sigma_s : \Sigma_s \rightarrow T_s^t(\Sigma_s)$. Note that, with this notation, for μ_s -almost every x , the differential $DT_s^t(x)$ is a linear map from the p -dimensional space $T_x \Sigma_s$ to the q -dimensional space $T_{T_s^t(x)}(T_s^t(\Sigma_s))$, $q \leq p$ (q possibly depending on x).

Remark 3.7 (A sufficient condition for the p -rectifiability of μ_t). The following sufficient condition for the p -rectifiability of the geodesic μ_t follows by combining the work of McCann-Pass-Warren [10, Theorem 1.2] with Lemma 3.3.

Given $p \in \{1, \dots, n\}$, let $\mu_0, \mu_1 \in \mathcal{P}_c(M, \mathcal{H}^p)$ with $\mu_i = \rho_i \mathcal{H}^p \llcorner \Sigma_i$, for some smooth p -dimensional submanifolds Σ_i , $i = 1, 2$. Consider the restriction of the quadratic cost function d^2 to the product $\Sigma_0 \times \Sigma_1$; if

$$(7) \quad \det \left[\left(\frac{\partial^2}{\partial x_i \partial y_j} d^2 \Big|_{\Sigma_0 \times \Sigma_1} \right)_{i,j=1,\dots,p} \right] \neq 0$$

and moreover

$$(8) \quad \Sigma_0 \cap \left(\bigcup_{x \in \Sigma_1} \text{Cut}(x) \right) = \emptyset \quad \text{and} \quad \Sigma_1 \cap \left(\bigcup_{x \in \Sigma_0} \text{Cut}(x) \right) = \emptyset,$$

where $\text{Cut}(x)$ is the cut locus of the point $x \in M$, then every W_2 -geodesic $\{\mu_t\}_{t \in [0,1]}$ between μ_0 and μ_1 satisfies that $\mu_t = \rho_t \mathcal{H}^p \llcorner \Sigma_t \in \mathcal{P}_c(M, \mathcal{H}^p)$ for every $t \in [0, 1]$ for a countably \mathcal{H}^p -rectifiable set $\Sigma_t \subset M$.

Indeed, calling π the L^2 -optimal coupling induced by the geodesic $\{\mu_t\}_{t \in [0,1]}$, by using (7) we can apply [10, Theorem 1.2] and get that π is supported on a p -dimensional Lipschitz submanifold S of $\Sigma_0 \times \Sigma_1 \subset M \times M$. Using now (8), we get that for every $(x, y) \in \Sigma_0 \times \Sigma_1$ there exists a unique geodesic $t \mapsto \gamma_t(x, y)$ from $x = \gamma_0(x, y) \in \Sigma_0$ to $y = \gamma_1(x, y) \in \Sigma_1$; moreover the map $\gamma_t(\cdot, \cdot) : \Sigma_0 \times \Sigma_1 \rightarrow M \times M$ is Lipschitz, for every fixed $t \in [0, 1]$. Calling $\Sigma_t := \gamma_t(S)$ we get that μ_t is concentrated on the p -rectifiable subset Σ_t . The fact that we can write $\mu_t = \rho_t \mathcal{H}^p \llcorner \Sigma_t$ for some density $\rho_t \in L^1(\mathcal{H}^p)$ follows then by Lemma 3.3.

Remark 3.8. For fixed s and t , pick a (resp. orthonormal) basis $(e_i)_{i=1, \dots, p}$ of $T_x \Sigma_s \subset T_x M$, and also a (resp. orthonormal) basis $(f_i)_{i=1, \dots, n}$ of $T_{T_s^t(x)} M$ such that $(f_i)_{i=1, \dots, q}$ is a basis of $T_{T_s^t(x)}(T_s^t(\Sigma_s))$. We can then see $DT_s^t(x)$ as a linear map from \mathbb{R}^p to \mathbb{R}^p (if $q < p$ just identify \mathbb{R}^q with $\{(x^1, \dots, x^p) : x^1 = \dots = x^{p-q} = 0\}$). Since the rank and the determinant are independent of the chosen basis, $\det[DT_s^t(x)]$ and the fact that $DT_s^t(x)$ is non-degenerate are then well defined concepts.

In the next lemma we show that the optimal transport map T_s^t is differentiable μ_s -a.e. on Σ_s and that at least an inequality holds in the Monge-Ampère equation (cf. [3]); this will be sufficient (and crucial) to our aims of characterizing curvature bounds in terms of optimal transport.

Lemma 3.9. *Let M be a complete Riemannian manifold and $\{\mu_t\}_{t \in [0,1]}$ a W_2 -geodesic with $\mu_t \ll \mathcal{H}^p \llcorner \Sigma_t \in \mathcal{P}_c(M, \mathcal{H}^p)$ for some countably \mathcal{H}^p -rectifiable subset $\Sigma_t \subset M$, for every $t \in [0, 1]$. For fixed $s \in (0, 1)$ and $t \in [0, 1]$, let T_s^t be the optimal transport map from μ_s to μ_t given in Remark 3.2.*

Then $T_s^t : \Sigma_s \rightarrow T_s^t(\Sigma_s) \subset M$ is differentiable μ_s -a.e. and the following Monge-Ampère inequality holds:

$$(9) \quad \rho_s(x) \leq \det[DT_s^t(x)] \rho_t(T_s^t(x)) \quad \mu_s\text{-a.e. } x, \forall s \in (0, 1), \forall t \in [0, 1].$$

In particular, $DT_s^t : \mathbb{R}^p \rightarrow \mathbb{R}^p$ is μ_s -a.e. non-degenerate. Moreover (9) holds with equality if $t, s \in (0, 1)$.

Let us stress that in the above lemma we do not claim that T_s^t is μ_s -a.e. differentiable as a map from M to M , but just as a map from Σ_s to its image, i.e. we claim differentiability with respect to infinitesimal variations which are *tangential* to Σ_s .

Proof. Step 1. Differentiability μ_s -a.e..

From Theorem 3.1 and Remark 3.2, we know that $T_s^t : \Sigma_s \rightarrow T_s^t(\Sigma_s)$ is a Lipschitz map; since by assumption Σ_s is countably \mathcal{H}^p -rectifiable, Rademacher Theorem implies that $T_s^t : \Sigma_s \rightarrow T_s^t(\Sigma_s)$ is differentiable \mathcal{H}^p -a.e. .

Step 2. Monge-Ampère inequality.

Since by construction $(T_s^t)_\# \mu_s = \mu_t$, it follows that for an arbitrary Borel subset $A \subset \Sigma_s$ it holds

$$(10) \quad \mu_s(A) \leq \mu_s((T_s^t)^{-1}(T_s^t(A))) = \mu_t(T_s^t(A)).$$

Equality holds for $s, t \in (0, 1)$ as the map T_s^t is μ_s -essentially injective. Recalling that $\mu_s = \rho_s \mathcal{H}^p \llcorner \Sigma_s$ and $\mu_t = \rho_t \mathcal{H}^p \llcorner \Sigma_t$, by the area formula we infer that

$$(11) \quad \begin{aligned} \mu_t(T_s^t(A)) &= \int_{T_s^t(A)} \rho_t d\mathcal{H}^p \llcorner \Sigma_t \\ &\leq \int_{T_s^t(A)} \rho_t(y) \mathcal{H}^0((T_s^t \llcorner A)^{-1}(y)) d\mathcal{H}^p \llcorner \Sigma_t(y) = \int_A \rho_t(T_s^t(x)) \det[DT_s^t(x)] d\mathcal{H}^p \llcorner \Sigma_s(x), \end{aligned}$$

with equality if $s, t \in (0, 1)$ as the map T_s^t is μ_s -essentially injective. The combination of (10) and (11) gives that for an arbitrary Borel subset $A \subset \Sigma_s$ it holds

$$\int_A \rho_s d\mathcal{H}^p = \mu_s(A) \leq \mu_t(T_s^t(A)) \leq \int_A \rho_t(T_s^t(x)) \det[DT_s^t(x)] d\mathcal{H}^p,$$

and the Monge-Ampère inequality (9) follows, with equality for $s, t \in (0, 1)$. \square

In order to have a more clear notation, in the next lemma we pick $s = 1/2$ and consider the Lipschitz map $T_{1/2}^t : \Sigma_{1/2} \rightarrow \Sigma_t$, $t \in [0, 1]$, but the same arguments hold for any fixed $s \in (0, 1)$. For $\mu_{1/2}$ -a.e. $x \in M$ let $\gamma_x \in \text{Geo}(M)$ be the geodesic defined by $[0, 1] \ni t \mapsto \gamma_x(t) := T_{1/2}^t(x)$ and, for $\mu_{1/2}$ -a.e. $x \in \Sigma_{1/2}$ let $v(x) \in T_x M$ be such that $\gamma_x(t) = \exp_x((t - \frac{1}{2})v(x))$, that is $v(x) = \dot{\gamma}_x(\frac{1}{2})$. Denote also with ∇_t the covariant derivative in M in the direction of $\dot{\gamma}_x(t)$ and with $D_t = \top \circ \nabla_t$ where $\top : T_{\gamma_x(t)} M \rightarrow T_{\gamma_x(t)} \Sigma_t$ is the orthogonal projection map.

Lemma 3.10. *The map $M \ni x \mapsto v(x) \in TM$ is well defined and differentiable $\mu_{1/2}$ -a.e. . As a consequence we can find a subset $N \subset \Sigma_{1/2}$ (independent of $t \in [0, 1]$) with $\mu_{1/2}(N) = 0$, such that for every $t \in [0, 1]$ the map $T_{1/2}^t : \Sigma_{1/2} \rightarrow \Sigma_t$ is differentiable at every $x \in \Sigma_{1/2} \setminus N$.*

Moreover, up to replacing N with a larger set of null $\mu_{1/2}$ -measure, for every $x \in \Sigma_{1/2} \setminus N$ the map $t \mapsto D_x T_{1/2}^t : T_x \Sigma_{1/2} \rightarrow T_{T_{1/2}^t(x)} \Sigma_t$ is differentiable at $t = 1/2$ and $D_t|_{t=1/2} D_x T_{1/2}^t : T_x \Sigma_{1/2} \rightarrow T_x \Sigma_{1/2}$ is self-adjoint.

Proof. Step 1: the map $v(\cdot) : \Sigma_{1/2} \setminus N \rightarrow TM$ is well defined.

In a first instance let $N \subset \Sigma_{1/2}$, with $\mu_{1/2}(N) = 0$, be such that for every $x \in \Sigma_{1/2} \setminus N$ the curve $t \mapsto \gamma_x(t) := T_{1/2}^t(x)$ is a well defined geodesic. In particular, the curve $t \mapsto \gamma_x(t)$ is C^1 and we can set $v(x) = \dot{\gamma}_x(\frac{1}{2})$; this is clearly well defined as a map from $\Sigma_{1/2} \setminus N$ to TM . Note moreover that, since by standing assumption μ_0 and μ_1 (and therefore all the measures μ_t) have compact support, we have

$$(12) \quad \mu_{1/2}\text{-ess sup}_{x \in \Sigma_{1/2}} |v(x)| \leq \frac{1}{2} \sup_{(x,y) \in \text{supp } \mu_0 \times \text{supp } \mu_1} d(x,y) =: C_{\mu_0, \mu_1} < \infty.$$

Step 2: the map $v(\cdot) : \Sigma_{1/2} \setminus N \rightarrow TM$ is differentiable.

First of all, note that there exists $\delta > 0$ small enough so that $T_x M \supset B_{C_{\mu_0, \mu_1}}(0) \ni w \mapsto \exp_x(tw)$ is a diffeomorphism onto its image for every $t \in (-\delta, \delta)$ and every $x \in \Sigma_{1/2}$. Fix $x_0 \in \Sigma_{1/2} \setminus N$. Since by Lemma 3.9 the map $T_{1/2}^{(1+\delta)/2}$ is differentiable $\mu_{1/2}$ -a.e., it follows that also the map

$$B_{\delta C_{\mu_0, \mu_1}}(x_0) \cap \Sigma_{1/2} \setminus N \rightarrow TM, \quad x \mapsto v(x) := \frac{2}{\delta} \exp_x^{-1} \left(T_{1/2}^{(1+\delta)/2}(x) \right)$$

is differentiable $\mu_{1/2}$ -a.e.. Therefore, up to redefining the $\mu_{1/2}$ -negligible set N , the claim is proved. Notice that in particular the map

$$(13) \quad T_{1/2}^t(x) = \exp_x(tv(x)) \quad \text{is differentiable everywhere on } \Sigma_{1/2} \setminus N \text{ for every } t \in \left(\frac{1-\delta}{2}, \frac{1+\delta}{2} \right).$$

Step 3: the map $T_{1/2}^t : \Sigma_{1/2} \setminus N \rightarrow \Sigma_t$ is differentiable for every $t \in [0, 1]$.

By construction we have that $T_{1/2}^t(x) = \exp_x(tv(x))$ and, using again that μ_0 and μ_1 have compact support, we know that there exists a compact subset $E \subset \subset M$ such that $T_{1/2}^t(\Sigma_{1/2}) \subset E$ for every $t \in [0, 1]$. In particular, there exists $\delta > 0$ small enough such that, for every $x_0 \in E$, the exponential map

$$\exp_{(\cdot)}(\cdot) : B_{\delta C_{\mu_0, \mu_1}}(x_0) \times B_{\delta C_{\mu_0, \mu_1}}(0) \rightarrow M$$

is smooth, where C_{μ_0, μ_1} was defined in (12).

Let $t_j := \frac{1}{2} + \frac{\delta}{2}j$, for $j = -\lfloor \frac{1}{\delta} \rfloor, \dots, 0, \dots, \lfloor \frac{1}{\delta} \rfloor$, be a $\frac{\delta}{2}$ -grid in $[0, 1]$ centered at $1/2$; for convenience choose $\delta \notin \mathbb{Q}$ so that $\frac{1}{2} + \frac{\delta}{2} \lfloor \frac{1}{\delta} \rfloor < 1$. By repeating the same argument of step 2 and replacing $1/2$ by t_j in (13), we get that for every $j = -\lfloor \frac{1}{\delta} \rfloor, \dots, 0, \dots, \lfloor \frac{1}{\delta} \rfloor$ there exists a subset $N_j \subset \Sigma_{t_j}$ with $\mu_{t_j}(N_j) = 0$ such that $T_{t_j}^t$ is differentiable everywhere on $\Sigma_{t_j} \setminus N_j$ for every $t \in (t_{j-1}, t_{j+1})$.

Since by Lemma 3.9 the maps $T_{t_i}^{t_{i+1}} : \text{supp } \mu_{t_i} \rightarrow \text{supp } \mu_{t_{i+1}}$ are bi-Lipschitz and since μ_t is equivalent to $\mathcal{H}^p \llcorner (\Sigma_t \cap \{\rho_t > 0\})$ for every $t \in [0, 1]$, we get that

$$N_+ := N_0 \cup \left[(T_{1/2}^{t_1})^{-1}(N_1) \right] \cup \left[(T_{t_1}^{t_2} \circ T_{1/2}^{t_1})^{-1}(N_2) \right] \cup \dots \cup \left[(T_{\lfloor \frac{1}{\delta} \rfloor}^{t_{\lfloor \frac{1}{\delta} \rfloor}} \circ \dots \circ T_{t_1}^{t_2} \circ T_{1/2}^{t_1})^{-1}(N_{\lfloor \frac{1}{\delta} \rfloor}) \right]$$

satisfies $\mu_{1/2}(N_+) = 0$. Defining analogously N_- by considering $t_j \leq \frac{1}{2}$ and setting $N = N_+ \cup N_-$ we get that $\mu_{1/2}(N) = 0$.

Fix now an arbitrary $t \in [1/2, 1]$ and let $j_0 := \max\{j : t_j \leq t\}$. Since we can write $T_{1/2}^t = T_{t_{j_0}}^t \circ \dots \circ T_{t_1}^{t_2} \circ$

$T_{1/2}^{t_1}$, it follows that $T_{1/2}^t : \Sigma_{1/2} \setminus N \rightarrow \Sigma_t$ is differentiable everywhere, as composition of differentiable functions. The argument for $t \in [-1, 1/2]$ is completely analogous, so the lemma is proved.

Step 4: the map $t \mapsto D_x T_{1/2}^t : T_x \Sigma_{1/2} \rightarrow T_{T_{1/2}^t(x)} \Sigma_t$ is differentiable at $t = 1/2$ and $D_t|_{t=1/2} D_x T_{1/2}^t : T_x \Sigma_{1/2} \rightarrow T_x \Sigma_{1/2}$ is self-adjoint.

Recall that the transport geodesics $\gamma_x(t) := T_{1/2}^t(x)$ are precisely the gradient flow curves of the corresponding Hamilton-Jacobi shift $\phi_t = H_t \phi$ of a d^2 -Kantorovich potential ϕ from μ_0 to μ_1 . Note in particular that, for $t \in (0, 1]$, ϕ_t is semi-concave. Moreover, $\phi_{1/2}$ is differentiable in x and admits a gradient $\nabla \phi_{1/2}$ in the classical sense with $\nabla \phi_{1/2}(x) = \dot{\gamma}_x(1/2)$ (page 61 in [13]). In particular, $\phi_{1/2}$ is differentiable $\mu_{1/2}$ -a.e. .

By construction, $\nabla \phi_{1/2}$ coincides $\mu_{1/2}$ -a.e. with the vector field v defined above and v is also differentiable on $\Sigma_{1/2} \setminus N$ as vector field along $\Sigma_{1/2}$. For every $x \in \Sigma_{1/2} \setminus N$ denote $A_x : T_x \Sigma_{1/2} \rightarrow T_x \Sigma_{1/2}$, $A_x w := D_w v(x) := (\nabla_w v(x))^\top$. For any fixed $x \in \Sigma_{1/2} \setminus N$, thanks to the semi-concavity of $\phi_{1/2}$ and the differentiability of $v : \Sigma_{1/2} \setminus N \rightarrow TM$ we can follow the proof of [14, Theorem 2.8] to show that the second derivatives of $\phi_{1/2}$ in x tangential to $\Sigma_{1/2}$ exist, and the following Taylor expansion holds for every curve $\lambda(t) \in \Sigma_{1/2}$ with $\lambda(0) = x$ and $\dot{\lambda}(0) = w \in T_x \Sigma_{1/2}$:

$$(14) \quad \phi_{1/2}(\lambda(t)) = \phi_{1/2}(\lambda(0)) + t \langle \nabla \phi_t(\lambda(0)), w \rangle + \frac{t^2}{2} \langle A_x w, w \rangle + o(t^2).$$

Though [14] only considers the case of convex functions in \mathbb{R}^n , it is clear that the proof works as well in the context of Riemannian manifolds and semi-concave functions. Finally, the Taylor expansion (14) implies that A_x must be self-adjoint.

In order to conclude the proof, observe that $(0, 1) \ni t \mapsto DT_{1/2}^t(x)$ is C^1 and that, by the symmetry of second order derivatives of distributions, for $\mu_{1/2}$ -a.e. $x \in \Sigma_{1/2}$ it holds

$$D_t|_{t=1/2} (DT_{1/2}^t(x)) = D(\nabla_t|_{t=1/2} T_{1/2}^t(x)) =: Dw := A_x.$$

□

Remark 3.11. Let $\{\mu_t\}_{t \in [0,1]}$ be a countably \mathcal{H}^p -rectifiable W_2 -geodesic, $s \in (0, 1)$, and $x \in \Sigma_s \setminus N$ where $N \subset \Sigma_s$ with $\mu_s(N) = 0$ is given by Lemma 3.10. Since a countably \mathcal{H}^p -rectifiable set has p -dimensional euclidean tangent spaces \mathcal{H}^p -a.e., without loss of generality we can assume that for every $x \in \Sigma_s \setminus N$ it holds $\dim T_x \Sigma_s = p$. Choose an orthonormal basis (e_1, \dots, e_p) of $T_x \Sigma_s$ and consider the vector fields $J_1, \dots, J_p : [0, 1] \rightarrow T_{\gamma_x(t)} M$ along the geodesic $\gamma_x : [0, 1] \rightarrow M$ defined by

$$J_i(t) := (DT_s^t(x)) [e_i] = \left(D(\exp_{(\cdot)}(tv(\cdot)))(x) \right) [e_i], \quad \forall i = 1, \dots, p, \quad \forall t \in [0, 1]$$

where $v(x)$ was defined before in Lemma 3.10. A standard computation of Riemannian geometry shows that the map $t \mapsto J_i(t)$ satisfies the Jacobi equation

$$\nabla_t \nabla_t J_i + R(\dot{\gamma}_x, J_i) \dot{\gamma}_x = 0, \quad \forall i = 1, \dots, p, \quad \text{on } [0, 1],$$

where ∇_t is the covariant derivative of vector fields along γ_x at the point $\gamma_x(t)$. In other words, J_i is a Jacobi field. We then set

$$(15) \quad B_x(t) : T_x \Sigma_s \rightarrow T_{\gamma_x(t)} M, \quad B_x(t) := DT_s^t(x) \quad \forall t \in [0, 1], \quad \forall x \in \Sigma_s \setminus N.$$

The combination of Lemma 3.9 and Lemma 3.10 yields that $B_x(t)$ is non-degenerate for every $t \in [0, 1]$ for μ_s -almost every $x \in \Sigma_s$. So in particular, for μ_s -a.e. x we have that $\dim \text{Im}[DT_s^t(x)] = p$ and $\{J_i(t)\}_{i=1, \dots, p}$ is a basis of $\text{Im}[DT_s^t(x)]$ for every $t \in [0, 1]$. We can (and will) consider $B_x(t)$ as a map from $T_x \Sigma_s$ to $T_{\gamma_x(t)} \Sigma_t$. Finally, we also proved that $D_t B_x(t)|_{t=s} : T_x \Sigma_s \rightarrow T_x \Sigma_s$ is self-adjoint for μ_s -almost every $x \in \Sigma_s$.

4. JACOBI FIELDS COMPUTATIONS

Let (M, g) be a complete Riemannian manifold without boundary, and let $\gamma : [0, 1] \rightarrow M$ be a minimizing, constant speed geodesic with $\gamma(0) = x$. Moreover, let $\{e_i\}_{i=1, \dots, p}$ be orthonormal vectors in $T_x M$, and let $J_{e_i} : [0, 1] \rightarrow TM$ be non-vanishing Jacobi fields along γ with $J_i(0) = e_i$ and $J_i'(0) = f_i$, for some

$f_i \in T_x M$ to be specified later. We denote with $T_{\gamma(t)\Sigma_t} \subset T_{\gamma(t)}M$ the span of $\{J_{e_i}(t)\}_{i=1,\dots,p}$ for each $t \in [0, 1]$, and with v^\top the orthogonal projection of a vector $v \in T_{\gamma(t)}M$ to the subspace $T_{\gamma(t)\Sigma_t}$. Similarly, v^\perp is its projection to the orthogonal complement $(T_{\gamma(t)\Sigma_t})^\perp$ of $T_{\gamma(t)\Sigma_t}$. We also denote with $\top : T_{\gamma(t)}M \rightarrow T_{\gamma(t)\Sigma_t}$ the orthogonal projection map.

Lemma 4.1. *Define the vector fields $E_i : [0, 1] \rightarrow T_{\gamma(t)\Sigma_t}$, $i = 1, \dots, p$, along γ with values in $\bigcup_{t \in [0,1]} T_{\gamma(t)\Sigma_t}$ as the solution of*

$$(16) \quad (\nabla_t E_i)^\top = 0, \quad \text{with } E_i(0) = e_i,$$

where ∇_t is the covariant derivative of vector fields along γ at the point $\gamma(t)$. Then $\{E_i(t)\}_{i=1,\dots,p}$ is an orthonormal basis for $T_{\gamma(t)\Sigma_t}$ for every $t \in [0, 1]$.

Proof. The existence and uniqueness of $E_i : [0, 1] \rightarrow T_{\gamma(t)\Sigma_t}$ solving (16) is standard as it corresponds to solve a system of first order linear homogeneous ODEs with Cauchy conditions. By definition of E_i , $i = 1, \dots, p$ we have

$$\frac{d}{dt} \langle E_i, E_j \rangle = \langle \nabla_t E_i, E_j \rangle + \langle E_i, \nabla_t E_j \rangle = \langle (\nabla_t E_i)^\top, E_j \rangle + \langle E_i, (\nabla_t E_j)^\top \rangle = 0.$$

Hence, $\langle E_i, E_j \rangle$ is constant along γ , and since $E_i(0) = e_i$, $i = 1, \dots, p$, is an orthonormal basis of $T_{\gamma_0}\Sigma_0$ the claim follows. \square

In the following we denote $D_t := \top \circ \nabla_t$. For E_i as in Lemma 4.1, by construction we have $D_t E_i = 0$.

Let $B(t) : T_{\gamma(0)\Sigma_0} \rightarrow T_{\gamma(t)}M$ be the 1-parameter family of linear maps defined via $B(t)e_i = J_{e_i}(t)$, and consider $\nabla_t B(t) : T_{\gamma(0)\Sigma_0} \rightarrow T_{\gamma(t)}M$ given by $(\nabla_t B(t))e_i = \nabla_t J_{e_i}$. If we consider $B(t)$ as a map from $T_{\gamma(0)\Sigma_0}$ to $T_{\gamma(t)\Sigma_t}$, its derivative $D_t B(t)$ defined by $[D_t B(t)]e_i = D_t J_{e_i}$ is a map from $T_{\gamma_0}\Sigma_0$ to $T_{\gamma(t)\Sigma_t}$ as well. Moreover, since $\{J_{e_i}\}_{i=1,\dots,p}$ are Jacobi fields in M , the Jacobi equation yields

$$(17) \quad \nabla_t \nabla_t B(t) + R(\dot{\gamma}_t, B(t))\dot{\gamma}_t = 0.$$

In the rest of the section we are going to work under the assumption that $B(t) : T_{\gamma(0)\Sigma_0} \rightarrow T_{\gamma(t)\Sigma_t}$ is invertible for all $t \in [0, 1]$, in fact that will be satisfied in the optimal transport application of the next section thanks to Lemma 3.9. It will be convenient to consider the operators:

$$(18) \quad \mathcal{U}(t) := (\nabla_t B(t))B(t)^{-1} : T_{\gamma(t)\Sigma_t} \rightarrow T_{\gamma(t)}M$$

$$(19) \quad \mathcal{U}^\top(t) := [\mathcal{U}(t)]^\top = (D_t B(t))B(t)^{-1} : T_{\gamma(t)\Sigma_t} \rightarrow T_{\gamma(t)\Sigma_t}.$$

$$(20) \quad \mathcal{U}^\perp(t) := [\mathcal{U}(t)]^\perp : T_{\gamma(t)\Sigma_t} \rightarrow (T_{\gamma(t)\Sigma_t})^\perp.$$

Lemma 4.2. *Let $J_i := J_{e_i}$ and E_i , $i = 1, \dots, p$ be as above. Then*

$$T_{\gamma(t)\Sigma_t}^\perp \ni \nabla_t E_i(t) = \mathcal{U}^\perp(t)E_i.$$

Proof. First, we write $J_i = \sum_{j=1}^p \langle J_i, E_j \rangle E_j$ and set $A_{ij} = \langle J_i, E_j \rangle$ where the matrix $A := (A_{ij})_{i,j} \in GL_n(\mathbb{R})$. Let A^{-1} be its inverse. We compute

$$\nabla_t J_i(t) = \sum_{j=1}^p \langle \nabla_t J_i(t), E_j(t) \rangle E_j(t) + \sum_{j=1}^p \langle J_i(t), \nabla_t E_j(t) \rangle E_j(t) + \sum_{j=1}^p \langle J_i(t), E_j(t) \rangle \nabla_t E_j(t),$$

where the second sum on the right hand side vanishes since $(\nabla_t E_j(t))^\top = 0$. Rearranging terms and multiplying by A^{-1} yields for $k = 1, \dots, p$

$$\begin{aligned} \sum_{i=1}^p (A^{-1})_{ki} (\nabla_t J_i(t))^\perp &= \sum_{i=1}^p (A^{-1})_{ki} \left[\nabla_t J_i(t) - \sum_{j=1}^p \langle \nabla_t J_i(t), E_j(t) \rangle E_j(t) \right] \\ &= \sum_{i=1}^p (A^{-1})_{ki} \left[\sum_{j=1}^p \langle J_i(t), E_j(t) \rangle \nabla_t E_j(t) \right] = \sum_{i=1}^p \sum_{j=1}^p (A^{-1})_{ki} A_{ij} \nabla_t E_j(t) = \nabla_t E_k(t). \end{aligned}$$

Now, we recall that $\nabla_t J_i = \nabla_t J_{E_i} = \nabla_t J_{B^{-1}(t)J_i} = \mathcal{U}(t)J_i$. Therefore

$$\nabla_t E_k(t) = \sum_{i=1}^p (A^{-1})_{ki} (\nabla_t J_i(t))^\perp = \sum_{i=1}^p (A^{-1})_{ki} [\mathcal{U}(t)J_i]^\perp = \left[\mathcal{U}(t) \left(\sum_{i=1}^p (A^{-1})_{ki} J_i \right) \right]^\perp = [\mathcal{U}(t)E_k(t)]^\perp,$$

as desired. \square

Lemma 4.3. *Let $B(t) : T_{\gamma(0)}\Sigma_0 \rightarrow T_{\gamma(t)}\Sigma_t$, $t \in [0, 1]$, be as above, and $B(t)^{-1} : T_{\gamma(t)}\Sigma_t \rightarrow T_{\gamma(0)}\Sigma_0$. Then*

$$(21) \quad D_t[B(t)^{-1}] = -B(t)^{-1}(D_t B(t))B(t)^{-1}.$$

Proof. Let $\{E_i\}_{i=1, \dots, p}$ be as in the previous lemma. Then, we obviously have $B(t)B(t)^{-1}E_i(t) = E_i(t)$ for any $i = 1, \dots, p$. Applying D_t yields

$$(D_t B(t))B(t)^{-1}E_i(t) + B(t)(D_t(B(t)^{-1}))E_i(t) + B(t)B(t)^{-1}(D_t E_i(t)) = D_t E_i(t) = 0.$$

Rearranging the terms and applying $B(t)^{-1}$ from the left of both sides yields the claim. \square

The next proposition expresses the “ p -dimensional volume distortion” along the geodesic γ in terms of the p -Ricci curvature and will be crucial for proving the characterization of lower curvature bounds in terms of optimal transport in the next section.

Proposition 4.4. *Let $\mathcal{U}(t), \mathcal{U}(t)^\top, \mathcal{U}(t)^\perp$ be defined in (18), (19), (20). Then it holds*

$$\nabla_t \mathcal{U}(t) + \mathcal{U}(t)\mathcal{U}^\top(t) + R(\dot{\gamma}_t, \cdot)\dot{\gamma}_t = 0.$$

Taking the trace along $T_{\gamma(t)}\Sigma_t$ yields

$$\text{tr}(D_t \mathcal{U}(t)) + \text{tr}((\mathcal{U}^\top(t))^2) + \text{Ric}_p(T_{\gamma(t)}\Sigma_t, \dot{\gamma}) = 0,$$

and moreover

$$(22) \quad \text{tr}(\mathcal{U}^\top(t))' + \text{tr}((\mathcal{U}^\top(t))^2) + \text{Ric}_p(T_{\gamma(t)}\Sigma_t, \dot{\gamma}(t)) = \|\mathcal{U}^\perp(t)\|^2.$$

If $D_t B(t)|_{t=0} : T_{\gamma(0)}\Sigma_0 \rightarrow T_{\gamma(0)}\Sigma_0$ is self-adjoint then $\mathcal{U}^\top(t) : T_{\gamma(t)}\Sigma_t \rightarrow T_{\gamma(t)}\Sigma_t$ is self-adjoint for all $t \in [0, 1]$ and, setting $y(t) = \log \det B(t)$, it holds

$$(23) \quad y''(t) + \frac{1}{p}y'(t)^2 + \text{Ric}_p(T_{\gamma(t)}\Sigma_t, \dot{\gamma}(t)) - \|\mathcal{U}^\perp(t)\|^2 \leq 0.$$

Remark 4.5. In case $p = \dim(M)$ then $\mathcal{U}^\top(t) = \mathcal{U}(t)$, $\mathcal{U}^\perp(t) = 0$ and $\text{Ric}_p(T_{\gamma(t)}\Sigma_t, \dot{\gamma}(t)) = \text{Ric}(\dot{\gamma}(t), \dot{\gamma}(t))$, so that Proposition 4.4 recovers the classical Jacobian estimates expressing the volume distortion along a geodesic in terms of Ricci curvature (see for instance [4, Lemma 3.1]).

Proof. First of all, there is a natural extension of $B(t)$ (and of $\nabla_t B(t)$) to maps from the whole $T_{\gamma(0)}M$ just by composing with the orthogonal projection into $T_{\gamma(0)}\Sigma_0$, i.e. for $v \in T_{\gamma(0)}M$ we consider $B(t)v^\top$. Differentiating the identity $\top \circ \top = \top$ gives $\nabla_t \top \circ \top + \top \circ \nabla_t \top = \nabla_t \top$; left and right composing with \top , yields $\top \circ \nabla_t \top \circ \top = 0$. Therefore, using (17) and (21), we get

$$\begin{aligned} \nabla_t [\mathcal{U}(t)] &= [\nabla_t \nabla_t B(t)] B(t)^{-1} + \nabla_t B(t)(\top \circ \nabla_t \top \circ \top)B(t)^{-1} + \nabla_t B(t)[\nabla_t B(t)^{-1}]^\top \\ &= -R(\dot{\gamma}(t), \cdot)\dot{\gamma}(t) + \nabla_t B(t)[D_t B(t)^{-1}] = -R(\dot{\gamma}(t), \cdot)\dot{\gamma}(t) - \nabla_t B(t)B(t)^{-1}D_t B(t)B(t)^{-1} \\ &= -R(\dot{\gamma}(t), \cdot)\dot{\gamma}(t) - \mathcal{U}(t)\mathcal{U}^\top(t). \end{aligned}$$

Taking the trace along $T_{\gamma(t)}\Sigma_t$ yields the second identity. To get the identity (22), observe that $\text{tr} \mathcal{U}^\top(t) = \sum_{i=1}^p \langle \mathcal{U}(t)E_i(t), E_i(t) \rangle$ and

$$\langle \mathcal{U}(t)E_i(t), E_i(t) \rangle' = \langle [D_t \mathcal{U}(t)] E_i(t), E_i(t) \rangle + \langle \mathcal{U}(t) [D_t E_i(t)], E_i(t) \rangle + \langle \mathcal{U}(t)E_i(t), \nabla_t E_i(t) \rangle.$$

Since $D_t E_i = (\nabla_t E_i)^\top = 0$ and $\nabla_t E_i = (\nabla_t E_i)^\perp$, we conclude that

$$(\text{tr} \mathcal{U}^\top(t))' = \text{tr}(D_t \mathcal{U}(t)) + \sum_{i=1}^p \langle \mathcal{U}(t)E_i(t), \nabla_t E_i(t) \rangle = \text{tr}(D_t \mathcal{U}(t)) + \sum_{i=1}^p \langle (\mathcal{U}(t)E_i(t))^\perp, \nabla_t E_i(t) \rangle.$$

In particular let us explicitly observe that, in general, $\text{tr}(D_t \mathcal{U}(t)) \neq \text{tr}(\mathcal{U}^\top(t))'$. The claimed identity (22) follows by observing that $\|\mathcal{U}(t)^\perp\|^2 = \sum_{i=1}^p \langle (\mathcal{U}(t)E_i(t))^\perp, \nabla_t E_i(t) \rangle$.

The rest of the proof is devoted to show (23). Setting $y(t) = \log \det B(t)$, we have that

$$\begin{aligned} y'(t_0) &= \frac{d}{dt} \Big|_{t=t_0} \log \det (B(t)B(t_0)^{-1}) = \frac{d}{dt} \Big|_{t=t_0} \log \det \left[\langle (B(t)B(t_0)^{-1}E_i(t), E_j(t)) \rangle_{i,j} \right] \\ &= \text{tr} \left[(D_t B(t))B(t_0)^{-1} \right] \Big|_{t=t_0} + 2 \sum_{i=1}^p \langle D_t E_i(t), E_i(t) \rangle \Big|_{t=t_0} \\ (24) \quad &= \text{tr} \left[(D_t B(t))B(t_0)^{-1} \right] \Big|_{t=t_0} = \text{tr}(\mathcal{U}^\top(t_0)), \end{aligned}$$

since by construction $D_t E_i(t) = 0$.

We next claim that, under the assumption that $D_t B(t)|_{t=0}$ is self-adjoint, then

$$(25) \quad \mathcal{U}^\top(t) : T_{\gamma(t)}\Sigma_t \rightarrow T_{\gamma(t)}\Sigma_t \quad \text{is self-adjoint for all } t \in [0, 1].$$

To this aim, calling $(\mathcal{U}^\top(t))^*$ the adjoint operator, we observe that

$$(26) \quad (\mathcal{U}^\top(t))^* - \mathcal{U}^\top(t) = (B(t)^*)^{-1} [(D_t B(t)^*)B(t) - B(t)^*(D_t B(t))] B(t)^{-1},$$

and that

$$(27) \quad D_t [(D_t B(t)^*)B(t) - B(t)^*(D_t B(t))] = (D_t^2 B(t)^*)B(t) - B(t)^*(D_t^2 B(t)).$$

Now, combining the Jacobi equation (17) with the identity $\top \circ \nabla_t \top \circ \top = 0$ proved at the beginning of the proof, we have

$$(28) \quad \begin{aligned} D_t^2 B(t) &= \top \nabla_t (\top \nabla_t B(t)) = \top (\nabla_t^2 B(t)) + \top (\nabla_t \top) \top \nabla_t B(t) = \top (\nabla_t^2 B(t)) \\ &= - (R(\dot{\gamma}(t), B(t))\dot{\gamma}(t))^\top = -\mathcal{R}(t)B(t), \end{aligned}$$

where

$$(29) \quad \mathcal{R}(t) : T_{\gamma(t)}\Sigma_t \rightarrow T_{\gamma(t)}\Sigma_t, \quad \mathcal{R}(t)[v] := [R(\dot{\gamma}(t), v)\dot{\gamma}(t)]^\top$$

is self-adjoint; indeed, in the orthonormal basis $\{E_i(t)\}_{i=1, \dots, p}$, it is represented by the symmetric matrix $\langle R(\dot{\gamma}(t), E_i(t))\dot{\gamma}(t), E_j(t) \rangle$. Plugging (28) into (27), we obtain that $(D_t B(t)^*)B(t) - B(t)^*(D_t B(t))$ is constant in t and thus vanishes identically, since by assumption $B(0) = \text{Id}$ and $D_t B(t)|_{t=0}$ is self-adjoint. Taking into account (26), this concludes the proof of the claim (25).

Using that $\mathcal{U}^\top(t)$ is a self-adjoint operator over a p -dimensional space, by Cauchy-Schwartz inequality, we have that

$$(30) \quad \text{tr} [(\mathcal{U}^\top(t))^2] \geq \frac{1}{p} (\text{tr} [\mathcal{U}^\top(t)])^2.$$

The desired estimate (23) then follows from the combination of (22), (24) and (30). \square

In the final part of the section we specialize to the case $p = 1$, giving the self-contained easier arguments.

Proposition 4.6. *Assume $p = 1$, let $J := J_{e_1}$ and $E := E_1$ be as above. In particular, $\dim T_{\gamma(t)}\Sigma_t = 1$ for every $t \in [0, 1]$, and $E = |J(t)|^{-1}J(t)$. Then*

$$(31) \quad \nabla_t E(t) = |J(t)|^{-1}(\nabla_t J(t))^\perp.$$

Proof. We compute $\nabla_t E$ as follows

$$\nabla_t E(t) = (|J(t)|^{-1})' J(t) + |J(t)|^{-1} \nabla_t J(t).$$

Since

$$(|J|^{-1})' = \left(\langle J, J \rangle^{-\frac{1}{2}} \right)' = -|J|^{-3} \langle J, \nabla_t J \rangle = -|J|^{-2} \langle E, \nabla_t J \rangle,$$

we get

$$\nabla_t E = -|J|^{-1} \langle E, \nabla_t J \rangle E + |J|^{-1} \nabla_t J = |J|^{-1} (\nabla_t J)^\perp.$$

\square

Corollary 4.7. *Assume $p = 1$, and consider $\mathcal{U}^\top(t)$ as above. Then, we have*

$$\langle \mathcal{U}^\top(t)E(t), E(t) \rangle' + \langle \mathcal{U}^\top(t)E(t), E(t) \rangle^2 + \text{Ric}_1(T_{\gamma(t)}\Sigma_t, \dot{\gamma}_t) = |\mathcal{U}(t)E(t)^\perp|^2.$$

Proof. Since

$$\text{tr}((\mathcal{U}^\top)^2) = \langle (\mathcal{U}^\top)^2(t)E(t), E(t) \rangle = \langle \mathcal{U}^\top(t)E(t), E(t) \rangle^2,$$

and, from (31), we have

$$(32) \quad \nabla_t E(t) = |J(t)|^{-1}(\nabla_t J(t))^\perp = |J(t)|^{-1}(\nabla_t J_{B(t)^{-1}J(t)})^\perp = |J(t)|^{-1}(\mathcal{U}(t)J(t))^\perp = (\mathcal{U}(t)E(t))^\perp,$$

the claim follows from (22). \square

5. OT CHARACTERIZATION OF SECTIONAL CURVATURE UPPER BOUNDS

Proposition 5.1. *Let M be a complete Riemannian manifold without boundary with $\text{Ric}_1 \leq K$, let $\{\mu_t\}_{t \in [0,1]}$ be a countably \mathcal{H}^1 -rectifiable W_2 -geodesic, and consider $B_x(t) : T_x \Sigma_0 \rightarrow T_{\gamma(t)} \Sigma_t$, $t \in [0, 1]$ as in Remark 3.11. Then, the function $[0, 1] \ni t \mapsto \mathcal{J}_x(t) := \det B_x(t) \in \mathbb{R}$ belongs to $C([0, 1]) \cap C^2((0, 1))$ and satisfies*

$$(33) \quad \mathcal{J}_x'' + K|\dot{\gamma}|^2 \mathcal{J}_x \geq 0 \quad \text{on } (0, 1).$$

In particular, if $t_0, t_1 \in [0, 1]$, $\tau(s) = s(t_1 - t_0) + t_0$, and $s \in [0, 1] \mapsto \varsigma_s = \gamma_{\tau(s)}$, we have for all $s \in [0, 1]$

$$(34) \quad \mathcal{J}_x(\tau(s)) \leq \sigma_{K,1}^{(1-s)}(|\dot{\varsigma}|) \mathcal{J}_x(t_0) + \sigma_{K,1}^{(s)}(|\dot{\varsigma}|) \mathcal{J}_x(t_1).$$

Proof. First note that, setting $t \in [0, 1] \mapsto y_x(t) := \log \mathcal{J}_x(t)$ we have that $y_x'(t) = \text{tr}((D_t B_x(t))B_x^{-1}(t)) = \text{tr} \mathcal{U}_x(t) = \text{tr} \mathcal{U}_x(t)^\top$. Then Corollary 4.7 yields that

$$y_x'' + (y_x')^2 + K|\dot{\gamma}_x|^2 \geq 0 \quad \text{on } (0, 1).$$

By computing $\mathcal{J}_x''(t) = (e^{y_x})''(t)$ this yields (33). Moreover, considering t_0, t_1, τ and ς as above we get

$$\frac{d^2}{ds^2} \mathcal{J}_x \circ \tau + K|\dot{\varsigma}|^2 \mathcal{J}_x \circ \tau \geq 0 \quad \text{on } (0, 1).$$

that is equivalent to (34) by classical comparison principle. \square

Theorem 5.2 (Curvature upper bounds). *Let (M, g) be a complete Riemannian manifold without boundary and let $K \geq 0$. Then the following statements (i) and (ii) are equivalent:*

- (i) $\text{Ric}_1 \leq K$ or, equivalently, $\text{Sec} \leq K$.
- (ii) *Let $\{\mu_t\}_{t \in [0,1]}$ be a countably \mathcal{H}^1 -rectifiable W_2 -geodesic, and let Π be the corresponding dynamical transport plan. Then, if $t_0, t_1 \in (0, 1)$ and $\tau(s) = (1-s)t_0 + st_1$, it holds*

$$\mathcal{H}^1(\text{supp } \mu_{\tau(s)}) \leq \int \left[\sigma_{K,1}^{(1-s)}(|\dot{\gamma} \circ \tau|) \rho_{t_0}(\gamma(t_0))^{-1} + \sigma_{K,1}^{(s)}(|\dot{\gamma} \circ \tau|) \rho_{t_1}(\gamma(t_1))^{-1} \right] d\Pi(\gamma), \quad \forall s \in [0, 1],$$

where ρ_t is the density of μ_t w.r.t. \mathcal{H}^1 .

In the case of $K = 0$ the inequality in (ii) becomes

$$\mathcal{H}^1(\text{supp } \mu_{\tau(s)}) \leq (1-s)\mathcal{H}^1(\text{supp } \mu_{t_0}) + s\mathcal{H}^1(\text{supp } \mu_{t_1}), \quad \forall s \in [0, 1].$$

Remark 5.3. Recall from Remark 2.3 that the condition $\text{Ric}_1 \leq K < 0$ is never satisfied as $\text{Ric}_1(\mathbb{R}v, v) = 0$ for every $v \in TM$; hence it makes sense just to assume a non-negative upper bound $K \geq 0$ and, in this case, $\text{Ric}_1 \leq K$ is equivalent to $\text{Sec} \leq K$.

Remark 5.4. In the assertion (ii) of Theorem 5.2, one cannot relax the assumption to $t_0, t_1 \in [0, 1]$. For instance, one can consider a cylinder $\mathbb{R} \times \mathbb{S}^1$ that is a space of zero (in particular non-positive) sectional curvature. Parametrize \mathbb{S}^1 by arclength on $[0, 2\pi]$, in particular 0 and π are two antipodal points in \mathbb{S}^1 . Then, the uniform distribution on the set of all geodesics connecting $(s, 0)$ and (s, π) for $s \in [0, 1]$ defines a countably \mathcal{H}^1 -rectifiable W_2 -geodesic $\{\mu_t\}_{t \in [0,1]}$ such that $\text{supp } \mu_0 = [0, 1] \times \{0\}$, $\text{supp } \mu_1 = [0, 1] \times \{\pi\}$, $\text{supp } \mu_{1/2} = [0, 1] \times \{\pi/2\} \cup [0, 1] \times \{3\pi/2\}$. Hence, we have $\mathcal{H}^1(\text{supp } \mu_{1/2}) = 2$, $\mathcal{H}^1(\text{supp } \mu_0) = \mathcal{H}^1(\text{supp } \mu_1) = 1$.

Proof. (i) \implies (ii). Let $\{\mu_t\}_{t \in [0,1]}$ be a countably \mathcal{H}^1 -rectifiable W_2 -geodesic, i.e. for every $t \in [0,1]$ the probability measure μ_t is concentrated on a countably \mathcal{H}^1 -rectifiable set $\Sigma_t \subset M$ and is $\mathcal{H}^1|_{\Sigma_t}$ -absolutely continuous. Also, thanks to Theorem 3.1 (see also Remark 3.2), the W_2 -geodesic $\{\mu_t\}_{t \in [0,1]}$ is given by Lipschitz optimal transport maps; more precisely there exist unique Lipschitz maps $T_{1/2}^t : \Sigma_{1/2} \rightarrow \Sigma_t$ such that $(T_{1/2}^t)_\# \mu_{1/2} = \mu_t$ and $\pi_{0,1} := (T_{1/2}^0, T_{1/2}^1)_\# \mu_{1/2}$ is an optimal coupling between μ_0 and μ_1 . Let $\gamma_x(t) := T_{1/2}^t(x) \in \text{Geo}(M)$ and $\gamma_x \circ \tau(s) =: \varsigma_x(s)$.

The map $x \mapsto (t \mapsto T_{1/2}^t(x)) =: \gamma_x \in \text{Geo}(X)$ yields a measurable map from M into the space of geodesics $\text{Geo}(X)$, and the push-forward of $\mu_{1/2}$ under this map is the associated optimal dynamical transport plan Π . In particular

$$(35) \quad \int f(\gamma) d\Pi(\gamma) = \int f(\gamma_x) d\mu_{1/2}(x)$$

for any non-negative measurable function $f : \text{Geo}(X) \rightarrow [0, \infty]$.

Setting $\mathcal{J}_x(t) := \det[DT_{1/2}^t(x)]$ for $\mu_{1/2}$ -a.e. $x \in \Sigma_{1/2}$ and making use of Lemma 3.9 and Proposition 5.1, we can compute for every $s \in [0,1]$:

$$\begin{aligned} \int_{\Sigma_{\tau(s)}} \rho_{\tau(s)}(x)^{-1} d\mu_{\tau(s)}(x) &= \int_{\Sigma_{1/2}} \rho_{\tau(s)}(T_{1/2}^{\tau(s)}(y))^{-1} d\mu_{1/2}(y) \\ &\stackrel{(9)}{=} \int_{\Sigma_{1/2}} \mathcal{J}_x(\tau(s)) d\mathcal{H}^1(x) \\ &\stackrel{(34)}{\leq} \int_{\Sigma_{1/2}} \left[\sigma_{K,1}^{(1-s)}(|\dot{\varsigma}_x|) \mathcal{J}_x(t_0) + \sigma_{K,1}^{(s)}(|\dot{\varsigma}_x|) \mathcal{J}_x(t_1) \right] d\mathcal{H}^1(x) \\ &\stackrel{(9)}{=} \int_{\Sigma_{1/2}} \left[\sigma_{K,1}^{(1-s)}(|\dot{\varsigma}_x|) \rho_{t_0}(T_{1/2}^{t_0}(x))^{-1} + \sigma_{K,1}^{(s)}(|\dot{\varsigma}_x|) \rho_{t_1}(T_{1/2}^{t_1}(x))^{-1} \right] \rho_{1/2}(x) d\mathcal{H}^1(x) \\ &\stackrel{(35)}{=} \int \left[\sigma_{K,1}^{(1-s)}(|\gamma \dot{\circ} \tau|) \rho_{t_0}(\gamma(t_0))^{-1} + \sigma_{K,1}^{(s)}(|\gamma \dot{\circ} \tau|) \rho_{t_1}(\gamma(t_1))^{-1} \right] d\Pi(\gamma). \end{aligned}$$

Note that the assumption $t_0, t_1 \in (0,1)$ was used above in order to apply (9) with equality.

(ii) \implies (i).

We argue by contradiction. Assume there exist $x_0 \in M$, a line $P \subset T_{x_0}M$ and $0 \neq v \in T_{x_0}M$ such that the 1-Ricci curvature of P in the direction of v satisfies

$$(36) \quad \text{Ric}_1(P, v) > (K + 3\epsilon)|v|^2,$$

for some $\epsilon > 0$. Let $\delta > 0$ be sufficiently small such that $\exp_{x_0}|_{B_\delta(0)}$ is a diffeomorphism onto its image. Then $\exp_{x_0}(P \cap B_\delta(0)) =: \Sigma_{\frac{1}{2}}$ is a smooth 1-dimensional submanifold. Let $\phi \in C_0^\infty(M)$ be a Kantorovich potential such that

$$(37) \quad \nabla\phi(x_0) = v \neq 0 \quad \text{and} \quad \nabla^2\phi(x_0) = 0.$$

By replacing ϕ with $\eta\phi$ for a sufficiently small number $\eta > 0$ we get that ϕ is a Kantorovich potential as well and $|\nabla\phi|(y)$ is smaller than the injectivity radius at y , for every $y \in \text{supp}(\phi) \subset M$. It is easily checked that for $\delta > 0$ small enough the map $y \mapsto T_t(y) = \exp_y(-t\nabla\phi(y))$ is a diffeomorphism from $B_\delta(0)$ onto its image for any $t \in [-\frac{1}{2}, \frac{1}{2}]$. Hence, $\Sigma_t := T_{t-\frac{1}{2}}(\Sigma_{\frac{1}{2}})$ for $t \in [0,1]$ is a 1-parameter family of smooth 1-dimensional submanifolds with finite 1-dimensional Hausdorff measure. We define $\mu_{\frac{1}{2}} := \mathcal{H}^1(\Sigma_{\frac{1}{2}})^{-1} \mathcal{H}^1 \llcorner \Sigma_{\frac{1}{2}}$; note that $\mu_t := (T_{t-\frac{1}{2}})_\# \mu_{\frac{1}{2}}$, with $t \in [0,1]$, is the unique L^2 -Wasserstein geodesic between μ_0 and μ_1 . Moreover, by construction, μ_t is a \mathcal{H}^1 -absolutely continuous probability measure concentrated on Σ_t .

Calling $\gamma_x(t) := T_{t-\frac{1}{2}}(x) = \exp_x(- (t - \frac{1}{2})\nabla\phi(x))$ for $x \in \Sigma_{\frac{1}{2}}$ the geodesic performing the transport, note that by continuity there exist $\delta, \sigma > 0$ small enough such that

$$(38) \quad \text{Ric}_1(T_{\gamma_x(t)}\Sigma_t, \dot{\gamma}_x(t)) > (K + 2\epsilon)|\dot{\gamma}_x(t)|^2, \quad \forall x \in \Sigma_{\frac{1}{2}} \subset B_\delta(x_0), \quad \forall t \in \left[\frac{1}{2} - \sigma, \frac{1}{2} + \sigma \right].$$

For every $x \in \Sigma_{\frac{1}{2}}$ note that $\gamma_x(t) := T_{t-\frac{1}{2}}(x)$ is a geodesic connecting $T_{-\frac{1}{2}}(x) \in \Sigma_0$ to $T_{\frac{1}{2}}(x) \in \Sigma_1$. Choose $e \in T_x \Sigma_{\frac{1}{2}}$, consider the Jacobi field $J : [0, 1] \rightarrow T_{\gamma_x(t)}M$ such that $J(\frac{1}{2}) = e$ and $J'(\frac{1}{2}) = [\nabla^2 \phi(x)]e$, and set $|J(t)|^{-1}J(t) = E(t)$. We introduce again the linear operator $\mathcal{U}^\top(t) = D_t B_x(t)B_x(1)^{-1}$ where $B_x(t) : T_x \Sigma_{\frac{1}{2}} \rightarrow T_{\gamma_x(t)}M$ by $B_x(t)e := J(t) = DT_{t-\frac{1}{2}}(x)e$. Then, as in Corollary 4.7, we get

$$(39) \quad \langle \mathcal{U}_x^\top(t)E(t), E(t) \rangle' + \langle \mathcal{U}_x^\top(t)E(t), E(t) \rangle^2 + \text{Ric}_1(T_{\gamma(t)}\Sigma_t, \dot{\gamma}_t) = |(\mathcal{U}_x(t)E(t))^\perp|^2 \leq |\mathcal{U}_x(t)E(t)|^2.$$

Since by construction $\mathcal{U}_{x_0}(0) := \nabla_t B_{x_0}(0)B_{x_0}^{-1}(0) = \nabla^2 \phi(x_0)|_{T_{x_0}\Sigma_{\frac{1}{2}}} = 0$ and $v \neq 0$, again by continuity we can choose $\delta, \sigma > 0$ even smaller so that

$$(40) \quad |\mathcal{U}_x(t)E(t)|^2 < \epsilon |\dot{\gamma}_x(t)|^2, \quad \forall x \in \Sigma_{\frac{1}{2}} \subset B_\delta(x_0), \quad \forall t \in \left[\frac{1}{2} - \sigma, \frac{1}{2} + \sigma \right].$$

The combination of (38), (39) and (40) then yields

$$0 > \langle \mathcal{U}^\top(t)E(t), E(t) \rangle' + \langle \mathcal{U}^\top(t)E(t), E(t) \rangle^2 + (K + \epsilon) |\dot{\gamma}_x(t)|^2, \quad \forall x \in \Sigma_{\frac{1}{2}}, \quad \forall t \in \left[\frac{1}{2} - \sigma, \frac{1}{2} + \sigma \right].$$

Observe that the affine reparametrization $t = g(s) = \frac{1}{2} - \sigma + 2\sigma s$, $g : [0, 1] \rightarrow [\frac{1}{2} - \sigma, \frac{1}{2} + \sigma]$, corresponds to consider the rescaled Kantorovich potential $2\sigma\phi$ in place of ϕ in the arguments above, and thus gives

$$0 > \langle \mathcal{U}^\top(g(s))E(g(s)), E(g(s)) \rangle' + \langle \mathcal{U}^\top(g(s))E(g(s)), E(g(s)) \rangle^2 + (K + \epsilon) |\dot{\gamma}_x(g(s))|^2 \\ \forall x \in \Sigma_{\frac{1}{2}}, \quad \forall s \in [0, 1].$$

Since g is affine, the restricted and rescaled curve $\{\tilde{\mu}_s := \mu_{g(s)}\}_{s \in [0, 1]}$ is still a W_2 -geodesic from $\tilde{\mu}_0 = \mu_{\frac{1}{2}-\sigma}$ to $\tilde{\mu}_1 = \mu_{\frac{1}{2}+\sigma}$. By repeating the arguments in the proof of (i) \implies (ii), with reversed inequalities and K replaced by $K + \epsilon$, we obtain

$$(41) \quad \int \left[\sigma_{K+\epsilon, 1}^{(\frac{1}{2})} (|\gamma \circ g|) \tilde{\rho}_0(\gamma(1/2-\sigma))^{-1} + \sigma_{K+\epsilon, 1}^{(\frac{1}{2})} (|\gamma \circ g|) \tilde{\rho}_1(\gamma(1/2+\sigma))^{-1} \right] d\tilde{\Pi}(\gamma) < \int_{\Sigma_{\frac{1}{2}}} \tilde{\rho}_{\frac{1}{2}}(y)^{-1} d\tilde{\mu}_{\frac{1}{2}}(y),$$

where $\tilde{\Pi}$ is the optimal plan induced by the W_2 -geodesic $\{\tilde{\mu}_s\}_{s \in [0, 1]}$. Using that the distortion coefficients $\sigma_{K, 1}^{(t)}(\theta)$ are monotone increasing in K , we arrive to contradict (ii) with $t = \frac{1}{2}$. \square

We remind the reader that there is a notion of upper curvature bounds for geodesic metric spaces (X, d) that goes under the name $\text{CAT}(K)$ for $K \in \mathbb{R}$ (see for instance [1, Chapter 9]). In case $K = 0$, the condition reduces to require 1-convexity of $\frac{1}{2}d(y, \cdot)^2$ for any $y \in X$. For Riemannian manifolds (M, g_M) the condition $\text{CAT}(K)$ for the induced metric space (M, d_M) implies an upper sectional curvature bound by K , moreover it also implies that geodesics are always extendible in case $K \leq 0$. The next corollary then follows.

Corollary 5.5. *Let (M, g) be a complete, simply connected Riemannian manifold without boundary. Then the following statements (i) and (ii) are equivalent:*

(i) (M, d_M) satisfies $\text{CAT}(0)$.

(ii) Let $\{\mu_t\}_{t \in [0, 1]}$ be a countably \mathcal{H}^1 -rectifiable W_2 -geodesic. Then

$$\mathcal{H}^1(\text{supp } \mu_t) \leq (1-t)\mathcal{H}^1(\text{supp } \mu_0) + t\mathcal{H}^1(\text{supp } \mu_1), \quad \forall t \in [0, 1].$$

Proof. The implication (i) \implies (ii) follows from the extendibility of geodesics.

The reverse implication follows from the reverse implication in Theorem 5.2. Indeed, the theorem implies that M has non-positive sectional curvature, therefore the $\text{CAT}(0)$ -condition holds locally. Then, since M is simply connected the condition globalizes by [1, Theorem 9.2.9]. \square

6. OT CHARACTERIZATION OF SECTIONAL, AND MORE GENERALLY p -RICCI, CURVATURE LOWER BOUNDS

Throughout the section, $\{\mu_t\}_{t \in [0,1]}$ is a countably \mathcal{H}^p -rectifiable W_2 -geodesic and Π is the corresponding dynamical optimal plan, i.e. $\mu_t = \rho_t \mathcal{H}^p \llcorner \Sigma_t$ where $\Sigma_t \subset M$ is a countably \mathcal{H}^p -rectifiable subset and $\rho_t \in L^1(M, \mathcal{H}^p)$. From Theorem 3.1 (see also Remark 3.2), we know that $\mu_t = (T_{1/2}^t)_\# \mu_{1/2}$ with $T_{1/2}^t : \Sigma_{1/2} \rightarrow \Sigma_t$ is Lipschitz. Lemma 3.10 (see also Remark 3.11) ensures the existence of a subset $N \subset \Sigma_{1/2}$, with $\mathcal{H}^p(N) = 0$, such that $T_{1/2}^t$ is differentiable for every $x \in \Sigma_{1/2} \setminus N$ and we set

$$B_x(t) : T_x \Sigma_{1/2} \rightarrow T_{\gamma_x(t)} \Sigma_t, \quad B_x(t) := DT_{1/2}^t(x) \quad \forall t \in [0, 1], \forall x \in \Sigma_{1/2} \setminus N.$$

Moreover $D_t B_x(t)|_{t=1/2} : T_x \Sigma_{1/2} \rightarrow T_x \Sigma_{1/2}$ is self-adjoint for every $x \in \Sigma_s \setminus N$.

Lemma 3.9 yields that $B_x(t)$ is invertible for every $t \in [0, 1]$ for every $x \in \Sigma_{1/2} \setminus N$, up to enlarging the subset N . Since $B_x(1/2) = \text{Id}$, it follows in particular that $\det[B_x(t)] > 0$ for all $t \in [0, 1]$. Now, for every $x \in \Sigma_{1/2} \setminus N$ and $t \in [0, 1]$, let $\gamma_x(t) := T_{1/2}^t(x)$ be the geodesic performing the transport and consider

$$\begin{aligned} \mathcal{U}_x(t) &:= (\nabla_t B_x(t)) B_x(t)^{-1} : T_{\gamma_x(t)} \Sigma_t \rightarrow T_{\gamma_x(t)} M, \\ \mathcal{U}_x^\top(t) &:= [\mathcal{U}_x(t)]^\top = (D_t B_x(t)) B_x(t)^{-1} : T_{\gamma_x(t)} \Sigma_t \rightarrow T_{\gamma_x(t)} \Sigma_t, \\ \mathcal{U}_x^\perp(t) &:= [\mathcal{U}_x(t)]^\perp : T_{\gamma_x(t)} \Sigma_t \rightarrow (T_{\gamma_x(t)} \Sigma_t)^\perp, \end{aligned}$$

where ∇_t denotes the covariant derivative along $\gamma_x(t)$ in M and $D_t := \top \circ \nabla_t$, \top being the orthogonal projection on $T_{\gamma_x(t)} \Sigma_t$ and \perp being the orthogonal projection on the orthogonal complement $(T_{\gamma_x(t)} \Sigma_t)^\perp$ of $T_{\gamma_x(t)} \Sigma_t$. For every $x \in \Sigma_{1/2} \setminus N$ such that $|\dot{\gamma}_x| \neq 0$, we define

$$(42) \quad \kappa_{\gamma_x} : [0, |\dot{\gamma}_x|] \rightarrow \mathbb{R}, \quad \kappa_{\gamma_x}(|\dot{\gamma}_x| t) |\dot{\gamma}_x|^2 := \|\mathcal{U}_x^\perp(t)\|^2, \quad \forall t \in [0, 1],$$

if $|\dot{\gamma}_x| = 0$, we set $\kappa_{\gamma_x}(0) = 0$. Observe that the map $[0, 1] \ni t \mapsto \kappa_{\gamma_x}(|\dot{\gamma}_x| t) \in \mathbb{R}$ is invariant under constant speed reparametrization of the geodesic γ_x .

We now introduce the generalized distortion coefficients σ_κ associated to a continuous function $\kappa : [0, \theta] \rightarrow \mathbb{R}$ (cf. [9]). First of all, the generalized sin-function associated to κ , denoted by \sin_κ , is defined as the unique solution $v : [0, \theta] \rightarrow \mathbb{R}$ of the equation

$$v'' + \kappa v = 0 \quad \& \quad v(0) = 0, \quad v'(\theta) = 1.$$

The generalized distortion coefficients $\sigma_\kappa^{(t)}(\theta)$, for $t \in [0, 1]$ and $\theta > 0$, are defined as

$$(43) \quad \sigma_\kappa^{(t)}(\theta) := \begin{cases} \frac{\sin_\kappa(t\theta)}{\sin_\kappa(\theta)} & \text{if } \sin_\kappa(s\theta) > 0 \text{ for all } s \in [0, 1], \\ \infty & \text{otherwise.} \end{cases}$$

Using Sturm-Picone comparison Theorem one can check that (see for instance [9, Proposition 3.4])

$$(44) \quad \kappa_1 \leq \kappa_2 \text{ on } [0, \theta] \quad \implies \quad \sigma_{\kappa_1}^{(t)}(\theta) \leq \sigma_{\kappa_2}^{(t)}(\theta) \quad \forall t \in [0, 1].$$

Moreover, by the strong maximum principle (see for instance [20, XVIII]), it holds

$$(45) \quad \kappa_1 < \kappa_2 \text{ on } (0, \theta) \quad \& \quad \sigma_{\kappa_1}^{(\cdot)}(\theta) \neq \infty \quad \implies \quad \sigma_{\kappa_1}^{(t)}(\theta) < \sigma_{\kappa_2}^{(t)}(\theta), \quad \forall t \in (0, 1).$$

It is convenient to also set $\sigma_\kappa^{(\cdot)}(0) \equiv 1$, $\kappa^-(t) := \kappa(\theta - t)$ and $\kappa^+(t) := \kappa(t)$.

If $v_0, v_1 \in [0, \infty)$, a straightforward computation gives that $v(t) := \sigma_{\kappa^-}^{(1-t)}(\theta) v_0 + \sigma_{\kappa^+}^{(t)}(\theta) v_1$ solves

$$(46) \quad v''(t) + \kappa(t\theta)\theta^2 v = 0, \quad \forall t \in (0, 1) \text{ with } v(0) = v_0 \quad \& \quad v(1) = v_1,$$

provided $t \in [0, 1] \mapsto \sigma_{\kappa^+}^{(t)}(\theta)$ (or, equivalently, $t \in [0, 1] \mapsto \sigma_{\kappa^-}^{(t)}(\theta)$) is real-valued.

By [9, Proposition 3.8], if $u : [0, 1] \rightarrow (0, \infty)$ with $u \in C^0([0, 1]) \cap C^2((0, 1))$ satisfies

$$(47) \quad u''(t) + \kappa(t\theta)\theta^2 u(t) \leq 0, \quad \forall t \in (0, 1) \text{ with } u(0) = v_0 \quad \& \quad u(1) = v_1 \quad \implies \quad u \geq v \text{ on } [0, 1].$$

It also convenient to consider a slightly different comparison function. To this aim we define the function $g : [0, 1] \times [0, 1] \rightarrow [0, 1]$ by

$$(48) \quad g(s, t) := \begin{cases} (1-s)t & \text{if } t \in [0, s], \\ s(1-t) & \text{if } t \in [s, 1], \end{cases}$$

so that for all $s \in (0, 1)$ one has

$$(49) \quad -\frac{\partial^2}{\partial t^2} g(s, t) = \delta_s \quad \text{in } \mathcal{D}'(0, 1), \quad g(s, 0) = g(s, 1) = 0.$$

Given $w_0, w_1 \in [0, \infty)$ and a continuous function $u : [0, 1] \rightarrow [0, \infty)$, a straightforward computation gives that $w(t) := (1-t)w_0 + tw_1 + \int_0^1 g(s, t) u(s) ds$ solves

$$(50) \quad w''(t) + u(t) = 0, \quad \forall t \in (0, 1) \text{ with } w(0) = w_0 \text{ \& } w(1) = w_1.$$

Theorem 6.1 (OT Characterization of curvature lower bounds). *Let (M, g) be a complete Riemannian manifold with $\partial M = \emptyset$ and let $K \in \mathbb{R}$. Then the following statements are equivalent:*

(i) $\text{Ric}_p \geq K$.

(ii) *Let $\{\mu_t\}_{t \in [0, 1]}$ be a countably \mathcal{H}^p -rectifiable W_2 -geodesic, and let Π be the corresponding dynamical optimal plan. Then, for any $p' \geq p$, it holds*

$$(51) \quad S_{p'}(\mu_t | \mathcal{H}^p) \leq - \int \left[\sigma_{(K - \kappa_\gamma^-)/p'}^{(1-t)}(|\dot{\gamma}|) \rho_0^{-\frac{1}{p'}}(\gamma(0)) + \sigma_{(K - \kappa_\gamma^+)/p'}^{(t)}(|\dot{\gamma}|) \rho_1^{-\frac{1}{p'}}(\gamma(1)) \right] d\Pi(\gamma), \quad \forall t \in [0, 1]$$

where κ_γ was defined in (42) and the generalized distortion coefficients σ are as in (43).

(iii) *Let $\{\mu_t\}_{t \in [0, 1]}$ and Π be as in (ii). Then*

$$\text{Ent}(\mu_t | \mathcal{H}^p) \leq (1-t) \text{Ent}(\mu_0 | \mathcal{H}^p) + t \text{Ent}(\mu_1 | \mathcal{H}^p) - \int \int_0^1 g(s, t) |\dot{\gamma}|^2 (K - \kappa_\gamma(s|\dot{\gamma}|)) ds d\Pi(\gamma), \quad \forall t \in [0, 1]$$

where $g(s, t)$ was defined in (48).

Remark 6.2. We emphasize that Theorem 6.1 is sharp. First of all, one can not omit the correction term κ_γ : even in \mathbb{R}^n , the convexity of S_p is not true in general. For instance consider \mathbb{R}^2 and the line segment $\{(t, \frac{1}{2}t) : t \in [0, 1]\} =: L_0$ and let $\mu_0 = \mathcal{H}^1|_{L_0}$; similarly, define $\mu_1 = \mathcal{H}^1|_{L_1}$ where $L_1 := \{(t, -\frac{1}{2}t) : t \in [0, 1]\}$. Then, it is easy to check that the optimal transport between μ_0 and μ_1 is supported on geodesics that connect $(t, \frac{1}{2}t)$ and $(t, -\frac{1}{2}t)$ and $\mu_{1/2}$ is exactly $\mathcal{H}^1|_{[0, 1] \times \{0\}}$. If Theorem 6.1 would hold with $K = 0$ and $\kappa_\gamma \equiv 0$, then the Brunn-Minkowski inequality (see Corollary 6.5 below) would contradict that the \mathcal{H}^1 -measure of $[0, 1] \times \{0\}$ is strictly smaller than the one of L_0 and L_1 .

Second, we stress that the arguments in the proof of Theorem 6.1 are sharp, since for this example all the inequalities become identities (for the details see Remark 6.4 after the proof).

The proof of Theorem 6.1 will make use of the next proposition.

Proposition 6.3. *Let M be a complete Riemannian n -dimensional manifold without boundary. Assume that $\text{Ric}_p \geq K$, for some $p \in \{1, \dots, n\}$ and $K \in \mathbb{R}$, and consider a countably \mathcal{H}^p -rectifiable W_2 -geodesic $\{\mu_t\}_{t \in [0, 1]}$.*

Then, using the notation recalled at the beginning of Section 6 and denoting $\mathcal{J}_x(t) := \det[B_x(t)]$, it holds

$$(52) \quad \frac{d^2}{dt^2} \mathcal{J}_x^{\frac{1}{p'}}(t) \leq -\frac{(K - \kappa_{\gamma_x(t)})}{p'} |\dot{\gamma}_x|^2 \mathcal{J}_x^{\frac{1}{p'}}(t), \quad \forall x \in \Sigma_{1/2} \setminus N, \mu_{1/2}(N) = 0, \forall p' \geq p, \forall t \in (0, 1),$$

and thus

$$(53) \quad \mathcal{J}_x^{\frac{1}{p'}}(t) \geq \sigma_{\frac{K - \kappa_{\gamma_x^-}}{p'}}^{(1-t)}(|\dot{\gamma}_x|) \mathcal{J}_x^{\frac{1}{p'}}(0) + \sigma_{\frac{K - \kappa_{\gamma_x^+}}{p'}}^{(t)}(|\dot{\gamma}_x|) \mathcal{J}_x^{\frac{1}{p'}}(1). \quad \forall x \in \Sigma_{1/2} \setminus N, \forall t \in [0, 1], \forall p' \geq p.$$

Proof. If we set $y_x(t) = \log \mathcal{J}_x(t) = \log \det B_x(t)$, from (23) in Proposition 4.4 we know that

$$y_x''(t) + \frac{1}{p} y_x'(t)^2 + \text{Ric}_p(T_{\gamma_x(t)} \Sigma_t, \dot{\gamma}_x(t)) - \|\mathcal{U}^\perp(t)\|^2 \leq 0, \quad \forall t \in (0, 1), \quad \forall x \in \Sigma_{1/2} \setminus N.$$

Plugging the assumption $\text{Ric}_p \geq K$ together with the definition (42) of κ_{γ_x} , we get

$$(54) \quad y_x''(t) + \frac{1}{p'} y_x'(t)^2 + (K - \kappa_{\gamma_x}(t)) |\dot{\gamma}_x(t)|^2 \leq 0, \quad \forall t \in (0, 1), \quad \forall x \in \Sigma_{1/2} \setminus N, \quad \forall p' \geq p,$$

which is equivalent to

$$(55) \quad (\mathcal{J}^{\frac{1}{p'}})''_x(t) + \frac{K - \kappa_{\gamma_x}(t)}{p'} |\dot{\gamma}_x(t)|^2 \mathcal{J}^{\frac{1}{p'}}(t) \leq 0, \quad \forall t \in (0, 1), \quad \forall x \in \Sigma_{1/2} \setminus N, \quad \forall p' \geq p.$$

The claimed (53) follows then by the comparison principle (47) and by Proposition 3.8 in [9]. More precisely, in step **3** of the proof of [9, Proposition 3.8] it is showed that if $\mathcal{J}^{\frac{1}{p'}}$ satisfies (55) and $\mathcal{J}^{\frac{1}{p'}}(t) > 0$ for some $t \in [0, 1]$ then $\sigma_{(K - \kappa_{\dot{\gamma}}^+)_/p'}^{(t)}(|\dot{\gamma}|) < \infty$; the desired (53) follows then from (47). \square

Proof of Theorem 6.1. (i) \implies (ii). Let $\{\mu_t\}_{t \in [0,1]}$ be a countably \mathcal{H}^p -rectifiable geodesic. Recall that for every $t \in [0, 1]$ it holds $\mu_t = \rho_t \mathcal{H}^p \llcorner \Sigma_t = (T_{1/2}^t)_\# \mu_{1/2}$. Let Π be the optimal dynamical plan associated to the W_2 -geodesic $\{\mu_t\}_{t \in [0,1]}$, i.e. $\mu_t = (e_t)_\# \Pi$.

Setting $\mathcal{J}_x(t) = \det B_x(t) = \det[DT_{1/2}^t(x)]$, for all $t \in (0, 1)$ and $p' \geq p$ we get:

$$\begin{aligned} \int_{\Sigma_t} \rho_t(y)^{-\frac{1}{p'}} d\mu_t(y) &= \int_{\Sigma_{1/2}} \rho_t(T_{1/2}^t(x))^{-\frac{1}{p'}} d\mu_{1/2}(x) \\ &\stackrel{(9)}{=} \int_{\Sigma_{1/2}} \mathcal{J}_x^{\frac{1}{p'}}(t) \rho_{1/2}(x)^{1-\frac{1}{p'}} d\mathcal{H}^p(x) \\ &\stackrel{(53)}{\geq} \int_{\Sigma_{1/2}} \left[\sigma_{(K - \kappa_{\gamma_x})_- / p'}^{(1-t)}(|\dot{\gamma}_x|) \mathcal{J}_x^{\frac{1}{p'}}(0) \rho_{1/2}(x)^{1-\frac{1}{p'}} + \sigma_{(K - \kappa_{\gamma_x})_+ / p'}^{(t)}(|\dot{\gamma}_x|) \mathcal{J}_x^{\frac{1}{p'}}(1) \rho_{1/2}(x)^{1-\frac{1}{p'}} \right] d\mathcal{H}^p(x) \\ &\stackrel{(9)}{\geq} \int_{\Sigma_{1/2}} \left[\sigma_{(K - \kappa_{\gamma_x})_- / p'}^{(1-t)}(|\dot{\gamma}_x|) \rho_0(T_{1/2}^0(x))^{-\frac{1}{p'}} + \sigma_{(K - \kappa_{\gamma_x})_+ / p'}^{(t)}(|\dot{\gamma}_x|) \rho_1(T_{1/2}^1(x))^{-\frac{1}{p'}} \right] \rho_{1/2}(x) d\mathcal{H}^p(x) \\ &\stackrel{(35)}{=} \int \left[\sigma_{(K - \kappa_\gamma)_- / p'}^{(1-t)}(|\dot{\gamma}|) \rho_0(\gamma(0))^{-\frac{1}{p'}} + \sigma_{(K - \kappa_\gamma)_+ / p'}^{(t)}(|\dot{\gamma}|) \rho_1(\gamma(1))^{-\frac{1}{p'}} \right] d\Pi(\gamma). \end{aligned}$$

This concludes the proof of (51) for $t \in (0, 1)$. In case $t = 0$ or $t = 1$ just observe that from the very definition (43) it holds $\sigma_{(K - \kappa_\gamma)_- / p'}^{(0)}(|\dot{\gamma}|) = 0$ and $\sigma_{(K - \kappa_\gamma)_- / p'}^{(1)}(|\dot{\gamma}|) = 1$, so the claim (51) is trivially satisfied.

(ii) \implies (i).

We argue by contradiction. Assume there exist $x_0 \in M$, a p -dimensional plane $P \subset T_{x_0}M$ and $0 \neq v \in T_{x_0}M$ such that the p -Ricci curvature of P in the direction of v satisfies

$$(56) \quad \text{Ric}_p(P, v) \leq (K - 4\epsilon)|v|^2,$$

for some $\epsilon > 0$. Let $\delta > 0$ be sufficiently small such that $\exp_{x_0}|_{B_\delta(0)}$ is a diffeomorphism onto its image. Then $\exp_{x_0}(P \cap B_\delta(0)) =: \Sigma_{\frac{1}{2}}$ is a smooth p -dimensional submanifold. Let $\phi \in C_0^\infty(M)$ be a Kantorovich potential such that

$$(57) \quad \nabla \phi(x_0) = v \neq 0 \quad \text{and} \quad \nabla^2 \phi(x_0) = 0.$$

By replacing ϕ with $\eta\phi$ for a sufficiently small number $\eta > 0$ we get that ϕ is a Kantorovich potential as well and $|\nabla \phi|(y)$ is smaller than the injectivity radius at y , for every $y \in \text{supp}(\phi) \subset M$. It is easily checked that for $\delta > 0$ small enough the map $y \mapsto T_t(y) = \exp_y(-t\nabla \phi(y))$ is a diffeomorphism from $B_\delta(0)$ onto its image for any $t \in [-\frac{1}{2}, \frac{1}{2}]$. Hence, $\Sigma_t := T_{t-\frac{1}{2}}(\Sigma_{\frac{1}{2}})$ for $t \in [0, 1]$ is a 1-parameter family of smooth p -dimensional submanifolds with finite p -dimensional Hausdorff measure. We define $\mu_{\frac{1}{2}} := \mathcal{H}^p(\Sigma_{\frac{1}{2}})^{-1} \mathcal{H}^p \llcorner \Sigma_{\frac{1}{2}}$; note that $\mu_t := (T_{t-\frac{1}{2}})_\# \mu_{\frac{1}{2}}$, with $t \in [0, 1]$, is the unique L^2 -Wasserstein geodesic between μ_0 and μ_1 . Moreover, by construction, μ_t is a \mathcal{H}^p -absolutely continuous probability

measure concentrated on Σ_t .

Calling $\gamma_x(t) := T_{t-\frac{1}{2}}(x) = \exp_x(- (t - \frac{1}{2})\nabla\phi(x))$ for $x \in \Sigma_{\frac{1}{2}}$ the geodesic performing the transport, note that by continuity there exist $\delta, \sigma > 0$ small enough such that

$$(58) \quad \text{Ric}_p(T_{\gamma_x(t)}\Sigma_t, \dot{\gamma}_x(t)) < (K - 3\epsilon)|\dot{\gamma}_x(t)|^2, \quad \forall x \in \Sigma_{\frac{1}{2}} \subset B_\delta(x_0), \quad \forall t \in \left[\frac{1}{2} - \sigma, \frac{1}{2} + \sigma\right].$$

The identity (22) proved in Proposition 4.4 reads as

$$(59) \quad \text{tr}[\mathcal{U}_x^\top(t)]' + \text{tr}[\mathcal{U}_x^\top(t)^2] + \text{Ric}_p(T_{\gamma_x(t)}\Sigma_t, \dot{\gamma}_x(t)) = \|\mathcal{U}_x^\perp(t)\|^2, \quad \forall x \in \Sigma_{\frac{1}{2}}, \quad \forall t \in [0, 1].$$

Since by construction $\mathcal{U}_{x_0}(0) := \nabla_t B_{x_0}(0)B_{x_0}^{-1}(0) = \nabla^2\phi(x_0)|_{T_{x_0}\Sigma_{\frac{1}{2}}} = 0$ and $v \neq 0$, again by continuity we can choose $\delta, \sigma > 0$ even smaller so that

$$(60) \quad \|\mathcal{U}_x^\perp(t)\|^2 + \text{tr}[(\mathcal{U}_x^\top(t))^2] = \sum_{i=1}^p [|\mathcal{U}^\perp(t)E_i|^2 + |\mathcal{U}^\top(t)E_i|^2] = \sum_{i=1}^p |\mathcal{U}(t)E_i|^2 = \|\mathcal{U}_x(t)\|^2 < \epsilon|\dot{\gamma}_x(t)|^2$$

for all $x \in \Sigma_{\frac{1}{2}} \subset B_\delta(x_0)$ and all $t \in [\frac{1}{2} - \sigma, \frac{1}{2} + \sigma]$. The combination of (58), (59) and (60) yields

$$(61) \quad 0 \leq \|\mathcal{U}_x^\perp(t)\|^2 < \text{tr}[\mathcal{U}_x^\top(t)]' + (K - 2\epsilon)|\dot{\gamma}_x(t)|^2 \leq \text{tr}[\mathcal{U}_x^\top(t)]' + \frac{1}{p}\text{tr}[\mathcal{U}_x^\top(t)]^2 + (K - 2\epsilon)|\dot{\gamma}_x(t)|^2,$$

for all $x \in \Sigma_{\frac{1}{2}}$ and all $t \in [\frac{1}{2} - \sigma, \frac{1}{2} + \sigma]$. Observe that the affine reparametrization $t = g(s) = \frac{1}{2} - \sigma + 2\sigma s$, $g : [0, 1] \rightarrow [\frac{1}{2} - \sigma, \frac{1}{2} + \sigma]$, corresponds to consider the rescaled Kantorovich potential $2\sigma\phi$ in place of ϕ in the arguments above, and thus gives

$$\begin{aligned} \text{tr}[\mathcal{U}_x^\top(g(s))]' + \frac{1}{p}\text{tr}[\mathcal{U}_x^\top(g(s))]^2 + (K - 2\epsilon)|\dot{\gamma}_x(g(s))|^2 &= 4\sigma^2 \left[\text{tr}[\mathcal{U}_x^\top(t)]' + \frac{1}{p}\text{tr}[\mathcal{U}_x^\top(t)]^2 + (K - 2\epsilon)|\dot{\gamma}_x(t)|^2 \right] \\ &> 0, \quad \forall x \in \Sigma_{\frac{1}{2}}, \quad \forall s \in [0, 1]. \end{aligned}$$

Arguing as in the proof of Proposition 6.3 (but with reversed inequalities), the last differential inequality gives

$$(62) \quad \mathcal{J}_x^{\frac{1}{p}}(g(s)) \leq \sigma \frac{(1-t)}{K-2\epsilon} (|\dot{\gamma}_x \circ g|) \mathcal{J}_x^{\frac{1}{p}}(0) + \sigma \frac{t}{K-2\epsilon} (|\dot{\gamma}_x \circ g|) \mathcal{J}_x^{\frac{1}{p}}(1), \quad \forall x \in \Sigma_{1/2} \setminus N, \quad \forall s \in [0, 1].$$

Note in particular that, since $\mathcal{J}_x^{\frac{1}{p}}(g(s)) > 0$ for all $s \in [0, 1]$, then $\sigma \frac{(\cdot)}{K-2\epsilon} (|\dot{\gamma}_x \circ g|) \neq 0$.

Since g is affine, the restricted and rescaled curve $\{\tilde{\mu}_s := \mu_{g(s)}\}_{s \in [0, 1]}$ is still a W_2 -geodesic from $\tilde{\mu}_0 = \mu_{\frac{1}{2}-\sigma}$ to $\tilde{\mu}_1 = \mu_{\frac{1}{2}+\sigma}$. By repeating the arguments in the proof of (i) \implies (ii), with reversed inequalities (note that (9) holds with equality since we are considering the interior of a geodesic, use (62) instead of (53), and replace $K - \kappa_\gamma$ by $K - 2\epsilon$), we obtain

$$(63) \quad \int_{\Sigma_{\frac{1}{2}}} \tilde{\rho}_{\frac{1}{2}}(y)^{-\frac{1}{p}} d\tilde{\mu}_{\frac{1}{2}}(y) \leq \int \left[\sigma \frac{(\frac{1}{2})}{(K-2\epsilon)/p} (|\dot{\gamma}|) \tilde{\rho}_0(\gamma(0))^{-\frac{1}{p}} + \sigma \frac{(\frac{1}{2})}{(K-2\epsilon)/p} (|\dot{\gamma}|) \tilde{\rho}_1(\gamma(1))^{-\frac{1}{p}} \right] d\tilde{\Pi}(\gamma),$$

where $\tilde{\Pi}$ is the dynamical optimal associated to $\{\tilde{\mu}_s = \tilde{\rho}_s \mathcal{H}^p\}_{s \in [0, 1]}$.

Observing that (60) gives $\kappa_\gamma \leq \epsilon$ and noting that $\|\mathcal{U}_x^\perp(t)\|^2$ correctly scales when we apply the reparametrization g , using (44) we get that (63) implies

$$\begin{aligned} \int_{\Sigma_{\frac{1}{2}}} \tilde{\rho}_{\frac{1}{2}}(y)^{-\frac{1}{p}} d\tilde{\mu}_{\frac{1}{2}}(y) &\leq \int \left[\sigma \frac{(\frac{1}{2})}{(K-\kappa_{\gamma_x}-\epsilon)/p} (|\dot{\gamma}|) \tilde{\rho}_0(\gamma(0))^{-\frac{1}{p}} + \sigma \frac{(\frac{1}{2})}{(K-\kappa_{\gamma_x}-\epsilon)/p} (|\dot{\gamma}|) \tilde{\rho}_1(\gamma(1))^{-\frac{1}{p}} \right] d\tilde{\Pi}(\gamma) \\ &\stackrel{(45)}{<} \int \left[\sigma \frac{(\frac{1}{2})}{(K-\kappa_{\gamma_x})/p} (|\dot{\gamma}|) \tilde{\rho}_0(\gamma(0))^{-\frac{1}{p}} + \sigma \frac{(\frac{1}{2})}{(K-\kappa_{\gamma_x})/p} (|\dot{\gamma}|) \tilde{\rho}_1(\gamma(1))^{-\frac{1}{p}} \right] d\tilde{\Pi}(\gamma). \end{aligned}$$

This contradicts (ii) for the geodesic $\{\tilde{\mu}_s = \tilde{\rho}_s \mathcal{H}^p\}_{s \in [0, 1]}$.

(i) \implies (iii).

For $t \in (0, 1)$ we have

$$(64) \quad \begin{aligned} \text{Ent}(\mu_t | \mathcal{H}^p) &= \int_{\Sigma_t} \log \rho_t(y) d\mu_t(y) = \int_{\Sigma_{1/2}} \log \rho_t(T_{1/2}^t(x)) d\mu_{1/2}(x) \stackrel{(9)}{=} \int_{\Sigma_{1/2}} \log[\rho_{1/2}(x) \mathcal{J}_t(x)^{-1}] d\mu_{1/2}(x) \\ &= \text{Ent}(\mu_{1/2} | \mathcal{H}^p) - \int_{\Sigma_{1/2}} y_x(t) d\mu_{1/2}(x), \end{aligned}$$

where $y_x(t) = \log(\mathcal{J}_t(x))$. Using (54) we obtain

$$(65) \quad \frac{d^2}{dt^2} \text{Ent}(\mu_t | \mathcal{H}^p) \geq \int_{\Sigma_{1/2}} (K - \kappa_{\gamma_x}(t)) |\dot{\gamma}_x(t)|^2 d\mu_{1/2}(x), \quad \forall t \in (0, 1).$$

We then get (iii) using (50) and standard comparison.

(iii) \implies (i).

Assume that by contradiction (58) holds and repeat verbatim the first part of the proof of (ii) \implies (i) to reach (61), i.e.

$$\text{tr}[\mathcal{U}_x^\top(t)]' + (K - 2\epsilon) |\dot{\gamma}_x(t)|^2 > 0, \quad \forall x \in \Sigma_{\frac{1}{2}}, \forall t \in \left[\frac{1}{2} - \sigma, \frac{1}{2} + \sigma \right].$$

Considering as above the affine reparametrization $t = g(s) = \frac{1}{2} - \sigma + 2\sigma s$, $g : [0, 1] \rightarrow [\frac{1}{2} - \sigma, \frac{1}{2} + \sigma]$ and recalling (24), we obtain

$$y_x(g(s))'' + (K - 2\epsilon) |\dot{\gamma}_x(g(s))|^2 > 0, \quad \forall x \in \Sigma_{\frac{1}{2}}, \forall s \in [0, 1].$$

Calling as above $\{\tilde{\mu}_s := \mu_{g(s)}\}_{s \in [0, 1]}$ the corresponding rescaled W_2 -geodesic, the combination of the last inequality with (64) gives

$$(66) \quad \frac{d^2}{ds^2} \text{Ent}(\tilde{\mu}_s | \mathcal{H}^p) < \int_{\Sigma_{1/2}} (K - 2\epsilon) |\dot{\gamma}_x(g(s))|^2 d\mu_{1/2}(x), \quad \forall s \in (0, 1).$$

Calling $\tilde{\Pi}$ the dynamical optimal plan associated to the geodesic $\{\tilde{\mu}_s := \mu_{g(s)}\}_{s \in [0, 1]}$, using (50) and standard comparison we get that

$$\text{Ent}(\tilde{\mu}_s | \mathcal{H}^p) > (1 - s) \text{Ent}(\tilde{\mu}_0 | \mathcal{H}^p) + s \text{Ent}(\tilde{\mu}_1 | \mathcal{H}^p) - \int_0^1 \int_0^1 g(t, s) |\dot{\gamma}|^2 (K - 2\epsilon) dt d\tilde{\Pi}(\gamma).$$

Observing now that (60) gives $\kappa_\gamma \leq \epsilon$ for $\tilde{\Pi}$ -a.e. γ , we obtain

$$\text{Ent}(\tilde{\mu}_s | \mathcal{H}^p) > (1 - s) \text{Ent}(\tilde{\mu}_0 | \mathcal{H}^p) + s \text{Ent}(\tilde{\mu}_1 | \mathcal{H}^p) - \int_0^1 \int_0^1 g(t, s) |\dot{\gamma}|^2 (K - \kappa_\gamma(t) |\dot{\gamma}| - \epsilon) dt d\tilde{\Pi}(\gamma),$$

which contradicts (iii) thanks to the strict positivity of g on $(0, 1) \times (0, 1)$. \square

Remark 6.4. In order to show that Theorem 6.1 is sharp, we show that equality is achieved in (51) for the example of Remark 6.2, $p' = 1$. First of all recall that, in euclidean spaces, the Jacobi fields are affine functions along the geodesics. The initial measure μ_0 is supported on the segment $\{(t, \frac{1}{2}t) : t \in [0, 1]\}$ that is generated by the unit vector $\frac{1}{\sqrt{5}}(2, 1) = e$, and the final measure is supported on the segment $\{(t, -\frac{1}{2}t) : t \in [0, 1]\}$ that is generated by $-\frac{1}{\sqrt{5}}(2, 1)$. Set $(\frac{2}{\sqrt{5}}, 0) = v$ and $(0, \frac{1}{\sqrt{5}}) = w$. We have

$$B_x(t) = J_e(t) = v + (1 - 2t)w = \left(\frac{2}{\sqrt{5}}, (1 - 2t)\frac{1}{\sqrt{5}} \right),$$

and $J'_e(t) = -2w = \left(0, -\frac{2}{\sqrt{5}} \right)$. Clearly, $u(t) = (t - \frac{1}{2}, 1)$ is orthogonal to $J_e(t)$ for every $t \in [0, 1]$. Thus,

$$\|(J'_e(t))^\perp\| = \frac{1}{\|u(t)\|} \langle J'_e(t), u(t) \rangle = -\frac{2}{\sqrt{5}(t^2 - t + \frac{5}{4})}.$$

Using the identity (32), we get that

$$\kappa_\gamma(t|\dot{\gamma}_x|)|\dot{\gamma}_x|^2 = \|(\mathcal{U}(t)E(t))^\perp\|^2 = \left\| \|J_e(t)\|^{-1} (J'_e(t))^\perp \right\|^2 = \frac{1}{(t^2 - t + \frac{5}{4})^2} = \kappa(t)$$

where $E(t) = \|J_e(t)\|^{-1} J_e(t)$. It follows that the coefficient $\kappa_\gamma(t|\dot{\gamma}|)|\dot{\gamma}|^2$ does not depend on γ . We thus get

$$(67) \quad \mathcal{H}^1(\text{supp } \mu_t) = \sigma_{-\kappa^-}^{(1-t)}(1) \mathcal{H}^1(\text{supp } \mu_0) + \sigma_{-\kappa^+}^{(t)}(1) \mathcal{H}^1(\text{supp } \mu_1).$$

Indeed, the \mathcal{H}^1 -measure of the support of μ_t is given by the length of the Jacobi field $J_e(t)$ with the normalisation $\alpha > 0$ such that $\alpha J_e(\frac{1}{2}) = (1, 0)$:

$$\mathcal{H}^1(\text{supp } \mu_t) = \sqrt{t^2 - t + \frac{5}{4}}.$$

A straightforward computation shows that $t \mapsto \mathcal{H}^1(\text{supp } \mu_t)$ solves the boundary value problem $f''(t) = \kappa(t)f(t)$, $f(1) = f(0) = \mathcal{H}^1(\text{supp } \mu_0) = \mathcal{H}^1(\text{supp } \mu_1) = \frac{\sqrt{5}}{2}$, thus (67) follows.

Note that, for this example, in the proof of Theorem 6.1 every inequality becomes an identity, showing the sharpness of the arguments.

As an application of Theorem 6.1 we establish a new Brunn-Minkowski type inequality involving the \mathcal{H}^p -measure and countably \mathcal{H}^p -rectifiable sets. The main novelty is about the measure: the standard Brunn-Minkowski inequality involves the top Hausdorff measure \mathcal{H}^n in an n -dimensional Riemannian manifold. A second refinement is that, in comparison with the standard Brunn-Minkowski inequality where one gives a lower bound on the measure of *all intermediate points*, here we give more sharply a lower bound on the measure of just the *t -intermediate points where the optimal transport is performed* (let us mention that this was already the case in [17] even if not explicitly stated, but there one considers the top dimensional Hausdorff measure).

Corollary 6.5 (*p -Brunn-Minkowski inequality*). *Let (M, g) be a complete n -dimensional Riemannian manifold without boundary. Assume that $\text{Ric}_p \geq K$ for some $p \in \{1, \dots, n\}$ and $K \in \mathbb{R}$. Let $A_0, A_1 \subset M$ be bounded p -rectifiable subsets with positive and finite \mathcal{H}^p -measure. Set $\mu_i = \mathcal{H}^p(A_i)^{-1} \mathcal{H}^p \llcorner A_i$ for $i = 0, 1$ and assume that there exists a W_2 -geodesic $\{\mu_t\}_{t \in [0,1]}$ such that for some $t_0 \in (0, 1)$ the measure μ_{t_0} is concentrated on a countably \mathcal{H}^p -rectifiable subset $\Sigma_{t_0} \subset M$.*

Then for every $t \in [0, 1]$ one has $\mu_t = \rho_t \mathcal{H}^p \in \mathcal{P}_c(M, \mathcal{H}^p)$ and it holds

$$(68) \quad \mathcal{H}^p(\{\rho_t > 0\})^{\frac{1}{p'}} \geq \int \left[\sigma_{(K-\kappa^-)/p'}^{(1-t)}(|\dot{\gamma}|) \rho_0^{-\frac{1}{p'}}(\gamma(0)) + \sigma_{(K-\kappa^+)/p'}^{(t)}(|\dot{\gamma}|) \rho_1^{-\frac{1}{p'}}(\gamma(1)) \right] d\Pi(\gamma), \quad \forall p' \geq p.$$

In particular, calling

$$A_t := \{\gamma(t) : \gamma \in \text{Geo}(X), \gamma(0) \in A_0, \gamma(1) \in A_1\}$$

the set of all t -intermediate points of geodesics with endpoints in A_0 and A_1 , it holds

$$(69) \quad \mathcal{H}^p(A_t)^{\frac{1}{p'}} \geq \int \left[\sigma_{(K-\kappa^-)/p'}^{(1-t)}(|\dot{\gamma}|) \rho_0^{-\frac{1}{p'}}(\gamma(0)) + \sigma_{(K-\kappa^+)/p'}^{(t)}(|\dot{\gamma}|) \rho_1^{-\frac{1}{p'}}(\gamma(1)) \right] d\Pi(\gamma), \quad \forall p' \geq p.$$

Proof. From Lemma 3.3 we know that for every $t \in [0, 1]$ it holds $\mu_t = \rho_t \mathcal{H}^p \in \mathcal{P}_c(M, \mathcal{H}^p)$. Moreover $\mathcal{H}^p(\{\rho_t > 0\} \setminus A_t) = 0$. Therefore, if we prove (68) then also (69) will follow.

In order to get (68), observe that from Theorem 6.1 the W_2 -geodesic $\{\mu_t\}_{t \in [0,1]}$ satisfies

$$(70) \quad \int_{\{\rho_t > 0\}} \rho_t(x)^{-\frac{1}{p'}} d\mu_t(x) \geq \int \left[\sigma_{(K-\kappa^-)/p'}^{(1-t)}(|\dot{\gamma}|) \rho_0^{-\frac{1}{p'}}(\gamma(0)) + \sigma_{(K-\kappa^+)/p'}^{(t)}(|\dot{\gamma}|) \rho_1^{-\frac{1}{p'}}(\gamma(1)) \right] d\Pi(\gamma), \quad \forall p' \geq p,$$

On the other hand, from Jensen inequality we get

$$\begin{aligned}
 \int_{\{\rho_t > 0\}} \rho_t(x)^{-\frac{1}{p'}} d\mu_t(x) &= \mathcal{H}^p(\{\rho_t > 0\}) \int_{\{\rho_t > 0\}} \rho_t^{1-\frac{1}{p'}} d\left(\frac{1}{\mathcal{H}^p(\{\rho_t > 0\})} \mathcal{H}^p \llcorner \{\rho_t > 0\}\right) \\
 &\leq \mathcal{H}^p(\{\rho_t > 0\}) \left(\int_{\{\rho_t > 0\}} \rho_t d\left(\frac{1}{\mathcal{H}^p(\{\rho_t > 0\})} \mathcal{H}^p \llcorner \{\rho_t > 0\}\right) \right)^{1-\frac{1}{p'}} \\
 (71) \qquad \qquad \qquad &= \mathcal{H}^p(\{\rho_t > 0\})^{\frac{1}{p'}}.
 \end{aligned}$$

The combination of (70) and (71) implies (68). \square

REFERENCES

- [1] D. Burago, Y. Burago, and S. Ivanov, A course in metric geometry, *Graduate Studies in Mathematics*, **33**, American Mathematical Society, Providence, RI, (2001).
- [2] P. Castillon, Submanifolds, isoperimetric inequalities and optimal transportation, *J. Funct. Anal.*, **259**, (2010), 79–103.
- [3] D. Cordero-Erausquin, R. J. McCann, and M. Schmuckenschläger, A Riemannian interpolation inequality á la Borell, Brascamp and Lieb, *Invent. Math.* **146**, (2), (2001), 219–257.
- [4] D. Cordero-Erausquin, R. J. McCann, and M. Schmuckenschläger, Prékopa-Leindler type inequalities on Riemannian manifolds, Jacobi fields, and optimal transport, *Ann. Fac. Sci. Toulouse Math.*, **6**, (15), (2006), 613–635.
- [5] W. Gangbo and R. McCann, Shape recognition via Wasserstein distance, *Quart. Appl. Math.* **58**, (2000), 705–737.
- [6] J. Kitagawa and M. Warren, Regularity for the optimal transportation problem with Euclidean distance squared cost on the embedded sphere, *SIAM J. Math. Anal.*, **44**, (2012), 2871–2887.
- [7] J. Lott, On tangent cones in Wasserstein space, *Proc. Amer. Math. Soc.*, **145**, 7, (2017), 3127–3136.
- [8] J. Lott and C. Villani, Ricci curvature for metric-measure spaces via optimal transport, *Ann. of Math. (2)* **169** (2009), 903–991.
- [9] C. Ketterer, On the geometry of metric measure spaces with variable curvature bounds, *J. Geom. Anal.* **27** (2017), no 3, 1951–1994.
- [10] R.J. McCann, B. Pass and M. Warren, Rectifiability of optimal transportation plans, *Canad. J. Math.* **64**, (2012) 924–934.
- [11] R. McCann and M. Sosio, Hölder continuity for optimal multivalued mappings, *SIAM J. Math. Anal.*, **43**, 4, (2011), 1855–1871.
- [12] P. Petersen and F. Wilhelm, On Frankel’s Theorem, *Canad. Math. Bull.*, **46**, (1), (2003), 130–139.
- [13] A. Petrunin, Alexandrov meets Lott-Villani-Sturm, *Münster J. Math.* **4**, (2011), 53–64.
- [14] R.T. Rockafellar, Second-order convex Analysis, *J. Nonlinear Convex Anal.*, **1**, (1), (2000), 1–16.
- [15] Z.M. Shen, On complete manifolds of nonnegative k th-Ricci curvature, *Trans. Amer. Math. Soc.*, **338**, (1), (1993), 289–310.
- [16] K.T. Sturm, On the geometry of metric measure spaces. I, *Acta Math.*, **196**, (2006), 133–177.
- [17] K.T. Sturm, On the geometry of metric measure spaces. II, *Acta Math.*, **196**, (2006), 133–177.
- [18] K.T. Sturm and M.K. von Renesse, Transport inequalities, gradient estimates, entropy, and Ricci curvature, *Comm. Pure Appl. Math.*, **58**, (2005), 923–940.
- [19] C. Villani, Optimal transport. Old and new, *Grundlehren der Mathematischen Wissenschaften*, **338**, Springer-Verlag, Berlin, (2009).
- [20] W. Walter, Ordinary differential equations, 6th Edition, *Graduate Texts in Mathematics* **182**, Springer-Verlag, (1998).
- [21] F. Wilhelm, On intermediate Ricci curvature and fundamental groups, *Illinois Journ. of Math.*, **41**, (3), (1997), 488–494.
- [22] H. Wu, Manifolds of partially positive curvature, *Indiana Univ. Math. J.*, **36**, (3), (1987), 525–548.
- [23] H.W. Xu and F. Ye, Differentiable sphere theorems for submanifolds of positive k -th Ricci curvature, *Manuscripta Math.*, **138**, (3-4), (2012), 529–543.