



# Descriptors for Pentane-2,4-dione and Its Derivatives

Michael H. Abraham<sup>1</sup>  · William E. Acree Jr.<sup>2</sup>

Received: 17 April 2017 / Accepted: 3 June 2017 / Published online: 19 August 2017  
© The Author(s) 2017. This article is an open access publication

**Abstract** We have used equations for partition coefficients of compounds from water and the gas phase to various solvents to obtain descriptors for pentane-2,4-dione and 21 of its derivatives. These descriptors can then be used to estimate further partition coefficients into a wide variety of solvents. The descriptors also yield information about the properties of pentane-2,4-dione and its derivatives. Pentane-2,4-dione and its alkyl derivatives are quite polar, with substantial hydrogen bond basicity but with no hydrogen bond acidity. In contrast 1,1,1-trifluoropentane-2,4-dione and hexafluoropentane-2,4-dione have significant hydrogen bond acidities.

**Keywords** Pentane-2,4-dione · Partition coefficients · Absolv descriptors · Acetylacetone · Linear free energy relationships

## 1 Introduction

The extraction of metal ions from aqueous solution into organic phases by the use of organic ligands is of great importance in the separation and purification of metals [1]. Quantitative studies of the extraction process require a knowledge of the water–solvent partition of the organic ligand, and so any method of estimating or predicting partition coefficients of organic ligands would be a considerable help in the design of new or novel extraction systems. We have set out a systematic method for the determination of properties or ‘descriptors’ of molecules [2–5], mostly using experimental values of water to

---

✉ Michael H. Abraham  
m.h.abraham@ucl.ac.uk

William E. Acree Jr.  
bill.acree@unt.edu

<sup>1</sup> Department of Chemistry, University College London, 20 Gordon St, London WC1H 0AJ, UK

<sup>2</sup> Department of Chemistry, University of North Texas, 1155 Union Circle Drive #305070, Denton, TX 76203-5017, USA

solvent partition coefficients. These descriptors, known as Abraham or Absolv descriptors can then be used to estimate partition coefficients for other water–solvent systems, as well as numerous physicochemical, environmental and biological properties. We have already used this method to determine descriptors for organophosphorus extractants [6]. Another well-known class of extracting agents is based on pentane-2,4-dione, or acetylacetone and its derivatives. Although we have preliminary descriptors for acetylacetone and some derivatives [7, 8], these were based on limited data, and so we have re-determined descriptors for acetylacetone itself, and have obtained new descriptors for 20 derivatives that have been used as extraction agents.

In solution, acetylacetone and its derivatives exist as a mixture of keto and enol forms, the proportion of which depends on the solvent. If the keto–enol equilibrium constant is known in a number of solvents for which the corresponding water–solvent partition coefficients are known, then it is possible to determine descriptors separately for the keto and enol forms [9]. For many of the compounds that we consider in this work, the keto–enol equilibrium constants are not known, and so we use the experimental water–solvent partition coefficients to obtain descriptors for the keto–enol mixture. These obtained descriptors can then be used to predict further experimental partition coefficients into a wide range of solvents.

## 2 Methodology

We start with our well-known linear free energy relationships, LFERs, Eqs. 1 and 2 [2–5] for the partition of neutral molecules (non-electrolytes) from water to another solvent or solvent system,

$$\log_{10}P = c + eE + sS + aA + bB + vV \quad (1)$$

$$\log_{10}K = c + eE + sS + aA + bB + lL \quad (2)$$

In Eq. 1, the dependent variable is  $\log_{10} P$ , where  $P$  is the water to solvent partition coefficient for a series of non-electrolytes in a given water to solvent system. In Eq. 2, the dependent variable is  $\log_{10} K$ , where  $K$  is the gas phase to solvent system partition coefficient. The independent variables are descriptors as described previously [2–5].  $E$  is the non-electrolyte (or solute) excess molar refractivity in units of  $(\text{cm}^3 \cdot \text{mol}^{-1})/10$ ,  $S$  is the solute dipolarity/polarizability,  $A$  and  $B$  are the overall or summation solute hydrogen bond acidity and basicity,  $V$  is the solute McGowan characteristic volume in units of  $(\text{cm}^3 \cdot \text{mol}^{-1})/100$ , and  $L$  is  $\log_{10} K16$ , where  $K16$  is the gas to hexadecane partition coefficient at 298 K. The use of Eqs. 1 and 2 has been reviewed [2–5]; the review of Clarke and Mallon [5] is particularly exhaustive.

In order to obtain descriptors we first need LFERs, based on Eq. 1 for partition from water to various solvent systems. The coefficients in Eq. 1 for partition from water into wet (water saturated) solvents are in Table 1 [6]. Then if we have  $\log_{10} P$  values for a given solute in systems for which we have descriptors, we can determine values of the descriptors in Eq. 1 by solution of a set of simultaneous equations. We usually have more equations than we have unknowns (i.e. the descriptors). In this case, use of Microsoft ‘Solver’ is a very convenient way of solving the set of equations by trial-and-error. The solution is the set of descriptors that gives the best fit of the dependent variable. The solution is greatly helped if we have prior knowledge of some of the descriptors. The  $E$ -descriptor can be

**Table 1** Coefficients in Eqs. 1 and 2 for partition of solutes from water and the gas phase to wet organic solvents at 298 K

Solvent	<i>c</i>	<i>e</i>	<i>s</i>	<i>a</i>	<i>b</i>	<i>l</i>	<i>v</i>
Coefficients in Eq. 1							
Butan-1-ol	0.376	0.434	−0.718	−0.097	−2.350	0.000	2.682
Pentan-1-ol	0.185	0.367	−0.732	0.105	−3.100	0.000	3.395
Hexan-1-ol	−0.006	0.460	−0.940	0.142	−3.284	0.000	3.792
Heptan-1-ol	0.041	0.497	−0.976	0.030	−3.438	0.000	3.859
Octan-1-ol	0.088	0.562	−1.054	0.034	−3.460	0.000	3.814
Nonan-1-ol	−0.041	0.562	−1.103	0.090	−3.540	0.000	3.922
Decan-1-ol	−0.136	0.542	−0.989	0.046	−3.722	0.000	3.996
Isobutanol	0.249	0.480	−0.639	−0.050	−2.284	0.000	2.758
Dichloromethane	0.319	0.102	−0.187	−3.058	−4.090	0.000	4.324
Trichloromethane	0.191	0.105	−0.403	−3.112	−3.514	0.000	4.395
Tetrachloromethane	0.199	0.523	−1.159	−3.560	−4.594	0.000	4.618
1,2-Dichloroethane	0.183	0.294	−0.134	−2.801	−4.291	0.000	4.180
Hexane	0.333	0.560	−1.710	−3.578	−4.939	0.000	4.463
Heptane	0.297	0.634	−1.755	−3.571	−4.946	0.000	4.488
Octane	0.241	0.690	−1.769	−3.545	−5.011	0.000	4.511
Decane	0.172	0.726	−1.750	−3.446	−4.496	0.000	4.489
Undecane	0.058	0.603	−1.661	−3.421	−5.120	0.000	4.619
Dodecane	0.114	0.668	−1.644	−3.545	−5.006	0.000	4.459
Hexadecane	0.087	0.667	−1.617	−3.587	−4.869	0.000	4.433
Cyclohexane	0.159	0.784	−1.678	−3.740	−4.929	0.000	4.577
Methylcyclohexane	0.246	0.782	−1.982	−3.517	−4.293	0.000	4.528
Isooctane	0.318	0.555	−1.737	−3.677	−4.864	0.000	4.417
Benzene	0.142	0.464	−0.588	−3.099	−4.625	0.000	4.491
Toluene	0.143	0.527	−0.720	−3.010	−4.824	0.000	4.545
Fluorobenzene	0.139	0.152	−0.374	−3.030	−4.601	0.000	4.540
Chlorobenzene	0.065	0.381	−0.521	−3.183	−4.700	0.000	4.614
Bromobenzene	−0.017	0.436	−0.424	−3.174	−4.558	0.000	4.445
Iodobenzene	−0.192	0.298	−0.308	−3.213	−4.653	0.000	4.588
Nitrobenzene	−0.152	0.525	0.081	−2.332	−4.494	0.000	4.187
Benzonitrile	0.097	0.285	0.059	−1.605	−4.562	0.000	4.028
Diethyl ether	0.248	0.561	−1.016	−0.226	−4.553	0.000	4.075
Diisopropylether	0.472	0.413	−0.745	−0.632	−5.251	0.000	4.059
Dibutylether	0.252	0.677	−1.506	−0.807	−5.249	0.000	4.815
Ethyl acetate	0.441	0.591	−0.699	−0.325	−4.261	0.000	3.666
Butyl acetate	−0.475	0.428	−0.094	−0.241	−4.151	0.000	4.046
Methyl isobutyl ketone	0.383	0.801	−0.831	−0.121	−4.441	0.000	3.876
Tributylphosphate	0.327	0.570	−0.837	−1.069	−4.333	0.000	3.919
Gas–water	−0.994	0.577	2.549	3.813	4.841	0.000	−0.869
Coefficients in Eq. 2							
Butan-1-ol	−0.095	0.262	1.396	3.405	2.565	0.523	0.000
Pentan-1-ol	−0.107	−0.001	1.188	3.614	1.671	0.721	0.000
Hexan-1-ol	−0.302	−0.046	0.880	3.609	1.785	0.824	0.000

**Table 1** continued

Solvent	<i>c</i>	<i>e</i>	<i>s</i>	<i>a</i>	<i>b</i>	<i>l</i>	<i>v</i>
Heptan-1-ol	-0.159	0.018	0.825	3.539	1.425	0.830	0.000
Octan-1-ol	-0.222	0.088	0.701	3.478	1.477	0.851	0.000
Nonan-1-ol	-0.197	0.141	0.694	3.616	1.299	0.827	0.000
Decan-1-ol	-0.302	0.233	0.741	3.531	1.177	0.835	0.000
Isobutanol	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Dichloromethane	0.192	-0.572	1.492	0.460	0.847	0.965	0.000
Trichloromethane	0.157	-0.560	1.259	0.374	1.333	0.976	0.000
Tetrachloromethane	0.217	-0.435	0.554	0.000	0.000	1.069	0.000
1,2-Dichloroethane	0.017	-0.337	1.600	0.774	0.637	0.921	0.000
Hexane	0.320	0.000	0.000	0.000	0.000	0.945	0.000
Heptane	0.284	0.000	0.000	0.000	0.000	0.950	0.000
Octane	0.219	0.000	0.000	0.000	0.000	0.960	0.000
Decane	0.159	0.000	0.000	0.000	0.000	0.972	0.000
Undecane	0.113	0.000	0.000	0.000	0.000	0.971	0.000
Dodecane	0.053	0.000	0.000	0.000	0.000	0.986	0.000
Hexadecane	0.000	0.000	0.000	0.000	0.000	1.000	0.000
Cyclohexane	0.163	-0.110	0.000	0.000	0.000	1.013	0.000
Methylcyclohexane	0.318	-0.215	0.000	0.000	0.000	1.012	0.000
Isooctane	0.264	-0.230	0.000	0.000	0.000	0.975	0.000
Benzene	0.107	-0.313	1.053	0.457	0.169	1.020	0.000
Toluene	0.121	-0.222	0.938	0.467	0.099	1.012	0.000
Fluorobenzene	0.181	-0.621	1.432	0.647	0.000	0.986	0.000
Chlorobenzene	0.064	-0.399	1.151	0.313	0.171	1.032	0.000
Bromobenzene	-0.064	-0.326	1.261	0.323	0.292	1.002	0.000
Iodobenzene	-0.171	-0.192	1.197	0.245	0.245	1.002	0.000
Nitrobenzene	-0.295	0.121	1.682	1.247	0.370	0.915	0.000
Benzonitrile	-0.075	-0.341	1.798	2.030	0.291	0.880	0.000
Diethylether	0.206	-0.169	0.873	3.402	0.000	0.882	0.000
Dipropylether	0.065	-0.202	0.776	3.074	0.000	0.948	0.000
Diisopropylether	0.114	-0.032	0.685	3.108	0.000	0.940	0.000
Dibutylether	0.369	-0.216	0.026	2.626	-0.499	1.124	0.000
Ethyl acetate	0.130	0.031	1.202	3.199	0.463	0.828	0.000
Butyl acetate	-0.664	0.061	1.671	3.373	0.824	0.832	0.000
Methyl isobutyl ketone	0.244	0.183	0.987	3.418	0.323	0.854	0.000
Tributylphosphate	0.097	-0.098	1.103	2.411	0.588	0.844	0.000
Gas–water	-1.271	0.822	2.743	3.904	4.814	-0.213	0.000

obtained from a refractive index at 293 K (for liquid solutes), or can be calculated from an estimated refractive index [10]. Both the available software programs [7, 8] for descriptors give calculated values of *E*. The *V*-descriptor can easily be calculated from its molecular formula [2, 11]. Thus we have three descriptors in Eq. 1 to determine (*S*, *A* and *B*).

However we can convert all values of  $\log_{10} P$  into corresponding gas–solvent partition coefficients, as  $\log_{10} K$ , through Eq. 3 where  $K_w$  is the gas to water partition coefficient, all partition coefficients being at 298 K. Note that  $K_w$  has no units. We take  $\log_{10} K_w$  as another unknown ‘descriptor’ and use both Eqs. 1 and 2 in our set of simultaneous equations. Coefficients in Eq. 2 are given in Table 1. Then even if we have a limited number of  $\log_{10} P$  values, say only five, we then have five equations in  $\log_{10} P$  and five equations in  $\log_{10} K$ . We also have two equations for  $\log_{10} K_w$ , see Table 1, making a total of 12 equations from which to derive five unknowns ( $S$ ,  $A$ ,  $B$ ,  $L$  and  $\log_{10} K_w$ ). This is the procedure we use for the determination of descriptors.

$$\log_{10} P = \log_{10} K - \log_{10} K_w \quad (3)$$

### 3 Results

The required values of  $\log_{10} P$  that we need to initiate the calculation of descriptors are known for acetylacetone and for a number of substituted compounds. Quite fortunately, Leo [12] has collected these  $\log_{10} P$  values, many of which are scattered over the literature, and lists them in his software program ‘BioLoom’. The  $\log_{10} P$  values that we use are nearly all from BioLoom. We start with acetylacetone itself. A value of  $E = 0.412$  from an experimental value of the refractive index [13] is available, and  $V = 0.8445$  [2, 11]. Partition coefficients into no less than 26 solvents are available [12]. Values of  $\log_{10} P$  into hexane, decane and butyl acetate were well out of line and were not used, leaving 23 data points. We have also 23 corresponding values of  $\log_{10} K$ , two equations in  $\log_{10} K_w$  and one equation for the NIST Kovats GC retention index, GCRI, leading to a total set of 49 equations. A value of GCRI = 790 for acetylacetone is listed in ChemSpider [13]. We have used NIST Kovats GC values to obtain Eq. 4:

$$\text{GCRI} = 69.6 + 12.1E + 76.3S + 200.0L \quad (4)$$

$$N = 286, SD = 46.4, R^2 = 0.992, F = 12079, PRESS = 634316, Q^2 = 0.992, PSD = 47.7$$

Here and elsewhere,  $N$  is the number of data points (compounds),  $SD$  is the regression standard deviation,  $R$  is the correlation coefficient,  $F$  is the  $F$ -statistic,  $PRESS$  and  $Q^2$  are the leave-one-out statistics and  $PSD$  is the predictive standard deviation [14]. In order not to bias the results, we use GCRI/100 in the set of simultaneous equations.

The total of 49 simultaneous equations were solved to yield the descriptors given in Table 2 with  $SD = 0.149 \log_{10}$  units. The 23 observed and calculated values of  $\log_{10} P$  are in Table 3. For this set of data the Absolute Error ( $AE$ ) = 0.018 and  $SD = 0.146 \log_{10}$  units. The descriptors for acetylacetone are not unusual, except that  $A = 0$ . It might be expected that acetylacetone would have some hydrogen bond acidity through the ‘active’  $\text{CH}_2$  group but we have enough data, with 49 equations, to be reasonably certain that  $A = 0$ .

For hexane-2,4-dione,  $\log_{10} P$  values into six solvents are known, and yield six corresponding  $\log_{10} K$  values. There are also two further equations in  $\log_{10} K_w$  and an equation in GCRI (890), giving a total of 15 equations. From a known refractive index,  $E = 0.381$  and  $V = 0.9854$ . The 15 equations were solved to yield the descriptors in Table 2 with an  $SD = 0.122 \log_{10}$  units.

**Table 2** Descriptors for pentane-2,4-dione and some of its derivatives

Compound	<i>E</i>	<i>S</i>	<i>A</i>	<i>B</i>	<i>V</i>	<i>L</i>	$\log_{10} K_w$
Pentane-2,4-dione	0.412	0.80	0.00	0.62	0.8445	3.347	3.59
Hexane-2,4-dione	0.381	0.78	0.00	0.65	0.9854	3.727	3.51
Heptane-2,4-dione	0.385	0.73	0.00	0.64	1.1263	4.204	3.22
Octane-2,4-dione	0.380	0.73	0.00	0.65	1.2672	4.780	3.14
Nonane-2,4-dione	0.380	0.76	0.00	0.66	1.4081	5.308	3.15
Heptane-3,5-dione	0.385	0.78	0.00	0.66	1.1263	4.217	3.45
Octane-3,5-dione	0.380	0.75	0.00	0.65	1.2672	4.787	3.19
Nonane-4,6-dione	0.399	0.77	0.00	0.66	1.4081	5.242	3.20
Undecane-5,7-dione	0.370	0.76	0.00	0.68	1.6899	6.223	3.03
Tridecane-6,8-dione	0.370	0.75	0.00	0.67	1.9717	7.371	2.70
5,5-Dimethylhexane-2,4-dione	0.380	0.81	0.00	0.73	1.2672	4.798	3.75
2,6-Dimethylheptane-3,5-dione	0.380	0.72	0.00	0.65	1.4081	5.206	2.99
2,8-Dimethylnonane-4,6-dione	0.370	0.77	0.00	0.72	1.6899	6.224	3.24
3-Methyl-2,4-pentanedione	0.380	0.77	0.00	0.65	0.9854	3.781	3.48
2,2,6,6-Tetramethylheptane-3,5-dione	0.360	0.78	0.00	0.72	1.6899	6.247	3.25
Benzoylacetone	1.000	1.06	0.00	0.60	1.3114	6.012	4.05
1,1,1-Trifluorobenzoylacetone	0.690	0.69	0.00	0.77	1.3645	5.494	3.72
Thenoylacetone	1.100	1.00	0.00	0.69	1.2361	5.810	4.46
1,1,1-Trifluoro-2-thenoylacetone	0.530	0.91	0.00	0.83	1.2892	5.210	4.58
Trifluoroacetylacetone	0.106	0.47	0.16	0.72	0.8976	2.878	3.60
Hexafluoroacetylacetone <sup>a</sup>	-0.217	0.07	0.32	0.80	0.9507	2.340	3.34

<sup>a</sup> Provisional only

There are only four partition coefficients available for heptane-2,4-dione, but these still yield 11 equations (GCRI = 989). A value for *E* (0.385) can be obtained from a known refractive index and *V* = 1.1263 and the 11 equations solved with *SD* = 0.089 to give the descriptors in Table 2.

In the case of octane-2,4-dione,  $\log_{10} P$  values are known only for partition into heptane and tetrachloromethane, leading to a total of seven equations. *E* was estimated as 0.380, *A* was taken as zero and *V* calculated as 1.2672. The seven equations were solved with *SD* = 0.031 to give the descriptors in Table 2.

A  $\log_{10} P$  value into tetrachloromethane is all that is available for nonane-2,4-dione. With GCRI = 1188 [13] we have only five equations. *E* was estimated as 0.380, *A* was taken as zero and *V* calculated as 1.4084. This is just enough to obtain the descriptors in Table 2.

Seven  $\log_{10} P$  values are listed for heptane-3,5-dione, and lead to 17 equations (GCRI = 989). An experimental value of the refractive index leads to *E* = 0.389, *V* is calculated as 1.1263, and the 17 equations can be solved with *SD* = 0.143 to give the descriptors in Table 2.

For octane-3,5-dione only two partition coefficients were available, so that we have seven equations. *E* was estimated as 0.380, *A* was taken as 0.00 and *V* calculated as 1.2672. The equations were solved to give the descriptors in Table 2 with an *SD* of 0.063  $\log_{10}$  units.

**Table 3** Calculated and observed values of  $\log_{10} P$  for water–solvent partition of 2,4-pentanedione (acac)

Solvent	Calc	Obs
Pentan-1-ol	0.70	0.43
Hexan-1-ol	0.60	0.35
Heptan-1-ol	0.59	0.32
Octan-1-ol	0.55	0.40
Nonan-1-ol	0.43	0.26
Dichloromethane	1.33	1.33
Trichloromethane	1.45	1.40
Tetrachloromethane	0.54	0.52
1,2-Dichloroethane	1.07	1.08
Heptane	−0.12	−0.10
Octane	−0.18	0.08
Nonane	−0.14	−0.01
Cyclohexane	−0.05	−0.05
Benzene	0.79	0.76
Toluene	0.66	0.62
<i>m</i> -Xylene	0.55	0.54
Chlorobenzene	0.79	0.75
Bromobenzene	0.75	0.79
Iodobenzene	0.67	0.79
Nitrobenzene	0.88	0.84
Dibutylether	0.14	0.04
Tributylphosphate	0.52	0.72
Methyl isobutylketone	0.57	0.80

There is more data for nonane-4,6-dione. We used seven  $\log_{10} P$  values, which translated into 17 equations (GCRI = 1188). A known refractive index gave  $E = 0.399$  and  $V = 1.4081$ . The set of equations was solved with an  $SD = 0.167$ , to give the descriptors in Table 2.

For undecane-5,7-dione we used five partition coefficients, leading to 12 equations. We estimated  $E$  as 0.37, calculated  $V$  as 1.6899 and solved the set of equations to yield the descriptors in Table 2 with an  $SD$  value of 0.150  $\log_{10}$  units.

As for undecane-5,7-dione we had only five partition coefficients for tridecane-6,8-dione. Taking  $E = 0.37$  and  $V = 1.9717$  we solved the set of 12 equations to obtain the descriptors in Table 2 with a rather large  $SD$  of 0.178  $\log_{10}$  units.

There are also a number of branched chain alkyl derivatives for which partition coefficients are available [12]. For 5,5-dimethylhexane-2,4-dione partition coefficients are known into five solvents, and with GCRI (1004) we have 13 equations. With  $E = 0.38$  and  $V = 1.2672$  we solved the set of equations to obtain the descriptors in Table 2;  $SD = 0.128 \log_{10}$  units.

Partition coefficients are available for 2,6-dimethylheptane-3,5-dione and with GCRI = 1060 we had 15 equations. The given experimental value [12] for partition into octanol,  $\log_{10} P = 2.22$ , was well out of line and was omitted. The resulting 13 equations with  $E = 0.38$  and  $V = 1.4081$  gave the descriptors in Table 2 with  $SD = 0.162 \log_{10}$  units.

The only partition coefficient that we could find for 2,8-dimethyl-4,6-dione was that of  $\log_{10} P = 4.05$  for partition into benzene [14], but a value of 1258 was available for GCRI. These yielded only five equations. We estimated  $E$  as 0.37, we know that  $V = 1.6899$ , and in order to solve the equations we also estimated that  $B = 0.72$ . Then solution of the equations gave  $SD = 0.076$  and the remaining descriptors as shown in Table 2.

There are two other alkyl derivatives of acac that have been used as complexing agents, 3-methylpentane-2,4-dione and 2,2,6,6-tetramethylheptane-3,5-dione. There is insufficient data on these compounds to yield a set of equations that can be solved to get descriptors, but from the results we have for the other alkyl derivatives, we estimate the descriptors as shown in Table 2.

A number of other derivatives of acetylacetone have been widely used as complexing agents; for several of these compounds, numerous values of  $\log_{10} P$  are known [12]. We start with benzoylacetone (1-phenylbutane-1,3-dione) for which partition into 20 solvents has been studied. A value of 1364 for GCRI is available [13] and so we have no less than 43 equations on the lines of Eqs. 1 and 2. We took  $E = 1.00$  from addition of fragments and also from calculations [7, 8] and  $V = 1.3114$ . The equations were solved to yield the descriptors in Table 2 with a very small value of  $SD = 0.086 \log_{10}$  units. The observed and calculated values of  $\log_{10} P$  are in Table 4. For the 20 solvents,  $AE = 0.01$  and  $SD = 0.086 \log_{10}$  units

Partitions into 11 solvents are known for 1,1,1-trifluorobenzoylacetone. The value of  $\log_{10} P$  into trichloromethane was quite out of line (obs. 2.73, calc. 3.28) and if this is left out we have 23 equations (GCRI = 1198). We estimated  $E = 0.69$  from values for pentane-2,4-dione, 1,1,1-trifluoropentane-2,4-dione and benzoylacetone, and calculated  $V = 1.3645$ . The set of equations were solved to give the descriptors in Table 2 with  $SD = 0.126 \log_{10}$  units.

**Table 4** Calculated and observed values of  $\log_{10} P$  for water–solvent partition of benzoylacetone

Solvent	Calc	Obs
Hexanol	2.46	2.48
Octanol	2.46	2.52
Nonanol	2.37	2.43
Decanol	2.37	2.38
Dichloromethane	3.44	3.48
Trichloromethane	3.52	3.50
Tetrachloromethane	2.79	2.77
1,2-Dichloroethane	3.24	3.23
Heptane	1.99	2.03
Octane	1.98	2.00
Nonane	1.97	1.97
Decane	1.95	2.00
Cyclohexane	2.21	2.11
Benzene	3.10	3.05
Toluene	2.96	2.91
m-Xylene	2.83	2.81
Chlorobenzene	3.13	3.13
Bromobenzene	3.06	3.36
Nitrobenzene	3.25	3.19
Dibutylether	2.50	2.36



For thenoylacetone (1-(2-thienyl)butane-1,3-dione) we have  $\log_{10} P$  values into hexane and benzene. Together with a value of 1385 for GCRI these gave seven equations. We estimated  $E = 1.10$  by addition of fragments, calculated  $V$  as 1.2361 and solved the equations to give the descriptors in Table 2 with  $SD = 0.039 \log_{10}$  units.

There are a large number of  $\log_{10} P$  values available for trifluorothenoylacetone (4,4,4-trifluoro-1-(2-thienyl)butane-1,3-dione). These include values for partition into numerous esters for which we have no coefficients in Eqs. 1 and 2. For partition into ethyl acetate and butyl acetate, however, the observed values of  $\log_{10} P$  are so far away from our calculated values that we suggest all the given  $\log_{10} P$  values into esters be used with caution. We were left with 12 values of  $\log_{10} P$ , together with a value of 1199 for GCRI, leading to 27 equations. A calculated refractive index [10] leads to  $E = 0.524$ , close to a calculated value for  $E$  of 0.53 [8]. We used the latter value and our calculated value of  $V = 1.2892$ , and solved the 27 equations to give the descriptors in Table 2 with an  $SD$  of  $0.101 \log_{10}$  units.

There are also  $\log_{10} P$  values for 2-furoyltrifluoroacetone and pivaloyltrifluoroacetone, but we could not obtain any reasonable set of descriptors for these two compounds.

Finally we deal with trifluoroacetylacetone (1,1,1-trifluoropentane-2,4-dione) and hexafluoroacetylacetone (1,1,1,3,3,3-hexafluoropentane-2,4-dione). For trifluoroacetylacetone we have  $\log_{10} P$  values into 15 solvents. The value of  $\log_{10} P$  into trichloromethane was considerably out of line (calc. 0.94, obs. 0.33) and was left out. With  $GCRI = 624$  this leaves 31 equations to solve. An experimental refractive index of 1.3890 [13] leads to  $E = 0.106$  and with  $V = 0.8976$  we obtained the descriptors in Table 2 with an  $SD$  of  $0.125 \log_{10}$  units. The calculated and observed values of  $\log_{10} P$  are in Table 5, and yield  $AE = 0.011$  and  $SD = 0.123 \log_{10}$  units (omitting trichloromethane). It is noteworthy that the  $A$ -descriptor is not zero, but with a set of 31 equations, we can be reasonably confident about this descriptor.

The position with hexafluoroacetylacetone is not straightforward. We have four values [15] of  $\log_{10} P$  into trichloromethane (−1.75), tetrachloromethane (−1.92), hexane (−2.04) and benzene (−1.91), and also a value of GCRI (459), leading to eleven equations.

**Table 5** Calculated and observed values of  $\log_{10} P$  for water–solvent partition of trifluoroacetylacetone

Solvent	Calc	Obs
Pentanol	0.72	0.52
Hexanol	0.67	0.62
Nonanol	0.49	0.59
Dichloromethane	0.70	0.40
Trichloromethane <sup>a</sup>	0.94	0.33
Tetrachloromethane	−0.01	0.02
Hexane	−0.52	−0.51
Heptane	−0.55	−0.57
Octane	−0.63	−0.62
Dodecane	−0.75	−0.76
Benzene	0.13	0.15
Toluene	0.04	0.13
Ethylbenzene	−0.10	−0.18
Nitrobenzene	0.10	0.30
Dibutylether	0.04	0.08

<sup>a</sup> Not used in the calculations of descriptors

The solution of this set of simultaneous equations yields completely unreasonable values for the descriptors. Stokely [16] has shown that hexafluoroacetylacetone decomposes in water. He measured a value for  $\log_{10} P$  into benzene of  $-0.42$  (in contrast to the value of  $-1.91$  [15]), and found that the partition coefficient decreased with time. We can obtain a value of  $-0.217$  for  $E$  from the refractive index and we can calculate  $V = 0.9507$ , but there is still not enough data to obtain a full set of descriptors. We can deduce that  $B = 0.80$  and  $L = 2.340$  by comparison to other compounds in Table 2, and from Absolv calculations [7]. Then with  $S = 0.07$  and  $A = 0.32$  we can reproduce Stokely's [16] value of  $-0.42$  for  $\log_{10} P$  into benzene, and the associated values of  $\log_{10} K$  into benzene and  $\log_{10} K$ , with the descriptors in Table 2. However, we caution that these results must be regarded as provisional only.

## 4 Discussion

We have managed to obtain descriptors for acetylacetone and 21 of its derivatives, as shown in Table 2. These can be combined with the equation coefficients in Table 1 to yield estimates of partition coefficients from water and the gas phase into all the listed solvents, and (hypothetical) partition coefficients into a large number of dry solvents for which we have also determined equation coefficients [17–19]. In addition we have determined equation coefficients for partition into water–ethanol [20, 21] and water–methanol mixtures [22], and so values of  $\log_{10} P$  and  $\log_{10} K$  into these solvent mixtures can also be estimated. In addition to the usual organic solvents, we have also studied ionic liquids [23], and partitions into these solvents can be estimated for the various acetylacetonates. Partitions or permeations in biological systems [24–26] can also be estimated from the descriptors listed in Table 2.

Inspection of the descriptors themselves shows that all the acetylacetonates are quite polar, with substantial values of the  $S$ -descriptor, and, as expected from the presence of the two carbonyl groups, are quite strong hydrogen bond bases, with  $B$ -values almost double those for simple aliphatic ketones which have  $B$ -values around 0.45 [7, 8]. Perhaps unexpectedly, the alkylsubstituted acetylacetonates all have zero hydrogen bond acidity, as do some of the trifluoro derivatives. Only with trifluoroacetylacetone, and with hexafluoroacetylacetone are significant values of the  $A$ -descriptor found.

The  $L$ -values form a very regular series, and can be taken to show the internal consistency of our set of descriptors. For the acetylacetonates with linear alkyl substituents we find Eq. 5, where CN is the number of carbon atoms.


$$L = 0.7132 + 0.5069 \text{ CN} \quad (5)$$

$$N = 10, SD = 0.059, R^2 = 0.998, F = 3734.0$$

$$PRESS = 0.06235, Q^2 = 0.995, PSD = 0.089$$

The branched chain substituents behave remarkably similarly to the linear chain substituents, and for all the alkyl substituted acetylacetonates we find Eq. 6.

$$L = 0.7516 + 0.5015 \text{ CN} \quad (6)$$



**Fig. 1** Plot of the descriptor  $L$  against the number of carbon atoms, CN, in alkyl pentane-2,4-diones, *open circle* linear alkylpentane-2,4-diones, *filled circle* branched chain alkylpentane-2,4-diones

**Table 6** Observed and calculated values of  $\log_{10} P$  for acetylacetonates

Compound	Obs	Calc			acd [10]	PHA [7]
		This work	[12]	[27]		
Pentane-2,4-dione	0.40	0.55	-0.46	0.05	0.34	0.58
Hexane-2,4-dione		0.99	0.07	0.54	0.87	0.91
Heptane-2,4-dione		1.62	0.60	1.03	1.40	1.28
Octane-2,4-dione		2.12	1.13	1.53	1.93	1.76
Nonane-2,4-dione		2.59	1.65	2.02	2.46	2.39
Heptane-3,5-dione		1.50	0.60	1.02	1.40	1.28
Octane-3,5-dione		2.09	1.13	1.53	1.93	1.76
Nonane-4,6-dione	2.23	2.59	1.65	2.02	2.46	2.24
Undecane-5,7-dione	3.56	3.59	2.71	3.00	2.99	3.53
Tridecane-6,8-dione	4.88	4.71	3.77	3.98	4.59	4.39
5,5-Dimethylhexane-4,6-dione	1.67	1.75	1.00	1.41	1.57	1.67
2,6-Dimethylheptane-3,5-dione	2.22	2.66	1.65	1.87	2.10	2.09
2,8-Dimethylnonane-4,6-dione		3.44	2.45	2.85	3.16	2.92
3-Methyl-2,4-pentanedione		1.00	0.07	0.47	0.69	0.92
2,2,6,6-Tetramethylheptane-3,5-dione		3.42	2.45	2.78	2.80	2.91
Benzoylacetone	2.52	2.46	1.09	0.91	2.52	1.85
1,1,1-Trifluorobenzoylacetone		2.29	1.65	1.02	4.17	2.54
Thenoylacetone		1.98	0.86	0.43	2.11	1.29
1,1,1-Trifluorothenoxyacetone	1.46	1.47	1.42	0.84	3.76	1.99
Trifluoroacetylacetone		0.59	0.10	0.47	2.09	1.32
Hexafluoroacetylacetone		0.76	0.66	0.88	3.84	1.69

**Table 7** Comparison of observed and calculated values of  $\log_{10} P$  for acetylacetonates

	This work	[12]	[27]	[10]	[7]
AE <sup>a</sup>	-0.10	0.76	0.60	-0.17	0.08
SD	0.23	0.91	0.80	0.91	0.38

<sup>a</sup> Obs - Calc

$$N = 15, SD = 0.053, R^2 = 0.998, F = 6169.3$$

$$PRESS = 0.0616440, Q^2 = 0.997, PSD = 0.069$$

The plots of  $L$  against CN are excellent, as shown in Fig. 1. Equation 5 or especially Eq. 6 could be used to estimate an  $L$ -value for any alkylsubstituted acetylacetonate.

Once we have descriptors for the acetylacetonates, we can then deduce the corresponding water–octanol partition coefficients, as  $\log_{10} P$ . These partition coefficients are of considerable interest, as they are often used as a measure of hydrophobicity of solutes, and they are the most commonly estimated of all water–solvent partition coefficients. We can compare our own calculated values with those calculated through four very common methods, the ClogP program of Leo [12], the EPI Suite TM [27], the ACD program in ChemSketch [10] and the ACD program that is part of the Absolv ADME Suite [7]. In addition, we can compare all the calculated values with the (few) observed values. Details are in Table 6. There are eight compounds for which observed values are available, and a comparison of these with the various calculated values, in terms of the average error and standard deviation, is in Table 7.

Inspection of Tables 6 and 7 suggests that where our descriptors are available, they yield estimates of water–octanol partition coefficients, as  $\log_{10} P$ , that are at least as good as those from standard calculation methods [7, 10, 12, 27]. In addition, use of our descriptors has the advantage that water–solvent partition coefficients can be estimated for a very large number of organic solvents. The deviations in observed and calculated values of  $\log_{10} P$  for the eight acetylacetonates are quite similar thus indicating that the errors in the descriptors for the various acetylacetonates are also quite close.

## 5 Conclusions

We have been able to obtain Abraham or Absolv descriptors for pentane-2,4-dione and 21 of its derivatives. These descriptors encode important chemical properties, and show that pentane-2,4-dione itself has no hydrogen bond acidity, but that the trifluoro- and hexafluoro-derivatives have substantial hydrogen bond acidity. The descriptors for the 22 compounds enable partition coefficients to be estimated for partition from water to a very large number of organic solvents. In the case of water–octanol partition coefficients we show that estimations through our descriptors are at least as good as the best calculations through well-known calculational programs.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution,

and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

1. Stevens, G.W., Perera, J.M., Grieser, F.: Metal ion extraction. *Curr. Opin. Colloid Interface. Sci.* **2**, 629–634 (1997)
2. Abraham, M.H.: Scales of hydrogen bonding: their construction and application to physicochemical and biochemical processes. *Chem. Soc. Rev.* **22**, 73–83 (1993)
3. Abraham, M.H., Ibrahim, A., Zissimos, A.M.: The determination of sets of solute descriptors from chromatographic measurements. *J. Chromatogr. A* **1037**, 29–47 (2004)
4. Poole, C.F., Atapattu, S.N., Poole, S.K., Bell, A.K.: Determination of solute descriptors by chromatographic methods. *Anal. Chim. Acta* **652**, 32–53 (2009)
5. Clarke, E.D., Mallon, L.: The determination of Abraham descriptors and their application to crop protection research. In: Jeschke, P., Kramer, W., Schirmer, U., Witschel, M. (eds.) *Modern Methods in Crop Protection Research*. Wiley, Weinheim (2012)
6. Abraham, M.H., Acree Jr., W.E.: Descriptors for the prediction of partition coefficients and solubilities of organophosphorus compounds. *Sep. Sci. Technol.* **48**, 884–897 (2013)
7. Absolv. ADME Suite 5.0, Advanced Chemistry Development, 110 Yonge Street, 14th Floor, Toronto, Ontario, M5C 1T4, Canada
8. Endo, S., Brown, T.N., Watanabe, N., Ulrich, N., Bronner, G., Abraham, M. H., Goss, K-U.: UFZ-LSER database v 3.1 [Internet], Leipzig, Germany, Helmholtz Centre for Environmental Research-UFZ. (2015). w.ufz.de/lserd, <http://www.ufz.de/lserd>. Accessed 9 July 2017
9. Abraham, M.H., Leo, A.J.: Partition between phases of a solute that exists as two interconverting species. *J. Chem. Soc. Perkin Trans.* **2**, 1839–1842 (1995)
10. ChemSketch, Advanced Chemistry Development, 110 Yonge Street, 14th Floor, Toronto, Ontario, M5C 1T4, Canada
11. Abraham, M.H., McGowan, J.C.: The use of characteristic volumes to measure cavity terms in reversed-phase liquid chromatography. *Chromatographia* **23**, 243–246 (1987)
12. Leo, A.J.: ‘BioLoom’, BioByte Corp, 201W. 4th Street, #204 Claremont, CA 91711–4707, USA
13. [www.chemspider.com](http://www.chemspider.com). Accessed 9 July 2017
14. Koshimura, H., Okubo, T.: Effect of substituents on the distribution coefficients of alkyl-substituted  $\beta$ -diketones and their copper and iron chelates. *Anal. Chim. Acta* **55**, 163–168 (1971)
15. Sekine, T., Hasegawa, Y., Ihara, N.: Acid dissociation and two-phase distribution constants of eight  $\beta$ -diketones in several solvent extraction systems. *J. Inorg. Nucl. Chem.* **35**, 3968–3970 (1973)
16. Stokely, J.R.: *Solvent Extraction and Gas Chromatography of Metal Fluoroacetylacetonates*. Ph.D. Thesis, Clemson University (1966)
17. Acree Jr., W.E., Horton, M.Y., Higgins, E., Abraham, M.H.: Abraham model linear free energy relationships as a means of extending solubility studies to include the estimation of solute solubilities in additional organic solvents. *J. Chem. Thermodyn.* **102**, 392–397 (2016)
18. Schmidt, A., Grover, D., Zetti, H., Koshevarova, V., Dai, C., Zhang, S., Hart, E., Brumfield, M., De La Rosa, J., Portillo, V., Pugh, A., Sanchez, A., Acree Jr., W.E., Abraham, M.H.: Determination of Abraham model solute descriptors for isophthalic acid from experimental solubility data in organic solvents at 298 K. *Phys. Chem. Liquids* **54**, 747–757 (2016)
19. Abraham, M.H., Acree Jr., W.E.: Descriptors for the prediction of partition coefficients for 8-hydroxyquinoline and its derivatives. *Sep. Sci. Technol.* **49**, 2135–2141 (2014)
20. Abraham, M.H., Acree Jr., W.E.: Partition coefficients and solubilities of compounds in the water–ethanol solvent system. *J. Solution Chem.* **40**, 1279–1290 (2011)
21. Abraham, M.H., Acree Jr., W.E.: Equations for the partition of neutral molecules, ions and ionic species from water to water–ethanol mixtures. *J. Solution Chem.* **41**, 730–740 (2012)
22. Abraham, M.H., Acree Jr., W.E.: Equations for the partition of neutral molecules, ions and ionic species from water to water–methanol mixtures. *J. Solution Chem.* **45**, 861–874 (2016)
23. Jang, B., Horton, M.Y., Acree Jr., W.E., Abraham, M.H.: Ion-specific equation coefficient version of the Abraham model for ionic liquid solvents: determination of coefficients for tributylethylphosphonium, 1-butyl-1-methylmorpholinium, 1-allyl-3-methylimidazolium and octyltriethylammonium cations. *Phys. Chem. Liquids* **55**, 358–385 (2017)

24. Zhang, K., Chen, M., Scriba, G.K.E., Abraham, M.H., Fahr, A., Lui, X.: Human skin permeation of neutral species and ionic species: extended linear free-energy relationship analysis. *J. Pharm. Sci.* **101**, 2034–2044 (2012)
25. Abraham, M.H., Gola, J.R.M., Ibrahim, A., Acree Jr., W.E., Liu, X.: The prediction of blood–tissue partitions, water–skin partitions and skin permeation for agrochemicals. *Pest Manag. Sci.* **70**, 1130–1137 (2014)
26. Abraham, M.H., Gola, J.R.M., Ibrahim, A., Acree Jr., W.E., Liu, X.: A simple method for estimating in vitro air–tissue and in vivo blood–tissue partition coefficients. *Chemosphere* **120**, 188–191 (2015)
27. EPI Suite TM, United States Environmental Protection Agency (2000–2012)