

<https://helda.helsinki.fi>

---

## The plastid genome of the invasive plant common ragweed (*Ambrosia artemisiifolia*, Asteraceae)

Amiryousefi, Ali

2017

---

Amiryousefi, A, Hyvönen, J & Poczai, P 2017, ' The plastid genome of the invasive plant common ragweed ( *Ambrosia artemisiifolia* , Asteraceae) ' , Mitochondrial DNA Part B, Resources , vol. 22 , pp. 753-754 . <https://doi.org/10.1080/23802359.2017.1390423>

---

<http://hdl.handle.net/10138/231051>

<https://doi.org/10.1080/23802359.2017.1390423>

---

cc\_by

publishedVersion


---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*



## Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <http://www.tandfonline.com/loi/tmdn20>

# The plastid genome sequence of the invasive plant common Ragweed (*Ambrosia artemisiifolia*, Asteraceae)

Ali Amiryousefi, Jaakko Hyvönen & Péter Poczai

To cite this article: Ali Amiryousefi, Jaakko Hyvönen & Péter Poczai (2017) The plastid genome sequence of the invasive plant common Ragweed (*Ambrosia artemisiifolia*, Asteraceae), *Mitochondrial DNA Part B*, 2:2, 753-754, DOI: [10.1080/23802359.2017.1390423](https://doi.org/10.1080/23802359.2017.1390423)

To link to this article: <http://dx.doi.org/10.1080/23802359.2017.1390423>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 18 Oct 2017.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Full Terms & Conditions of access and use can be found at  
<http://www.tandfonline.com/action/journalInformation?journalCode=tmdn20>

## The plastid genome sequence of the invasive plant common Ragweed (*Ambrosia artemisiifolia*, Asteraceae)

Ali Amiryousefi<sup>a,b</sup>, Jaakko Hyvönen<sup>b</sup>  and Péter Poczai<sup>a</sup> 

<sup>a</sup>Finnish Museum of Natural History (Botany), University of Helsinki, Helsinki, Finland; <sup>b</sup>Department of Bioscience (Plant Biology), Viikki Plant Science Centre, University of Helsinki, Helsinki, Finland

### ABSTRACT

In the current study, we present the complete chloroplast genome sequence of *Ambrosia artemisiifolia*. The genome is 152,223 bp long and consist of 83 protein coding genes, 38 tRNAs, and four rRNAs duplicated in the inverted repeat. Detected large single-copy (LSC) and small single-copy (SSC) regions separated with two inverted repeat regions (IR) of length 25,098. The phylogenetic hypotheses obtained based on the analyses of 18 cp genomes places common ragweed within the tribe Heliantheae of the Asteraceae.

### ARTICLE HISTORY

Received 5 October 2017  
Accepted 6 October 2017

### KEYWORDS

Chloroplast genome; *de novo* assembly; genome skimming; phylogenomics; plastid evolution


*Ambrosia artemisiifolia* L. (common ragweed) is one of the most prevalent invasive plants. The main concern regarding this plant is its large production of wind dispersed pollen that causes allergy for 15–60% of the European population (Taramarcz et al. 2005). One single plant can produce six billion pollen grains and thousands of seeds during its lifecycle (Kazinczi et al. 2008). Besides its major health impact, it is causing serious economical problems for crop yield in agriculture. Yield losses alone were estimated to be €130 million per year for certain countries (Kórníves et al. 2006). Ragweed is a plant of concern especially now in the era of warming climate because tests have shown that higher levels of carbon dioxide will greatly increase pollen production, and thus raising the number of people suffering of allergic reactions (Virág et al. 2016).

We extracted DNA according to Shi et al. (2012) from 20 g fresh ragweed leaves collected in Serbia (43.276149, 21.902976; voucher P.Poczai 0012867). Paired-end libraries of 2 × 150 bp were prepared with Illumina TruSeq DNA Sample prep kit and sequencing was carried out on an Illumina MiSeq platform. Raw reads were filtered with Trimmomatic (Bolger et al. 2014), and *de novo* assembly of the plastid genome was carried out with the Geneious R10 assembler platform (Kearse et al. 2012). We annotated the genome using Geneious and in-house scripts. Here, we report the complete chloroplast sequence of *Ambrosia artemisiifolia* to provide resources for taxonomic studies and invasive weed biology.

The complete chloroplast genome of *Ambrosia artemisiifolia* (GenBank accession MG019037) has a total length of 152,223 bp which is divided by two IR regions of 25,098 bp. This genome comprises of 125 genes and has 38% overall GC content. The genes are classified into 38 tRNA, 4 rRNA and 83 coding-protein genes.

Using the RAxMLv8.0 (Stamatakis 2014) the best scoring ML tree with 10,000 bootstrap replicates was calculated under GTR-GAMMA after running jModelTest2 (Darriba et al. 2012) including 16 representative species of the Heliantheae tribe (Asteraceae) and two outgroup species of Apiales (Figure 1). We also made phylogenetic analysis using parsimony as an optimality criterion and obtained similar topology. The same matrix was analyzed also with parsimony as an optimality criterion using WinClada (Nixon 2002) and TNT (Goloboff et al. 2008). Prior to the analysis we used the WinClada command 'Mop uninformative characters' to exclude parsimony uninformative characters. This resulted in a matrix with 30,216 characters and due to its small size, we were able to perform analyses using implicit enumeration of the TNT that ensures finding optimal tree(s). Phylogenetic hypothesis obtained by our study supported previous topologies (Panero and Funk 2008).

We expect this sequence to clarify the taxonomic status of the *Ambrosia* genus within the separate Ambrosiinae subtribe, and provide additional genomic resources for invasive plant genomics.



**Figure 1.** The ML tree of 18 selected chloroplast genome sequences and *Ambrosia artemisiifolia*. The values on the node show the bootstraps of 10,000 replicates and scale is substitution per site.

## Acknowledgements


We thank staff and colleagues of the Viikki Biocenter who kindly contributed reagents, materials and analyses tools for our study.

## Disclosure statement

The authors claim no conflict of interest.

## ORCID

Jaakko Hyvönen  <http://orcid.org/0000-0001-7559-8295>

Péter Poczai  <http://orcid.org/0000-0002-0107-1068>

## References

- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. 30:2114–2120.
- Darriba D, Taboada GL, Doallo R, Posada D. 2012. jModelTest 2: more models, new heuristics and parallel computing. *Nat Methods*. 9:772.
- Goloboff PA, Farris JS, Nixon KC. 2008. TNT, a free program for phylogenetic analysis. *Cladistics*. 24:774–786.
- Kazinczi G, Beres I, Novak R, Birò K, Pathy Z. 2008. Common ragweed (*Ambrosia artemisiifolia*): a review with special regards to the results in Hungary. I. Taxonomy, origin and distribution, morphology, life cycle and reproduction strategy. *Herbologia*. 9:55–91.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, et al. 2012. Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*. 28:1647–1649.
- Kömíves T, Béres I, Reisinger P, Lehoczky É, Berke J, Tamás J, Páldy A, Csornai G, Nádor G, Kardeván P, et al. 2006. New strategic programme of the integrated ragweed control. *Hungarian Weed Res Technol*. 7:5–51.
- Nixon KC. 2002. WinClada 1.00.08. [accessed 2017 Mar 10]. <http://www.cladistics.com>
- Panero JL, Funk VA. 2008. The value of sampling anomalous taxa in phylogenetic studies: major clades of the Asteraceae revealed. *Mol Phylogenet Evol*. 47:757–782.
- Shi C, Hu N, Huang H, Gao J, Zhao Y-J, Gao L-Z. 2012. An improved chloroplast DNA extraction procedure for whole plastid genome sequencing. *PLoS One*. 7:e31468.
- Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*. 30:1312–1313.
- Tamarcaz P, Lambelet C, Clot B, Keimer C, Hauser C. 2005. Ragweed (*Ambrosia*) progression and its health risks: will Switzerland resist this invasion? *Swiss Med Wkly*. 135:538–548.
- Virág E, Hegedűs G, Barta E, Nagy E, Mátyás K, Kolics B, Tallér J. 2016. Illumina sequencing of common (short) ragweed (*Ambrosia artemisiifolia* L.) reproductive organs and leaves. *Front Plant Sci*. 7:1506.