

1 **Fine mapping of genes determining extrafusal fiber properties in murine soleus**
2 **muscle**

3 Carroll AM^{1,2}, Cheng R³, Collie-Duguid ESR^{1,4}, Meharg C⁵, Scholz ME¹, Fiering S⁶, Fields
4 JL⁶, Palmer AA⁷, Lionikas A¹.

5

6 Affiliation:

7 ¹School of Medicine, Medical Sciences and Nutrition, University of Aberdeen, Foresterhill
8 Aberdeen, AB25 2ZD, UK

9 ²The New Zealand Institute for Plant & Food Research Limited, Batchelar Road, Private Bag
10 11600, Palmerston North 4442, New Zealand

11 ³Research School of Biology, The Australian National University, 134 Linnaeus Way, Acton,
12 ACT 2601, Australia

13 ⁴Centre for Genome Enabled Biology and Medicine, University of Aberdeen, Old Aberdeen,
14 Aberdeen, AB24 3RY, UK

15 ⁵Institute for Global Food Security, Queen's University Belfast, 18-30 Malone Road, Belfast,
16 BT9 5BN, UK

17 ⁶Department of Microbiology/immunology, Geisel School of Medicine at Dartmouth,
18 Hanover, NH 03755, USA

19 ⁷Department of Psychiatry, University of California San Diego, 9500 Gilman Drive, Mail Code
20 0667, BRF2 3A24, La Jolla, CA 92093, USA

21

22

23

24

25

26

27 **Abstract**

28 **Introduction.** Muscle fiber cross-sectional area (CSA) and proportion of different fiber types
29 are important determinants of muscle function and overall metabolism. Genetic variation
30 plays a substantial role in phenotypic variation of these traits, however, the underlying genes
31 remain poorly understood.

32 **Aims.** This study aimed to map quantitative trait loci (QTL) affecting differences in soleus
33 muscle fiber traits between the LG/J and SM/J mouse strains.

34 **Methods.** Fiber number, CSA, and proportion of oxidative type I fibers were assessed in the
35 soleus of 334 genotyped female and male mice of the F₃₄ generation of advanced intercross
36 lines (AIL) derived from the LG/J and SM/J strains. To increase the QTL detection power,
37 these data were combined with 94 soleus samples from the F₂ intercross of the same
38 strains. Transcriptome of the soleus muscle of LG/J and SM/J females was analysed using
39 microarray.

40 **Results.** Genome-wide association analysis mapped 4 QTL (genome-wide $p < 0.05$)
41 affecting the properties of muscle fibers to Chromosome 2, 3, 4 and 11. A 1.5-LOD QTL
42 support interval ranged between 2.36 Mb and 4.67 Mb. Based on the genomic sequence
43 information, functional and transcriptome data, candidate genes were identified for each of
44 these QTL.

45 **Conclusion.** Combination of analyses in F₂ and F₃₄ AIL populations with transcriptome and
46 genomic sequence data in the parental strains is an effective strategy for refining QTL and
47 nomination of the candidate genes.

48

49 Key words: skeletal muscle, muscle fiber types, genetic variation

50

51 **Introduction**

52 Skeletal muscle plays a broad range of biological functions including locomotion,
53 thermoregulation, respiration, postural support, protection of bones and viscera; as well as
54 serving as a source of amino acids in times of starvation or disease. Muscle tissue in
55 livestock also provides an essential source of dietary proteins. In humans, there is more
56 than a 2-fold difference in muscle mass between individuals of similar age and same sex (3,
57 33). This is the outcome of variability in the number of muscle fibers and their size (51).
58 These differences are of clinical relevance. Variability in muscle mass significantly impacts
59 energy expenditure (58), influencing preponderance to obesity. In addition, individuals with
60 lower muscle mass may be more vulnerable to impairment of these vital functions due to
61 aging and/or disease related muscle loss. It has recently been reported that there is a
62 positive association between muscle mass and longevity in older adults (66).

63
64 Human skeletal muscles are mainly comprised of a mixture of type I, IIA and IIX muscle
65 fibers (62). The number of fibers, their size and varying proportions of the fiber types affect
66 morphological and functional properties of the muscle (6). A larger diameter of the fibers
67 and higher number of fibers typically leads to augmented muscular strength and power (25,
68 28). The proportion of type I muscle fibers is a factor determining success in endurance
69 sporting events (15, 18) and overall metabolism in humans (24, 29, 44, 74). In livestock,
70 proportion of oxidative type I fibers is associated with meat quality (65).

71
72 In humans, genetic factors account for around half of the variation in strength (19, 24, 74)
73 and the upper limit heritability is even greater (over 0.9) for muscle mass (26). Heritability
74 estimates of proportion of type I fibers is also high, ranging between 0.4 and 0.9, indicating
75 that genetic factors play an important role in determining muscle fiber properties (37, 63).
76 Effects of genetic factors on muscle fibers have also been demonstrated in mouse (20, 22,

77 59), cattle (68), sheep (10, 38) and pig (71). However, the specific genes underlying these
78 effects remain largely elusive.

79

80 Attempts at mapping the polygenic architecture of muscle fiber properties in mouse (11), pig
81 (17, 43, 52, 55, 77), cattle (1) and carp (80) have been made. A number of QTL have been
82 identified in these studies. However, the resolution achieved in the F_2 population is not
83 adequate for reliable nomination of the candidate genes in the majority of the QTLs of
84 polygenic traits. The mouse soleus muscle (primarily consists of type I and IIA fiber types),
85 closely resembles the fiber type composition of human skeletal muscles (primarily comprised
86 of type I, IIA and IIX fiber types), and is therefore a particularly interesting experimental
87 model. In our previous study, we mapped soleus muscle fiber traits in an F_2 intercross
88 between the LG/J and SM/J laboratory mouse strains (11). These strains differ in a number
89 of muscular phenotypes, with the LG/J strain displaying a greater proportion of type I fibers,
90 and a greater cross-sectional area (CSA) of type I and IIA muscle fibers. We identified in
91 that study three significant QTLs contributing to the difference in the CSA of muscle fibers
92 between LG/J and SM/J strains (11). Regions of conserved synteny from the identified loci
93 were also implicated in fiber phenotypes in pig supporting the importance of these genomic
94 regions in determining muscle fiber properties. However, the exact genes underlying their
95 effects remain to be determined.

96

97 Integration of advanced study populations, high throughput gene expression technology and
98 increasing availability of knockout models aid identification of the causative genes.

99 Nomination of the genes underlying QTL effects can be facilitated by improving the mapping
100 resolution, and by utilising genomic sequence and transcriptome information. Advanced
101 intercross lines (AIL) have been proposed as a powerful population for mapping QTLs (16).
102 It has been demonstrated recently that a joint F_2 and AIL analysis can combine the

103 advantages of both mapping populations by increasing the power to detect QTLs and
104 achieving a higher mapping resolution of various traits in mice (13, 47). Additionally, testing
105 for differences in specific gene expression has led to several nominations of quantitative trait
106 genes (30, 35). For validation of such candidate genes, phenotypic effects of relevant
107 alleles can be examined in experimental populations where these alleles segregate albeit on
108 a different genetic background. In addition, available knockout models offers particularly
109 attractive option for validation experiments.

110 In the present study we aimed to fine-map QTL and nominate candidate genes affecting the
111 CSA and proportion of oxidative type I fibers in the soleus muscle in a combined analysis of
112 F₂ and F₃₄ AIL mice, and by cross referencing QTL data with soleus transcriptome profiles in
113 the parental strains. Further filtering of the emerged candidates was carried out in an
114 independent AIL and a knockout model.

115

116 **Methods**

117 **Muscle Samples**

118 This study was carried out on soleus muscles dissected from females and males of the F₃₄
119 advanced intercross lines (AIL) of the LG/J and SM/J inbred strains. Animals were
120 maintained as previously described (13) and sacrificed at 94 ± 4 days. All procedures were
121 approved by the Institutional Animal Care and Use Committee of the University of Chicago.
122 Soleus muscle samples from F₃₄ AIL mice described in our previous study (47) were
123 subjected to histological analyses. The final sample size used in the present study was 334
124 F₃₄ mice, 142 females and 192 males, after discarding samples of poor tissue quality. A set
125 of 94 F₂ samples (38 females and 56 males) described in our previous study (11) was also
126 used in order to increase the QTL detection power.

127 In addition, we also analysed soleus muscle samples for two hypothesis driven studies
128 aimed at testing the effects of identified candidate genes on percentage of oxidative, type I
129 fibers. First, we examined solei samples from the *Chd6* ATPase knockout (n=6),
130 heterozygous (n=4) and wild type (n=4) females. The generation of the *Chd6* mutant mice
131 has been previously reported (40). Briefly, the genetic manipulation generated an allele with
132 the ATPase domain of *Chd6* (exon 12) flanked by loxP sites so that the action of Cre
133 recombinase would delete this domain. The mice were mated to a germline Cre-expressing
134 strain (Jackson lab strain 003465) to delete both exon 12 and the neomycin resistance
135 marker used for the targeting. Subsequently breeding generated the *Chd6* ATPase
136 knockout mice utilized in the present study. Second, solei of the advanced intercross mice
137 (generations F₉-F₁₂), all homozygous carries of the C57BL/6J (n=22) or DBA/2J (n=23)
138 alleles at the region harbouring the *Alad* gene were selected from the tissue bank of our
139 previous study (9).

140 **Phenotype assessment**

141 The soleus muscles were frozen in isopentane cooled in liquid nitrogen. Transverse
142 sections from the belly of the muscle were cut at a thickness of 10 μm with a cryotome
143 (Leica CM1850UV) at -20°C . The muscle sections were subjected to ATPase staining (acid
144 pre-incubation, pH 4.47) to distinguish between fiber types (8). Microscopic images of
145 stained sections were taken at x5 and x20 magnification.

146 The following phenotypes were assessed: muscle fiber number (type I and IIA) and percent
147 of type I muscle fibers, cross-sectional area (CSA) of type I and type IIA fibers (**Figure 1**).
148 Muscle fiber traits were manually analysed using ImageJ software (NIH-version 1.43). 25
149 measurements of each fiber type were taken using the freehand selection tool at x20
150 magnification to obtain a value representing the mean CSA of type I or type IIA fibers for that
151 muscle. This was deemed as a representative sample by empirical testing as described
152 previously (11). Total number of type I and type IIA muscle fibers were counted using the
153 ImageJ cell counter plugin on x5 magnification images. As all fibers in mouse soleus pass
154 through the belly of the muscle (69), this method provides an accurate estimate of the
155 number of fibers constituting the muscle. Total number of type I fibers and total number of
156 type IIA fibers were counted, permitting derivation of percentage of type I fibers. Over the
157 course of the study ~200,000 muscle fibers were counted and ~6,700 fibers measured for
158 CSA.

159 Statistical analyses

160 The GraphPad Prism version 5.0 statistical package was used (GraphPad software, La Jolla,
161 CA). Data are presented as mean \pm SD, unless otherwise stated. The CSA of type I and
162 type IIA fibers were analysed using a two-way (sex and fiber type) paired-measures (type I
163 and type IIA fibers) ANOVA.

164 Genotyping and QTL mapping

165 Mice were genotyped using a custom designed SNP array that included 4,610 polymorphic
166 SNPs that were approximately evenly distributed across the genome, as described

167 previously (13). The genome-wide association analysis was performed in the combined
168 population of the F₃₄ and recently published F₂ intercrosses (11) using the R package
169 QTLRel (12). This software accounted for the complex relationships (e.g., sibling, half-
170 sibling, cousins) among the F₃₄ mice by using a mixed model, as previously described (12,
171 13). Due to the sex differences in muscle mass in these mice (47), and the discovery of sex
172 specific QTL in other studies (45, 46), we included sex as an additive and interacting
173 covariate. Threshold of significance was estimated by 1000 permutations (14). We defined
174 the support interval for each QTL as the 1.5-LOD drop off on either side of the peak marker.
175 This interval was expressed in physical map position (Mb) by using the nearest genotyped
176 SNP that flanked the support interval, based on the mouse genome build GRCm38.p3.

177 Transcriptome analysis

178 Soleus muscle tissues from 92-day old LG/J and SM/J females (n=3 of each strain) were
179 used. RNA was isolated using TRIzol (Invitrogen Life Technologies, Carlsbad, CA) followed
180 by purification and DNase digestion using RNeasy minikits (Qiagen, Venlo, Netherlands)
181 according to the manufacturer's instructions. Quantification of total RNA was performed on
182 a NanoDrop spectrophotometer (Thermo Scientific) and quality tested on an Agilent
183 Tapestation with R6K Screentapes (RIN ≥7.3). Generation of sense strand cDNA from
184 purified total RNA (Ambion[®] WT expression kit, Ambion, Austin, Texas) followed by
185 fragmentation and labelling (GeneChip WT labelling kit, Affymetrix, Santa Clara, CA) were
186 performed according to the manufacturer's instructions. Hybridisation, washing, staining and
187 scanning of microarrays were carried out on Affymetrix Mouse Gene 2.0 ST microarrays
188 according to the manufacturer's standard protocols using a GeneChip Fluidics station 450
189 and GCS3000 scanner (Affymetrix[®], Santa Clara, CA). Microarray data are available in the
190 ArrayExpress database (www.ebi.ac.uk/arrayexpress) under accession number E-MTAB-
191 5290.

192

193 Data pre-processing and quality control analysis was performed using Affymetrix®
194 Genechip® Expression Console v1.2. Probe cell intensity data on the Mouse Gene 2.0 ST
195 array (CEL files) were processed using the RMA16 algorithm (Affymetrix, Santa Clara, CA,
196 USA) which fits a robust linear model at probe level by employing background correction,
197 quantile normalisation of log₂ transformed data and summarisation to probe level data (CHP
198 files, 41,345 probe sets).

199

200 Data was analysed for differentially expressed genes in Partek® Genomics Suite® version
201 6.6, build 6.15.0730 (Partek Inc., St Louis, MO) using a *Mus musculus* build mm10
202 annotation file for Mouse Gene 2.0 ST microarrays (MoGene-2_0-st-v1.na35.mm10). CEL
203 files (Expression Console v 1.2, Affymetrix, Santa Clara, CA) were imported to Partek
204 Genomics Suite v 6.6 and processed using RMA normalisation with background correction
205 of log₂ transformed data and probe set summarisation by median polish. Differential
206 expression analysis between the LG/J and SM/J strains of all genes (n=41,345 transcript
207 clusters) was determined by 1-way ANOVA with Storey's FDR, and q-value ≤0.05
208 considered significant (n=819 genes differentially expressed ≥ 1.2 fold; see **Supplementary**
209 **Table 1**).

210 To assess transcription of positional candidate genes in each strain, a hypothesis driven
211 analysis of differential gene expression was performed between the LG/J and SM/J strains
212 on all genes mapping to the support interval defined for each QTL in the GWAS described
213 above. Using Partek Genomics Suite v.6.6, a total of 159 genes that were represented on
214 the mouse Gene 2.0ST microarray, were identified in *Mus musculus* genome build
215 GRCm38, mm10 within mapping co-ordinates Chr2:158908559–162608559 (26 genes),
216 Chr3:33308451–35708451 (15 genes), Chr4:57605946–62913639 (77 genes) or Chr
217 11:27900000–31500000 (41 genes). 1-way ANOVA identified differentially expressed genes
218 between the LG/J and SM/J strains (P<0.05). Fold change was calculated using the
219 geometric mean of samples in each group.

220 Candidate genes

221 Nomination of the candidate genes was based on the following three criteria. First, we
222 scrutinized polymorphisms in positional candidates between the LG/J and SM/J strains. The
223 emphasis was on the indels and SNPs that would affect the coding sequence and lead to
224 changes in amino acids. To assess whether amino acid substitution would influence the
225 function of a protein, evolutionary conservation at the site of substitution and properties of
226 substituted amino acids were considered using three different bioinformatics tools as
227 described by Nikolskiy and colleagues (56). Second, we examined expression of positional
228 candidates across a panel of over ninety mouse tissues and cell types available in BioGPS
229 GeneAtlas MOE430, gcrma dataset (79). This analysis permits a quantitative comparison of
230 transcript abundance of a gene between tissues. We considered that an abundant
231 expression in skeletal muscle lineage, i.e. muscle tissue and/or C2C12 myogenic cell line,
232 implies functional and/or structural relevance of a gene in this tissue. Third, we compared
233 gene expression levels in the soleus muscle between the two strains as described in the
234 previous section. Expression difference in this analysis might point at the strain-specific,
235 genotype-dependent mechanism underlying the phenotypic difference.

236

237 **Results**

238 Phenotypic analyses

239 **CSA.** Cross section analysis of soleus muscle fibers were done on mice of both sexes from
240 the F₃₄ cohort. For muscle fiber cross-sectional area, we observed a statistically significant
241 sex by fiber type interaction (P<0.0001). In the female F₃₄ mice there was no significant
242 difference between type I and type IIA muscle fiber areas (913 ± 229 μm², n=140; and 952 ±
243 242 μm², n=140 respectively; P=0.2). However, there was a significant difference within the
244 males, with the type I muscle fiber area being smaller than IIA fiber area (1084 ± 238 μm²,

245 n=187; and $1215 \pm 294 \mu\text{m}^2$, n=187 respectively; $P < 0.0001$). Muscle fiber area was lower
246 in females than males for type I CSA, ($P < 0.0001$) and type IIA CSA ($P < 0.0001$).

247 **Percentage of type I fibers.** The number of type I fibers as a percentage of total fibers
248 varied substantially between individuals, ranging from 30% to 67% in females, and from 26%
249 to 59% in males (**Figure 1**) and was greater in females than males ($46 \pm 8\%$, n=142; and 39
250 $\pm 6\%$, n=189; respectively; $P < 0.0001$).

251 **Total fiber number.** No difference was observed in the total soleus fiber number between
252 females and males (646 ± 102 , n=120, and 667 ± 105 , n=177, respectively; $P = 0.0979$).

253

254 QTL analyses

255 Muscle fiber traits approximated the normal distribution in both the F_2 and F_{34} population
256 (**Supplementary Figure 1**). We identified significant QTL (at the 1% or 5% level of genome-
257 wide statistical significance) (39) for CSA of type I and type IIA fibers and the percentage of
258 type I fibers. We also identified chromosome-wide significant QTL for CSA of type I and type
259 IIA fibers, the percentage of type I fibers and total fiber number (**Table 1**). The size of the
260 support interval of these QTL ranged from 0.4-40.7 Mb, with a median of 4.6 Mb.

261 The QTL at the genome-wide level of significance for CSA of type I and type IIA fibers on
262 chromosome 3 was named *Mfq5*. The QTL at the genome-wide level of significance for the
263 percentage of type I fibers on chromosome 2 and 4 were named *Mfq4* and *Mfq6*,
264 respectively. The SM/J allele conferred a greater percentage of type I fibers at *Mfq4*, and a
265 greater CSA at *Mfq5*. The LG/J allele conferred a greater percentage of type I fibers at *Mfq6*
266 locus.

267 A significant QTL affecting CSA of type I and type IIA fibers was also detected on
268 chromosome 11 (**Figure 2**) within the same region as locus *Mfq3*, previously identified in the
269 F_2 intercross of the same parental strains (11). The QTL exhibited male-specificity in both

270 type I and IIA fibers of the F₃₄ mice (**Figure 3**). Because this QTL recapitulated properties of
271 the *Mfq3* locus, which we also found to be male specific in the F₂ population, we concluded
272 that the same locus has been refined in F₃₄ and did not assign a new name for this QTL.
273 Earlier reported *Mfq2* locus has been refined in a similar manner; a QTL on chromosome 6
274 affecting CSA of type I and type IIA fibers (at 1% chromosome specific threshold) was
275 engulfed by the support interval of *Mfq2* and also replicated its increasing allele, LG/J, in
276 both females and males (not shown).

277 Gene expression analyses

278 We hypothesized that each identified QTL harbours one or more genetic variants that drive
279 phenotypic differences by means of differential gene expression. Hypothesis driven analysis
280 of differential expression in soleus muscle was performed between LG/J and SM/J strains for
281 the genes in the most robust QTLs affecting fiber CSA or % Type I fibers (*Mfq3*, *Mfq4*, *Mfq5*
282 and *Mfq6*). The Mouse Gene 2.0 ST expression array contains 159 genes residing within
283 the support intervals of these QTLs (**Supplementary Table 2**). Twenty genes (**Table 2**)
284 showed evidence of differential expression (ANOVA, p≤0.05), 2 of which, *Alad* and *Hdhd3*,
285 were significant after correction for the multiple testing problem (Storey's FDR q≤0.05).
286 Compared to other tissues and cell types, expression of differentially expressed genes *Mafb*,
287 *Acyp2* and *Mtif2* (**Table 2**), is particularly enriched in skeletal muscle (BioGPS, Mouse
288 MOE430 gene expression data).

289 Genomic analyses

290 Positional candidates with non-synonymous polymorphisms provide a plausible genetic
291 cause for the phenotypic differences. Based on the genomic sequence of the LG/J and
292 SM/J strains (56), we identified 21 genes in the QTL regions with non-synonymous
293 polymorphisms predicted to affect protein function by at least one out of three algorithms
294 used in the analysis (**Supplementary Table 3**). Four of those genes (*Mfq3*: *Mtif2*, *Rtn4*,
295 *Psme4*; *Mfq5*: *Dnajc19*) are prioritized further because of their preferential expression in

296 muscle lineage (differentiated muscle and/or C2C12 myoblasts) compared to other tissues
297 and cell types. Among those, the *Mtif2* gene differs by 3 (rs26871496, rs26871494,
298 rs29436813) and *Rtn4* by 9 (rs29473364, rs29469198, rs13463765, rs29465940,
299 rs26857726, rs26857725, rs29474377, rs26857722, rs26857721) amino acids between the
300 two strains. At all SNPs the SM/J strains carries reference while the LG/J strain the
301 alternative allele.

302 Candidate gene analyses

303 The *Chd6* gene emerged as a differentially expressed positional candidate for the *Mfq4*
304 locus affecting percentage of type I fibers (**Table 2**). To test its effect we examined soleus
305 muscles of *Chd6* knockout, heterozygous and wild type littermates. This analysis however
306 revealed that the genotype of the animals did not have a significant effect ($P=0.30$) on the
307 percentage of type I fibers (**Figure 4**).

308 The *Alad* gene emerged as a candidate for another locus affecting proportion of type I fibers,
309 *Mfq6*. In the animals of an advanced intercross between the C57BL/6J and DBA/2J strains
310 (these strains carry one or three copies of *Alad*, respectively (3)), we examined if percentage
311 of type I fibers was genotype-dependent. The analysis revealed no difference in the
312 percentage of type I fibres between the carriers of the C57BL/6J and the DBA/2J alleles, 42
313 $\pm 7\%$ and $42 \pm 8\%$, respectively.

314 **Discussion**

315 A previous study on muscle weight in LG/J and SM/J strains identified a two-fold difference
316 in soleus muscle size (47). We then explored the cellular and genetic mechanisms
317 contributing to this phenomenon, finding that the difference was largely due to the CSA of
318 muscle fibers and we mapped QTL affecting muscle fiber traits in an F₂ intercross between
319 the LG/J and SM/J strains (11). The present study, which utilizes the F₃₄ advanced
320 intercross, verified, refined and expanded our earlier findings.

321 A number of studies have previously reported the effects of *Stat5a* and *Stat5b* (36), *Pgc-1α*
322 (42), *Ky* (4), myostatin (54), leptin (61), calcineurin (76), *Sod1* (5), alpha-actinin-3 (50),
323 dystrophin (7), *Tbx15* (41) and IIB myosin heavy chains (2) genes on muscle fiber area in
324 knockout or mutant models. In addition, *Pgc-1α* (75), calcineurin (76), *Foxo1* (34) and
325 myostatin (20) are reported to affect the proportion of muscle fiber types. However, the
326 genomic positions of these genes have not been linked to muscle fiber differences between
327 the LG/J and SM/J strains, implicating involvement of novel genes.

328 **Muscle fiber number.** The number of fibers is an important determinant of muscle size and
329 functional properties. It is set during embryogenesis and the first post-natal week in mice
330 (78). The number of muscle fibers in males (667 ± 105) and females (646 ± 102) of the F₃₄
331 population was comparable to that observed in the soleus of the F₂ population (645 ± 102
332 and 595 ± 107, respectively), and within the range of the fiber count observed in solei of a
333 variety of different strains of mice ~250~900 fibers (32, 49, 57, 70, 72).

334 From these data it emerged that males and females are born with a similar number of fibers
335 in soleus muscle, and that the sex difference in muscle weight (males have approximately
336 30% larger soleus than females) is due to the difference in fiber size. Comparison of the
337 parental strains also revealed a similar number of fibers (11), despite the 2-fold difference in
338 soleus weight (47), demonstrating that size rather than number of fibers determines variation
339 in muscle weight between the LG/J and SM/J strains.

340 **Fiber area.** The CSA of muscle fibers in the LG/J strain is 49% to 90% greater than the
341 corresponding fibers in the SM/J strain, indicating that this variable accounts for a large
342 portion of the muscle mass difference between the strains (47).

343 The area of type I ($1084 \pm 238 \mu\text{m}^2$ and $913 \pm 229 \mu\text{m}^2$ for males and females, respectively)
344 and type IIA ($1215 \pm 294 \mu\text{m}^2$ and $952 \pm 242 \mu\text{m}^2$, respectively) of the F_{34} mice was
345 comparable to the corresponding fiber area of the F_2 mice of the same lineage (11) and it is
346 within the range reported for the type I, between $920 \mu\text{m}^2$ and $1780 \mu\text{m}^2$ (32, 57, 70), and
347 type IIA fiber area, between $700 \mu\text{m}^2$ and $1400 \mu\text{m}^2$ (32, 70), in various inbred mouse
348 strains.

349 **Percentage of type I fibers.** The percentage of type I fibers in male ($39 \pm 6\%$) and female
350 ($47 \pm 8\%$) F_{34} mice were also within the range of previous studies, which showed the
351 percentage of type I fibers in the soleus muscle fluctuates between ~25 and ~66% (32, 57,
352 70).

353 In the F_{34} mice we replicated our observation in the F_2 population that the percentage of type
354 I fibers was significantly greater in females than males. This sex difference is also observed
355 in various human muscles where, in general, women have a higher percentage of type I
356 muscle fibers than males (27, 53, 60, 64, 67). The phenomenon is likely to be explained, at
357 least partly, by the effect of androgens; castration leads to a higher percentage of type I
358 fibers in the soleus of male mice (73).

359 **Validation and refinement of genetic architecture.** In the present study, we validated and
360 refined the genetic architecture of muscle fibers identified in an F_2 intercross between the
361 same parental strains (11). In order to increase QTL detection power, we increased sample
362 size by combining the F_{34} and F_2 data. The median mapping resolution of 4.6 Mb for muscle
363 fiber QTLs was comparable with 3.7 Mb of muscle weight QTLs obtained in the same
364 population albeit using ~1,600 fewer genetic markers than in the present analysis (47). A
365 genome-wide significant QTL identified in the present study between 27.9 Mb and 31.4 Mb

366 on chromosome 11 (**Table 1**) overlapped with a significant QTL, *Mfq3*, mapped in the F₂
367 population (11). In addition to the chromosomal location, the increasing allele of this locus
368 (LG/J) and its male-specific effect (**Figure 3**) were also replicated in F₃₄, suggesting that the
369 same gene(s) were involved in two different populations and permitting us to refine the *Mfq3*
370 locus from 51.6 Mb to 3.57 Mb. The presence of two satellite QTL proximal of the refined
371 *Mfq3* (**Table 1**) suggests that the QTL observed in the F₂ population (11) might have been
372 an outcome of up to three linked loci.

373 The recently reported “mini-muscle” locus, mapped to 67.1–70.2 Mb on chromosome 11,
374 affects muscle fiber area and proportion of fiber types (21-23). However, the mutation
375 responsible for the “mini-muscle” phenotype maps to an intron of *Myh4* gene located at 67.2
376 Mb (31), between the support intervals of two adjacent QTLs affecting fiber type between the
377 LG/J and SM/J strains (**Table 1**). Together, these data suggest that a number of genes
378 residing on chromosome 11 might be involved in the regulation of muscle fiber phenotypes.

379 The QTL affecting the CSA of type I and type IIA fibers on chromosome 6, albeit at 1%
380 chromosome-wide threshold of significance (**Table 1**), overlapped with the *Mfq2* locus found
381 in the F₂ population, characterized by the same increasing allele, LG/J. Thus, the support
382 interval of *Mfq2* could be considered to be 5.18 Mb rather than the previously reported 56.5
383 Mb. Importantly, the immediate proximity of the refined region (Chr 6: 110.8-116.0 Mb) to
384 the syntenic region (Chr 6:116.0-118.0 Mb) implicated in the QTL affecting the diameter of
385 pig IIA fibers (17) suggest that the same genes could be underlying the effects of these
386 QTLs in mice.

387 A QTL affecting percentage of type I fibers (at 10% chromosome-wide threshold) on
388 chromosome 1 (67.6to 70.8 Mb) overlapped with *Mfq1* locus which influenced the CSA of
389 type I and type IIA fiber area in the F₂ population (11). However, because the CSA and
390 percentage of type I fibers are poorly correlated traits both in the F₃₄ (**Supplementary Table**

391 4) and the F₂ mice (11), it is likely that different genes are underlying the *Mfq1* locus and the
392 QTL identified in the F₃₄ population. Further studies are required to clarify this observation.

393 **Transcriptome analysis**

394 In the present study, the expressed transcriptome in soleus muscle of the parental strains
395 was examined in order to facilitate nomination of the candidate genes within the refined QTL.
396 We hypothesized that if the phenotypic effect of the QTL was brought about by the allele
397 specific abundance of transcripts encoded by genes within the QTL, such genes would be
398 differentially expressed in the transcriptome between the parental strains. Comparison of
399 expression of the genes within the four most robust QTLs identified *Alad* and *Hdhd3* genes
400 as potential candidates for the *Mfq6* locus, which affects the proportion of type I fibers.
401 Transcripts of both genes are more abundant in the LG/J compared to the SM/J strain. This
402 is consistent with our findings in the TA muscle of the same strains (48). Of these two
403 identified candidate genes, transcripts of *Alad* are ~20 times more abundant in the mouse
404 muscle than *Hdhd3*, regardless of strain. In addition, *Alad* may play a role during
405 myogenesis as its expression in C2C12 myogenic cells is 5-fold higher compared to
406 differentiated muscle (79).

407 **Candidate genes.**

408 The support intervals of four most robust QTLs harbor 159 genes (**Supplementary Table 2**).
409 These regions were scrutinized further for the genes fulfilling one of the following criteria:
410 presence of the functional variants (i.e. non-synonymous SNPs predicted to alter function of
411 encoded protein); abundance of transcript in muscle lineage, particularly in comparison to
412 other tissues and cell types; differential expression in the soleus of the two strains; and by
413 comparing genomic sequence between the LG/J and SM/J strains a list of 21 genes was
414 highlighted (**Supplementary Table 3**) with the strain-specific functional variants. Using
415 bioinformatics, 4 genes abundantly and/or preferentially expressed in skeletal muscle
416 compared to other types of tissues and cells were identified. Our own analysis of gene

417 expression in soleus muscle highlighted a set of 20 genes differentially expressed between
418 the two strains (**Table 2**). Intersection of all these lists permitted us to prioritise nine
419 candidate genes which appeared on more than one of these lists and/or for which
420 independent and accessible validation models were available (i.e. *Chd6* and *Alad*). Because
421 neither the *Chd6* (**Figure 4**) nor *Alad* genes were found to affect proportion of type I fibres in
422 the way predicted by the QTL analyses, the list of prioritised candidates was reduced to 7
423 genes annotated in **Supplementary Table 5**. Three out of four QTLs contain one (*Mfq6*) or
424 more candidate genes. All candidates are abundantly transcribed in muscle lineage with
425 *Psme4*, *Acyp2* and *Mafb* showing the highest level of expression in skeletal muscle
426 compared to other tissues and cells. None of the seven candidates have been previously
427 implicated to affect properties of skeletal muscle fibres although some of them have been
428 implicated in cardiomyopathy or function as transcription factors (**Supplementary Table 5**).
429 Thus, genomic and gene expression analyses permitted focusing on a limited number of
430 positional candidates in the future validation studies for establishing the causative genes.

431 **Conclusion**

432 In conclusion, we have refined the genetic architecture affecting cross sectional area of
433 soleus muscle fibers and proportion of type I fibers in the LG/J and SM/J derived lineage.
434 Integrating QTL mapping, genomic and transcriptome data from homologous muscle
435 highlighted several candidate genes that may underpin muscle phenotypes critical to health
436 and disease and worthy of follow up analyses.

437

438

439

440

441 Grants

442 The study was supported by NIH R01AR056280 grant (AL), NIH R01DA024845 (AAP) and
443 Marie Curie International Reintegration Grant 249156 (AL).

444

445 Acknowledgements

446 The authors wish to acknowledge technical assistance from Diane Stewart, Jasmine C.
447 Berry and Jake T. Brown. We acknowledge the Dartmouth Mouse Modelling Shared
448 Resource of the Norris Cotton Cancer Center (support grant P30CA023108-27), for
449 providing *Chd6* knockout mouse samples.

450

451 Author contributions

452 A.L. conceived and supervised the study, A.M.C. phenotyped muscle samples, A.A.P.
453 provided genotypes and oversaw the QTL analyses, R.C. designed the QTLRel software
454 used in the QTL analysis, A.M.C. carried out QTL mapping, M.E.S, C.M. and E.C.D. did
455 transcriptome analysis, S.N.F. and J.L.F. generated and provided *Chd6* knockout samples,
456 A.L. and A.M.C. wrote the manuscript with input from all co-authors.

457

458

459 **Figure legends**

460 **Figure 1. Individual variability in proportion of oxidative fibers.** Representative images
461 of F₃₄ female soleus cross-sections following myosin ATPase staining (acid pre-incubation).
462 Dark fibers type I, pale fibers type IIA.

463

464 **Figure 2. Type I fiber cross-sectional area QTL on chromosome 11.** Analyses were
465 carried out in the F₂ intercross and in the combined F₂ and F₃₄ populations. X-axis indicates
466 the relative position in the linkage map in centimorgan (cM). The thresholds are at the level
467 of 0.05 genome wise significance for the F₂ output (dotted line) and combined output (solid
468 line).

469

470 **Figure 3. Sex specificity of *Mfq3* locus on cross-sectional area (CSA) of soleus type I
471 and IIA fibers in the F₃₄ intercross.** Mean and SEM. Genotype at the peak marker: LG,
472 homozygous for LG/J allele; H, heterozygous; SM, homozygous for SM/J allele.

473

474

475 **Figure 4. Percentage of type I fibers in the soleus muscle of 4 month old *Chd6*
476 knockout (KO), heterozygous (HET) and wild-type (WT) females.** There is no difference
477 in percentage of type I muscle fibers in the soleus muscle between knockout, heterozygotes
478 and wild-type groups (P=0.3041). Each data point is from a single mouse, horizontal lines
479 represent group mean.

480

481 **Table 1.** Characteristics of muscle fiber QTL in combined analyses of the F₂ and F₃₄
 482 intercrosses derived from the LG/J and SM/J strains.

Chr	Thr**	Level***	Start Mb [†]	End Mb	Size Mb	Trait	Locus¥
1	C	0.1	67.6	70.7	3.1	% Type I	
1	C	0.1	193.9	194.3	0.4	% Type I & CSA2A	
2	C	0.1	92.4	104.8	12.4	% Type I	
2	C	0.05	139.6	145.6	6.0	% Type I	
2	G	0.01	158.8	162.5	3.7	% Type I	Mfq4 (SM)
3	G	0.05	33.6	40.0	6.4	CSA1 & CSA2A	Mfq5 (SM)
4	G	0.05	57.7	62.7	5.0	% Type I	Mfq6 (LG)
4	C	0.05	103.9	106.1	2.2	% Type I	
6	C	0.05	81.9	84.1	2.2	CSAIIA	
6	C	0.01	110.8	116.0	5.2	CSA1 & CSA2A	Mfq2* (LG)
7	C	0.05	138.4	140.0	1.6	% Type I	
8	C	0.1	7.4	12.4	5.0	% Type I	
8	C	0.05	89.0	92.4	3.4	TOTAL	
8	C	0.01	121.9	128.6	6.7	TOTAL	
10	G	0.1	120.7	121.3	0.6	% Type I	
11	C	0.1	12.4	17.2	4.8	CSAIIA	
11	C	0.1	19.1	23.1	4.0	CSAIIA	
11	G	0.01	28.0	31.5	3.5	CSA1 & CSA2A	Mfq3* (LG)
11	C	0.1	62.5	64.2	1.7	% Type I	
11	C	0.1	70.6	76.2	5.6	% Type I	
12	C	0.1	27.6	29.3	1.7	CSA1	
13	C	0.01	5.3	9.9	4.6	% Type I	
13	C	0.05	71.5	74.0	2.5	CSAIIA	
14	C	0.05	93.6	102.3	8.7	CSAIIA	
15	C	0.1	12.1	20.3	8.2	TOTAL	
16	C	0.05	68.9	75.1	6.2	CSA1 & CSA2A	
X	C	0.01	11.8	52.5	40.7	TOTAL	

483 * refined previously identified QTL in the LG/J and SM/J F₂ intercross (47).

484 ** C – chromosome-wide threshold, G- genome-wide threshold

485 *** Level of significance

486 ¥ LG –increasing allele is LG/J, SM- increasing allele is SM/J

487 † Genomic positions based on GRCm38.p3.

488

489 **Table 2.** Positional candidate genes differentially expressed between LG/J and SM/J soleus muscles.

Chr	QTL	Probe set ID	Gene	p-value*	Fold-Change**	Gene name***		
2	<i>Mfq4</i>	17393868	<i>Mafb</i>	0.033	-1.77	v-maf musculoaponeurotic fibrosarcoma oncogene family, protein B (avian)		
		17393910	<i>Chd6</i>	0.042724	-1.13	chromodomain helicase DNA binding protein 6		
		17404652	<i>Gm24780</i>	0.032914	-1.98	Predicted gene Gm24780, predicted protein is B4HDV3.		
3	<i>Mfq5</i>	17396801	<i>Ttc14</i>	0.033387	-1.14	tetratricopeptide repeat domain 14		
		17396876	----	0.0219376	-1.80	There are no assigned mRNA sequences for this probe set. The probe set lies within lincRNA Sox2ot (Sox2 overlapping transcript, non-protein coding)		
		17425606	<i>Gm12526</i>	0.046791	-1.17	predicted gene 12526		
		17414380	<i>Gm24277</i>	0.00525644	-2.07	Gm24277 a known snRNA. The probeset also lies within an intronic region of RefSeq gene Pakap (PALM2-AKAP2), a read through transcript on chromosome 4		
		17425701	<i>Mir3095</i>	0.0298235	-1.78	Mir3095 (Entrez ID 100526502; EST ENSMUST00000175552).		
		17426097	<i>Mup3</i>	0.00304	-1.49	major urinary protein 3		
		17426126	<i>Fkbp15</i>	0.018038	-1.10	FK506 binding protein 15		
4	<i>Mfq6</i>	17414545	<i>Slc31a1</i>	0.049799	1.12	solute carrier family 31, member 1		
		17426166	<i>Cdc26</i>	0.021407	-1.29	cell division cycle 26		
		17426198	<i>Hdhd3</i>	0.000869	1.86	haloacid dehalogenase-like hydrolase domain containing 3		
		17426206	<i>Alad</i>	9.75E-05	1.91	aminolevulinate, delta-, dehydratase		
		17414600	<i>Rgs3</i>	0.023096	1.08	regulator of G-protein signaling 3		
		17248064	<i>Mtif2</i>	0.015517	-1.16	mitochondrial translational initiation factor 2		
		17261285	<i>LOC102637613</i>	0.00432682	1.68	linc RNA [AK084560 (EST)/ Gm12092 (predicted gene)].		
		17248127	----	0.00187422	-1.46	There are no assigned mRNA sequences for this transcript. The probe set lies within an intron of <i>Sptbn1</i> .		
		11	<i>Mfq3</i>	17261393	<i>Acyp2</i>	0.011777	1.26	acylphosphatase 2, muscle type
				17248196	<i>Asb3</i>	0.014221	-1.15	ankyrin repeat and SOCS box-containing 3

491 * ANOVA p-value for strain effect; ** Fold change uses SM/J as baseline (negative values indicate LG/J expression is down compared to SM/J,
492 positive values LG/J expression up compared to SM/J); bold indicates that gene is predominantly and/ or strongly expressed in skeletal muscle
493 tissue (79). *** For probe sets not designed against an annotated gene, genes at the genomic loci of the Affymetrix probeset were identified in
494 UCSC genome browser using mouse genome build GRCm38.

495

496 **References**

497

- 498 1. **Abe T, Saburi J, Hasebe H, Nakagawa T, Kawamura T, Saito K, Nade T, Misumi S, Okumura**
499 **T, Kuchida K, Hayashi T, Nakane S, Mitsuhasi T, Nirasawa K, Sugimoto Y, and Kobayashi E.** Bovine
500 quantitative trait loci analysis for growth, carcass, and meat quality traits in an F2 population from a
501 cross between Japanese Black and Limousin. *Journal of animal science* 86: 2821-2832, 2008.
- 502 2. **Allen DL, Sartorius CA, Sycuro LK, and Leinwand LA.** Different pathways regulate expression
503 of the skeletal myosin heavy chain genes. *The Journal of biological chemistry* 276: 43524-43533,
504 2001.
- 505 3. **Bishop TR, Miller MW, Wang A, and Dierks PM.** Multiple copies of the ALA-D gene are
506 located at the Lv locus in *Mus domesticus* mice. *Genomics* 48: 221-231, 1998.
- 507 4. **Blanco G, Coulton GR, Biggin A, Grainge C, Moss J, Barrett M, Berquin A, Marechal G,**
508 **Skykker M, van Mier P, Nikitopoulou A, Kraus M, Ponting CP, Mason RM, and Brown SD.** The
509 kyphoscoliosis (ky) mouse is deficient in hypertrophic responses and is caused by a mutation in a
510 novel muscle-specific protein. *Human molecular genetics* 10: 9-16, 2001.
- 511 5. **Bordet T, Schmalbruch H, Pettmann B, Hagege A, Castelnau-Ptakhine L, Kahn A, and Haase**
512 **G.** Adenoviral cardiostrophin-1 gene transfer protects pmn mice from progressive motor
513 neuronopathy. *The Journal of clinical investigation* 104: 1077-1085, 1999.
- 514 6. **Bottinelli R, and Reggiani C.** Human skeletal muscle fibres: molecular and functional
515 diversity. *Progress in biophysics and molecular biology* 73: 195-262, 2000.
- 516 7. **Briguet A, Courdier-Fruh I, Foster M, Meier T, and Magyar JP.** Histological parameters for
517 the quantitative assessment of muscular dystrophy in the mdx-mouse. *Neuromuscular disorders* :
518 *NMD* 14: 675-682, 2004.
- 519 8. **Brooke MH, and Kaiser KK.** Muscle fiber types: how many and what kind? *Archives of*
520 *neurology* 23: 369-379, 1970.
- 521 9. **Carbonetto P, Cheng R, Gyekis JP, Parker CC, Blizard DA, Palmer AA, and Lionikas A.**
522 Discovery and refinement of muscle weight QTLs in B6 × D2 advanced intercross mice. *Physiological*
523 *Genomics* 46: 571-582, 2014.
- 524 10. **Carpenter CE, Rice OD, Cockett NE, and Snowden GD.** Histology and composition of muscles
525 from normal and callipyge lambs. *Journal of animal science* 74: 388-393, 1996.
- 526 11. **Carroll AM, Palmer AA, and Lionikas A.** QTL Analysis of Type I and Type IIA Fibers in Soleus
527 Muscle in a Cross between LG/J and SM/J Mouse Strains. *Frontiers in genetics* 2: 99, 2011.
- 528 12. **Cheng R, Abney M, Palmer AA, and Skol AD.** QTLRel: an R package for genome-wide
529 association studies in which relatedness is a concern. *BMC genetics* 12: 66, 2011.
- 530 13. **Cheng R, Lim JE, Samocho KE, Sokoloff G, Abney M, Skol AD, and Palmer AA.** Genome-wide
531 association studies and the problem of relatedness among advanced intercross lines and other
532 highly recombinant populations. *Genetics* 185: 1033-1044, 2010.
- 533 14. **Cheng R, and Palmer AA.** A Simulation Study of Permutation, Bootstrap, and Gene Dropping
534 for Assessing Statistical Significance in the Case of Unequal Relatedness. *Genetics* 193: 1015-1018,
535 2013.
- 536 15. **Costill DL, Daniels J, Evans W, Fink W, Krahenbuhl G, and Saltin B.** Skeletal muscle enzymes
537 and fiber composition in male and female track athletes. *Journal of applied physiology* 40: 149-154,
538 1976.
- 539 16. **Darvasi A, and Soller M.** Advanced intercross lines, an experimental population for fine
540 genetic mapping. *Genetics* 141: 1199-1207, 1995.
- 541 17. **Estelle J, Gil F, Vazquez JM, Latorre R, Ramirez G, Barragan MC, Folch JM, Noguera JL, Toro**
542 **MA, and Perez-Enciso M.** A quantitative trait locus genome scan for porcine muscle fiber traits
543 reveals overdominance and epistasis. *Journal of animal science* 86: 3290-3299, 2008.

- 544 18. **Fink WJ, Costill DL, and Pollock ML.** Submaximal and maximal working capacity of elite
545 distance runners. Part II. Muscle fiber composition and enzyme activities. *Annals of the New York*
546 *Academy of Sciences* 301: 323-327, 1977.
- 547 19. **Frederiksen H, Gaist D, Petersen HC, Hjelmborg J, McGue M, Vaupel JW, and Christensen K.**
548 Hand grip strength: a phenotype suitable for identifying genetic variants affecting mid- and late-life
549 physical functioning. *Genetic epidemiology* 23: 110-122, 2002.
- 550 20. **Girgenrath S, Song K, and Whittemore LA.** Loss of myostatin expression alters fiber-type
551 distribution and expression of myosin heavy chain isoforms in slow- and fast-type skeletal muscle.
552 *Muscle & nerve* 31: 34-40, 2005.
- 553 21. **Guderley H, Houle-Leroy P, Diffie GM, Camp DM, and Garland T, Jr.** Morphometry,
554 ultrastructure, myosin isoforms, and metabolic capacities of the "mini muscles" favoured by
555 selection for high activity in house mice. *Comparative biochemistry and physiology Part B,*
556 *Biochemistry & molecular biology* 144: 271-282, 2006.
- 557 22. **Guderley H, Joanisse DR, Moka S, Bilodeau GM, and Garland T, Jr.** Altered fibre types in
558 gastrocnemius muscle of high wheel-running selected mice with mini-muscle phenotypes.
559 *Comparative biochemistry and physiology Part B, Biochemistry & molecular biology* 149: 490-500,
560 2008.
- 561 23. **Hartmann J, Garland T, Jr., Hannon RM, Kelly SA, Munoz G, and Pomp D.** Fine mapping of
562 "mini-muscle," a recessive mutation causing reduced hindlimb muscle mass in mice. *The Journal of*
563 *heredity* 99: 679-687, 2008.
- 564 24. **Hernelahti M, Tikkanen HO, Karjalainen J, and Kujala UM.** Muscle fiber-type distribution as
565 a predictor of blood pressure: a 19-year follow-up study. *Hypertension* 45: 1019-1023, 2005.
- 566 25. **Hubal MJ, Gordish-Dressman H, Thompson PD, Price TB, Hoffman EP, Angelopoulos TJ,**
567 **Gordon PM, Moyna NM, Pescatello LS, Visich PS, Zoeller RF, Seip RL, and Clarkson PM.** Variability in
568 muscle size and strength gain after unilateral resistance training. *Medicine and science in sports and*
569 *exercise* 37: 964-972, 2005.
- 570 26. **Huygens W, Thomis MA, Peeters MW, Vlietinck RF, and Beunen GP.** Determinants and
571 upper-limit heritabilities of skeletal muscle mass and strength. *Canadian journal of applied*
572 *physiology = Revue canadienne de physiologie appliquee* 29: 186-200, 2004.
- 573 27. **Jaworowski A, Porter MM, Holmback AM, Downham D, and Lexell J.** Enzyme activities in
574 the tibialis anterior muscle of young moderately active men and women: relationship with body
575 composition, muscle cross-sectional area and fibre type composition. *Acta physiologica Scandinavica*
576 176: 215-225, 2002.
- 577 28. **Kadi F, Eriksson A, Holmner S, Butler-Browne GS, and Thornell LE.** Cellular adaptation of the
578 trapezius muscle in strength-trained athletes. *Histochemistry and cell biology* 111: 189-195, 1999.
- 579 29. **Karjalainen J, Tikkanen H, Hernelahti M, and Kujala UM.** Muscle fiber-type distribution
580 predicts weight gain and unfavorable left ventricular geometry: a 19 year follow-up study. *BMC*
581 *cardiovascular disorders* 6: 2, 2006.
- 582 30. **Karp CL, Grupe A, Schadt E, Ewart SL, Keane-Moore M, Cuomo PJ, Kohl J, Wahl L,**
583 **Kuperman D, Germer S, Aud D, Peltz G, and Wills-Karp M.** Identification of complement factor 5 as
584 a susceptibility locus for experimental allergic asthma. *Nature immunology* 1: 221-226, 2000.
- 585 31. **Kelly SA, Bell TA, Selitsky SR, Buus RJ, Hua K, Weinstock GM, Garland T, Pardo-Manuel de**
586 **Villena F, and Pomp D.** A Novel Intronic Single Nucleotide Polymorphism in the Myosin heavy
587 polypeptide 4 Gene Is Responsible for the Mini-Muscle Phenotype Characterized by Major Reduction
588 in Hind-Limb Muscle Mass in Mice. *Genetics* 195: 1385-1395, 2013.
- 589 32. **Kilikevicius A, Venckunas T, Zelniene R, Carroll AM, Lionikaite S, Ratkevicius A, and**
590 **Lionikas A.** Divergent physiological characteristics and responses to endurance training among
591 inbred mouse strains. *Scandinavian journal of medicine & science in sports* 23: 657-668, 2013.
- 592 33. **Kim J, Wang Z, Heymsfield SB, Baumgartner RN, and Gallagher D.** Total-body skeletal
593 muscle mass: estimation by a new dual-energy X-ray absorptiometry method. *The American journal*
594 *of clinical nutrition* 76: 378-383, 2002.

- 595 34. **Kitamura T, Kitamura YI, Funahashi Y, Shawber CJ, Castrillon DH, Kollipara R, DePinho RA,**
596 **Kitajewski J, and Accili D.** A Foxo/Notch pathway controls myogenic differentiation and fiber type
597 specification. *The Journal of clinical investigation* 117: 2477-2485, 2007.
- 598 35. **Klein RF, Allard J, Avnur Z, Nikolcheva T, Rotstein D, Carlos AS, Shea M, Waters RV,**
599 **Belknap JK, Peltz G, and Orwoll ES.** Regulation of bone mass in mice by the lipoxigenase gene
600 Alox15. *Science (New York, NY)* 303: 229-232, 2004.
- 601 36. **Klover P, Chen W, Zhu BM, and Hennighausen L.** Skeletal muscle growth and fiber
602 composition in mice are regulated through the transcription factors STAT5a/b: linking growth
603 hormone to the androgen receptor. *FASEB journal : official publication of the Federation of American*
604 *Societies for Experimental Biology* 23: 3140-3148, 2009.
- 605 37. **Komi PV, Viitasalo JH, Havu M, Thorstensson A, Sjodin B, and Karlsson J.** Skeletal muscle
606 fibres and muscle enzyme activities in monozygous and dizygous twins of both sexes. *Acta*
607 *physiologica Scandinavica* 100: 385-392, 1977.
- 608 38. **Koohmaraie M, Shackelford SD, Wheeler TL, Lonergan SM, and Doumit ME.** A muscle
609 hypertrophy condition in lamb (callipyge): characterization of effects on muscle growth and meat
610 quality traits. *Journal of animal science* 73: 3596-3607, 1995.
- 611 39. **Lander E, and Kruglyak L.** Genetic dissection of complex traits: guidelines for interpreting
612 and reporting linkage results. *Nature genetics* 11: 241-247, 1995.
- 613 40. **Lathrop MJ, Chakrabarti L, Eng J, Harker Rhodes C, Lutz T, Nieto A, Denny Liggitt H, Warner**
614 **S, Fields J, Stöger R, and Fiering S.** Deletion of the Chd6 exon 12 affects motor coordination.
615 *Mammalian Genome* 21: 130-142, 2010.
- 616 41. **Lee KY, Singh MK, Ussar S, Wetzell P, Hirshman MF, Goodyear LJ, Kispert A, and Kahn CR.**
617 Tbx15 controls skeletal muscle fibre-type determination and muscle metabolism. *Nature*
618 *communications* 6: 8054, 2015.
- 619 42. **Leick L, Hellsten Y, Fentz J, Lyngby SS, Wojtaszewski JF, Hidalgo J, and Pilegaard H.** PGC-
620 1alpha mediates exercise-induced skeletal muscle VEGF expression in mice. *American journal of*
621 *physiology Endocrinology and metabolism* 297: E92-103, 2009.
- 622 43. **Li WB, Ren J, Zhu WC, Guo BL, Yang B, Liu LT, Ding NS, Ma JW, Li L, and Huang LS.** Mapping
623 QTL for porcine muscle fibre traits in a White Duroc x Erhualian F(2) resource population. *Journal of*
624 *animal breeding and genetics = Zeitschrift fur Tierzucht und Zuchtungsbiologie* 126: 468-474,
625 2009.
- 626 44. **Lillioja S, Young AA, Culter CL, Ivy JL, Abbott WG, Zawadzki JK, Yki-Jarvinen H, Christin L,**
627 **Secomb TW, and Bogardus C.** Skeletal muscle capillary density and fiber type are possible
628 determinants of in vivo insulin resistance in man. *The Journal of clinical investigation* 80: 415-424,
629 1987.
- 630 45. **Lionikas A, Blizard DA, Gerhard GS, Vandenberg DJ, Stout JT, Vogler GP, McClearn GE, and**
631 **Larsson L.** Genetic determinants of weight of fast- and slow-twitch skeletal muscle in 500-day-old
632 mice of the C57BL/6J and DBA/2J lineage. *Physiological genomics* 21: 184-192, 2005.
- 633 46. **Lionikas A, Blizard DA, Vandenberg DJ, Glover MG, Stout JT, Vogler GP, McClearn GE, and**
634 **Larsson L.** Genetic architecture of fast- and slow-twitch skeletal muscle weight in 200-day-old mice
635 of the C57BL/6J and DBA/2J lineage. *Physiological genomics* 16: 141-152, 2003.
- 636 47. **Lionikas A, Cheng R, Lim JE, Palmer AA, and Blizard DA.** Fine-mapping of muscle weight QTL
637 in LG/J and SM/J intercrosses. *Physiological genomics* 42a: 33-38, 2010.
- 638 48. **Lionikas A, Meharg C, Derry JM, Ratkevicius A, Carroll AM, Vandenberg DJ, and Blizard**
639 **DA.** Resolving candidate genes of mouse skeletal muscle QTL via RNA-Seq and expression network
640 analyses. *BMC genomics* 13: 592, 2012.
- 641 49. **Luff AR, and Goldspink G.** Total number of fibers in muscles of several strains of mice.
642 *Journal of animal science* 30: 891-893, 1970.
- 643 50. **MacArthur DG, Seto JT, Chan S, Quinlan KG, Raftery JM, Turner N, Nicholson MD, Kee AJ,**
644 **Hardeman EC, Gunning PW, Cooney GJ, Head SI, Yang N, and North KN.** An Actn3 knockout mouse

645 provides mechanistic insights into the association between alpha-actinin-3 deficiency and human
646 athletic performance. *Human molecular genetics* 17: 1076-1086, 2008.

647 51. **MacDougall JD, Sale DG, Alway SE, and Sutton JR.** Muscle fiber number in biceps brachii in
648 bodybuilders and control subjects. *Journal of applied physiology: respiratory, environmental and*
649 *exercise physiology* 57: 1399-1403, 1984.

650 52. **Malek M, Dekkers JC, Lee HK, Baas TJ, Prusa K, Huff-Lonergan E, and Rothschild MF.** A
651 molecular genome scan analysis to identify chromosomal regions influencing economic traits in the
652 pig. II. Meat and muscle composition. *Mammalian genome : official journal of the International*
653 *Mammalian Genome Society* 12: 637-645, 2001.

654 53. **Mannion AF, Dumas GA, Cooper RG, Espinosa FJ, Faris MW, and Stevenson JM.** Muscle
655 fibre size and type distribution in thoracic and lumbar regions of erector spinae in healthy subjects
656 without low back pain: normal values and sex differences. *Journal of anatomy* 190 (Pt 4): 505-513,
657 1997.

658 54. **McPherron AC, Lawler AM, and Lee SJ.** Regulation of skeletal muscle mass in mice by a new
659 TGF-beta superfamily member. *Nature* 387: 83-90, 1997.

660 55. **Nii M, Hayashi T, Mikawa S, Tani F, Niki A, Mori N, Uchida Y, Fujishima-Kanaya N, Komatsu**
661 **M, and Awata T.** Quantitative trait loci mapping for meat quality and muscle fiber traits in a
662 Japanese wild boar x Large White intercross. *Journal of animal science* 83: 308-315, 2005.

663 56. **Nikolskiy I, Conrad DF, Chun S, Fay JC, Cheverud JM, and Lawson HA.** Using whole-genome
664 sequences of the LG/J and SM/J inbred mouse strains to prioritize quantitative trait genes and
665 nucleotides. *BMC genomics* 16: 415, 2015.

666 57. **Nimmo MA, Wilson RH, and Snow DH.** The inheritance of skeletal muscle fibre composition
667 in mice. *Comparative biochemistry and physiology A, Comparative physiology* 81: 109-115, 1985.

668 58. **Pourhassan M, Bosy-Westphal A, Schautz B, Braun W, Gluer CC, and Muller MJ.** Impact of
669 body composition during weight change on resting energy expenditure and homeostasis model
670 assessment index in overweight nonsmoking adults. *The American journal of clinical nutrition* 99:
671 779-791, 2014.

672 59. **Rehfeldt C, Ott G, Gerrard DE, Varga L, Schlote W, Williams JL, Renne U, and Bunger L.**
673 Effects of the compact mutant myostatin allele Mstn (Cmpt-dl1Abc) introgressed into a high growth
674 mouse line on skeletal muscle cellularity. *Journal of muscle research and cell motility* 26: 103-112,
675 2005.

676 60. **Roepstorff C, Donsmark M, Thiele M, Vistisen B, Stewart G, Vissing K, Schjerling P, Hardie**
677 **DG, Galbo H, and Kiens B.** Sex differences in hormone-sensitive lipase expression, activity, and
678 phosphorylation in skeletal muscle at rest and during exercise. *American journal of physiology*
679 *Endocrinology and metabolism* 291: E1106-1114, 2006.

680 61. **Sainz N, Rodriguez A, Catalan V, Becerril S, Ramirez B, Gomez-Ambrosi J, and Fruhbeck G.**
681 Leptin administration favors muscle mass accretion by decreasing FoxO3a and increasing PGC-1alpha
682 in ob/ob mice. *PLoS one* 4: e6808, 2009.

683 62. **Schiaffino S.** Fibre types in skeletal muscle: a personal account. *Acta physiologica (Oxford,*
684 *England)* 199: 451-463, 2010.

685 63. **Simoneau JA, and Bouchard C.** Genetic determinism of fiber type proportion in human
686 skeletal muscle. *FASEB journal : official publication of the Federation of American Societies for*
687 *Experimental Biology* 9: 1091-1095, 1995.

688 64. **Simoneau JA, and Bouchard C.** Human variation in skeletal muscle fiber-type proportion and
689 enzyme activities. *The American journal of physiology* 257: E567-572, 1989.

690 65. **Sosnicki A.** Histopathological observation of stress myopathy in M. longissimus in the pig
691 and relationships with meat quality, fattening and slaughter traits. *Journal of animal science* 65: 584-
692 596, 1987.

693 66. **Srikanthan P, and Karlamangla AS.** Muscle Mass Index As a Predictor of Longevity in Older
694 Adults. *The American Journal of Medicine* 127: 547-553, 2014.

- 695 67. **Staron RS, Hagerman FC, Hikida RS, Murray TF, Hostler DP, Crill MT, Ragg KE, and Toma K.**
696 Fiber type composition of the vastus lateralis muscle of young men and women. *The journal of*
697 *histochemistry and cytochemistry : official journal of the Histochemistry Society* 48: 623-629, 2000.
- 698 68. **Stavaux D, Art T, McEntee K, Reznick M, and Lekeux P.** Muscle fibre type and size, and
699 muscle capillary density in young double-muscled blue Belgian cattle. *Zentralblatt fur*
700 *Veterinarmedizin Reihe A* 41: 229-236, 1994.
- 701 69. **Timson BF, Bowlin BK, Dudenhoeffer GA, and George JB.** Fiber number, area, and
702 composition of mouse soleus muscle following enlargement. *Journal of applied physiology*
703 *(Bethesda, Md : 1985)* 58: 619-624, 1985.
- 704 70. **Totsuka Y, Nagao Y, Horii T, Yonekawa H, Imai H, Hatta H, Izaikai Y, Tokunaga T, and Atomi**
705 **Y.** Physical performance and soleus muscle fiber composition in wild-derived and laboratory inbred
706 mouse strains. *Journal of applied physiology (Bethesda, Md : 1985)* 95: 720-727, 2003.
- 707 71. **Van den Maagdenberg K, Stinckens A, Lefaucheur L, Buys N, and De Smet S.** The effect of
708 mutations in the insulin-like growth factor-II and ryanodine receptor-1 genes on biochemical and
709 histochemical muscle fibre characteristics in pigs. *Meat science* 79: 757-766, 2008.
- 710 72. **van der Laarse WJ, Diegenbach PC, and Maslam S.** Quantitative histochemistry of three
711 mouse hind-limb muscles: the relationship between calcium-stimulated myofibrillar ATPase and
712 succinate dehydrogenase activities. *The Histochemical journal* 16: 529-541, 1984.
- 713 73. **Vaughan HS, Aziz U, Goldspink G, and Nowell NW.** Sex and stock differences in the
714 histochemical myofibrillar adenosine triphosphatase reaction in the soleus muscle of the mouse. *The*
715 *journal of histochemistry and cytochemistry : official journal of the Histochemistry Society* 22: 155-
716 159, 1974.
- 717 74. **Wade AJ, Marbut MM, and Round JM.** Muscle fibre type and aetiology of obesity. *Lancet*
718 335: 805-808, 1990.
- 719 75. **Wang YX, Zhang CL, Yu RT, Cho HK, Nelson MC, Bayuga-Ocampo CR, Ham J, Kang H, and**
720 **Evans RM.** Regulation of muscle fiber type and running endurance by PPARdelta. *PLoS biology* 2:
721 e294, 2004.
- 722 76. **Wilkins BJ, Dai YS, Bueno OF, Parsons SA, Xu J, Plank DM, Jones F, Kimball TR, and**
723 **Molkentin JD.** Calcineurin/NFAT coupling participates in pathological, but not physiological, cardiac
724 hypertrophy. *Circulation research* 94: 110-118, 2004.
- 725 77. **Wimmers K, Fiedler I, Hardge T, Murani E, Schellander K, and Ponsuksili S.** QTL for
726 microstructural and biophysical muscle properties and body composition in pigs. *BMC genetics* 7: 15,
727 2006.
- 728 78. **Wirtz P, Loermans HM, Peer PG, and Reintjes AG.** Postnatal growth and differentiation of
729 muscle fibres in the mouse. I. A histochemical and morphometrical investigation of normal muscle.
730 *Journal of anatomy* 137 (Pt 1): 109-126, 1983.
- 731 79. **Wu C, Orozco C, Boyer J, Leglise M, Goodale J, Batalov S, Hodge CL, Haase J, Janes J, Huss**
732 **JW, 3rd, and Su AI.** BioGPS: an extensible and customizable portal for querying and organizing gene
733 annotation resources. *Genome biology* 10: R130, 2009.
- 734 80. **Zhang Y, Xu P, Lu C, Kuang Y, Zhang X, Cao D, Li C, Chang Y, Hou N, Li H, Wang S, and Sun X.**
735 Genetic linkage mapping and analysis of muscle fiber-related QTLs in common carp (*Cyprinus carpio*
736 L.). *Marine biotechnology (New York, NY)* 13: 376-392, 2011.

A



B



500 μ m



Chromosome 11





