



# *Effect of composting and soil type on dissipation of veterinary antibiotics in land-applied manures*

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Chen, C., Ray, P., Knowlton, K. F., Pruden, A. and Xia, K. (2018) Effect of composting and soil type on dissipation of veterinary antibiotics in land-applied manures. *Chemosphere*, 196. pp. 270-279. ISSN 0045-6535 doi: <https://doi.org/10.1016/j.chemosphere.2017.12.161> Available at <http://centaur.reading.ac.uk/74979/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.chemosphere.2017.12.161>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

1 **Effect of composting and soil type on dissipation of veterinary**  
2 **antibiotics in land-applied manures**

3 **Chaoqi Chen<sup>\*1</sup>, Partha Ray<sup>2,4</sup>, Katharine F. Knowlton<sup>2</sup>, Amy Pruden<sup>3</sup>, and Kang Xia<sup>1</sup>**

4 <sup>1</sup>Department of Crop and Soil Environmental Sciences, Virginia Tech, Blacksburg, Virginia  
5 24061, United States

6 <sup>2</sup>Department of Dairy Science, Virginia Tech, Blacksburg, Virginia 24061, United States

7 <sup>3</sup>Department of Civil & Environmental Engineering, Virginia Tech, Blacksburg, Virginia 24061,  
8 United States

9 <sup>4</sup>Division of Animal, Dairy & Food Chain Sciences, School of Agriculture, Policy and  
10 Development, Univ. of Reading, Reading RG6 6AR, UK

11

12

13

14

15

## 16 Abstract

17 The objective of this study was to determine the fate of commonly used veterinary antibiotics in  
18 their naturally excreted form when manure-based amendments are applied to soil. Beef cattle  
19 were administered sulfamethazine, tylosin, and chlortetracycline and dairy cows were treated  
20 with pirlimycin according to standard animal production practice. The resulting manure was  
21 composted for 42 days under static or turned conditions and applied at agronomic N rates to  
22 sandy, silt, and silty clay loam soils and compared with amendment with corresponding raw  
23 manures in sacrificial microcosms over a 120-day period. Antibiotic dissipation in the raw  
24 manure-amended soils followed bi-phasic first order kinetics. The first phase half-lives for  
25 sulfamethazine, tylosin, chlortetracycline, and pirlimycin ranged from 6.0 to 18 days, 2.7 to 3.7  
26 days, 23 to 25 days, and 5.5 to 8.2 days, respectively. During the second phase, dissipation of  
27 sulfamethazine was negligible, while the half-lives for tylosin, chlortetracycline, and pirlimycin  
28 ranged from 41 to 44 days, 75 to 144 days, and 87 to 142 days, respectively. By contrast,  
29 antibiotic dissipation in the compost-amended soils followed single-phase first order kinetics  
30 with negligible dissipation of sulfamethazine and half-lives of tylosin and chlortetracycline  
31 ranging from 15 to 16 days and 49 to 104 days, respectively. Pirlimycin was below the detection  
32 limit in the compost-amended soils. After incubating 120-days, antibiotics in compost-amended  
33 soils (up to  $3.1 \mu\text{g kg}^{-1}$ ) were significantly lower than in the manure-amended soils (up to  $19 \mu\text{g}$   
34  $\text{kg}^{-1}$ ;  $p < 0.0001$ ), with no major effect of soil type on the dissipation. Risk assessment suggested  
35 that manure composting can reduce antibiotic resistance selection potential in manure-amended  
36 soils.

37 Keywords: environmental fate, antibiotics, dairy, beef, soil, risk assessment

38

## 39 Highlights

- 40 • Antibiotic dissipation follows bi-phasic 1<sup>st</sup>-order kinetics in manure-amended soils
- 41 • Antibiotic dissipation obeys single-phase 1<sup>st</sup>-order kinetics in compost-amended soils
- 42 • Manure-borne antibiotics persist in soils at low concentrations
- 43 • Soil type had negligible effect on dissipation kinetics of manure-borne antibiotics
- 44 • Manure composting can reduce antibiotic resistance selection potential in manure-
- 45 amended soils

46

47 1. Introduction

48 Antibiotics are widely used in livestock for therapeutic and sub-therapeutic uses (Chee-Sanford et  
49 al., 2009). About 80% of the 13.5 million kilograms of antibiotics sold yearly in the USA is used  
50 in animal production (Done et al., 2015), among which macrolides are categorized as “critically  
51 important” while sulfonamides, tetracyclines and cephalosporins are categorized as “highly  
52 important” in human medicine by World Health Organization (Collignon et al., 2016).  
53 Administered antibiotics are not fully metabolized, leading to their excretion in animal manure  
54 (Chiesa et al., 2015). In the environment, soil is the primary receiver of antibiotics used in animal  
55 production, mainly through land application of manure (Jechalke et al., 2014). A wide range of  
56 antibiotics has been detected worldwide in various manure and manure-amended soils, at  
57 concentrations up to several thousand  $\mu\text{g kg}^{-1}$  (Martinez-Carballo et al., 2007; Ho et al., 2014; Li  
58 et al., 2015; Yang et al., 2016).

59 Although composting has been proposed as a mean to reduce antibiotic levels in manure before  
60 land application, recent studies have indicated that it cannot completely remove antibiotics  
61 (Dolliver et al., 2008; Cessna et al., 2011; Ray et al., 2017). There is concern that antibiotics  
62 released into soils could create a selective pressure on the native microbial community (Cleary et  
63 al., 2016; Nordenholt et al., 2016) and enrich resistant bacteria and resistance genes (Jechalke et  
64 al., 2014).

65 Previous studies demonstrated that antibiotic dissipation in soils is influenced by environmental  
66 factors, such as temperature, pH, and soil physicochemical properties, as well as the initial  
67 concentrations of antibiotics and their associated matrixes (Otker and Akmehmet-Balcioglu, 2005;  
68 Cengiz et al., 2010; Srinivasan and Sarmah, 2014). Chemical properties of the antibiotics can also

69 influence their interactions with soil minerals and organic matter (Otker and Akmehmet-Balcioglu,  
70 2005; Wegst-Uhrich et al., 2014). To date, investigations on antibiotic fate in soils have largely  
71 been conducted on antibiotic-spiked soils or soils mixed with antibiotic-spiked manure (Carlson  
72 and Mabury, 2006; Fang et al., 2014; Pan and Chu, 2016). Although such approaches are in  
73 accordance with the US EPA guidelines for pesticides (USEPA., 2008a, b), they ignore effects of  
74 passing through the animal gut and subsequent manure management on the interactions of  
75 antibiotics with manure matrix before land application. Such influences are important because: *1)*  
76 antibiotics typically enter the soil via a manure matrix, whereas pesticides are normally directly  
77 applied to soils; *2)* passage through the animal gut will influence their sorption within the manure  
78 matrix, partially biodegrade antibiotics, and alter microbial communities involved; and *3)* manure  
79 amendment alters soil physical, chemical, and biological properties, all of which will influence the  
80 fate of antibiotics.

81 Although there is limited information comparing the environmental fate of manure-borne versus  
82 spiked antibiotics, studies of biosolids have demonstrated significantly faster dissipation rates of  
83 the antimicrobial triclosan when spiked into soils compared to when it is biosolids-borne (Kwon  
84 and Xia, 2012). Decreased diffusion, increased sorption, and reduced bioavailability when  
85 biosolids-borne could contribute to prolonged persistence, compared to predicted values and those  
86 determined using spiked soils.

87 The objective of this study was to determine the effect of composting and soil type on the  
88 dissipation of manure-borne antibiotics following land-application, using manure derived from  
89 antibiotic-administrated dairy cows and beef cattle. The results help to evaluate whether benefits  
90 of composting using FDA FSMA guidelines (FDA, 2014) extend towards reducing antibiotics and

91 their potential impacts in soils, particularly within the USDA National Organic Program's  
92 recommended 120-day waiting period between raw manure application and harvest of crops that  
93 are in contact with soil (USDA, 2012).

## 94 2. Materials and methods

### 95 *2.1 Raw manure and compost*

96 Raw manure was collected from sulfamethazine, tylosin and chlortetracycline-treated beef cattle  
97 or pirlimycin-treated dairy cows ([Table S1](#) and [Text S1 of Supplemental Information \(SI\)](#)). After  
98 laboratory-scale composting for 42-days, using static and turned techniques as recommended by  
99 the FDA (FDA, 2014) and described in a previous study (Ray et al., 2017), compost was collected  
100 from composting tumblers ([Text S1, SI](#)). The physical properties and antibiotic concentrations in  
101 raw manure and the compost are listed in [Table S2](#).

### 102 *2.2 Soil microcosms*

103 The sandy loam, silt loam, and silty clay loam soils were top soils (0-5 cm) collected from three  
104 locations in Virginia ([Text S2, SI](#) and [Table S3](#)).

105 For each microcosm unit, 15 g air-dried soil sieved through 2-mm was added to a 150 mL  
106 pre-washed (70% ethanol) and air-dried glass jar. Calculated based on the typical agronomic  
107 nitrogen application rate in Virginia (Evanylo, 2009), 2.44 g raw manure or 1.60 g compost was  
108 then added to a glass jar and mixed by hand shaking and stirring. Ultrapure water was added to the  
109 manure-soil mixture to bring the soil moisture to 50% of its field moisture capacity and maintained  
110 by recording the total weight of each microcosm jar. Each jar was covered loosely with an  
111 aluminum foil sheet to reduce moisture loss while maintaining aerobic conditions. All microcosms



112 were kept in the dark at room temperature (20°C) with soil moisture adjusted weekly by adding  
113 ultrapure water to bring the total weight of each jar to its recorded weight. Soils were collected in  
114 triplicate on days 0, 1, 3, 7, 29, 57, 90 and 120 via destructive sampling, freeze dried, and stored  
115 at -20°C for antibiotics analysis.

116 Antibiotics were analyzed using SPE-UPLC-MS/MS method with small modifications (Text S3,  
117 SI) (Ray et al., 2014; Ray et al., 2017). The samples were extracted using sonication with methanol:  
118 McIlvaine buffer (50:50, v/v) or methanol: phosphate buffer (70:30, v/v). The extracts were  
119 cleaned up using OASIS HLB Cartridges (Waters, Milford, MA). After the sample preparation,  
120 the antibiotics were analyzed using Agilent 1290 UPLC coupled with Agilent 6490 Triple Quad  
121 tandem mass spectrometry (Agilent Technologies Inc., Santa Clara, CA ). To estimate the method  
122 detection limits (MDLs), series dilutions of antibiotic standards were spiked into manure amended  
123 soils. The spiked samples were processed through the entire analytical method. The MDLs were  
124 then determined using seven samples spiked near the lowest concentration that was detected. The  
125 MDLs for sulfamethazine, tylosin, chlortetracycline, and pirlimycin were 0.13, 0.25, 0.59, and  
126 1.54  $\mu\text{g kg}^{-1}$ , respectively. The method quantification limits (3.3 time MDLs) for sulfamethazine,  
127 tylosin, chlortetracycline, and pirlimycin were 0.43, 0.83, 1.95, and 1.78  $\mu\text{g kg}^{-1}$ , respectively. The  
128 recoveries for sulfamethazine, tylosin, chlortetracycline, and pirlimycin ranged from 90% to 118%,  
129 81% to 121%, 58% to 76%, and 89% to 91% in manure amended soils, respectively.

### 130 *2.3 Data graphing and statistical analysis*

131 Bi-phasic or single phasic first order kinetic models were used to fit the  $C_t/C_o$  vs  $t$  curves. The  
132 rate constants were acquired based on the slope of the curve fit of  $\ln(C_t/C_o)$  vs  $t$  to the respective

133 model and the half-lives were derived from the calculated rate constants. If all triplicate samples  
134 were non-detect, then the  $C_t/C_o$  point was noted as below the detection limit (BDL).

135 All statistical analyses were carried out using JMP (JMP<sup>®</sup>, Version 12.0. SAS Institute Inc.,  
136 Cary, NC, 1989-2007). For non-detect samples, half of the method detection limit was used for  
137 statistical purposes. Two-way analysis of variance (ANOVA) was carried out at the 95%  
138 confidence level to test the effects of amendment type (raw manure, static compost, or turned  
139 compost), soil type, and their interactions on antibiotic dissipation. One-way ANOVA was used  
140 to test the effect of amendment type on the final antibiotic concentrations in soils at the end of  
141 microcosm study. Multiple paired comparisons were conducted using the Tukey-Kramer method  
142 (Tukey, 1949).

#### 143 *2.4 Assessment of antibiotic resistance selection potential*

144 Risk quotient (RQ) values were applied to assess the potential impact of antibiotic residue on  
145 antibiotic resistance selection potential in soils at day 0 and 120 days after manure application.  
146 The RQ value was calculated as the ratio of the measured concentrations in the soils before and  
147 after incubation (Table 1) to the predicted no-effect concentrations for antibiotic resistance  
148 selection in soils ( $PNEC_{soil}$ ). The predicted no-effect concentration in soil for an antibiotic  
149 ( $PNEC_{soil}$ ) can be calculated using the following equations (Thomaidi et al., 2016):

$$150 \quad PNEC_{soil} = PNEC_{water} \times k_d = PNEC_{water} \times k_{oc} \times f_{oc}$$

151 Where  $PNEC_{water}$  is the predicted no-effect concentration for resistance selection in water ( $\mu\text{g L}^{-1}$ ),  
152  $k_d$  is the soil-water partition coefficient of antibiotics ( $\text{L kg}^{-1}$ ),  $k_{oc}$  is the soil organic carbon-water  
153 partitioning coefficient of an antibiotic. The  $PNEC_{soil}$  for sulfamethazine, tylosin, chlortetracycline,

154 and pirlimycin were estimated (Text S4, SI) and ranged from 58-100, 7.0-12, 0.68-1.2, and 10-18  
155  $\mu\text{g kg}^{-1}$ , respectively (Table S8). Similar to the classification used for ecological risk evaluation  
156 (Ho et al., 2015), RQ values  $<0.1$ ,  $0.1-1$ , and  $>1$  are categorized to three levels, as “low”, “medium”,  
157 and “high” antibiotic resistance selection potentials, respectively.

### 158 3. Results and discussion

#### 159 3.1 Antibiotic dissipation patterns in the soil microcosms

160 The term “dissipation” is used here to refer to the collective effects of biodegradation,  
161 transformation, sorption, loss of extractability, and other processes that contribute to a net decrease  
162 in measured antibiotic. Limited field studies have reported the dissipation of manure-borne  
163 antibiotics (Halling-Sorensen et al., 2005; Heuer et al., 2008). While such field studies are of value  
164 to gain a general sense of antibiotic behavior in the real world, it is not possible to isolate the  
165 effects of various environmental processes in governing their fate. For example, transport, plant  
166 uptake, and photodegradation will all contribute to some extent to the persistence of antibiotics in  
167 soils at field-scale and cannot be distinguished from other processes, such as biodegradation. In  
168 contrast, microcosms provide the advantage of a closed system with a limited number of variables.

##### 169 3.1.1 Raw manure-amended soils

170 The initial concentrations (dry weight basis) of sulfamethazine, tylosin, and chlortetracycline in  
171 the three cattle manure-amended soils ranged from 29-47, 8.4-10, and 46-80  $\mu\text{g kg}^{-1}$ , respectively  
172 (Table 1). These concentrations were more relevant to real-world conditions compared to levels  
173 typically spiked to soils (100 to 1,000,000  $\mu\text{g kg}^{-1}$ ) (Accinelli et al., 2007; Pan and Chu, 2016).

174 Dissipation of sulfamethazine was rapid in all three soils within the first 7 days, with 43 to 77%  
175 remaining at day 7 (Fig. 1), but slowed significantly thereafter. Sulfamethazine remained constant  
176 at levels of 32 to 45% in the three soils until day 120 (Fig. 1). Our observations are consistent with  
177 a prior field study examining the dissipation of five sulfonamides, including sulfamethazine, in  
178 manure-amended soils (Stoob et al., 2007), where the dissipation was initially fast, but slowed  
179 down considerably after 14 days. Similar results were also reported when examining the  
180 sulfamethazine dissipation in a swine manure-amended sandy loam soil with an initial spiked  
181 concentration of 100,000  $\mu\text{g kg}^{-1}$  (Lertpaitoonpan et al., 2015). Other studies of structurally-related  
182 sulfonamides also yielded similar results (Wang et al., 2006; Liu et al., 2010; Fang et al., 2014).  
183 For example, Fang et al. (2014) observed rapid dissipation of sulfadiazine within 7 days, followed  
184 by dramatically slower dissipation in manure-amended soils with an initial spiked concentration  
185 of 20,000  $\mu\text{g kg}^{-1}$  (Fang et al., 2014). This study supports the overall conclusion that  
186 sulfamethazine becomes more persistent and less bioavailable with time.

187 Similar to the case with sulfamethazine, initial dissipation of tylosin was rapid in raw manure-  
188 amended soils (Fig. 1). By day 7, 44% to 58% of tylosin remained in the three soils. However, in  
189 contrast to sulfamethazine, the dissipation of tylosin continued through 120 days, although at a  
190 slower rate. By 120 days, only 5.9 to 8.4% of tylosin remained in the three soils, with final  
191 concentrations ranging 0.51 to 0.87  $\mu\text{g kg}^{-1}$  (Fig. 1, Table 1). Continuous dissipation of tylosin in  
192 soils were observed in previous studies (Halling-Sorensen et al., 2005; Carlson and Mabury,  
193 2006; Schlusener and Bester, 2006; Hu and Coats, 2007; Sassman and Lee, 2007; Liu et al.,  
194 2010). A field study using manure-borne antibiotics showed that the concentrations of tylosin

195 declined from 50  $\mu\text{g kg}^{-1}$  to 10  $\mu\text{g kg}^{-1}$  in a sandy loam soil and from 25 to 3  $\mu\text{g kg}^{-1}$  in a sand soil  
196 within 155 days (Halling-Sorensen et al., 2005).

197 The trend in the dissipation of chlortetracycline in the raw manure-amended soils was similar to  
198 that of tylosin; however, initial dissipation rates were markedly lower (Fig. 1). At day 7, 62% to  
199 72% of chlortetracycline remained in the three soils, while around 50% of tylosin was  
200 transformed by day 7 (Fig. 1). Continuous dissipation of chlortetracycline in soils has also been  
201 observed in laboratory and field experiments (Halling-Sorensen et al., 2005; Carlson and  
202 Mabury, 2006; Zhang and Zhang, 2010; Fang et al., 2014). In a field study using manure-borne  
203 antibiotics, 50% reduction of chlortetracycline was observed in both a sandy loam and sandy soil  
204 within 20-34 days and 28-42 days, respectively (Halling-Sorensen et al., 2005).

205 In contrast to the above antibiotic dissipation patterns, the levels of pirlimycin in all raw manure-  
206 amended soils first increased 1.98-2.70 times from day 0 to day 3, rapidly decreased from day 3  
207 to day 29, and then remained relatively constant thereafter until day 120 (Fig. 1). The initial  
208 spike in pirlimycin concentration was most notable in the sandy loam soil (Table 1). Previous  
209 research has shown that pirlimycin administered to dairy cows can be conjugated in the liver  
210 and gastrointestinal tract to form pirlimycin-sulfoxide, pirlimycin-sulfone, pirlimycin-adenylate,  
211 pirlimycin-uridylate, and pirlimycin sulfoxide-adenylate (Hornish et al., 1998). These conjugates  
212 are subsequently excreted into the feces and urine at substantial levels, up to 50% of the total  
213 secreted pirlimycin. Therefore, we hypothesize that the initial observed rise in concentration was  
214 likely due to deconjugation of pirlimycin conjugates back to pirlimycin.

### 215 3.1.2 Compost-amended soils

216 Although a significant proportion (62%-99%) of manure-borne antibiotics can be transformed  
217 during composting (Ray et al., 2017), to the best of our knowledge no prior study has examined  
218 whether the residual antibiotics in finished compost (e.g., [Table S2](#)) are subject to further  
219 dissipation after application to soil.

220 The initial concentrations of sulfamethazine in the three soils ranged from 1.1-1.6  $\mu\text{g kg}^{-1}$  and  
221 from 0.35 to 0.80  $\mu\text{g kg}^{-1}$ , after amending with static and turned composts, respectively ([Table](#)  
222 [1](#)). One prior study indicated that the highest concentration of chlortetracycline in soils nearby a  
223 swine manure composting facility was 0.85  $\mu\text{g kg}^{-1}$  (Awad et al., 2014). This highlights the  
224 importance of understanding the fate of low concentrations of antibiotics in compost-amended  
225 soils. In contrast to raw manure-amended soils, no dissipation of sulfamethazine was observed in  
226 the compost-amended soils over the 120-day period ([Fig. 1](#)), indicating that it may not have been  
227 bioavailable to soil microorganisms. This was consistent with the observation that no dissipation  
228 of sulfamethazine occurred after day 7 in raw manure-amended soil ([Fig. 1](#)). Strong adsorption to  
229 compost could be a key factor limiting the bioavailability of sulfamethazine and contributing to  
230 its observed persistence.

231 The initial concentrations of tylosin in soils after application with static compost and turned  
232 compost ranged from 1.3 to 1.8  $\mu\text{g kg}^{-1}$  and from 0.27 to 1.2  $\mu\text{g kg}^{-1}$ , respectively ([Table 1](#)).  
233 Rapid dissipation was observed within the first 7 days of microcosm incubation, resulting in 48-  
234 54% and 28-68% of the initially added tylosin remaining in all three soils for static compost-  
235 amended soil and turned compost-amended soil, respectively ([Fig. 1](#)). By day 57, tylosin was  
236 below detection in all soils applied with compost.

237 Initial concentrations of chlortetracycline ranged from 7.5 to 11  $\mu\text{g kg}^{-1}$  and from 4.7 to 6.4  $\mu\text{g}$   
238  $\text{kg}^{-1}$  (Table 1) after application of static compost and turned compost to soil, respectively.  
239 Compared to tylosin, chlortetracycline dissipation was much slower during the first week. By  
240 day 7, 82-92% and 82-106% of the initially added chlortetracycline remained in all three soils for  
241 static compost-amended soils and turned compost-amended soils, respectively (Fig. 1). After 120  
242 days, chlortetracycline was still above the detection limit, with concentrations ranging from 1.8  
243 to 3.1  $\mu\text{g kg}^{-1}$  and from 1.1 to 2.4  $\mu\text{g kg}^{-1}$  in static and turned compost-amended soil, respectively  
244 (Table 1).

## 245 3.2 Antibiotic dissipation rates in the soil microcosms

### 246 3.2.1 Raw manure-amended soils

247 Single-phase first order kinetics did not adequately describe the dissipation of the target  
248 antibiotics in raw manure-amended soils in this study, as the coefficients of determination ( $R^2$ )  
249 varied largely from 0.27 to 0.93 upon fitting the data to a single-phase first order kinetic model.  
250 When fitting their dissipation using bi-phasic first order kinetics using the Hockey-Stick model  
251 (Sarmah and Rohan, 2011),  $R^2$  values ranged from 0.94 to 0.99. This model consists of two  
252 sequential first-order kinetics with the integrated equation shown below:

$$253 C_t = C_o e^{-k_1 t_b} \text{ for } t < t_b, C_t = C_o e^{-k_1 t_b} e^{-k_2 (t - t_b)}, \text{ for } t > t_b$$

254 Where  $C_t$  is the compound concentration ( $\mu\text{g kg}^{-1}$ ) at time  $t$  (d) after application,  $C_o$  is the initial  
255 concentration ( $\mu\text{g kg}^{-1}$ ),  $k_1$  is the rate constant ( $\text{d}^{-1}$ ) until  $t = t_b$ . The time at which rate constant  
256 changes from  $k_1$  to  $k_2$  is denoted by  $t_b$  (breakpoint). The breakpoints for sulfamethazine, tylosin,  
257 and chlortetracycline were 7 days, 3 days, and 29 days, respectively. For pirlimycin, due to the

258 initial deconjugation of its conjugates, its  $C_o$  is defined as the peak concentration detected at day  
259 3, with a corresponding breakpoint of 29 days (26 days after day 3).

260 The antibiotic dissipation rate constants in the raw manure-amended soils are shown in [Table 2](#).  
261 Because the dairy manure matrix is distinct from that of beef cattle manure, the dissipation of  
262 pirlimycin is discussed separately. For the three soils, the first phase dissipation rate constants  
263 ( $k_1$ ) ranked in the order of tylosin > sulfamethazine > chlortetracycline ([Table 2](#)). It has been  
264 shown that antibiotic degradation is typically catalyzed by different extracellular hydrolytic  
265 enzymes (protease, lipase, and cellulase) released by microorganisms, mainly in the aqueous  
266 phase of soil systems (Thiele-Bruhn, 2003). Therefore, the overall dissipation rate of antibiotics  
267 is largely affected by their hydrophilicity (Otker and Akmeahmet-Balcioglu, 2005; Wegst-Uhrich  
268 et al., 2014). Accordingly, the observed order of dissipation rate constants was consistent with  
269 the order of the water solubility of these three compounds: tylosin ( $5000 \text{ mg L}^{-1}$ ) >  
270 sulfamethazine ( $1500 \text{ mg L}^{-1}$ ) > chlortetracycline ( $600 \text{ mg L}^{-1}$ ) ([Table S1](#)).

271 The dissipation rate constants of these three antibiotics in the second phase ( $k_2$ ) were much lower  
272 than those in the first phase ( $k_1$ ) ([Table 2](#)). Availability-adjusted first order kinetic models  
273 assume that antibiotic availability in soils decreases exponentially with time, largely due to  
274 sorption, and have been applied in prior studies with decreasing dissipation rates (Wang et al.,  
275 2006; Stoob et al., 2007; Pan and Chu, 2016). Sorption of antibiotics is an important factor  
276 affecting the dissipation rate (Otker and Akmeahmet-Balcioglu, 2005; Wegst-Uhrich et al., 2014)  
277 and thus the partitioning coefficient ( $k_d$ ) is a key parameter used in estimating the migration  
278 potential of aqueous-phase contaminants in contact with solid soil components. Median  $k_d$  values  
279 of sulfamethazine and tylosin reported in prior literature were 3 and  $100 \text{ L kg}^{-1}$ , respectively



280 (Wegst-Uhrich et al., 2014), while the  $k_d$  values of chlortetracycline ranged from 1208-2386 L  
281  $\text{kg}^{-1}$  (Sarmah et al., 2006). This suggests that soil sorption tendency (loss of availability) follows  
282 the order of chlortetracycline > tylosin > sulfamethazine. In the current study, the rate constants  
283 for the second phase ( $k_2$ ) were in the order of tylosin > chlortetracycline > sulfamethazine, which  
284 is not consistent with the assumption of loss of bioavailability of the antibiotics in soils due to  
285 adsorption. Antibiotics examined in previous studies were spiked into the systems (Wang et al.,  
286 2006; Stoob et al., 2007; Pan and Chu, 2016), while our study utilized manure from antibiotic-  
287 treated animals. Partitioning coefficients of antibiotics in manure are likely different from those  
288 described for soils (Loke et al., 2002). Also, in prior studies spiking antibiotics, it is likely that  
289 the interactions of antibiotics with manure or soil components did not achieve a steady state  
290 before dissipation began. Therefore, the bioavailability of the antibiotics in the soils could  
291 decrease over time. On the other hand, the antibiotics in our study are more likely to have  
292 reached equilibrium with the manure matrix after passing through the digestion system. After  
293 application of manure to the soils, it is likely that desorption begins to dominate sorption in the  
294 second dissipation phase, at which point aqueous antibiotic dissipation is near completion. The  
295 desorption of chlortetracycline could be retarded due to the lowest dissipation rate of the released  
296 chlortetracycline (smallest  $k_1$ ). By contrast, the desorption of tylosin could be accelerated due to  
297 the fastest dissipation rate of the released tylosin (highest  $k_1$ ). Lack of dissipation of  
298 sulfamethazine in the second phase may due to a fraction of sulfamethazine that is irreversibly  
299 sorbed to the manure.

300 Pirlimycin dissipation rate constants in all raw manure-amended soils were higher during the  
301 first phase (day 3 to 29) compared to the second phase (day 29 to 120) (Table 2). Similarly,

302 dissipation of clindamycin (a lincosamide antibiotic) in biosolids followed a biphasic pattern,  
303 with faster dissipation during the first phase followed by relatively stabilized second phase (Wu  
304 et al., 2009).

### 305 *3.2.2 Compost-amended soils*

306 Since sulfamethazine concentrations remained stable in compost-amended soils over the 120  
307 days, no rate constants could be estimated. By contrast, tylosin was below detection limit by day  
308 57. Therefore, for compost-amended soil, a concentration at half of the detection limit ( $0.12 \mu\text{g}$   
309  $\text{kg}^{-1}$ ) of tylosin was assumed beyond day 57, with simple first order kinetics used to fit the curve  
310 from day 0 to day 57, with  $R^2$  ranged from 0.67-0.99. . For chlortetracycline, simple first order  
311 kinetics were applied to fit the curves for soils applied with both compost types, with  $R^2$  ranging  
312 from 0.88 to 0.97. In compost-amended soils, the rate constants of tylosin in all three soils were  
313 greater than those for chlortetracycline, which is consistent with the trend for these two  
314 antibiotics in raw manure-amended soils. Curve fitting was not conducted for pirlimycin because  
315 it was below the detection limit in all compost-amended soils over the duration of the study  
316 (Table 2).

### 317 *3.3 Antibiotic dissipation half-lives in the soil microcosms*

318 The half-lives of the four target antibiotics in the current study are shown in [Table 3](#). In the raw  
319 manure amended soils, the half-lives of tylosin, chlortetracycline, and pirlimycin in the second  
320 phase were 14, 4.6, and 16 times as long as those in the first phase ([Table 3](#)), respectively,  
321 indicating that manure-borne antibiotics could persist in soil at low concentrations for a long  
322 period of time. The observed bi-phasic dissipation patterns suggest that a portion of the manure-

323 borne antibiotics is immediately bioavailable and transformed rapidly after manure application.  
324 The remaining portion appears to be released slowly from the manure as dissipation continues.  
325 The BIOWIN model in EPI Suite™ (USEPA., 2012) was applied to predict half-lives  
326 specifically with respect to primary biodegradation, estimating values of 8.67 days for  
327 sulfamerazine and 15 days for tylosin and chlortetracycline. The EPI Suite™-predicted half-lives  
328 are similar to the dissipation half-lives measured for the raw-manure amended soils, while, for  
329 most of the cases, significantly shorter than the second-phase dissipation half-lives and the  
330 single-phase half-lives for the compost-amended soils (Table 3). This suggests that the initial  
331 antibiotics are more bioavailable in the raw manure-amended soil and their dissipation is most  
332 likely biologically-driven. By contrast, in the later phase or in the compost-added soils these  
333 compounds became more recalcitrant and their dissipation is more likely affected by a complex  
334 array of biological, chemical, and physical factors.

335 Varied half-lives of antibiotics had been reported in the literature (Table 3) Lertpaitoonpan  
336 (2008) noted that longer half-lives of sulfamethazine were observed with higher initial spiked  
337 concentrations and suggested that microbial activity may be inhibited by higher antibiotic  
338 concentrations (Lertpaitoonpan, 2008). However, the highest concentration of sulfamethazine  
339 determined in the present study,  $47 \mu\text{g kg}^{-1}$  (Table 3), is far below these previous studies. An  
340 effective concentration ( $\text{EC}_{10}$  values) of  $13,000 \mu\text{g kg}^{-1}$  sulfamethazine was required to influence  
341 microbial respiration in rice paddy soils (Liu et al., 2009). Besides, even dissipation of antibiotics  
342 within the same class can vary. For example, six commonly used antibiotics were spiked into a  
343 sandy loam soils with an initial concentration of  $2,000 \mu\text{g kg}^{-1}$  for a 120-day microcosm study to  
344 examine their dissipation (Martinez-Carballo et al., 2007). Among them, the half-lives for four

345 structurally-related macrolides, including tylosin, erythromycin, oleandomycin, and  
346 roxithromycin ranged from 8 days to >120 days.

347 The half-life of pirlimycin in 0.1 N NaOH (pH 12.5) solution and in pure water (under UV  
348 exposure) were 5 and 6.7 days and thus comparable to those estimated for the first phase of  
349 dissipation in the raw manure amended soil current study.

350 In summary, the half-lives of antibiotics in soils reported in the literature appears system-specific  
351 and guidance may need to be system specific and incorporate safety factors, assuming the  
352 longest observed dissipation rates.

### 353 *3.4 Effect of amendment and soil type on dissipation of manure-borne antibiotics in soils*

354 Potential interactive effects of manure amendment type and soil type on antibiotic dissipation  
355 were examined, but none were found (Table 4). Overall, composting appears to be a promising  
356 approach for reducing antibiotic input to soils before manure land application. At the same  
357 nitrogen application rates, the initial antibiotic concentrations were much lower in compost than  
358 in manure-amended soils and remained low throughout the study period, with a much lower end-  
359 point concentration (Table 1). However, lower initial concentrations can translate to slower  
360 subsequent dissipation rates, as was observed for the compost-amended soils relative to the first  
361 phase dissipation rates in the raw manure-amended soils (Table 2). As suggested by the  
362 comparison of the EPI Suite™-predicted half-lives and the measured half-lives, antibiotics in  
363 compost are less bioavailable comparing to raw manure because most of the available fraction is  
364 transformed and the residual fraction becomes more recalcitrant during composting. Our prior  
365 study observed decreasing dissipation rate of antibiotics during manure composting (Ray et al.,

366 2017). Notably, static versus turned compost did not result in significantly different dissipation  
367 patterns or rates in soils (Table 3), which may be related to the high similarity of the small-scale  
368 compost conditions (Ray et al., 2017).

369 Statistically significant differences were not observed for dissipation of manure-borne antibiotics  
370 among different types of soil receiving manure application (Table 3 and Table S4). Soil  
371 properties, such as pH, organic matter content, and clay content theoretically could affect the  
372 partition coefficient of antibiotics (Gao and Pedersen, 2010; Wegst-Uhrich et al., 2014) and,  
373 therefore, affect the dissipation of antibiotics in soils. In particular, hydrophobic interactions  
374 between chemicals and the organic matter is considered to be a predominant mechanism of  
375 sorption (Zhang et al., 2010). However, these interactions and factors might not be applicable for  
376 manure-borne antibiotics because different from antibiotics that are spiked into soil systems, the  
377 manure-borne antibiotics enter into the soils are likely in various complexed forms with manure  
378 matrixes, most likely with the organic matter in manure. As a result, soil physic-chemical  
379 properties might become less important, as observed by others (Sassman et al., 2007; Bailey et  
380 al., 2016).

### 381 *3.5 Environmental implication*

382 More so than toxicity, a main concern regarding land application of antibiotic-containing manure  
383 is the potential to select for antibiotics resistance and gene transfer, resulting in accumulation in  
384 soils (Knapp et al., 2010; Knapp et al., 2011). Selection pressure has been reported to occur at  
385 very low antibiotic concentrations, as suggested by susceptible/resistant bacteria competition  
386 tests (Gullberg et al., 2011; Sandegren, 2014). Minimal selective antibiotic concentrations  
387 (MSCs), which could be several hundred-fold below the minimal inhibitory concentrations

388 (MICs) of susceptible bacteria, have been reported to be capable of enriching for resistant  
389 bacteria (Gullberg et al., 2011). Here, antibiotic resistance selection potential was assessed for the  
390 initial and final 120-day concentrations of antibiotic residues in the microcosms. Using the  
391 method described by Bengtsson-Palme and Larsson (2016) (Bengtsson-Palme and Larsson,  
392 2016) which assume the concentrations of antibiotics that inhibit growth of some bacteria will by  
393 consequence have selective effects on the community level, the estimated MSCs of targeted  
394 antibiotics instead of toxicity thresholds were applied to standard risk quotients.

395 The initial sulfamethazine levels were at the upper end of “medium” in raw manure-amended soils  
396 (Fig. 2). After 120 days, although sulfamethazine RQ values in the raw-manure amended soils  
397 decreased, the concentrations levels were still in the “medium” category for antibiotic resistant  
398 selection. In composted-amended soils, sulfamethazine RQ values remained  $<1$  throughout the  
399 120-day incubation period (Fig 2). The initial tylosin levels were at the lower end of “high” in raw  
400 manure-amended soils and at or close to “medium” in compost amended-soils, with RQ values  
401 ranging from 0.1-1 or close to 0.1, respectively (Fig 2). In contrast to sulfamethazine, the potential  
402 for tylosin to select for antibiotic resistance decreased from initial “high” or “medium” levels to  
403 “low” for all soils after 120-day incubation. The potential for chlortetracycline to select for  
404 antibiotic resistance remained “high” for all the soils during the 120-day incubation. Pirlimycin  
405 was detectable only in manure-amended soils. Similar to tylosin, the potential of pirlimycin for  
406 antibiotic resistance selection decreased from the lower end of “high” or upper end of “medium”  
407 levels to “low” after 120 days (Fig 2).

408 The result from this study suggest that composting manure reduces the potential for antibiotic  
409 resistance selection relative to raw manure application to soils. Further, the results support the

410 conceptual benefits of a wait period prior to harvest, especially for raw manure-amended soil.  
411 However, 120 days may not be sufficient for some antibiotics to reduce their potential to a “low”  
412 risk level for antibiotic resistance selection potential. After incubation for 120 days, the  
413 concentrations of antibiotics in raw manure-amended soils were still significantly higher than those  
414 in the compost-amended soils ( $p < 0.001$ ) (Table S5 and Table S6). The persistence of antibiotics  
415 in manure-amended soils and their potential for resistance selection imply that identification of  
416 appropriate manure management practice prior to land application warrants attention.

#### 417 4. Conclusions

418 The study employed a controlled, replicated microcosm approach to understand the effect of  
419 composting and soil type on the dissipation of manure-borne antibiotics in soils amended with  
420 raw manure or compost. Manure-borne antibiotics, including sulfamethazine and  
421 chlortetracycline, can persist in soils at low concentrations for extended periods (120-day).  
422 Extended persistence of these antibiotics in soils indicates the possibility of antibiotic  
423 accumulation in soils with repeated input of antibiotics with manure application over time.  
424 Dissipation of antibiotics in raw manure-amended soils was significantly faster than in compost-  
425 amended soils, but composting reduced initial inputs of antibiotics and generally resulted in  
426 lower levels by 120 days. Soil type did not have a measurable influence on the fate of manure-  
427 borne antibiotics, likely because the complex interactions between antibiotics and manure  
428 components in the animals’ digestive system and during composting reduce the relevance of soil  
429 properties in affecting antibiotic fate. Thus, manure management practices for reducing antibiotic  
430 inputs may be widely applicable to various soil types. Further, composting may be advantageous  
431 for reducing antibiotic inputs to soil systems, while enforcing a wait period prior to crop harvest

432 may provide additional benefits for reducing the chances of contributing to selection and spread  
433 of resistant bacteria.

## 434 Acknowledgements

435 This study was supported by the USDA NIFA Competitive Grants no. 2014-05280, no 2013-  
436 67019-21355, and no. 2013-67021-21140.

## 437 References

438

- 439 Accinelli, C., Koskinen, W.C., Becker, J.M., Sadowsky, M.J., 2007. Environmental fate of two  
440 sulfonamide antimicrobial agents in soil. *J Agric Food Chem* 55, 2677-2682.
- 441 Awad, Y.M., Kim, S.-C., Abd El-Azeem, S.A.M., Kim, K.-H., Kim, K.-R., Kim, K., Jeon, C.,  
442 Lee, S.S., Ok, Y.S., 2014. Veterinary antibiotics contamination in water, sediment, and soil near  
443 a swine manure composting facility. *Environ Earth Sci* 71, 1433-1440.
- 444 Bailey, C., Spielmeyer, A., Hamscher, G., Schüttrumpf, H., Frings, R.M., 2016. The veterinary  
445 antibiotic journey: comparing the behaviour of sulfadiazine, sulfamethazine, sulfamethoxazole  
446 and tetracycline in cow excrement and two soils. *J Soil Sediments* 16, 1690-1704.
- 447 Bengtsson-Palme, J., Larsson, D.G., 2016. Concentrations of antibiotics predicted to select for  
448 resistant bacteria: Proposed limits for environmental regulation. *Environ Int* 86, 140-149.
- 449 Carlson, J.C., Mabury, S.A., 2006. Dissipation kinetics and mobility of chlortetracycline, tylosin,  
450 and monensin in an agricultural soil in Northumberland County, Ontario, Canada. *Environ*  
451 *Toxicol Chem* 25, 1-10.
- 452 Cengiz, M., Balcioglu, I.A., Oruc, H.H., Cengiz, T.G., 2010. Evaluation of the interaction  
453 between soil and antibiotics. *J Environ Sci Health B* 45, 183-189.
- 454 Cessna, A.J., Larney, F.J., Kuchta, S.L., Hao, X., Entz, T., Topp, E., McAllister, T.A., 2011.  
455 Veterinary antimicrobials in feedlot manure: dissipation during composting and effects on  
456 composting processes. *J Environ Qual* 40, 188-198.
- 457 Chee-Sanford, J.C., Mackie, R.I., Koike, S., Krapac, I.G., Lin, Y.F., Yannarell, A.C., Maxwell,  
458 S., Aminov, R.I., 2009. Fate and transport of antibiotic residues and antibiotic resistance genes  
459 following land application of manure waste. *J Environ Qual* 38, 1086-1108.
- 460 Chiesa, L., Nobile, M., Arioli, F., Britti, D., Trutic, N., Pavlovic, R., Panseri, S., 2015.  
461 Determination of veterinary antibiotics in bovine urine by liquid chromatography-tandem mass  
462 spectrometry. *Food Chem* 185, 7-15.
- 463 Cleary, D.W., Bishop, A.H., Zhang, L., Topp, E., Wellington, E.M., Gaze, W.H., 2016. Long-  
464 term antibiotic exposure in soil is associated with changes in microbial community structure and  
465 prevalence of class 1 integrons. *FEMS Microbiol Ecol* 92.
- 466 Collignon, P.C., Conly, J.M., Andremont, A., McEwen, S.A., Aidara-Kane, A., World Health  
467 Organization Advisory Group, B.M.o.I.S.o.A.R., 2016. World Health Organization Ranking of  
468 Antimicrobials According to Their Importance in Human Medicine: A Critical Step for



469 Developing Risk Management Strategies to Control Antimicrobial Resistance From Food  
470 Animal Production. *Clin Infect Dis* 63, 1087-1093.

471 Dolliver, H., Gupta, S., Noll, S., 2008. Antibiotic degradation during manure composting. *J*  
472 *Environ Qual* 37, 1245-1253.

473 Done, H.Y., Venkatesan, A.K., Halden, R.U., 2015. Does the Recent Growth of Aquaculture  
474 Create Antibiotic Resistance Threats Different from those Associated with Land Animal  
475 Production in Agriculture? *AAPS J* 17, 513-524.

476 Evanylo, G.K., 2009. Agricultura Land Application of Biosolids in Virginia: Managing Biosolids  
477 for Agricultural use. Virginia Cooperative Extension.

478 Fang, H., Han, Y., Yin, Y., Pan, X., Yu, Y., 2014. Variations in dissipation rate, microbial  
479 function and antibiotic resistance due to repeated introductions of manure containing  
480 sulfadiazine and chlortetracycline to soil. *Chemosphere* 96, 51-56.

481 FDA, 2014. Food Safety Modernization Act Facts: Biological soil amendments: Subpart F, 2014.  
482 Accessed online at:  
483 <http://www.fda.gov/downloads/Food/GuidanceRegulation/FSMA/UCM359281.pdf>.

484 Gao, J., Pedersen, J.A., 2010. Sorption of sulfonamide antimicrobial agents to humic acid-clay  
485 complexes. *J Environ Qual* 39, 228-235.

486 Gullberg, E., Cao, S., Berg, O.G., Ilback, C., Sandegren, L., Hughes, D., Andersson, D.I., 2011.  
487 Selection of resistant bacteria at very low antibiotic concentrations. *PLoS Pathog* 7, e1002158.

488 Halling-Sorensen, B., Jacobsen, A.M., Jensen, J., Sengelov, G., Vaclavik, E., Ingerslev, F., 2005.  
489 Dissipation and effects of chlortetracycline and tylosin in two agricultural soils: a field-scale  
490 study in southern Denmark. *Environ Toxicol Chem* 24, 802-810.

491 Heuer, H., Focks, A., Lamshöft, M., Smalla, K., Matthies, M., Spiteller, M., 2008. Fate of  
492 sulfadiazine administered to pigs and its quantitative effect on the dynamics of bacterial  
493 resistance genes in manure and manured soil. *Soil Biol Biochem* 40, 1892-1900.

494 Ho, Y.B., Zakaria, M.P., Latif, P.A., Saari, N., 2014. Occurrence of veterinary antibiotics and  
495 progesterone in broiler manure and agricultural soil in Malaysia. *Sci Total Environ* 488-489,  
496 261-267.

497 Ho, Y.B., Zakaria, M.P., Latif, P.A., Saari, N., 2015. Environmental Risk Assessment for  
498 Veterinary Antibiotics and Hormone in Malaysian Agricultural Soil. *Iran J Public Health* 43, 67-  
499 71.

500 Hornish, R.E., Roof, R.D., Wiest, J.R., 1998. Pirlimycin residue in bovine liver--a case of  
501 reverse metabolism. *Analyst* 123, 2463-2467.

502 Hu, D., Coats, J.R., 2007. Aerobic degradation and photolysis of tylosin in water and soil.  
503 *Environ Toxicol Chem* 26, 884-889.

504 Jechalke, S., Heuer, H., Siemens, J., Amelung, W., Smalla, K., 2014. Fate and effects of  
505 veterinary antibiotics in soil. *Trends Microbiol* 22, 536-545.

506 Knapp, C.W., Dolfing, J., Ehlert, P.A., Graham, D.W., 2010. Evidence of increasing antibiotic  
507 resistance gene abundances in archived soils since 1940. *Environ Sci Technol* 44, 580-587.

508 Knapp, C.W., McCluskey, S.M., Singh, B.K., Campbell, C.D., Hudson, G., Graham, D.W.,  
509 2011. Antibiotic resistance gene abundances correlate with metal and geochemical conditions in  
510 archived Scottish soils. *PLoS One* 6, e27300.

511 Kwon, J.-W., Xia, K., 2012. Fate of triclosan and triclocarban in soil columns with and without  
512 biosolids surface application. *Environ. Toxicol. Chem.* 31, 262-269.

513 Lertpaitoonpan, W., 2008. Sorption, degradation, and transport of sulfamethazine in soils and  
514 manure-amended soils. Graduate Theses and Dissertations, Paper 11180.

515 Lertpaitoonpan, W., Moorman, T.B., Ong, S.K., 2015. Effect of Swine Manure on  
516 Sulfamethazine Degradation in Aerobic and Anaerobic Soils. *Water Air Soil Poll* 226, 81.

517 Li, C., Chen, J., Wang, J., Ma, Z., Han, P., Luan, Y., Lu, A., 2015. Occurrence of antibiotics in  
518 soils and manures from greenhouse vegetable production bases of Beijing, China and an  
519 associated risk assessment. *Sci Total Environ* 521-522, 101-107.

520 Liu, F., Ying, G.-G., Yang, J.-F., Zhou, L.-J., Tao, R., Wang, L., Zhang, L.-J., Peng, P.-A., 2010.  
521 Dissipation of sulfamethoxazole, trimethoprim and tylosin in a soil under aerobic and anoxic  
522 conditions. *Environ Chem* 7, 370-376.

523 Liu, F., Ying, G.G., Tao, R., Zhao, J.L., Yang, J.F., Zhao, L.F., 2009. Effects of six selected  
524 antibiotics on plant growth and soil microbial and enzymatic activities. *Environ Pollut* 157,  
525 1636-1642.

526 Loke, M.L., Tjornelund, J., Halling-Sorensen, B., 2002. Determination of the distribution  
527 coefficient (log K<sub>d</sub>) of oxytetracycline, tylosin A, olaquinox and metronidazole in manure.  
528 *Chemosphere* 48, 351-361.

529 Martinez-Carballo, E., Gonzalez-Barreiro, C., Scharf, S., Gans, O., 2007. Environmental  
530 monitoring study of selected veterinary antibiotics in animal manure and soils in Austria.  
531 *Environ Pollut* 148, 570-579.

532 Nordenholt, R.M., Goyne, K.W., Kremer, R.J., Lin, C.H., Lerch, R.N., Veum, K.S., 2016.  
533 Veterinary Antibiotic Effects on Atrazine Degradation and Soil Microorganisms. *J Environ Qual*  
534 45, 565-575.

535 Otker, H.M., Akmehmet-Balcioglu, I., 2005. Adsorption and degradation of enrofloxacin, a  
536 veterinary antibiotic on natural zeolite. *J Hazard Mater* 122, 251-258.

537 Pan, M., Chu, L.M., 2016. Adsorption and degradation of five selected antibiotics in agricultural  
538 soil. *Sci Total Environ* 545-546, 48-56.

539 Ray, P., Chen, C., Knowlton, K.F., Pruden, A., Xia, K., 2017. Fate and Effect of Antibiotics in  
540 Beef and Dairy Manure during Static and Turned Composting. *J Environ Qual* 46, 45-54.

541 Ray, P., Knowlton, K.F., Shang, C., Xia, K., 2014. Method development and validation: solid  
542 Phase extraction-ultra performance liquid chromatography-tandem mass spectrometry  
543 quantification of pirlimycin in bovine feces and urine. *J AOAC Int* 97, 1730-1736.

544 Sandegren, L., 2014. Selection of antibiotic resistance at very low antibiotic concentrations.  
545 *Upsala J Med Sci* 119, 103-107.

546 Sarmah, A.K., Meyer, M.T., Boxall, A.B., 2006. A global perspective on the use, sales, exposure  
547 pathways, occurrence, fate and effects of veterinary antibiotics (VAs) in the environment.  
548 *Chemosphere* 65, 725-759.

549 Sarmah, A.K., Rohan, M., 2011. Evaluation of four mathematical models to describe dissipation  
550 kinetics of 4-n-nonylphenol and bisphenol-A in groundwater-aquifer material slurry. *J Environ*  
551 *Monit* 13, 157-166.

552 Sassman, S.A., Lee, L.S., 2007. Sorption and degradation in soils of veterinary ionophore  
553 antibiotics: monensin and lasalocid. *Environ Toxicol Chem* 26, 1614-1621.

554 Sassman, S.A., Sarmah, A.K., Lee, L.S., 2007. Sorption of tylosin A, D, and A-aldol and  
555 degradation of tylosin A in soils. *Environ Toxicol Chem* 26, 1629-1635.

556 Schlusener, M.P., Bester, K., 2006. Persistence of antibiotics such as macrolides, tiamulin and  
557 salinomycin in soil. *Environ Pollut* 143, 565-571.

558 Srinivasan, P., Sarmah, A.K., 2014. Dissipation of sulfamethoxazole in pasture soils as affected  
559 by soil and environmental factors. *Sci Total Environ* 479-480, 284-291.

560 Stoob, K., Singer, H.P., Mueller, S.R., Schwarzenbach, R.P., Stamm, C.H., 2007. Dissipation  
561 and transport of veterinary sulfonamide antibiotics after manure application to grassland in a  
562 small catchment. *Environ Sci Technol* 41, 7349-7355.

563 Thiele-Bruhn, S., 2003. Pharmaceutical antibiotic compounds in soils - a review. *J Plant Nutr*  
564 *Soil Sc* 166, 145-167.

565 Thomaidi, V.S., Stasinakis, A.S., Borova, V.L., Thomaidis, N.S., 2016. Assessing the risk  
566 associated with the presence of emerging organic contaminants in sludge-amended soil: A  
567 country-level analysis. *Sci Total Environ* 548-549, 280-288.

568 Tukey, J.W., 1949. Comparing individual means in the analysis of variance. *Biometrics* 5, 99-  
569 114.

570 USDA, 2012. Guide for Organic Crop Producers. Accessed online at  
571 <https://www.ams.usda.gov/sites/default/files/media/GuideForOrganicCropProducers.pdf>.

572 USEPA., 2008a. Fate, Transport and Transformation Test Guidelines: OPPTS 835.0001  
573 Principles and Strategies Related to Biodegradation Testing of Organic Chemicals under the  
574 Toxic Substances Control Act (TSCA)

575 USEPA., 2008b. Fate, Transport and Transformation Test Guidelines: OPPTS 835.1230  
576 Adsorption/Desorption (Batch Equilibrium).

577 USEPA., 2012. Estimation Programs Interface Suite™ for Microsoft® Windows, v 4.11]. United  
578 States Environmental Protection Agency, Washington, DC, USA. Accessed online at  
579 <https://www.epa.gov/tsca-screening-tools/epi-suitetm-estimation-program-interface#citing>.

580 Wang, Q., Guo, M., Yates, S.R., 2006. Degradation kinetics of manure-derived sulfadimethoxine  
581 in amended soil. *J Agric Food Chem* 54, 157-163.

582 Wegst-Uhrich, S.R., Navarro, D.A., Zimmerman, L., Aga, D.S., 2014. Assessing antibiotic  
583 sorption in soil: a literature review and new case studies on sulfonamides and macrolides. *Chem*  
584 *Cent J* 8, 5.

585 Wu, C., Spongberg, A.L., Witter, J.D., 2009. Sorption and biodegradation of selected antibiotics  
586 in biosolids. *J Environ Sci Health A Tox Hazard Subst Environ Eng* 44, 454-461.

587 Yang, Y., Owino, A.A., Gao, Y., Yan, X., Xu, C., Wang, J., 2016. Occurrence, composition and  
588 risk assessment of antibiotics in soils from Kenya, Africa. *Ecotoxicology* 25, 1194-1201.

589 Zhang, G.X., Liu, X.T., Sun, K., Zhao, Y., Lin, C.Y., 2010. Sorption of tetracycline to sediments  
590 and soils: assessing the roles of pH, the presence of cadmium and properties of sediments and  
591 soils. *Front Environ Sci En* 4, 421-429.

592 Zhang, M., Zhang, H., 2010. Thermal Degradation of Chloroteracycline in Animal Manure and  
593 Soil. in: Xu, J., Huang, P.M. (Eds.). *Molecular Environmental Soil Science at the Interfaces in*  
594 *the Earth's Critical Zone*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 229-231.

595 Table 1. Initial ( $C_0$ ) and final (120-day) concentrations of antibiotics ( $\mu\text{g kg}^{-1}$ ) in three different soils amended with raw manure, static  
 596 compost, or turned compost.

Amendment types	Soil types	Sulfamethazine		Tylosin		Chlortetracycline		Pirlimycin		
		Initial	Final	Initial	Final	Initial	Final	Initial	Day 3 <sup>†</sup>	Final
Raw manure	sandy loam	47±9.9	18±3.5	8.4±2.6	0.70±0.07	57±14	17±2.2	4.6±0.64	12±1.4	0.79±0.08
	silt loam	29±4.8	12±2.3	8.6±1.1	0.51±0.03	80±13	19±2.4	10±1.8	19±1.9	0.47±0.21
	silty clay loam	32±7.3	13±3.6	10±2.3	0.87±0.07	46±11	9.3±1.0	13±1.6	26±1.7	0.63±0.05
Static compost	sandy loam	1.6±0.35	1.5±0.03	1.8±0.35	BDL	11±0.38	3.1±0.14	BDL	BDL	BDL
	silt loam	1.1±0.22	1.2±0.05	1.8±0.40	BDL	7.5±0.52	1.8±0.11	BDL	BDL	BDL
	silty clay loam	1.5±0.58	1.4±0.06	1.3±0.20	BDL	8.9±0.50	1.8±0.16	BDL	BDL	BDL
Turned compost	sandy loam	0.80±0.12	0.78±0.38	0.27±0.14	BDL	4.7±1.2	1.1±0.52	BDL	BDL	BDL
	silt loam	0.35±0.03	0.43±0.03	0.63±0.30	BDL	6.4±1.9	2.4±1.5	BDL	BDL	BDL
	silty clay loam	0.35±0.08	0.39±0.08	1.2±0.45	BDL	5.9±0.57	2.1±0.50	BDL	BDL	BDL

597 <sup>†</sup>The initial concentrations of pirlimycin in the raw manure amended soils are the peak concentrations determined at day 3 of the  
 598 microcosm incubation.

599 BDL: concentrations which are below method detection limits (0.13, 0.25, 0.59, and 0.54  $\mu\text{g kg}^{-1}$  for sulfamethazine, tylosin,  
 600 chlortetracycline, and pirlimycin, respectively)

601 Table 2. Dissipation rate constants ( $k$ ) and goodness of curve fitting of different antibiotics in three soils amended with raw manure,  
 602 static compost, or turned compost during the 120-day microcosm incubation study

Amendment types	Soil types	Sulfamethazine			Tylosin			Chlortetracycline			Pirlimycin		
		$k_1$	$k_2$	$R^2$	$k_1$	$k_2$	$R^2$	$k_1$	$k_2$	$R^2$	$k_1$	$k_2$	$R^2$
Raw manure†	sandy loam	0.116±0.029	ND	0.94	0.261±0.054	0.016±0.001	0.96	0.028±0.005	0.005±0.002	0.95	0.085±0.004	0.007±0.001	0.99
	silt loam	0.092±0.017	ND	0.96	0.192±0.038	0.023±0.001	0.98	0.030±0.004	0.006±0.002	0.94	0.104±0.010	0.008±0.001	0.90
	silty clay loam	0.039±0.021	ND	0.94	0.238±0.052	0.016±0.001	0.96	0.028±0.005	0.009±0.002	0.95	0.127±0.001	0.005±0.001	0.99
Static compost‡	sandy loam	ND		NA	NA		0.99	0.012±0.001		0.94	NA		
	silt loam	ND		NA	NA		0.97	0.013±0.001		0.96	NA		
	silty clay loam	ND		NA	NA		0.97	0.014±0.001		0.98	NA		
Turned compost‡	sandy loam	ND		NA	0.011±0.005§		0.85	0.011±0.001		0.96	NA		

	silt loam	ND	NA	0.019±0.011§	0.69	0.007±0.001	0.88	NA
	silty clay loam	ND	NA	0.032±0.008§	0.67	0.008±0.001	0.94	NA

603

604 † Dissipation of antibiotics followed bi-phasic first order kinetics in raw manure-amended soils

605 ‡ Dissipation of antibiotics followed single phase first order kinetics in the compost-amended soils

606 § The dissipation curves of tylosin in the compost-amended soil are fitted to a single phase first kinetic from day 0 to day 57, half of  
607 the method detection limit ( $0.12 \mu\text{g kg}^{-1}$ ] are used to represent the concentrations of tylosin at day 57.

608 ND: no dissipation (k values are close to 0];

609 NA: not available due to below detection limit of pirlimycin in the compost-amended soils.

610 Table 3. The half-lives of antibiotics in soils in this study and literatures

Antibiotics	Initial concentrations ( $\mu\text{g kg}^{-1}$ )	Samples	Half-lives (days)	References
Sulfamethazine	29-47 <sup>†</sup>	manure amended-sandy loam/-silt loam/-silty clay loam	6-18 (1 <sup>st</sup> phase) ND (>120, 2 <sup>nd</sup> phase)	This study
	1.1-1.5 <sup>†</sup>	static compost amended-sandy loam/-silt loam/-silty clay loam	ND (>120)	
	0.35-0.80 <sup>†</sup>	turned compost amended-sandy loam/-silt loam/-silty clay loam	ND (>120)	
	1,000 to 1,000,000 <sup>‡</sup>	silt loam/sandy loam	18.6	(Accinelli et al., 2007)
	500 to 100,000 <sup>‡</sup>	sandy loam	1.3-5.9	(Lertpaitoonpan, 2008)
	500 to 100,000 <sup>‡</sup>	manure amended-sandy loam	1.2-6.6	
	200 <sup>‡</sup>	manure amended-sandy loam/-clay loam	ND (>28)	(Bailey et al., 2016)
	100 <sup>‡</sup>	clay loam soil	24.8	(Pan and Chu, 2016)
Tylosin	8.4-10 <sup>†</sup>	manure amended-sandy loam/-silt loam/-silty clay loam	2.7-3.7 (1 <sup>st</sup> phase) 41-44 (2 <sup>nd</sup> phase)	This study
	1.3-1.8 <sup>†</sup>	static compost amended-sandy loam/-silt loam/-silty clay loam	15-17	

	0.27-1.2†	turned compost amended-sandy loam/-silt loam/-silty clay loam	22-63	
	1142‡	sandy loam	6.1	(Carlson and Mabury, 2006)
	1408‡	manure amended-sandy loam	4.5	
	2000‡	sandy loam	8	(Schlusener and Bester, 2006)
	50000‡	sandy loam	7	(Hu and Coats, 2007)
	50†	manure amended-loamy sand	67	(Halling-Sorensen et al., 2005)
	25†	manure amended-sandy	49	
Chlortetracycline	57-80†	manure amended-sandy loam/-silt loam/-silty clay loam	23-25 (1 <sup>st</sup> phase) 75-144 (2 <sup>nd</sup> phase)	This study
	7.5-11†	static compost amended-sandy loam/-silt loam/-silty clay loam	49-58	
	4.7-6.4†	turned compost amended-sandy loam/-silt loam/-silty clay loam	61-104	
	754‡	sandy loam	21	(Carlson and Mabury, 2006)
	705‡	manure amended-sandy loam	24	
	5000‡	loam	31.9	(Zhang and Zhang, 2010)
	5000‡	manure amended-loam	37.3	
		20000‡	silt loam	5.5



	20-30†	manure amended-loamy sand	25	(Halling-Sorensen et al., 2005)
	20-30†	manure amended-sandy	34	
Pirlimycin	12-26‡	manure amended-sandy loam/-silt loam/-silty clay loam	5.5-8.2 87-142	This study

611 † Anitibotics in naturally excreted form when manure-based amendments are applied to soil

612 ‡ Antibiotics directly spiked into the soil systems

613 Table 4. *P* values of two-way ANOVA of the effect of manure amendment type and soil type on antibiotics dissipation and multiple  
 614 pair comparisons of effect of manure amendment types on antibiotic dissipation

Factors	Sulfamethazine	Tylosin	Chlortetracycline	Pirlimycin
Amendment type	<b>&lt;0.0001</b>	0.53	<b>0.01</b>	NA
Soil Type	0.05	0.36	0.72	0.18
Amendment Type × Soils	0.12	0.11	0.81	NA
Pair comparisons of amendment type	Sulfamethazine	Tylosin	Chlortetracycline	Pirlimycin
Raw manure vs. Static compost	<b>&lt;0.0001</b>	0.95	0.17	NA
Raw manure vs. Turned compost	<b>&lt;0.0001</b>	0.51	<b>0.01</b>	NA
Static compost vs. Turned compost	0.06	0.71	0.44	NA

615 NA: not available due to below detection limit of pirlimycin in the compost-amended soils

616 Figure captions


617

618 Figure 1. Dissipation of sulfamethazine, tylosin, chlortetracycline, and pirlimycin in sandy, silt,  
619 and silty clay loam soils amended with raw manure, static compost, and turned compost. Error  
620 bars represent standard deviations from replicate microcosms (n=3].

621

622 Figure 2. The antibiotic resistant selection potential risk quotient (RQ] values of sulfamethazine,  
623 tylosin, chlortetracycline, and pirlimycin in soils applied with raw manure, static compost, and  
624 turned compost

625 Figure 1.



626

627

