



# *Effect of selected plant species within biodiverse pasture on in vitro fatty acid biohydrogenation and tissue fatty acid composition of lamb*

Article

Accepted Version

Kliem, K. E., Thomson, A. L., Crompton, L. A. and Givens, D. I. (2018) Effect of selected plant species within biodiverse pasture on in vitro fatty acid biohydrogenation and tissue fatty acid composition of lamb. *Animal*. ISSN 1751-732X doi: <https://doi.org/10.1017/S1751731118000265> Available at <http://centaur.reading.ac.uk/75153/>

It is advisable to refer to the publisher's version if you intend to cite from the work.

To link to this article DOI: <http://dx.doi.org/10.1017/S1751731118000265>

Publisher: Cambridge University Press

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

1 **Effect of selected plant species within biodiverse pasture on *in vitro* fatty acid**  
2 **biohydrogenation and tissue fatty acid composition of lamb**

3

4 K. E. Kliem<sup>1</sup>, A. L. Thomson<sup>1</sup>, L. A. Crompton<sup>1</sup>, D. I. Givens<sup>2</sup>

5

6 <sup>1</sup>Animal Dairy and Food Chain Sciences, School of Agriculture, Policy and  
7 Development, University of Reading, Reading, RG6 6AR, United Kingdom.

8 <sup>2</sup>Institute for Food Nutrition and Health, University of Reading, Reading, RG6 6AR,  
9 United Kingdom.

10

11

12 Corresponding author: K. E. Kliem Email: [k.e.kliem@reading.ac.uk](mailto:k.e.kliem@reading.ac.uk)

13

14 Short title: Fatty acid profiles of biodiverse forage species

15

16 **Abstract**

17 The effect of botanical diversity on supply of polyunsaturated fatty acids (PUFA) to  
18 ruminants *in vitro*, and the fatty acid (FA) composition of muscle in lambs was  
19 investigated. Six plant species, commonly grown as part of UK herbal ley mixtures  
20 (*Trifolium pratense*, *Lotus corniculatus*, *Achillea millefolium*, *Centaurea nigra*, *Plantago*  
21 *lanceolata* and *Prunella vulgaris*), were assessed for FA profile, and *in vitro*  
22 biohydrogenation of constituent PUFA, to estimate intestinal supply of PUFA available  
23 for absorption by ruminants. Modelling the *in vitro* data suggested that *L. Corniculatus*  
24 and *P. Vulgaris* had the greatest potential to increase 18:3 n-3 supply to ruminants,  
25 having the highest amounts escaping *in vitro* biohydrogenation . Biodiverse pastures  
26 were established using the six selected species, under-sown in a perennial ryegrass-

27 based sward. Lambs were grazed (~50 days) on biodiverse or control pastures and  
28 the effects on the FA composition of *m. longissimus thoracis* (lean and subcutaneous  
29 fat) and *m. semimembranosus* (lean) were determined. Biodiverse pasture increased  
30 18:2 n-6 and 18:3 n-3 contents of *m. semimembranosus* (+14.8 and +7.2 mg/100g  
31 tissue respectively) and the subcutaneous fat of *m. l. thoracis* (+158 and +166 mg/100g  
32 tissue respectively) relative to feeding a perennial ryegrass pasture. However, there  
33 was no effect on total concentrations of saturated FA in the tissues studied. It was  
34 concluded that enhancing biodiversity had a positive impact on muscle FA profile  
35 reflected by increased levels of total PUFA.

36

37 **Keywords:** Biodiversity; multispecies swards; fatty acids; biohydrogenation; lamb  
38 muscle

39

#### 40 **Implications**

41 The improvement of muscle fatty acid (FA) profile in lambs through increased  
42 polyunsaturated fatty acid (PUFA) concentration achieved in the present study adds to  
43 the growing body of evidence supporting the replacement of monoculture pasture with  
44 biodiverse mixtures. By including a greater proportion of species that were found to  
45 promote PUFA supply to ruminant tissues, such as selfheal and birdsfoot trefoil,  
46 biodiverse seed mixtures could be formulated to optimise the FA profile of resulting  
47 ruminant food products. However, the seemingly low persistence of these species  
48 within a competitive mixed sward remains a challenge to commercial uptake.

49

#### 50 **Introduction**

51 There is increasing interest in low-input biodiverse pasture as a sustainable forage for  
52 grass-based ruminant production systems (Luescher *et al.*, 2014). However, to-date,  
53 little work has focused on the ability of individual plants within biodiverse pastures to  
54 beneficially modify the fatty acid (FA) profile of ruminant food products, with the aim of  
55 increasing mono- and polyunsaturated fatty acids (MUFA, PUFA) and decreasing  
56 saturated fatty acid (SFA) concentrations (Elgersma, 2015). Certain plant species  
57 (Asteraceae, Apiaceae, Rosaceae, Cyperaceae) have been positively correlated with  
58 PUFA in milk (Collomb *et al.*, 2002). In ruminant muscle, biodiverse systems have been  
59 associated with enhanced PUFA concentrations in lambs (Whittington *et al.*, 2006;  
60 Campidonico *et al.*, 2016) and Lourenco *et al.* (2007) reported increased  
61 concentrations of docosahexaenoic acid (DHA; a very long chain n-3 PUFA) in the  
62 intramuscular fat of lambs grazing a biodiverse pasture compared with an intensive  
63 ryegrass pasture. However, few studies have attempted to relate species composition  
64 within a biodiverse sward to the FA supplied by the plants and their potential to alter  
65 the composition of ruminant food products. Identifying which species have the most  
66 potential for improving the resulting FA composition in ruminant products could aid in  
67 designing targetted seed mixtures for this purpose. For example, red clover (*Trifolium*  
68 *pratense*) and certain other perennial forage species contain enhanced levels of  
69 polyphenol oxidase, which can prevent lipolysis and subsequent rumen  
70 biohydrogenation of plant PUFA (Dewhurst *et al.*, 2006; Lee *et al.*, 2014). In addition,  
71 plants containing condensed tannins may protect PUFA from biohydrogenation, and  
72 thus enhance PUFA concentration in ruminant products (Campidonico *et al.*, 2016;  
73 Girard *et al.*, 2016a). However, the degree of PUFA protection can be influenced by  
74 the concentration, chemical structure, and degree of polymerisation of the condensed  
75 tannins, which can vary both within and between plant species (Azuhwi *et al.*, 2013).

76 This may explain why certain tannin containing species appear to affect FA profile to a  
77 greater extent than others. Girard *et al.* (2016b) observed that sainfoin (*Onobrychis*  
78 *viciifolia*) raised PUFA concentration in cheese to a greater extent than birdsfoot trefoil  
79 (*Lotus corniculatus*), with both providing similar amounts of alpha-linolenic acid (18:3  
80 n-3) to the animal. Therefore, the objectives of the present study, were to (i) determine  
81 the impact of a range of species on PUFA biohydrogenation *in vitro*, and (ii) determine  
82 the impact of increasing pasture botanical biodiversity by including selected plant  
83 species, on the FA profile of lamb meat.

84

## 85 **Material and methods**

86 *Experiment 1: In vitro biohydrogenation of selected species*

87 *Sample collection.* Six “candidate” plant species - birdsfoot trefoil (*Lotus corniculatus*),  
88 knapweed (*Centaurea nigra*), ribwort plantain (*Plantago lanceolate*), red clover  
89 (*Trifolium pratense*), selfheal (*Prunella vulgaris*) and yarrow (*Achillea millefolium*) -  
90 were selected from a larger group of species, due to containing relatively high  
91 concentrations of 18:2 n-6 and 18:3 n-3, and being relatively easy to establish (Kliem  
92 *et al.*, 2006). The entire above-ground plant material of each plant was collected from  
93 one of several already established plots on four separate occasions during the growing  
94 season (Kliem *et al.*, 2006), mixed well and transported to the laboratory. Samples  
95 were stored at -20°C before being lyophilised and milled (<1 mm).

96

97 *In vitro biohydrogenation.* To Wheaton flasks (capacity 125 ml), 1.0 g (+/- 0.01 g) of  
98 each freeze-dried and milled sample was accurately weighed in triplicate, and 90 ml of  
99 a reduced anaerobic buffer (Theodorou *et al.*, 1994) added. Flasks were warmed to  
100 39°C prior to inoculation with 10 ml strained bovine rumen fluid collected approximately

101 2 h post-feeding from two lactating dairy cows receiving a total mixed diet comprising  
102 50:50 forage:concentrate (DM basis), with the forage portion being predominantly  
103 maize silage. Flasks were loosely stoppered and vented via a needle. Flasks were  
104 incubated at 39°C with regular mixing by agitation of the bottles. Three flasks per plant  
105 species were removed following 0, 3, 6, 9, 12, 24 and 48 h incubation, flask contents  
106 were frozen at -20°C and then lyophilised. Lyophilised residue was mixed and stored  
107 at -20°C before subsequent FA analysis.

108

109 *Fatty acid analysis.* FA analysis of the whole plant material prior to *in vitro*  
110 biohydrogenation was performed on triplicate sub-samples of freeze-dried plant  
111 material, using a method based on Sukhija and Palmquist (1988) with toluene for  
112 extraction and 2% (v/v) sulphuric acid in methanol for methylation. Resulting FA methyl  
113 esters (FAME) were analysed on a Varian 3400 CX Gas Chromatograph equipped  
114 with a flame-ionization detector, using a temperature programme (Shingfield *et al.*,  
115 2003). Identification of FAME peaks was completed using a known external standard  
116 (GLC463, Nu-Check Prep., MN, USA). Individual FA concentrations were normalised  
117 according to the total lipid content, determined as ether extract (MAFF, 1986). The  
118 contents of individual FA were reported on an oven DM basis following measurement  
119 of the residual DM content of the freeze-dried samples (after oven drying at 100°C for  
120 18 h).

121

122 Biohydrogenation residues were analysed for FA composition using a method based  
123 on Folch *et al.* (1957) and methylated using a bi-methylation method (base-catalysed  
124 then acid-catalysed) derived from Kramer and Zhou (2001). A known amount of  
125 internal standard (Heneicosanoic acid methyl ester, H3265, Sigma-Aldrich, UK) was

126 added prior to methylation in order to quantify FAME. Extracted FAME were analysed  
127 as described above. The FA profiles were expressed as total mg/flask. Results for 18:2  
128 n-6 and 18:3 n-3 were used to calculate the extent of *in vitro* biohydrogenation for  
129 each plant based on the disappearance of 18:2 n-6 and 18:3 n-3.

130

131 *Data analysis.* Flask contents of selected fatty acids were analysed for effects of plant,  
132 time and their interaction by means of the Mixed model in SAS (v9.4, SAS Institute,  
133 Cary, NC, US), which included within sampling time comparison of plant least squares  
134 means (analysed using the PDIFF function). Results were considered significantly  
135 different when  $P < 0.05$ . Curves (constructed using the mean of three flasks over the  
136 entire incubation period from 0 to 48 h) describing the rate and extent of *in vitro*  
137 biohydrogenation (disappearance of 18:2 n-6 and 18:3 n-3) were fitted to the  
138 exponential model of Ørskov and McDonald (1979) using SigmaPlot (Systat Software  
139 Inc., London). Hydrogenation of FA was described by the equation  $P_t = x + ye^{-zt}$ , where  
140  $P_t$  is the amount (mg) of FA present at incubation time  $t$ ,  $x$  is the non-hydrogenated FA  
141 fraction (mg),  $y$  is the hydrogenated fraction (mg) and  $z$  is the fractional rate of  
142 disappearance of  $y$  (/h). Curve parameters were compared as in Boufaïed *et al.*  
143 (2003); effective disappearance (*ED*) and rumen bypass (*BP*) of 18:2 n-6 and 18:3 n-  
144 3 were calculated using a rumen fractional passage rate ( $k$ ) of 0.03/h (Alcaide *et al.*,  
145 2000). This rate describes the passage of small particulate matter in sheep.

146

147 *Experiment 2: Fatty acid profile of lamb*

148 *Plant species and establishment of biodiverse pastures.* The same six species  
149 assessed *in vitro* in experiment 1 were established within a permanent, perennial  
150 ryegrass-based sward at the University of Reading. In the previous five years the sward



151 had been used to graze sheep and had received approximately 100 kg fertiliser  
152 nitrogen/ha/year. The site was divided into ten plots (5 x 2 arrangement; each 60 m x  
153 29 m) allocated in a paired block design to either the biodiverse or a control (no  
154 additional species sown) treatment. Blocking was completed to account for potential  
155 variation in background conditions. The biodiverse plots were power harrowed prior to  
156 under-sowing at ~5 kg seed/ha, twice the recommended seed rate (DEFRA, 2004).  
157 The weight of each species within the seed mixture was as follows: birdsfoot trefoil (19  
158 %), knapweed (24 %), ribwort plantain (32 %), red clover (13 %), selfheal (10 %) and  
159 yarrow (2 %; Emorsgate Seeds, Norfolk, UK). These proportions were used so that the  
160 same number of seeds per g was included of each species. Establishment of the six  
161 'sown' species was completed using 0.75 and 0.25 of the total seed amount (5 kg) in  
162 spring and autumn, respectively. Owing to poor establishment of the biodiverse  
163 pastures a further ~5 kg seed/ha was applied in late autumn. After sowing the  
164 biodiverse plots were rolled and then left undisturbed for at least six weeks. The control  
165 pastures received 100 kg fertiliser nitrogen/ha in the first year but no additional fertiliser  
166 was applied to the biodiverse pastures.

167

168 Immediately prior to the start of the grazing study in the following spring, both the  
169 biodiverse and control plots were assessed for species richness as determined by the  
170 number and abundance of different sown and unsown plant species, and assessment  
171 of contribution to the overall biomass. This was achieved by estimating the number of  
172 different species and percentage ground cover of vascular plant species in 12  
173 randomly positioned 50 x 50 cm quadrats within each plot (areas within 1 m of the  
174 fences were excluded from the sampling). Simultaneously, ten random samples per  
175 plot were obtained by harvesting the above-ground plant matter that were pooled within

176 plots, frozen (-20°C), freeze-dried and milled, and stored at -20°C and subsequently  
177 analysed for fatty acid analysis, as per the process described for whole plants in  
178 Experiment 1.

179

180 *Experimental animals and the grazing study.* Fifty greyface mule x Texel castrated  
181 male lambs from an early lambing flock were weaned in April of the grazing year and  
182 given a forage-based diet until the start of the grazing study in mid-May. The lambs  
183 were weighed prior to the study (mean weight  $\pm$ SEM 26.8 kg  $\pm$  0.39), and five lambs  
184 were randomly allocated to each plot to ensure a similar mean live-weight within each  
185 plot and across the two treatments (26.8 and 26.7 kg for biodiverse and control  
186 pastures, respectively). Lambs had access to water *ad libitum*, and were weighed  
187 weekly, with live-weights recorded. The grazing period continued for a minimum of 50  
188 d (mean  $\pm$  s.e.m. control 64.7  $\pm$  0.93 days, biodiverse 64.3  $\pm$  0.93 days) after which  
189 time animals reaching the target weight of 45 kg or attaining optimum body condition  
190 score by palpation of the loin area were selected for slaughter. A total of three lambs  
191 from each plot were slaughtered. Animals were transported to the University of Bristol  
192 for slaughter, which occurred according to European Union Welfare guidelines. On  
193 arrival animals were stunned by captive bolt followed by abrupt exsanguination.  
194 Carcasses were prepared and graded, and and tissue samples were taken for study  
195 from *musculus longissimus thoracis* and *musculus semimembranosus*, and  
196 subcutaneous fat from above *m. l. thoracis*. Samples were stored frozen at -20°C until  
197 required for FA analysis.

198

199 *Fatty acid analysis.* Prior to analysis tissue samples were partially defrosted at room  
200 temperature for approximately 30 minutes and prepared by cutting into ~ 1 cm<sup>3</sup> pieces,

201 and blended to a homogeneous paste in a food processor within 2 minutes.  
202 Subsequently, FA in samples were extracted using the Folch *et al.* (1957) method  
203 followed by a base-catalysed methylation as described for *in vitro* samples in  
204 Experiment 1. For FA extraction, 2.0 g of each tissue (in duplicate) were homogenised  
205 in chloroform/methanol (2:1, v/v) using an IKA® Ultra-Turrax dispersal tool (IKA®-  
206 Werke GmbH & Co.. Staufen, Germany). After washing the extract with saline solution,  
207 the solvent was removed under vacuum at 40°C using a rotary evaporator and the  
208 remaining lipid extract was re-suspended in hexane. FAME were analysed as outlined  
209 previously, FA contents and profiles were obtained for each sample, and were  
210 expressed as mg/100 g fresh tissue.

211

212 *Data analysis.* Live-weight was analysed using the Mixed procedure of SAS (SAS  
213 version 9.4; SAS Institute), with a model that included fixed effects of time, treatment  
214 and time by treatment interaction (including time as a repeated measurement), and  
215 random effects of plot and lamb within plot. Pasture total lipid, FA content and species  
216 richness were analysed using a two-way ANOVA, with fixed effects of treatment and  
217 block. Tissue FA were analysed using the Mixed procedure of SAS with a model  
218 including fixed effects of treatment, block, and treatment by block interaction. Results  
219 were considered significantly different where  $P < 0.05$ , and tendencies were reported  
220 where  $P$  was between 0.05 and 0.1.

221

## 222 **Results**

### 223 *Experiment 1*

224 Of the six plant species, selfheal contained the highest amount of total FA, and ribwort  
225 plantain the least (Table 1). Yarrow was particularly high in 18:2 n-6, and Selfheal

226 contained the greatest quantity of 18:3 n-3 (Table 1). The effect of the six selected  
227 plant species on *in vitro* flask contents of selected FA are reported in Table 2. There  
228 were effects ( $P<0.001$ ) of plant, time and plant by time interaction for all FA presented  
229 in Table 2. At 0 h incubation, all flasks contained similar amounts of 18:0 ( $P=0.124$ ),  
230 but over time flask contents increased ( $P<0.001$ ). The interaction between plant and  
231 time for 18:0 reflected a lag in 18:0 accumulation for knapweed, and the greatest  
232 ( $P<0.05$ ) 18:0 accumulation at 48 hours for selfheal and birdsfoot trefoil. For *cis*-9 18:1,  
233 at 0 h incubation there was a difference ( $P<0.001$ ) between plants, most probably due  
234 to the high content in selfheal (Table 2). Over time this decreased ( $P<0.001$ ) for all  
235 flasks, but again the rate of disappearance varied between plants, with this being  
236 lowest for red clover after 3 hours of incubation. Flask contents of *trans*-11 18:1 were  
237 similar across all plants at time 0 ( $P=0.542$ ), but over time contents increased  
238 ( $P<0.001$ ) to a peak between 6 and 12 hours before decreasing again. The greatest  
239 amount of *trans*-11 18:1 was measured in flasks containing birdsfoot trefoil.

240

241 There were differences ( $P<0.001$ ) between plants for both 18:2 n-6 and 18:3 n-3, at 0  
242 h incubation. Over time both decreased ( $P<0.001$ ) but at different rates. According to  
243 the disappearance curves, knapweed contained the highest amount of non-  
244 hydrogenatable 18:2 n-6, and selfheal the lowest (Table 3). The effective  
245 disappearance of the hydrogenatable fraction was highest for yarrow, with yarrow and  
246 knapweed containing the highest amount of rumen bypass 18:2 n-6. Selfheal  
247 contained the highest amount of hydrogenatable 18:3 n-3, but had the lowest rate of  
248 18:3 n-3 disappearance of all plants. Due to this the ruminal bypass 18:3 n-3 was  
249 highest for selfheal.

250

251 *Experiment 2*

252 *Pasture botanical composition and lamb performance.* The number of different species  
253 (both sown and unsown) was higher in the biodiverse than the control pastures when  
254 expressed per quadrat ( $P<0.016$ ) and per plot ( $P<0.019$ ). However, birdsfoot trefoil  
255 was not recorded in any of the biodiverse pastures. The mean contribution of the sown  
256 and un-sown (non-grass species) plant species to the overall biomass was 25.4% in  
257 the biodiverse pastures. This contribution was largely comprised of ribwort  
258 plantain. The total lipid and FA composition of the conventional and biodiverse pastures  
259 immediately prior to start of the lamb grazing study is presented in Table 4. There were  
260 no statistically significant ( $P>0.05$ ) differences in the individual FA contents of the two  
261 pasture types. The predominant FA was 18:3 n-3 and accounted for approximately  
262 50% of the total FA.

263

264 Live-weight change of the lambs grazing the conventional and biodiverse pastures is  
265 summarised in Figure 1 and demonstrates an effect of time ( $P<0.001$ ) but no effect  
266 ( $P=0.717$ ) of treatment, with no interaction ( $P=0.773$ ). Overall mean live-weight gains  
267 were 10.2 and 10.0 kg ( $\pm 1.91$  SEM) over the grazing period for conventional and  
268 biodiverse groups, respectively.

269

270 *Fatty acid composition of tissues.* The summary of the amounts of key FA groups in all  
271 three tissues analysed are reported in Table 5 (for full details of FA content, see  
272 Supplementary tables S1, S2 and S3). The total FA content of *m. l. thoracis* was similar  
273 for the lambs grazing the biodiverse and control pastures: mean 1573 and 1648  
274 mg/100 g tissue respectively. The amount of 18:2 n-6 tended to be higher ( $P=0.060$ )  
275 in *m. l. thoracis* from the lambs grazing the biodiverse pasture, however, other

276 differences in FA content and profile were small. A significant block and block x  
277 treatment effect ( $P < 0.05$ ) was recorded for 22:2 *cis*-13, *cis*-16, due to the higher level  
278 of this FA in one of the blocks.

279

280 The total FA content was numerically higher in the *m. semimembranosus* than in the  
281 *m. l. thoracis* tissue (Table 5). A lower ( $P < 0.04$ ) content of *trans*-11 18:1 was found in  
282 *m. semimembranosus* from lambs grazing the biodiverse pasture (Table 5). At the  
283 same time 18:2 n-6 and 18:3 n-3 concentrations were higher (both  $P < 0.02$ ), resulting  
284 in a higher total n-3 and n-6 PUFA content in tissue from lambs grazing biodiverse  
285 pasture. Block x treatment effects were recorded for a number of FA, mainly due to  
286 some blocks having different mean values from the remaining blocks, which magnified  
287 any subtle treatment differences.

288

289 Subcutaneous fat contained 47,261 and 46,723 mg total FA/100 g tissue from lambs  
290 grazed control and biodiverse pastures respectively (Table 5). The content of *trans*-10  
291 18:1, *trans*-12 18:1, 19:0, 18:2 *cis*-9, *cis*-12, 18:3 n-3, 20:3 n-6, 24:0/20:5 n-3, 22:5 n-  
292 3 and total n-3, n-6, and very long chain n-3 PUFA were all higher ( $P < 0.05$ ) in  
293 subcutaneous fat from lambs grazed on biodiverse pasture compared with control  
294 pasture (Supplementary table S3).

295

## 296 **Discussion**

### 297 *In vitro* biohydrogenation

298 It has been suggested (Dewhurst *et al.*, 2001) that the proportion of leaf in the whole  
299 plant DM is an important determinant of FA concentration due to forage lipids being  
300 predominantly of leaf origin (Harfoot, 1981). Differences in leaf:stem ratios between

301 plant species may explain some of the differences in plant FA contents observed. The  
302 *in vitro* biohydrogenation characteristics were similar for the six plant species studied.  
303 Selfheal and birdsfoot trefoil displayed the greatest accumulation of 18:0 but these  
304 plants contained the greatest initial amounts of total PUFA. Rate of disappearance of  
305 18:2 n-6 was similar for all plants apart from knapweed (mean of plants excluding  
306 knapweed, 0.12 mg/h, knapweed 0.09 mg/h). This may indicate that knapweed exerts  
307 some other effect on rumen microbes and/or their enzymes, either by inhibiting the  
308 initial lipolysis or biohydrogenation itself. Indeed, knapweed resulted in a lower  
309 accumulation of both *trans*-11 18:1 and 18:0 which suggests less biohydrogenation.  
310 Kumarasamy *et al.* (2003) found that serotonin conjugates extracted from the seeds of  
311 knapweed had antimicrobial activity. These conjugates may also be present in other  
312 fractions of knapweed and therefore conferring potential antimicrobial effects.

313

314 For 18:3 n-3, both selfheal and birdsfoot trefoil displayed similar high values for ED  
315 compared with the other plants, and yet the accumulation of 18:0 for birdsfoot trefoil  
316 did not increase at the same rate as that of selfheal. This may be due to birdsfoot trefoil  
317 inhibiting the intermediary pathways of biohydrogenation, through, for example, the  
318 presence of condensed tannins. After ingestion some condensed tannins from  
319 birdsfoot trefoil remain free and unbound that may inhibit the extracellular enzyme  
320 action of certain bacteria (Barry and Manley, 1986). Min *et al.* (2002) found that  
321 including birdsfoot trefoil in the diet of sheep decreased the population of the rumen  
322 bacteria *Butyrivibrio proteoclasticus*, which is one of the few bacterial species that  
323 conducts the final step of rumen biohydrogenation of 18:3 n-3 (converting *trans*-11  
324 18:1 to 18:0). However, if condensed tannins affect bacterial biohydrogenation in this  
325 way, no reduction in the initial rate of disappearance of both 18:2 n-6 and 18:3 n-3 was

326 observed in the present study. No estimate of the tannin content or that of other  
327 polyphenols was completed in the present study but it is highlighted as an area of  
328 future study. When a rumen passage rate of 0.03/h (rate at which small particles leave  
329 the rumen of sheep) was applied, the amount of 18:3 n-3 by-passing hydrogenation  
330 was numerically higher for selfheal and birdsfoot trefoil. However, this observation is  
331 likely to reflect the higher initial concentration of 18:3 n-3 in these plants.

332

333 Compared with previous *in vitro* research, red clover did not appear to perform better  
334 than other species in terms of effective disappearance and by-pass of 18:2 n-6 and  
335 18:3 n-3. Van Ranst *et al.* (2013) reported lower lipolysis and biohydrogenation of 18:3  
336 n-3 and 18:2 n-6 with silages containing increasing amounts of red clover, which may  
337 have been due to the presence of polyphenol oxidase (PPO) within the red clover. In  
338 the present study however, red clover was being compared not with ryegrass but with  
339 other species which may have exerted similar biohydrogenation-inhibiting effects. An  
340 *in vivo* study reported higher concentrations of 18:3 n-3 in rumen fluid following the  
341 feeding of a 50:50 grass:red clover silage to lambs, compared with a 100% grass  
342 silage, suggesting PPO as a possible reason (Campidonico *et al.*, 2016). However  
343 there was no difference between the grass:red clover silage and grass:sainfoin silage  
344 treatments, with sainfoin suggested as having a different mechanism of action for  
345 inhibiting biohydrogenation.

346

347 *Using biodiverse pasture to alter the fatty acid profile of tissues*

348 To accelerate the assembly of a species-rich community within grassland, deliberate  
349 under-sowing permanent pasture with selected plant species is required.  
350 Establishment of the biodiverse pastures over approximately 12 months significantly



351 increased the number of different 'sown' and unsown species present in the  
352 experimental plots as compared with the control pastures. On average the biodiverse  
353 pastures were shown to contain an average of 16 different plant species per plot,  
354 compared with 9 for control pastures. However, despite the higher species richness,  
355 the most abundant species, and concomitant contributor to the overall plant biomass,  
356 was ribwort plantain. The abundance of the remaining species was low and therefore  
357 made a substantially smaller contribution to the available biomass available for  
358 grazing. Other studies carried out with the aim of introducing different species to create  
359 biodiverse pastures have been, for example, four years in duration (e.g. Hopkins *et al.*,  
360 1999; Pywell *et al.*, 2002). Therefore a longer period of establishment is required to  
361 enable some slower growing species to proliferate following initial sowing in order to  
362 create a truly biodiverse pasture.

363

364 The *m. semimembranosus* total FA concentrations were lower than those measured  
365 by Whittington *et al.* (2006) comparing different biodiverse systems with a control  
366 pasture. The *m. l. thoracis* total FA concentration was similar to that observed by  
367 Lourenço *et al.* (2007) comparing an intensive ryegrass pasture with an established  
368 biodiverse pasture. There were few differences in the FA profile between the pasture  
369 treatments for *m. l. thoracis*. Lourenço *et al.* (2007) observed a number of differences  
370 in this tissue between animals grazing biodiverse or intensive *lolium perenne*-based  
371 pastures, including a higher 18:2 n-6 resulting in an increased n-6:n-3 ratio for the  
372 biodiverse treatment. In the study of Lourenço *et al.* (2007) the lambs grazed the  
373 biodiverse pastures for a period of 12 weeks (84 days) compared to 50 days in the  
374 present study. This shorter grazing period may reflect some of the differences in the  
375 results recorded although a minimum of 50 days grazing is generally recommended in

376 order to detect differences in muscle phospholipids (Wood, personal communication).  
377 Another reason for the lack of effect in the present study is low establishment of  
378 biodiverse species.

379

380 *M. semimembranosus* contains a higher amount of phospholipids, which have a higher  
381 PUFA content (De Smet *et al.*, 2004). The differences in n-6 PUFA content observed  
382 in the present study for the lambs grazing biodiverse pastures are similar to those  
383 observed by Whittington *et al.* (2006), however these authors did not observe  
384 increases in n-3 FA that were recorded in the present study with the biodiverse  
385 treatment. These differences in intramuscular FA concentrations suggest an increased  
386 availability of both 18:2 n-6 and 18:3 n-3 for tissue incorporation. This may reflect a  
387 reduction in rumen biohydrogenation of these dietary FA for the lambs grazing  
388 biodiverse pastures. The lower *trans*-11 18:1 concentration further illustrates this point,  
389 as this is a key intermediate of the biohydrogenation of both PUFA. There are several  
390 possible explanations for this. Inhibition of initial lipolysis of plant lipids prior to rumen  
391 biohydrogenation (this may have been the mechanism of action observed during *in*  
392 *vitro* biohydrogenation of knapweed and selfheal) may have contributed to this effect.  
393 In addition, inhibition of biohydrogenation prior to the hydrogenation step that  
394 synthesises *trans*-11 18:1 and/or increased rate of passage for animals consuming  
395 biodiverse pasture may have resulted in greater amounts of PUFA escaping rumen  
396 biohydrogenation. However, the mechanism(s) underlying the finding of the present  
397 study are unclear and should be an area of further investigation.

398

399 Total FA concentration was greatest in the subcutaneous fat. Subcutaneous fat total  
400 FA content was lower than that observed by Enser *et al.* (1996; 70,572 mg/100 g

401 tissue) and Lourenço *et al.* (2007; 60,900 – 66,900 mg/100 g tissue). This might reflect  
402 the lighter carcass weight and therefore level of finish, but may also reflect slight  
403 difficulty in separating the subcutaneous fat from muscle. Subcutaneous fat was more  
404 susceptible to dietary change, due to the higher total FA concentration when compared  
405 with muscle. *Trans*-10 18:1 and *trans*-12 18:1 tend to arise following biohydrogenation  
406 of 18:2 n-6 (Jouany *et al.*, 2007). The reason for greater amounts of these FA in  
407 subcutaneous fat of lambs grazing the biodiverse pasture is unclear, especially as we  
408 hypothesise that biohydrogenation of dietary PUFA may have been lower with  
409 biodiverse pastures. It may reflect FA differences for deposition into subcutaneous  
410 tissue. Subcutaneous fat from biodiverse treatment lambs contained higher amounts  
411 of both 18:2 n-6 and 18:3 n-3 than control lambs, as well as a higher very long chain  
412 n-3 FA. There is evidence to suggest that in ruminant animals, 18:2 n-6 is preferentially  
413 deposited in phospholipids compared to 18:3 n-3 (De Smet *et al.*, 2004), which would  
414 suggest a lower n-6:n-3 ratio in subcutaneous fat than the lean muscle tissues.  
415 However in the present study the ratio of n-6:n-3 were similar for all the tissues studied.  
416  
417 Increasing human consumption of 18:3 n-3 has been suggested as a means of  
418 increasing synthesis of very long chain (VLC) n-3 FA through tissue elongation and  
419 desaturation. Burdge and Calder (2005) concluded that due to poor efficiency of  
420 conversion, 18:3 n-3 appears to be a limited source of VLC n-3 FA in humans, and  
421 consumption of preformed VLC n-3 FA is a more efficient means of attaining  
422 recommended intake levels. The efficiency of conversion of 18:3 n-3 to VLC n-3 FA in  
423 ruminant meat has not been measured, but increasing 18:3 n-3 consumption by  
424 ruminants has lead to increased amounts of VLC n-3 FA in lean tissues (Scollan *et al.*,  
425 2001; Wachira *et al.*, 2002). In the present study, the only tissue to display an increase

426 in VLC n-3 FA when lambs grazed biodiverse pasture was subcutaneous fat, which is  
427 likely to be consumed in variable amounts, according to consumer preference,  
428 alongside muscle tissue.

429

430 In conclusion, the results of the present study suggest that it is possible to manipulate  
431 the FA concentration and profile of muscle and subcutaneous fat in lamb by grazing  
432 biodiverse pastures. Grazing lambs on the biodiverse pastures established within our  
433 project increased overall PUFA content (~30 mg/100 g tissue) of lamb muscle. The  
434 three tissues analysed had varying responses to diet reflecting the presence of  
435 different lipid classes in each of the tissues. Differences reported from the *in vivo* study  
436 may have been more pronounced if the biodiverse species had established at the  
437 expected density, especially as the more promising species from the *in vitro* study were  
438 not present within the *in vivo* study pastures.

439

#### 440 **Acknowledgements**

441 This study was supported by the Rural Economy and Land Use programme (joint  
442 collaboration between ESRC, BBSRC and NERC), project “Implications of a nutrition  
443 driven food policy for land use and the rural environment” (RES-224-25-0073). The  
444 authors gratefully acknowledge staff at the Centre for Dairy Research Meat and Growth  
445 Research Unit for their technical contributions and care of the experimental animals.

446

#### 447 **References**

448 Alcaide EM, García AIM, and Aguilera JF, 2000. A comparative study of nutrient digestibility,  
449 kinetics of degradation and passage and rumen fermentation pattern in goats and sheep  
450 offered good quality diets. *Livestock Production Science* 64, 215 – 223.

451 Azuhwi BN, Boller B, Dohme-Meier F, Hess HD, Kreuzer M, Stringano E and Mueller-  
452 Harvey I, 2013. Exploring variation in proanthocyanidin composition and content of  
453 sainfoin (*Onobrychis viciifolia*). *Journal of the Science of Food and Agriculture* 93, 2102-  
454 2109.

455 Barry TN, and Manley TR, 1986. Interrelationships between the concentrations of total  
456 condensed tannin, free condensed tannin and lignin in lotus sp. and other possible  
457 consequences in ruminant nutrition. *Journal of the Science of Food and Agriculture* 37, 248  
458 – 254.

459 Boufaïed H, Chouinard PY, Tremblay GF, Petit HV, Michaud R, and Bélanger G, 2003. Fatty  
460 acids in forages. II. In vitro ruminal biohydrogenation of linolenic and linoleic acids from  
461 timothy. *Canadian Journal of Animal Science* 83, 513 – 522.

462 Burdge GC, and Calder PC, 2005. Alpha-linolenic acid metabolism in humans: the effects of  
463 gender and age on conversion to longer-chain polyunsaturated fatty acids. *European*  
464 *Journal of Lipid Science and Technology* 107, 426 – 439.

465 Campidonico L, Toral PG, Priolo A, Luciano G, Valenti B, Hervas G, Frutos P, Copani G,  
466 Ginane C and Niderkorn V, 2016. Fatty acid composition of ruminal digesta and  
467 longissimus muscle from lambs fed silage mixtures including red clover, sainfoin, and  
468 timothy. *Journal of Animal Science* 94, 1550-1560.

469 Collomb M, Bütikofer U, Sieber R, Jeangros B, and Bosset JO, 2002. Correlation between fatty  
470 acids in cows' milk fat produced in the lowlands, mountains and highlands of Switzerland  
471 and botanical composition of the fodder. *International Dairy Journal* 12, 661 – 666.

472 DEFRA (2004) Sward enhancement: diversifying grassland by oversowing and slot seeding.  
473 Rural Development Service Technical Advice Note 29.

474 De Smet S, Raes K, and Demeyer D, 2004. Meat fatty acid composition as affected by fatness  
475 and genetic factors: a review. *Animal Research* 53, 81 – 98.

476 Dewhurst RJ, Scollan ND, Youell SJ, Tweed JKS, and Humphreys MO, 2001. Influence of  
477 species, cutting date and cutting interval on the fatty acid composition of grasses. *Grass*  
478 *and Forage Science* 56, 68 – 74.

479 Dewhurst RJ, Shingfield KJ, Lee MRF, and Scollan ND, 2006. Increasing the concentrations  
480 of beneficial polyunsaturated fatty acids in milk produced by dairy cows in high-forage  
481 systems. *Animal Feed Science and Technology*, 131, 168-206.

482 Elgersma A, 2015. Grazing increases the unsaturated fatty acid concentration of milk from  
483 grass-fed cows: A review of the contributing factors, challenges and future perspectives.  
484 *European Journal of Lipid Science and Technology* 117, 1345-1369.

485 Enser M, Hallett K, Hewitt B, Fursey GAJ, and Wood JD, 1996. Fatty acid content and  
486 composition of English beef, lamb and pork at retail. *Meat Science* 42, 443 – 456.

487 Folch J, Lees M, and Stanley GH, 1957. A simple method for the isolation and purification of  
488 total lipids from animal tissues. *Journal of Biological Chemistry* 226, 497 – 509.

489 Girard M, Dohme-Meier F, Silacci P, Kragten SA, Kreuzer M and Bee G, 2016a. Forage  
490 legumes rich in condensed tannins may increase n-3 fatty acid levels and sensory quality  
491 of lamb meat. *Journal of the Science of Food and Agriculture* 96, 1923-1933.

492 Girard M, Dohme-Meier F, Wechsler D, Goy D, Kreuzer M and Bee G, 2016b. Ability of 3  
493 tanniferous forage legumes to modify quality of milk and Gruyere-type cheese. *Journal of*  
494 *Dairy Science* 99, 205-220.

495 Harfoot CG, 1981. Lipid metabolism in the rumen. In *Lipid Metabolism in Ruminant Animals*,  
496 pp 21 – 55 [W. W. Christie, editor]. Oxford, UK: Pergamon Press,

497 Hopkins A, Pywell RF, Peel S, Johnson RH, and Bowling PJ. 1999. Enhancement of botanical  
498 diversity of permanent grassland and impact on hay production in Environmentally Sensitive  
499 Areas in the UK. *Grass and Forage Science* 54, 163 – 173.

500 Jouany JP, Lassalas B, Doreau M, and Glasser F, 2007. Dynamic features of the rumen  
501 metabolism of linoleic acid, linolenic acid and linseed oil measured in vitro. *Lipids* 42, 351 –  
502 360.

503 Kliem KE, Deaville ER, Morgan R and Givens DI, 2006. Do biodiverse pastures have the  
504 potential to improve the fatty acid profile of ruminant meat? *Proceedings of the 8th British*  
505 *Grassland Society Research Conference, Cirencester, UK*, pp81.

506 Kramer JKG, and Zhou J, 2001. Conjugated linoleic acid and octadecenoic acids: extraction  
507 and isolation of lipids. *European Journal of Lipid Science and Technology* 103, 594 – 600.

508 Kumarasamy Y, Middleton M, Reid RG, Nahar L, and Sarker SD, 2003. Biological activity of  
509 serotonin conjugates from the seeds of *Centaurea nigra*. *Fitoterapia* 74, 609 – 612.

510 Lee MRF, Theobald VJ, Gordon N, Leyland M, Tweed JKS, Fychan R and Scollan, ND,  
511 2014. The effect of high polyphenol oxidase grass silage on metabolism of  
512 polyunsaturated fatty acids and nitrogen across the rumen of beef steers. *Journal of*  
513 *Animal Science* 92, 5076-5087.

514 Lourenço M, Van Ranst G, De Smet S, Raes K, and Fievez V, 2007. Effect of grazing pastures  
515 with different botanical composition by lambs on rumen fatty acid metabolism and fatty acid  
516 pattern of longissimus muscle and subcutaneous fat. *Animal* 1, 537 – 545.

517 Luescher A, Mueller-Harvey I, Soussana JF, Rees RM and Peyraud JL, 2014. Potential of  
518 legume-based grassland-livestock systems in Europe: a review. *Grass and Forage*  
519 *Science* 69, 206-228.

520 MAFF 1986. *The Analysis of Agricultural Materials*. Reference Book 427. HMSO London

521 Min BR, Barry TN, Attwood GT, and McNabb WC, 2002. The effect of condensed tannins from  
522 *Lotus corniculatus* on the proteolytic activities and growth of rumen bacteria. *Journal of*  
523 *Animal Science* 80 (Supplement 1), 1602.

524 Ørskov ER, and McDonald I, 1979. The estimation of protein degradability in the rumen from  
525 incubation measurements weighted according to rate of passage. *Journal of Agricultural*  
526 *Science* 92, 499 – 503.

527 Pywell RF, Bullock JM, Hopkins A, Walker KJ, Sparks TH, Burke MJW, and Peel S, 2002.  
528 Restoration of species-rich grassland on arable land: assessing the limiting processes using  
529 a multi-site experiment. *Journal of Applied Ecology* 39, 294 – 309.

530 Scollan ND, Choi NJ, Kurt E, Fisher AV, Enser M, Wood JD, 2001. Manipulating the fatty acid  
531 composition of muscle and adipose tissue in beef cattle. *British Journal of Nutrition* 85: 115  
532 – 124.

533 Shingfield KJ, Ahvenjärvi S, Toivonen V, Ärölä A, Nurmela KVV, Huhtanen P, and Griinari JM,  
534 2003. Effect of dietary fish oil on biohydrogenation of fatty acids and milk fatty acid content  
535 in cows. *Animal Science* 77, 165 - 179.

536 Sukhija PS and Palmquist DL, 1988. Rapid method for determination of total fatty acid content  
537 and composition of feedstuffs and feces. *Journal of Agricultural and Food Chemistry* 36,  
538 1202 – 1206.

539 Theodorou MK, Williams BA, Dhanoa MS, McAllan AB, France J 1994. A simple gas  
540 production method to determine the fermentation kinetics of ruminant feeds. *Animal Feed*  
541 *Science and Technology* 48, 185-197.

542 Van Ranst G, Vandewalle M, Gadeyne F, De Riek J and Fievez V 2013. Lipid Metabolism in  
543 mixtures of red clover (*Trifolium repens*) and perennial ryegrass (*Lolium perenne*) in lab  
544 scale silages and in vitro rumen incubations. *Animal* 7, 1454-1463.

545 Wachira AM, Sinclair LA, Wilkinson RG, Enser M, Wood JD, and Fisher AV 2002. Effects of  
546 dietary fat source and breed on the carcass composition, n-3 polyunsaturated fatty acid and  
547 conjugated linoleic acid content of sheep meat and adipose tissue. *British Journal of*  
548 *Nutrition* 88, 697 – 709.

549 Whittington FM, Dunn R, Nute GR, Richardson RI, and Wood JD, 2006. Effect of pasture type  
550 on lamb product quality. In *New developments in sheepmeat quality*. 9<sup>th</sup> Annual Langford  
551 Food Industry Conference, University of Bristol, UK, pp 27 – 31

552



553 **Table 1** Selected fatty acid content of six plants selected for in vitro biohydrogenation  
 554 incubations (mixture of four different sampling times; mg/g dry matter)

Fatty acid	Plant <sup>1</sup>					
	BT	K	RP	RC	S	Y
16:0	5.27	4.83	4.20	4.13	5.16	4.25
18:0	0.62	0.59	0.58	0.77	1.01	0.45
18:1 <i>cis</i> -9	1.34	1.75	2.28	1.47	5.69	2.44
18:2 <i>cis</i> -9, <i>cis</i> -12	4.85	6.84	5.25	4.78	5.64	8.95
18:3 n-3	8.94	6.28	5.70	4.59	11.9	4.94
Total 18:2 <i>cis</i> -9, <i>cis</i> -12 + 18:3 n-3	13.8	13.1	11.0	9.4	17.5	11.4
Total fatty acids	26.6	24.8	21.7	19.8	32.9	24.7

555 <sup>1</sup> Where BT – Birdsfoot trefoil; K – Knapweed; RP – Ribwort plantain; RC – Red clover; S –  
 556 Selfheal; Y – Yarrow.

557

558

559

560

561

562

563

564

565

566

567

568

569 **Table 2** Flask content (mg) of selected fatty acids over a 48 h *in vitro* incubation.

Fatty acid	Time (h)	Plant <sup>1</sup>						s.e.m.	P <sup>2</sup> (plant)
		BT	K	RP	RC	S	Y		
18:0	0	22.4	21.4	21.9	21.3	22.4	20.6	0.52	0.124
	3	23.0 <sup>b</sup>	22.7 <sup>b</sup>	23.4 <sup>b</sup>	23.9 <sup>ab</sup>	25.3 <sup>a</sup>	24.0 <sup>ab</sup>		0.013
	6	26.9 <sup>bc</sup>	24.6 <sup>d</sup>	25.4 <sup>cd</sup>	25.5 <sup>cd</sup>	29.0 <sup>a</sup>	27.1 <sup>b</sup>		<0.001
	9	29.6 <sup>b</sup>	26.2 <sup>d</sup>	27.4 <sup>cd</sup>	27.6 <sup>cd</sup>	32.1 <sup>a</sup>	28.0 <sup>c</sup>		<0.001
	12	29.8 <sup>b</sup>	26.3 <sup>d</sup>	28.3 <sup>c</sup>	28.0 <sup>c</sup>	33.4 <sup>a</sup>	29.3 <sup>bc</sup>		<0.001
	24	32.2 <sup>b</sup>	30.3 <sup>cd</sup>	29.5 <sup>cd</sup>	29.3 <sup>d</sup>	36.3 <sup>a</sup>	30.5 <sup>c</sup>		<0.001
	48	35.3 <sup>b</sup>	30.7 <sup>d</sup>	31.8 <sup>cd</sup>	31.5 <sup>cd</sup>	37.4 <sup>a</sup>	32.3 <sup>c</sup>		<0.001
18:1 <i>cis</i> -9	0	3.04 <sup>c</sup>	3.05 <sup>c</sup>	3.70 <sup>b</sup>	2.91 <sup>c</sup>	6.96 <sup>a</sup>	3.73 <sup>b</sup>	0.068	<0.001
	3	2.46 <sup>c</sup>	2.59 <sup>c</sup>	3.08 <sup>b</sup>	2.61 <sup>c</sup>	5.63 <sup>a</sup>	3.19 <sup>b</sup>		<0.001
	6	2.39 <sup>c</sup>	2.39 <sup>c</sup>	2.89 <sup>b</sup>	2.31 <sup>c</sup>	4.67 <sup>a</sup>	2.82 <sup>b</sup>		<0.001
	9	2.20 <sup>d</sup>	2.23 <sup>d</sup>	2.80 <sup>b</sup>	2.24 <sup>d</sup>	4.14 <sup>a</sup>	2.53 <sup>c</sup>		<0.001
	12	1.88 <sup>d</sup>	1.87 <sup>d</sup>	2.56 <sup>b</sup>	1.98 <sup>cd</sup>	3.31 <sup>a</sup>	2.14 <sup>c</sup>		<0.001
	24	1.33 <sup>d</sup>	1.32 <sup>d</sup>	1.81 <sup>b</sup>	1.39 <sup>cd</sup>	2.26 <sup>a</sup>	1.55 <sup>c</sup>		<0.001
	48	0.91 <sup>d</sup>	1.03 <sup>cd</sup>	1.23 <sup>bc</sup>	1.05 <sup>cd</sup>	1.73 <sup>a</sup>	1.16 <sup>bc</sup>		<0.001
18:1 <i>trans</i> -11	0	1.62	1.70	1.67	1.72	1.79	1.79	0.074	0.542
	3	2.47 <sup>b</sup>	3.03 <sup>a</sup>	2.47 <sup>b</sup>	2.51 <sup>b</sup>	2.52 <sup>b</sup>	3.02 <sup>a</sup>		<0.001
	6	3.69 <sup>a</sup>	3.06 <sup>c</sup>	2.63 <sup>d</sup>	3.05 <sup>c</sup>	2.95 <sup>c</sup>	3.29 <sup>b</sup>		<0.001
	9	3.91 <sup>a</sup>	3.28 <sup>bc</sup>	2.61 <sup>d</sup>	3.15 <sup>c</sup>	3.44 <sup>b</sup>	3.33 <sup>bc</sup>		<0.001
	12	3.78 <sup>a</sup>	3.04 <sup>c</sup>	2.53 <sup>d</sup>	2.95 <sup>c</sup>	3.40 <sup>b</sup>	3.14 <sup>c</sup>		<0.001
	24	3.00 <sup>b</sup>	2.62 <sup>d</sup>	2.47 <sup>d</sup>	2.59 <sup>d</sup>	3.41 <sup>a</sup>	2.89 <sup>c</sup>		<0.001
	48	2.91 <sup>ab</sup>	2.45 <sup>cd</sup>	2.36 <sup>d</sup>	2.57 <sup>c</sup>	3.11 <sup>a</sup>	2.87 <sup>b</sup>		<0.001
18:2 <i>cis</i> -9, <i>cis</i> -12	0	5.57 <sup>c</sup>	6.03 <sup>b</sup>	4.16 <sup>d</sup>	4.99 <sup>d</sup>	5.75 <sup>c</sup>	7.94 <sup>a</sup>	0.078	<0.001
	3	3.86 <sup>d</sup>	4.47 <sup>b</sup>	3.54 <sup>e</sup>	3.57 <sup>e</sup>	4.24 <sup>c</sup>	5.57 <sup>a</sup>		<0.001
	6	3.13 <sup>bc</sup>	3.89 <sup>b</sup>	2.94 <sup>cd</sup>	2.82 <sup>d</sup>	3.22 <sup>b</sup>	4.47 <sup>a</sup>		<0.001
	9	2.51 <sup>b</sup>	3.53 <sup>a</sup>	2.63 <sup>b</sup>	2.47 <sup>b</sup>	2.60 <sup>b</sup>	3.51 <sup>a</sup>		<0.001
	12	1.86 <sup>c</sup>	2.90 <sup>a</sup>	2.17 <sup>b</sup>	2.01 <sup>bc</sup>	1.89 <sup>c</sup>	2.81 <sup>a</sup>		<0.001
	24	1.26 <sup>bc</sup>	2.05 <sup>a</sup>	1.34 <sup>bc</sup>	1.39 <sup>b</sup>	1.13 <sup>c</sup>	1.87 <sup>a</sup>		0.002
	48	0.87 <sup>BC</sup>	1.54 <sup>A</sup>	0.96 <sup>BC</sup>	1.08 <sup>B</sup>	0.85 <sup>C</sup>	1.34 <sup>A</sup>		0.055
18:3 n-3	0	7.60 <sup>b</sup>	3.36 <sup>d</sup>	3.18 <sup>d</sup>	3.75 <sup>c</sup>	9.28 <sup>a</sup>	3.33 <sup>d</sup>	0.075	<0.001
	3	4.18 <sup>b</sup>	1.73 <sup>d</sup>	1.43 <sup>e</sup>	1.98 <sup>c</sup>	6.27 <sup>a</sup>	1.45 <sup>e</sup>		<0.001

6	2.67 <sup>b</sup>	1.39 <sup>cd</sup>	1.19 <sup>de</sup>	1.41 <sup>c</sup>	4.18 <sup>a</sup>	1.12 <sup>e</sup>	<0.001
9	1.97 <sup>b</sup>	1.24 <sup>c</sup>	1.05 <sup>cd</sup>	1.12 <sup>c</sup>	3.00 <sup>a</sup>	0.89 <sup>d</sup>	<0.001
12	1.32 <sup>b</sup>	1.00 <sup>c</sup>	0.89 <sup>cd</sup>	0.87 <sup>cd</sup>	1.83 <sup>a</sup>	0.73 <sup>d</sup>	<0.001
24	0.84 <sup>a</sup>	0.79 <sup>ab</sup>	0.60 <sup>bc</sup>	0.61 <sup>bc</sup>	0.95 <sup>a</sup>	0.56 <sup>c</sup>	<0.001
48	0.60 <sup>ab</sup>	0.63 <sup>ab</sup>	0.47 <sup>b</sup>	0.52 <sup>b</sup>	0.76 <sup>a</sup>	0.48 <sup>b</sup>	<0.001

570

571 <sup>1</sup> Where BT – Birdsfoot trefoil; K – Knapweed; RP – Ribwort plantain; RC – Red clover; S – Selfheal; Y – Yarrow.

572 <sup>2</sup> Significance of the effect of plant within sampling time. There were effects ( $P<0.001$ ) of plant, time and plant by time interaction for all fatty  
573 acids presented.

574 Values within rows with different superscripts are significantly different ( $P<0.050$ )

575

576

577

578

579

580

581 **Table 3.** Curve fit parameters for the disappearance of 18:2 n-6 and 18:3 n-3 over time.

	Plant <sup>1</sup>					
	BT	K	RP	RC	S	Y
18:2 <i>cis</i> -9, <i>cis</i> -12						
$x^2$	0.94	1.58	1.04	1.16	0.84	1.45
$y^2$	4.55	4.29	3.95	3.75	4.91	6.39
$z^2$	0.13	0.09	0.11	0.13	0.12	0.13
Curve fit <sup>3</sup>	0.992	0.982	0.976	0.991	0.998	0.995
ED <sup>4</sup>	3.67	3.22	3.12	3.03	3.93	5.18
BP <sup>5</sup>	1.81	2.64	1.86	1.87	1.81	2.66
18:3 n-3						
$x$	0.75	0.81	0.68	0.62	0.70	0.61
$y$	6.78	2.49	2.45	3.08	8.64	2.68
$z$	0.21	0.26	0.30	0.23	0.15	0.33
Curve fit	0.996	0.961	0.944	0.987	0.998	0.975
ED	5.94	2.23	2.22	2.73	7.22	2.46
BP	1.60	1.06	0.90	0.97	2.12	0.84

582 <sup>1</sup> Where BT – Birdsfoot trefoil; K – Knapweed; RP – Ribwort plantain; RC – Red clover; S –

583 Selfheal; Y – Yarrow.

584 <sup>2</sup> using the equation  $P_t = x + ye^{-zt}$ , where  $P_t$  is the amount (mg) of 18:2 n-6 or 18:3 n-3 present

585 in the flasks at time  $t$ ,  $x$  is the non-hydrogenatable fraction (mg),  $y$  is the hydrogenatable

586 fraction (mg),  $z$  is the rate of disappearance of fraction  $y$  (/h), and  $t$  is incubation time (h;

587 Ørskov & McDonald, 1979)

588 <sup>3</sup> R-squared value for the curve fit

589 <sup>4</sup> ED - Effective disappearance (mg/g DM) of 18:2 n-6 or 18:3 n-3 using a ruminal rate of

590 passage (k) of 0.03 (Alcaide *et al.*, 2000)

591 <sup>5</sup> BP – Potential ruminal bypass (mg/g DM) of 18:2 n-6 or 18:3 n-3.

592

593

594

595

596

597

598

599

600 **Table 4** Mean fatty acid contents (mg/g dry matter) of the control and biodiverse pastures prior  
601 to the commencement of the lamb grazing study.

Fatty acid	Pasture type		s.e.m.	<i>P</i> <sup>1</sup>
	Control	Biodiverse		
16:0	4.68	4.11	0.295	0.216
18:0	0.33	0.29	0.022	0.260
18:1 total	0.80	0.63	0.065	0.099
18:2 <i>cis</i> -9, <i>cis</i> -12	4.02	3.95	0.198	0.793
18:3 n-3	17.8	15.3	0.90	0.094
Total lipid	30.1	26.4	1.67	0.156

602

603 <sup>1</sup> Significance of the effect of pasture type

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621 **Table 5.** Fatty acid composition (mg/100g tissue) of tissues from lambs grazing control and biodiverse pasture.

	Forage type		SEM	<i>P</i> <sup>1</sup>		
	Conventional	Biodiverse		Treatment	Block	Treatment x block
<i>M. Longissimus Thoracis</i>						
18:1 <i>trans</i> -11	45.7	39.6	5.23	0.421	0.306	0.457
18:2 <i>cis</i> -9, <i>cis</i> -12	58.0	72.7	5.18	0.060	0.358	0.426
18:3 n-3	21.8	27.3	2.35	0.112	0.470	0.334
Total fatty acids	1573	1648	168.1	0.758	0.600	0.189
Total SFA <sup>2</sup>	748	776	81.9	0.808	0.650	0.197
Total <i>cis</i> -MUFA <sup>3</sup>	569	592	65.5	0.799	0.585	0.166
Total <i>trans</i> -MUFA <sup>4</sup>	76.2	73.6	8.40	0.835	0.386	0.361
n-3 PUFA <sup>5</sup>	37.7	44.5	3.24	0.155	0.495	0.389
n-6 PUFA <sup>6</sup>	92.5	111	7.10	0.078	0.349	0.399
Total PUFA <sup>7</sup>	130	156	10.2	0.094	0.398	0.389
Total CLA <sup>8</sup>	23.0	21.5	2.86	0.716	0.502	0.502
n-6:n-3	2.5	2.5	0.07	0.511	0.278	0.651
VLC n-3 <sup>9</sup>	21.1	22.8	1.37	0.389	0.504	0.682
<i>M. Semimembranosus</i>						
18:1 <i>trans</i> -11	75.4	61.5	4.40	0.037	0.187	0.018
18:2 <i>cis</i> -9, <i>cis</i> -12	70.1	84.9	2.90	0.002	0.201	0.099
18:3 n-3	28.4	35.6	2.01	0.020	0.286	0.250
Total fatty acids	2416	2315	147.5	0.634	0.695	0.058
Total SFA	1171	1119	76.4	0.636	0.798	0.079
Total <i>cis</i> -MUFA	865	819	55.3	0.557	0.648	0.048
Total <i>trans</i> -MUFA	134	115	9.4	0.170	0.311	0.100
n-3 PUFA	47.2	55.2	2.42	0.032	0.252	0.206
n-6 PUFA	111	128	4.2	0.010	0.220	0.104
Total PUFA	158	183	6.2	0.012	0.302	0.110
Total CLA	39.0	32.9	2.88	0.150	0.375	0.111
n-6:n-3	2.4	2.3	0.08	0.748	0.078	0.954
VLC n-3	24.4	25.6	0.68	0.254	0.204	0.204
Sub-cutaneous fat						
18:1 <i>trans</i> -11	1638	1494	81.7	0.228	0.077	0.615

18:2 <i>cis</i> -9, <i>cis</i> -12	707	865	37.1	0.007	0.059	0.532
18:3 n-3	344	510	30.6	0.002	0.169	0.777
Total fatty acids	47261	46723	1368.6	0.784	0.165	0.036
Total SFA	23260	23378	808.0	0.919	0.894	0.138
Total <i>cis</i> -MUFA	17681	16744	866.5	0.454	0.177	0.065
Total <i>trans</i> -MUFA	2950	2923	87.7	0.826	0.006	0.367
n-3 PUFA	445	631	32.8	0.001	0.207	0.834
n-6 PUFA	939	1126	51.6	0.018	0.071	0.598
Total PUFA	1384	1757	72.9	0.002	0.096	0.762
Total CLA	969	840	64.0	0.170	0.073	0.103
n-6:n-3	2.1	1.9	0.11	0.086	0.186	0.672
VLC n-3	111	135	6.1	0.011	0.985	0.913

622 <sup>1</sup> Significance of the effect of; T - treatment; B - block; T\*B, treatment\*block interaction

623 <sup>2</sup> SFA - saturated fatty acids. Sum of 12:0, 13:0, 14:0, 15:0, 16:0, 17:0, 18:0, 18:0, 19:0, 20:0, 22:0, 24:0.

624 <sup>3</sup> MUFA - mono-unsaturated fatty acids. Sum of *cis*-9 12:1, *cis*-9 14:1, *cis*-9 15:1, *cis*-9 16:1, *cis*-10 17:1, *cis*-9 18:1, *cis*-11 20:1, *cis*-13 22:1, *cis*-  
625 15 24:1

626 <sup>4</sup> Sum of *trans*-9 16:1, *trans*-6-8 18:1, *trans*-9 18:1, *trans*-10 18:1, *trans*-11 18:1, *trans*-12 18:1, *trans*-13-14 18:1

627 <sup>5</sup> PUFA – polyunsaturated fatty acids. Sum of 18:3 n-3, 20:5 n-3, 22:3 n-3, 22:5 n-3, 22:6 n-3

628 <sup>6</sup> Sum of *trans*-9, *trans*-12 18:2, *cis*-9, *cis*-12 18:2, 20:2 n-6, 20:3 n-6, 22:2 n-6, 22:4 n-6.

629 <sup>7</sup> Sum of n-3 and n-6 PUFA.

630 <sup>8</sup> CLA – conjugated linoleic acid

631 <sup>9</sup> VLC – very long chain

632

633 **Figure captions**

634

635 **Figure 1.** The mean liveweight of lambs grazing either a control or biodiverse pasture over a  
636 60 d study period. Mixed model analysis concluded an effect ( $P<0.001$ ) of time but no effect of  
637 treatment ( $P=0.717$ ) or time by treatment interaction ( $P=0.773$ ).

638