

University of New England
DUNE: DigitalUNE

Environmental Studies Faculty Publications

Environmental Studies Department

2-2009

The Functions And Values Of Fringing Salt Marshes In Northern New England, USA

Pamela A. Morgan

University of New England, pmorgan@une.edu

David M. Burdick

University of New Hampshire

Frederick T. Short

University of New Hampshire

Follow this and additional works at: http://dune.une.edu/env_facpubs

 Part of the [Environmental Health and Protection Commons](#), [Environmental Indicators and Impact Assessment Commons](#), [Environmental Monitoring Commons](#), [Marine Biology Commons](#), and the [Systems Biology Commons](#)

Recommended Citation

Morgan, Pamela A.; Burdick, David M.; and Short, Frederick T., "The Functions And Values Of Fringing Salt Marshes In Northern New England, USA" (2009). *Environmental Studies Faculty Publications*. Paper 4.

http://dune.une.edu/env_facpubs/4

This Article is brought to you for free and open access by the Environmental Studies Department at DUNE: DigitalUNE. It has been accepted for inclusion in Environmental Studies Faculty Publications by an authorized administrator of DUNE: DigitalUNE. For more information, please contact bkenyon@une.edu.

Left running head: P. Morgan et al.

Right running head: Fringing salt marshes

The Functions and Values of Fringing Salt Marshes in Northern New England, USA

Pamela A. Morgan^{1,*}, David M. Burdick² and Frederick T. Short²

¹*Department of Environmental Studies, University of New England, Biddeford, Maine, 04005, USA*

²*Jackson Estuarine Laboratory, University of New Hampshire, 85 Adams Point Road, Durham, New Hampshire, 03824, USA*

**Corresponding author; tele: 207 602 2227; fax: 207 602 5926; e-mail: pmorgan@une.edu*

1 ABSTRACT:

2 Although large salt marshes of the northeastern United States have been studied
3 extensively, very little is known about the smaller, fringing marshes in this area, despite the fact
4 that they are a common habitat type. We compared the functions and values of five fringing salt
5 marshes (FM) to those of five meadow marshes (MM) along the southern Maine/New Hampshire
6 coast. Specifically we compared their primary production, soil organic matter content, plant
7 diversity, sediment trapping ability and wave dampening properties. Also explored were the
8 relationships between these functions and several physical characteristics at each site, including
9 soil salinity, percent surface slope, elevation and size.

10 Differences between fringing salt marshes and meadow marshes included their soil
11 organic matter content (12% FM; 42% MM), plant species richness (8.8 FM; 15.6 MM) and
12 density of plant species in 1m² samples (2.8 FM; 4.5 MM). Although our results suggest that
13 fringing marshes trap more sediment per unit area than meadow marshes, this difference was not
14 significant, probably due to the great variability among sites. Traps located 1 m from the edge
15 trapped an average of 21.6 ±18.6 g m⁻² d⁻¹ (FM) and 3.2 ±0.8 g m⁻² d⁻¹ (MM), and traps randomly
16 distributed on the marsh surface trapped an average of 1.6 ±0.7 g m⁻² d⁻¹ (FM) and 0.6 ±0.2 g m⁻²
17 d⁻¹ (MM).

18 Similarities between fringing salt marshes and meadow marshes included aboveground
19 (285±52 g m⁻² FM; 274±22 g m⁻² MM) and belowground (1379±200 g m⁻² FM; 1884±360 g m⁻²
20 MM) peak season biomass. Also, both marsh types reduced the height of waves coming onto the
21 marsh surface by approximately 63% only 7 m into the marsh. The plant diversity of fringing
22 marsh and meadow marsh sites as measured by the Evenness Index was also found to be similar
23 (0.557 FM; 0.536 MM).

1 The results of this study indicate that despite their small size, fringing salt marshes are
2 valuable components of estuaries, performing many ecological functions to the same degree as
3 nearby meadow marshes. More effort should be made to include them in regional efforts to
4 conserve and restore coastal habitats.

5

6

7 Key Words: Fringing salt marsh; meadow marsh; functions and values; Gulf of Maine

1 INTRODUCTION:

2 Our current understanding of salt marsh ecology comes from studies of large marsh
3 systems, especially those along the Eastern coast of the United States. These salt marshes
4 typically have a distinct zonation of plant communities, which reflect their surface elevation and
5 the effects of tidal flooding (Miller and Egler 1950; Niering and Warren 1980). They are valued
6 for a number of reasons, including their role as nursery grounds for finfish and shellfish, their
7 ability to accrete sediments and counter the effects of sea level rise, their role in storm surge
8 protection, and their recreational and aesthetic values (Teal 1986; Short et al. 2000). However
9 not all salt marshes are large meadow marshes. Particularly in the northeastern United States,
10 many of the salt marshes that line the edges of bays and rivers are quite narrow in width and
11 small in size (Roman et al. 2000, Morgan 2002). In the state of Maine, Jacobson et al. (1987)
12 found that nearly half of the coastal salt marsh area was comprised of marshes 0.2 ha (0.49 acres)
13 or smaller. These smaller marshes are often referred to as fringing salt marshes. Despite the fact
14 that they can be quite extensive, little is known about the ecology or the functions and values of
15 fringing salt marshes. Studies in Massachusetts and Rhode Island have examined the role that
16 fringing salt marshes may play as transformers of nitrogen that enters the estuary from adjacent
17 uplands (Lyons et al. 1995; Tobias et al. 2001b; Davis et al. 2004), but other aspects of fringing
18 marsh ecology have been studied little or not at all.

19 Fringing salt marshes are in need of study not just because of the paucity of information
20 available about them, but also because they are particularly susceptible to environmental impacts.
21 On their landward borders they are often abutted by residential and commercial development,
22 and on their seaward borders they are exposed to the erosive force of waves. Because they are
23 narrow, impacts to the borders of a fringing marsh have proportionately large effects on the entire

1 marsh. Also because they are narrow, fringing marshes provide convenient access to open water
2 for fishermen and boaters, who may impact them unintentionally.

3 The purpose of this study was to compare the functions and values of fringing salt
4 marshes to those of larger, meadow salt marshes and in some cases, to shoreline areas where no
5 marsh was present. We chose to use a functions and values approach to studying salt marshes for
6 two reasons. First, indicators of salt marsh functions can be measured objectively with
7 repeatable and quantitative methods. Second, functions can be linked to values, to which the
8 general public can relate. The distinction between functions and values is an important one.
9 Functions are ecosystem activities or processes that occur over time and do not depend on
10 societal perceptions; that is, they continue to occur whether or not people care about them
11 (Brinson and Rheinhardt 1996). Values are things that people care about because they are
12 “worthy, desirable or useful to humans” (Mitsch and Gosselink 1993). Citizens can more easily
13 understand the concept of wetland values than the concept of wetland functions, and values often
14 weigh heavily in decisions concerning the future of coastal resources. Based on past experience
15 and the scientific literature we developed a list of the functions and values of New England’s salt
16 marshes (Short et al. 2000). We then selected several of these for study in salt marshes in
17 southern Maine and New Hampshire.

18 Our specific objectives were to: (1) measure several of the ecological functions of
19 fringing salt marshes and meadow salt marshes, including (a) primary production, (b) soil
20 organic matter accumulation, (c) filtration/trapping of sediments, (d) maintenance of plant
21 biodiversity, and (e) dissipation of physical forces of waves; (2) compare these ecological
22 functions in fringing salt marshes and meadow salt marshes; and (3) determine how marsh
23 physical characteristics (size, elevation, surface slope and soil salinity) are related to these marsh
24 functions.

1

2 MATERIALS AND METHODS:

3 Five fringing marshes and five meadow marshes were selected for study from the Saco
4 River, Maine, south to the Great Bay Estuary in New Hampshire (Fig. 1; Table 1). The mean
5 tidal range within the area is approximately 9 ft, with semi-diurnal tides. The fringing marshes
6 chosen are all located along the edges of rivers, bays or coves, and the meadow marshes are
7 found behind barrier beaches (Fig. 2). All of the meadow marshes are naturally divided into
8 sections by large creeks or rivers, so we selected one of these sections for study in each meadow
9 marsh.

10 Nine sample stations were established on each marsh site using a stratified random
11 sampling design, according to the proportion of high marsh area to low marsh area. In fringing
12 marshes, an x-y coordinate system was used to locate the random points. In meadow marshes, a
13 latitude/longitude grid was placed over a base map of the marsh area and then nine random
14 points were chosen from the grid. These points were then located in the field using global
15 positioning system technology (GPS). Physical data (porewater salinity, surface elevation,
16 surface slope) and biological data (primary productivity, soil organic matter accumulation and
17 plant diversity) were then collected at these sample stations.

18

19 *Physical Characteristics of Marsh Study Sites*

20 Physical characteristics were measured at each of the nine stations per site. Previous
21 studies have shown that the physical characteristics of salt marsh sites can influence their
22 ecological functions (e.g. Jacobson and Jacobson 1989; Knutson et al. 1982; Osgood and Zieman
23 1993; Warren and Niering 1993; Kastler and Wiberg 1996). We therefore measured several

1 physical characteristics that might influence the functions we were investigating and looked for
2 correlations between them. We measured soil porewater salinity, surface elevation, percent
3 surface slope, and the distance of each sample point from the seaward edge of the marsh. Soil
4 porewater salinity was determined in July and August with a temperature-corrected optical
5 refractometer after extracting water from 10-15 cm depth with a soil sipper. Elevations of all
6 sample locations were determined using a Meridian L6-20 level and stadia pole. The relative
7 elevations of all stations on a site were first measured by surveying from the station points to a
8 relative benchmark nearby each site. These relative benchmarks were then tied into a high tide
9 elevation on one date, which allowed for comparison of elevations between all sites. To
10 determine the high tide line, three stakes painted with water-soluble paint were placed in each of
11 the ten marsh sites before high tide (11.9' MHHW) on a windless day. Following high tide the
12 water line on each stake was marked and then tied into the relative benchmark elevation at each
13 site. The elevations of all the quadrats on all the sites were then calculated relative to 0' tide
14 elevation. Surface slope was measured at each sample station in a direction perpendicular to the
15 water's edge, across a horizontal distance of 1 m. The distance from each sample station to the
16 water's edge was measured using a 50 m tape or a Lytespeed 400 rangefinder. The area of each
17 site was determined with NIH Image 1.47, from U.S.G.S. topographic maps, aerial photographs
18 (1" = 200') and field measurements.

19 Means and standard errors of the nine data points for each of the physical characteristics
20 above were calculated for each marsh site. Means of the ten sites were compared using Analysis
21 of Variance (ANOVA) and then pairwise comparisons were made with Student-Newman-Keuls
22 or Scheffe's S tests, as appropriate. The overall means and standard errors for meadow and
23 fringing marsh types were also determined and compared using ANOVA.

24

1 *Ecological Functions of Marsh Study Sites*

2 Five salt marsh functions at each of the ten salt marsh sites were evaluated using a variety
3 of indicators, each developed based on knowledge of the literature and previous experience. The
4 functions we assessed and the corresponding indicators we used to measure them are listed in
5 Table 2.

6 Primary production of vascular plants at each site was evaluated by measuring the annual
7 standing crop, including both live aboveground and live belowground plant biomass. Samples
8 were collected from the nine random sample points (stratified by proportion of low and high
9 marsh) at each marsh site at the end of the growing season (late August) by clipping all
10 vegetation in a 0.25 m² quadrat. Live plants were washed, then separated from dead material of
11 previous years and dried at 60°C for 48 hr and weighed. A sediment core (20 cm deep, 3.5 cm
12 diameter) was taken from each quadrat, then washed on a 2 mm screen. Live roots and rhizomes
13 were separated from dead material and then dried at 60°C for 48 hr and weighed to determine
14 belowground biomass (Gross et al. 1991).

15 To determine soil percent organic matter content, a core (15 cm deep; 3.5 cm diameter)
16 was taken from each of the nine sample stations, and percent organic matter in the sediment
17 determined from weight loss upon ignition in a muffle furnace (400°C) for 4 hr (Craft et al.
18 1991).

19 Sediment filtration and trapping was assessed by measuring the amount of sediment
20 accumulated on sediment traps (discs) over a ten day period in mid-August. Sediment traps were
21 designed after those of Reed (1989), and consisted of a pre-weighed mylar disc (8 cm diameter)
22 attached to a piece of sheet metal with plastic coated clips and held onto the marsh surface by 6
23 in long metal sod staples. Five sediment traps were distributed randomly at five of the sample

1 stations at each site, stratified by proportion of low to high marsh. Five additional traps were
2 placed on the marsh surface 1 m from the water's edge. These traps were randomly distributed
3 along the seaward edge of each marsh site. Three traps were also placed in areas where no marsh
4 was present, adjacent to the five fringing marsh sites. Discs were collected after two weeks,
5 dried at 60°C for 48 hr and weighed. In addition, the surface elevation and the distance between
6 each trap and the seaward edge of the marsh were measured. The number of plant stems and the
7 percent cover of plant species present in a 1 m² quadrat around each trap on the marsh surface
8 were also recorded. The suspended sediment concentration in the water flooding the marsh
9 surface at high tide was determined by collecting water as it flooded all sites on the same spring
10 tide night. Samples were later filtered through pre-weighed 0.45 µm glass fiber filters, then dried
11 at 60°C for 48 hr and weighed.

12 The species richness and relative abundance of higher plants were assessed once at each
13 site, in late July. Percent cover of all species in 1 m² quadrats was estimated visually using the
14 following cover classes: 0%, 0% < x ≤ 1%; 1% < x ≤ 5%; 5% < x ≤ 10%; 10% < x ≤ 20%; and
15 continuing above 20% in 10% increments up to 100%. Total percent cover per quadrat did not
16 exceed 100%. The number of quadrats sampled on fringing and meadow marshes was based on
17 preliminary sampling and calculation of running averages for small and large marsh sites. The
18 results of this initial analysis indicated that the minimum number of quadrats needed on fringing
19 and meadow marsh sites was 10 and 30, respectively, in order to include the majority of plant
20 species on the two different sized sites. These quadrats were then distributed in a stratified
21 random manner, according to the proportion of high and low marsh at each site.

22 To assess how well fringing and meadow marshes dissipate wave energy, we looked at
23 the difference between wave heights at the marsh/water edge and 5 m and 7 m into the marsh,

1 along a transect perpendicular to the water's edge. We surveyed three of the fringing marsh sites,
2 three of the meadow marsh sites, and three areas where no marsh was present. Three transects
3 were laid out at each site, evenly spaced along the marsh/water edge. Stakes with meter sticks
4 attached were placed at 0 m, 5 m and 7 m along each transect. Waves were generated by the
5 wake of a 17' aluminum boat and then videotaped simultaneously at 0 m and 5 m, then at 0 m
6 and 7 m. Waves from the boat were filmed three times (takes) at 0 m and 5 m and three times at
7 0 m and 7 m along each transect. Videotapes were later viewed frame-by-frame (30 frames s⁻¹)
8 and wave peaks and troughs were recorded for each take at 0 m and at 5 m or 7 m. The
9 maximum trough to peak height was determined for each take, as were the two wave heights
10 following the maximum wave. The percent reduction in maximum wave heights from 0 m and 5
11 m, and from 0 m to 7 m was calculated for the three takes at each transect and then averaged.
12 The mean height of three waves (maximum and two following) per take was also calculated and
13 then the percent reduction in this 'three wave mean' height was determined from 0 m and 5 m,
14 and from 0 m to 7 m. Finally, percent wave height reduction values (maximum and three wave)
15 obtained for the three transects were averaged to determine means for each fringing marsh,
16 meadow marsh and 'no marsh' site. In addition, the depth of the water at the time of filming was
17 also recorded at the 0 m, 5 m and 7 m points along transects.

18 Before comparing fringing marsh and meadow marsh functions, the possible relationships
19 between each of the functions and the physical characteristics measured at the sites were
20 explored. Scatterplots were drawn comparing the quantitative assessment for each function with
21 each of the physical characteristics investigated for that function. Correlation coefficients were
22 then calculated for each function-physical characteristic pair. Results of these correlations aided
23 in the choice of which variables to use as covariates in the means comparisons described below.

1 For each of the functions in Table 2 and their associated metrics, the means and standard
2 errors of the five fringing marshes and the five meadow marshes were calculated. Means were
3 also calculated for the areas where no marsh was present when assessing sediment filtration and
4 trapping, and the dissipation of wave energy. Analysis of Variance (ANOVA) or Analysis of
5 Covariance (ANCOVA) was then employed to compare the mean values from the fringing marsh
6 sites with those of the meadow marsh sites for each function. If data were collected at 'no marsh'
7 sites, these were included in the means comparisons as well.

8 Data collected to assess the function of maintenance of plant diversity were analyzed to
9 compare the number of species per site, species density, and evenness (E) of vascular plant
10 species in meadow and fringing marshes. The percent covers of *Spartina alterniflora* and of the
11 dominant high marsh species *Juncus gerardii*, *Puccinellia maritima* and *Spartina patens* were
12 also calculated for each marsh site. Average values for each of these plant community attributes
13 were then calculated and compared using ANOVA. Calculations were based on ten random
14 quadrats per marsh site when calculating plant species evenness (E), as this diversity parameter
15 requires an equal sample size to compare two communities. For the other indicators of plant
16 species diversity, fringing marsh means are based on ten quadrats and meadow marsh means on
17 thirty quadrats, although the means of ten randomly selected meadow marsh quadrats are also
18 presented.

19

20 RESULTS:

21 *Physical Characteristics of Marsh Study Sites*

22 Soil porewater salinity, surface elevation, marsh area and distance to the water's edge of
23 sample points were all significantly less in fringing salt marshes than in meadow marshes

1 (calculated as means of five sites; $p < 0.05$) (Fig. 3a-d). However surface slope was significantly
2 greater in fringing marshes than in meadow marshes ($p < 0.01$)(Fig. 3e). To see how each marsh
3 contributed to these differences, Figure 3 also shows means and standard errors of the nine
4 samples at each marsh site.

6 *Comparison of Fringing Marsh Functions to Meadow Marsh Functions*

7 Primary production

8 We found no difference in the production of aboveground or belowground biomass
9 between fringing and meadow marshes (Fig. 4). The slope of the marsh surface may affect
10 productivity in fringing marshes, as it was highly correlated with both aboveground ($r = 0.941$, p
11 < 0.05) and belowground ($r = 0.951$, $p < 0.05$) biomass.

12

13 Soil organic matter accumulation

14 The organic matter content of meadow marsh soils was significantly greater than that of
15 the fringing marsh soils (ANOVA; $p < 0.001$) (Fig. 5a). The five fringing marsh sites had lower
16 surface elevations than the five meadow marshes, and two were significantly lower in elevation
17 (Fig. 3g). Soil percent organic matter content was discovered to correlate with marsh surface
18 elevation ($r = 0.801$, $p < 0.05$) (Fig. 5b), and in an ANCOVA of marsh type and elevation,
19 organic matter in meadow marsh soils was greater than that in fringing marsh soils, even after the
20 variability due to elevation was accounted for ($p < 0.001$).

1

2 Sediment filtration and trapping

3 Although there was on average more sediment deposited on the traps randomly
4 distributed on the surface of the fringing marshes than on the surface of the meadow marsh sites,
5 this difference was not significant (Fig. 6a). Areas where no marsh was present had an even
6 greater amount of sediment deposited per unit area. However the variance around the mean was
7 extremely high for 'no marsh' areas, with the standard deviation (6.84) greater than the mean
8 ($4.24 \text{ g m}^{-2} \text{ d}^{-1}$). A comparison of the means for meadow, fringing and 'no marsh' areas showed
9 no significant difference in the amount of sediment deposited on these three site types, even after
10 removing the variance associated with elevation, which was a significant ($p < 0.001$) covariate in
11 the model (ANCOVA; $p = 0.374$, log transformed data). If elevation was not included in the
12 model, then $p = 0.134$ (log transformed data). Sediment deposition was less at sites with a higher
13 mean elevation, at both fringing and meadow marsh sites.

14 Traps placed just one meter in from the edge of the marsh sites collected more sediment
15 than those that were distributed randomly, as expected. Once again there was no significant
16 difference in the mean amount of sediment deposited on fringing, meadow and 'no marsh' sites,
17 as determined by ANCOVA with elevation used as a covariate ($p = 0.120$, log transformed
18 data)(Fig. 6b). If the variability due to elevation was not removed, $p = 0.721$ (log transformed
19 data). It should be noted that one trap at site DIM had an unusually large amount of sediment
20 deposited on it ($1847.89 \text{ g m}^{-2} \text{ d}^{-1}$, compared to the next highest value of $0.92 \text{ g m}^{-2} \text{ d}^{-1}$). This
21 was attributed to the presence of a nearby culvert, which greatly increased the velocity of the
22 water moving through the area, most likely causing large amounts of sediment to be resuspended
23 and deposited. This data point was therefore considered as an outlier and was discarded.

1 We had expected that suspended sediment concentration of the tidal water moving onto
2 the marsh surface would influence the amount of sediment deposited on the sediment traps, but
3 this is not what we observed in meadow marshes (Fig. 7). Although significant in fringing
4 marshes, the relationship in Figure 7 is driven by observations at a single site ($r = 0.999$, $p <$
5 0.001). We did find that vegetative cover may influence the amount of sediment deposited on the
6 marsh surface, however. The greater the percent cover of plants around sediment traps, the less
7 the amount of sediment deposited ($r = -0.732$, $p < 0.05$).

8

9 Maintenance of plant communities

10 Several measures of plant diversity showed no significant difference between fringing
11 marsh and meadow marsh plant communities. The Shannon-Weiner Index (H'), Evenness (E),
12 and species richness were all similar in the two marsh types (Table 3). However, if the full
13 sample of thirty quadrats on meadow marshes was used, then species richness of meadow
14 marshes was greater than that of fringing marshes (ANOVA, $p < 0.01$). Although the percent
15 cover of *Spartina alterniflora* in fringing marshes was almost double what it was in meadow
16 marshes, this difference was also not significant, in large part due to the short form of this species
17 occurring in high marsh areas of meadow marshes. Species density, however, was less in
18 fringing marshes than in meadow marshes (ANOVA, $p < 0.01$). The total percent cover of the
19 dominant high marsh plants, including *Spartina patens*, *Juncus gerardii*, *Distichlis spicata* and
20 *Puccinellia maritima* was also less in fringing marshes (ANOVA, $p < 0.05$).

1

2 Dissipation of physical forces of waves

3 An example of the wave profiles generated from videotaping passing waves at 0 m (the
4 marsh edge) and at 5 m can be seen in Figure 8. Along all transects at all sites, the heights of the
5 largest waves at 0 m ranged from 3.5 cm to 27.3 cm, averaging 12 cm tall. The ‘three wave
6 mean’ height (mean height of the maximum and next two waves) at 0 m ranged from 2.7 cm to
7 21.2 cm, with an average of 7.8 cm.

8 It should be noted that the waves used to calculate percent height reductions along each
9 transect were not shallow water waves. We determined this by measuring wavelengths of
10 suspect waves on the video screen and comparing them to water depths at those points. The
11 water depth was always significantly greater than 1/20 of the wavelength (Denny 1988).

12 In both fringing and meadow marshes, the heights of the largest waves traveling 7 m
13 across the marsh surface were reduced by more than 60% (Fig. 9a). Where no marsh was
14 present, wave heights were reduced by only 33%. This difference between marsh and ‘no marsh’
15 areas was statistically significant (ANOVA, $p < 0.05$, square root transformed data; Student-
16 Neuman-Keuls test, $p < 0.05$). The percent reduction in wave height across 7 m was less when
17 we considered the ‘three wave mean’ height (55% in fringing and 52% in meadow marshes,
18 compared to 28% in ‘no marsh’ areas), and again the difference between marsh and ‘no marsh’
19 areas was statistically significant (ANOVA $p < 0.05$, square root transformed data; Student-
20 Neuman-Keuls test, $p < 0.05$) (Fig. 9b).

1

2 DISCUSSION:

3 Although there have been some studies of fringing marsh ecology (e.g. Kastler and
4 Wiberg 1996; Tobias et al. 2001a; Davis et al. 2004; Bozeck and Burdick 2005), few studies
5 have considered fringing salt marshes as unique habitats, distinct from larger, meadow salt
6 marshes. To clarify the role of fringing salt marshes in estuaries along the southern Maine/New
7 Hampshire coast, we studied how they function compared to large meadow marshes in the same
8 area. We discovered that fringing salt marshes are diverse in terms of their physical
9 characteristics and that this diversity is sometimes reflected in their ecological functions. We
10 also found that despite this diversity, fringing marshes as a group often function at levels similar
11 to meadow marshes.

12

13 *Physical Characteristics of Fringing and Meadow Marsh Study Sites*

14 In general, fringing marshes are narrower, steeper, and have lower mean surface
15 elevations than meadow marshes. They are also more variable than meadow marshes in terms of
16 their elevations and surface slopes (Fig. 3). This variability is an important property of fringing
17 salt marshes, in part because elevation and slope may influence marsh function, as will be
18 discussed. In addition, their variability makes it difficult to describe a “typical” fringing salt
19 marsh. For example, some fringing salt marshes contain predominantly low marsh plant
20 communities, but others have extensive high marsh areas. Although the fringing marsh study
21 sites had statistically lower mean soil porewater salinity than the meadow marsh sites, this

1 difference was primarily due to the very low soil salinity at one fringing marsh site (Fig. 3a,f).
2 We found no strong correlations between salinity and any of the functions we studied.

3

4 *Comparison of Fringing Marsh Functions to Meadow Marsh Functions*

5 Primary production

6 Our results demonstrate that the primary productivity of fringing marshes is as great as
7 that of meadow marshes, indicating that they are important contributors to estuarine food webs.
8 Mean aboveground production in fringing marshes was similar to that in meadow marsh sites,
9 and although the mean belowground production in meadow marsh sites was greater than that in
10 fringing marsh sites, this difference was not significant (Fig. 4). It should be noted that although
11 harvesting the peak season standing crop as a measure of aboveground production is a commonly
12 used method, it underestimates true aboveground net production by 10-15% (Nixon and Oviatt
13 1973). Comparing the aboveground biomass values we obtained to those of other studies is
14 difficult because of the variety of sample methods that have been employed to measure
15 aboveground production (Marinucci 1982). Nevertheless, our values are within the range of
16 those found in studies of other Maine and New Hampshire salt marshes (Lindthurst and Reimold
17 1978; Gross et al. 1991).

18 Studies of salt marsh belowground biomass production are few in number compared to
19 studies of aboveground biomass production, due to the difficulty of sampling and processing
20 belowground tissues (Gross et al. 1991). However investigating the belowground component of
21 production is important, as it can be 4-7 times greater than that of aboveground production
22 (Marinucci 1982). The belowground to aboveground biomass ratio was 4.8 in the fringing

1 marshes we sampled and 6.9 in the meadow marshes. Also, our values for belowground
2 production agree with what others have found in New England marshes (Lindthurst and Reimold
3 1978; Gross et al. 1991).

4 The positive relationship we observed between above and belowground production and
5 surface slope ($r = 0.941$ and $p = 0.951$, respectively) could be attributed to the “streamside
6 effect.” In general, marsh surfaces are more steeply sloped where they are adjacent to tidal
7 waters, either along the edge of a creek or along the seaward edge of the marsh, and aboveground
8 primary production is greater here (Gallagher and Kibby 1981; Burdick et al. 1989). Soils in
9 areas exposed to tidal waters more often are typically more well-drained, and sediment oxidation
10 rates are higher, so gas exchange between roots and the surrounding soils can take place more
11 rapidly than in waterlogged areas (Burdick et al. 1989). Although differences in belowground
12 biomass production in *Spartina* marshes have not been well studied, Gallagher and Kibby (1981)
13 found that streamside plants had greater recoverable underground reserves than back marsh
14 plants in a *Carex lyngbyei* tidal marsh. Ellison et al. (1986) also found that belowground
15 production in a Massachusetts salt marsh was greater at the marsh edge than on other parts of the
16 marsh.

17

18 Soil organic matter accumulation

19 Our results show that the percent organic matter content of meadow marsh soils is more
20 than three times that of fringing marsh soils (Fig. 5a). The meadow marshes we sampled had
21 greater surface elevations than the fringing marshes (Fig. 3b), and there was a positive correlation
22 between elevation and soil organic matter content (Fig. 5b). Schmitt et al. (1998) also found an

1 increase in the amount of organic matter deposited in the sediment and on the marsh surface with
2 increasing elevation in a Massachusetts salt marsh.

3 If salt marshes are to keep pace with rising sea level, they must be able to accrete at a rate
4 equal to or greater than that of sea level rise (Donnelly and Bertness 2001, Patrick and DeLaune
5 1990). Vertical accretion relies on two sources of sediment; one from waters that flood the marsh
6 surface, and the other from above and belowground plant biomass which does not completely
7 decompose, contributing organic material to marsh soils (Redfield 1972; Nixon 1982). The build
8 up of organic matter in marsh soils appears to be most important in the high marsh zone. In a
9 study of five Rhode Island salt marshes, Bricker-Urso et al. (1989) found that the contribution of
10 organic matter to accretion on the high marsh was more than twice that of inorganic sediments,
11 but in the low marsh the contribution of inorganic and organic sediments was equal. In addition,
12 Ellison et al. (1986) found that the decomposition rate of live roots and rhizomes was slower in
13 the high marsh zone than at the marsh edge. Lower decomposition rates in interior, poorly
14 drained high marsh soils may result in organic matter accumulation. The distance that sample
15 points are from the marsh edge has also been observed to correlate with soil percent organic
16 matter content. The percent organic matter in sediments of two Virginia salt marshes was lowest
17 at the water's edge and increased along a 30 m transect into their interiors (Kastler and Wiberg
18 1996). In this study, we also found that the soil organic matter content correlated with the
19 distance the sample points were from the edge of the marsh ($r = 0.704$, $p < 0.05$). Our results
20 indicate that meadow marshes along the southern Maine/NH coast rely more on soil organic
21 matter accumulation for accretion than fringing marshes do. We conclude that the salt marsh
22 function of soil organic matter accumulation is performed to a greater extent in meadow marshes
23 than in fringing marshes. If this is the case, then to keep pace with sea level rise, fringing

1 marshes must rely to a greater extent on the trapping of inorganic sediments as their predominant
2 mechanism of accretion.

3

4 Sediment filtration and trapping

5 Reed (1989) first developed the technique of trapping sediment on filter paper discs
6 attached to the marsh surface. Due to the activity of green crabs in our area, we modified her
7 design and used discs made of Mylar, which crabs do not find so appetizing. In one study of
8 sediment deposition on Louisiana tidal marshes, Reed (1989) found rates of $2.9 \text{ g m}^{-2} \text{ d}^{-1}$
9 (excluding winter storm events, when sedimentation rates were much higher). We obtained
10 similar values for sediment deposition, with marsh site means ranging from $0.44\text{-}4.31 \text{ g m}^{-2} \text{ d}^{-1}$
11 for traps randomly distributed on fringing marshes and $0.20\text{-}1.51 \text{ g m}^{-2} \text{ d}^{-1}$ for traps randomly
12 distributed on meadow marshes.

13 We observed that sediment deposition rates decreased with increasing elevation ($r = -$
14 0.732 , $p < 0.05$), probably because tidal waters cover marsh areas at higher elevations less
15 frequently and for a shorter period of time. Negative correlations between elevation and
16 sediment deposition have also been observed in Massachusetts (Schmitt et al. 1998) and North
17 Carolina (Leonard 1997) salt marshes.

18 Intertidal areas where no marsh vegetation was present (designated as 'no marsh' areas)
19 showed a greater rate of sediment deposition ($0.62\text{-}16.44 \text{ g m}^{-2} \text{ d}^{-1}$) when compared to fringing
20 marshes or meadow marshes, but these differences were not significant (Fig. 6a). Local
21 resuspension of surface sediment on 'no marsh' sites may have contributed to greater deposition
22 rates. We had expected to see reduced rates of sediment deposition on 'no marsh' areas, as the

1 presence and density of marsh vegetation has been observed to correlate positively with sediment
2 trapping (Gleason et al. 1979; Stumpf 1983).

3 We also expected greater sediment deposition per unit area on fringing salt marshes than
4 on meadow marshes because, considering the length of marsh bordering tidal waters, fringing
5 marshes have a greater edge: area ratio than meadow marshes, so a finite sediment supply would
6 appear as more sediment deposited per unit area on fringing marshes. We did observe higher
7 rates of deposition on fringing marshes than on meadow marshes, but this difference was not
8 significant (Fig. 6a), likely due to the high variability in deposition rates between sites within
9 each site type. To eliminate any effect of the greater edge: area ratio of fringing marshes, we
10 placed traps at fringing and meadow marsh sites just 1 m from the water's edge. Again, the rates
11 of sediment deposition were greater on fringing marshes, but this difference was not significant
12 (Fig. 6b).

13 Our results suggest that fringing marshes trap greater amounts of sediment per unit area
14 than meadow marshes, although the variability between sites was too high and our sample size
15 was too small to confirm this. With slower rates of organic matter accumulation than occur in
16 meadow marshes (Figure 5a), the trapping of inorganic sediments is an important mechanism of
17 accretion for fringing marshes. Whether fringing marshes in this area are performing the
18 function of sediment filtration and trapping at levels sufficient to ensure that their rates of
19 accretion will keep pace with sea level rise is an important question that deserves further
20 investigation. The use of feldspar clay marker horizons to measure marsh accretion over several
21 years could help in answering this question (Cahoon and Turner 1989).

1

2 Maintenance of plant diversity

3 Fringing salt marshes in New England are often thought to have very low plant diversity,
4 and to be comprised primarily of *Spartina alterniflora* (Cook et al. 1993; Bryan et al. 1997). In
5 contrast to this, we found that fringing salt marshes often have very developed high marsh
6 communities. All of the fringing marshes we studied contained *Spartina patens*, a typical high
7 marsh plant, and at two sites, *S. patens* was more abundant than *S. alterniflora*. So although the
8 proportion of high marsh to low marsh in the fringing marshes we studied was less (0.7:1) than in
9 meadow marshes (3.4:1), high marsh species are an important component of fringing marsh plant
10 communities.

11 Our results did confirm that meadow marshes are more species rich than fringing salt
12 marshes in this part of New England (Table 3). This is likely due to the “area effect” (the
13 number of species sampled increases with increasing sample size (Magurran 1988)); species
14 richness was correlated with marsh area ($r = 0.818$, $p < 0.05$). Species density (number of species
15 present per m^2) was also greater in meadow marshes than in fringing marshes, and correlated
16 with marsh area ($r = 0.676$, $p < 0.05$).

17 The low marsh zone in New England salt marshes is typically dominated by *Spartina*
18 *alterniflora*, and the sites we studied fit this pattern. However the high marsh zone of the
19 majority of the meadow marshes in our study did not fit the pattern commonly observed in other
20 New England marshes, where there are distinct bands of *S. patens* and *Juncus gerardii* (Miller
21 and Egler 1950; Niering and Warren 1980). Instead, the high marsh often contained large areas of
22 forbs (broad leaved plants), in a mosaic of patches of *S. patens*, *J. gerardii* and other dominant
23 high marsh grasses. Ewenchuck and Bertness (2004) also noted the occurrence of forb pannes in

1 a Wells, ME marsh (one of our study sites), and attributed the occurrence of these forb patches to
2 the waterlogged, anoxic soils found in northern New England salt marshes. Because northern
3 New England salt marshes have not been as extensively ditched as marshes farther south, they
4 tend to be wetter (Ewenchuck and Bertness, 2004). Jacobson and Jacobson (1989) found mosaic
5 patterns of vegetation in a number of the Maine salt marshes they sampled, which they
6 hypothesized was due to microrelief in high marsh areas.

7 Although meadow marshes have greater species richness and density, their plant
8 communities are comparable to those of fringing marshes in terms of two other measures of plant
9 diversity, the Shannon-Weiner index (H') and Evenness index (E) (Table 3). The Evenness index
10 we employed is the ratio of observed diversity to maximum diversity, $E = H' / H_{\max} = H' / \ln S$
11 (Magurran 1988). Values for E describe how close the set of species abundances for a marsh site
12 is to having maximum diversity, where the relative abundances for all species are equal. Our
13 results show that the relative abundances of species were similar in the fringing and meadow
14 marsh sites we sampled. It should be noted that these results are based on ten quadrats sampled in
15 both fringing and meadow marshes, as equal sample sizes must be used when calculating H' and
16 E .

17

18 Dissipation of physical forces of waves

19 Previous studies have shown that salt marshes reduce the height and energy of incoming
20 waves, helping to protect the adjacent upland from erosion (Knutson et al. 1982; Moeller et al.
21 1996). In addition, salt marshes reduce wave velocity, resulting in increased sediment deposition
22 on the marsh surface and decreased sediment erosion (Leonard and Luther 1995). We were

1 interested in knowing if marsh type (fringing or meadow) or other characteristics (vegetation,
2 slope) affected a marsh's ability to reduce the height (energy) of incoming waves.

3 Our results demonstrate that marsh type does not affect a site's ability to reduce the
4 height of incoming waves, with fringing and meadow marshes both causing waves to lose energy
5 as they traveled 7 m across the marsh surface (Fig. 9). The maximum wave height was reduced
6 62% in fringing marshes and 64% in meadow marshes after traveling 7 m across the marsh
7 surface. These values are similar to those obtained by Knutson et al. (1982), who found wave
8 heights reduced by 57% five meters into a *S. alterniflora* marsh, and 65% at ten meters. Leonard
9 and Luther (1995) found a 65% reduction in the turbulent energy of water coming onto the marsh
10 after it had traveled just 3 m in from the marsh edge.

11 Areas where no marsh was present were much less effective at reducing the height of
12 maximum waves (33% over 7 m), as expected. In Moeller et al.'s (1996) study of a *S.*
13 *alterniflora* marsh in England, they found that low marsh areas absorbed 2-3 times as much wave
14 energy as adjacent sand flats.

15 Our results demonstrate that for waves up to 27 cm in height (typical of boat or wind
16 generated waves), even narrow fringing marshes are capable of reducing wave energy by almost
17 two thirds, helping to protect adjacent shorelines from the erosive forces of waves.

18

19 Summary and Conclusions

20 *Fringing Salt Marshes Defined*

21 Found along the edges of bays and rivers, the fringing salt marshes of New England have
22 been described as relatively long and narrow in shape and dominated by *Spartina alterniflora*

1 (Bryan et al. 1997; Roman et al. 2000). We can now more clearly define fringing marshes as
2 having steeper slopes, lower elevations and soils with less organic matter than those of larger
3 marshes. In addition, their plant communities usually include both low marsh and high marsh
4 zones, although in more equal proportions than is seen in larger marshes, where the high marsh
5 dominates. It is also important to note that fringing salt marshes are quite diverse in terms of their
6 physical characteristics (width, length, slope, elevation, soils).

7

8 *Functions and Values of Fringing Salt Marshes*

9 Fringing salt marshes have important functions and values that had not been investigated
10 prior to this study. They are as productive on a per unit area basis as meadow marshes, making
11 valuable contributions to detrital and grazing food webs. Their ability to filter and trap sediments
12 from the water column improves water quality and adds to marsh accretion, helping fringing
13 marshes keep pace with sea level rise. By dampening the energy of incoming waves, fringing
14 marshes help protect the adjacent shoreline from erosion. This is especially important because in
15 New England, these narrow marshes are often the only buffer between the erosive forces of
16 waves and valuable upland coastal property. Fringing marshes are important in maintaining plant
17 biodiversity in the estuary, as they contain distinct low and high marsh zones dominated by the
18 same plant species found in meadow marshes. And finally, the importance of connectivity
19 between habitat patches is well known in conservation biology (Hunter and Gibbs 2007).
20 Fringing salt marshes serve the function of connecting larger salt marshes in New England's
21 estuaries. Lining the edges of rivers and bays, and distributed between the larger meadow
22 marshes, they serve an important role in the dispersal and colonization of salt marsh plant and
23 animal species.

1 In many New England estuaries, fringing salt marshes are the dominant marsh type. And
2 yet regional efforts aimed at marsh conservation and restoration still focus on larger, meadow
3 marshes (Konisky et al. 2006, Taylor 2008). With an improved understanding of the ecological
4 functions of fringing marshes and of their value to coastal communities, we can do a better job of
5 protecting these important resources.

6
7

8 Acknowledgements

9

10 We thank Thomas Lee, William B. Bowden and Michelle Dionne for their advice during this
11 project. Thanks also to the many field and laboratory assistants from the Jackson Estuarine Lab
12 and the University of New England. Special thanks to Ryan Davis, Jeff Gaeckle, and Blaine
13 Kopp for their support and critique. Funding for this project came from the University of New
14 England, University of New Hampshire and National Oceanic and Atmospheric Administration,
15 Grant or cooperative agreement no. NA77OR0235. This is Jackson Estuarine Laboratory
16 contribution number xxx.

17

18

LITERATURE CITED:

- Bozeck, C.M. and D.M. Burdick. 2005. Impacts of seawalls on saltmarsh plant communities in the Great Bay Estuary, New Hampshire USA. *Wetlands Ecology and Management* 13(5):553-568.
- Bricker-Urso, S., S. W. Nixon, J. K. Cochran, D. J. Hirschberg, and C. Hunt. 1989. Accretion rates and sediment accumulation in Rhode Island salt marshes. *Estuaries* 12:300-317.
- Brinson, M. M. and R. Rheinhardt. 1996. The role of reference wetlands in functional assessment and mitigation. *Ecological Applications* 6:69-76.
- Bryan, R. R., M. Dionne, R. A. Cook, J. Jones, and A. Goodspeed. 1997. Maine citizens guide to evaluating, restoring, and managing tidal marshes. Maine Audubon Society, Falmouth, ME.
- Burdick, D. M., I. A. Mendelssohn, and K. L. McKee. 1989. Live standing crop and metabolism of the marsh grass *Spartina patens* as related to edaphic factors in a brackish, mixed marsh community in Louisiana. *Estuaries* 12(3):195-204.
- Cook, R. A., A. J. Lindley Stone, and A. P. Amman. 1993. Method for the evaluation and inventory of vegetated tidal marshes in New Hampshire. Audubon Society of New Hampshire.
- Cahoon, D. R., and Turner, R. E. 1989. Accretion and canal impacts in a rapidly subsiding wetland; II. Feldspar marker horizon technique. *Estuaries* 12(4):260-268.
- Craft, C. B., E. D. Seneca, and S. W. Broome. 1991. Loss on ignition and Kjeldahl digestion for estimating organic carbon and total nitrogen in estuarine marsh soils: Calibration with dry combustion. *Estuaries* 14:175-179.

- Davis, J., B. Nowicki, and C. Wigand. 2004. Denitrification of fringing salt marshes of Narragansett Bay, Rhode Island, USA. *Wetlands* 24(4):870-878.
- Denny, M. W. 1988. *Biology and the mechanics of the wave-swept environment*. Princeton, N.J.: Princeton University Press.
- Donnelly, J. P. and M. D. Bertness. 2001. Rapid shoreward encroachment of salt marsh cordgrass in response to accelerated sea-level rise. *Proceedings of the National Academy of Sciences* 98(25):14218-14223.
- Ellison, A. M., M. D. Bertness, and T. Miller. 1986. Seasonal patterns in the belowground biomass of *Spartina alterniflora* (Gramineae) across a tidal gradient. *American Journal of Botany* 73(11):1548-1554.
- Ewenchuck, P. J. and M. D. Bertness. 2004. Structure and organization of a northern New England salt marsh plant community. *Journal of Ecology* 92:72-85.
- Gallagher, J. L. and H. V. Kibby. 1981. The streamside effect in a *Carex lyngbyei* estuarine marsh: the possible role of recoverable underground reserves. *Estuarine, Coastal and Shelf Science* 12:451-460.
- Gleason, M. L., D. Elmer, and N. C. Pien. 1979. Effects of stem density upon sediment retention by salt marsh cord grass, *Spartina alterniflora* Loisel. *Estuaries* 2:271-273.
- Gross, M. F., M. A. Hardisky, P. L. Wolf, and V. Klemas. 1991. Relationship between aboveground and belowground biomass of *Spartina alterniflora* (smooth cordgrass). *Estuaries* 14:180-191.
- Hunter, M. L. and J.P. Gibbs. 2007. *Fundamentals of conservation biology*. Malden, MA: Blackwell Publishing.
- Jacobson, H. A. and G. L. J. Jacobson. 1989. Variability of vegetation in tidal marshes of Maine, U.S.A. *Canadian Journal of Botany* 67:230-238.

- Jacobson, H. A., G. L. J. Jacobson, and J. T. Kelley. 1987. Distribution and abundance of tidal marshes along the coast of Maine. *Estuaries* 10:126-131.
- Kastler, J. A. and P. L. Wiberg. 1996. Sedimentation and boundary changes of Virginia salt marshes. *Estuarine, Coastal, and Shelf Science* 42:683-700.
- Knutson, P. L., R. A. Brochu, W. N. Seelig, and M. Inskeep. 1982. Wave dampening in *Spartina alterniflora* marshes. *Wetlands* 2:87-104.
- Konisky, R.A., D.M. Burdick, M. Dionne and H.A. Neckles. 2006. A regional assessment of salt marsh restoration and monitoring in the Gulf of Maine. *Restoration Ecology* 14(4): 516–525.
- Leonard, L. 1997. Controls of sediment transport and deposition in an incised mainland marsh basin, southeastern North Carolina. *Wetlands* 17(2):263-274.
- Leonard, L. A. and M. E. Luther. 1995. Flow hydrodynamics in tidal marsh canopies. *Limnology and Oceanography* 40:1474-1484.
- Lindthurst, R. A. and R. J. Rheimold. 1978. An evaluation of methods for estimating the net aerial primary productivity of estuarine angiosperms. *Journal of Applied Ecology* 15:919-31.
- Lyons, J., J. Ahern, J. McClelland, and I. Valiela. 1995. Macrophyte abundances in Waquoit Bay estuaries subject to different nutrient loads and the potential role of fringing salt marsh in groundwater nitrogen interception. *Biological Bulletin* 189:255-256.
- Magurran, A. C. 1988. *Ecological diversity and its measurement*. Princeton, NJ: Princeton University Press.
- Marinucci, A. C. 1982. Trophic importance of *Spartina alterniflora* production and decomposition to the marsh-estuarine ecosystem. *Biological Conservation* 22:35-58.

- Miller, W. B. and F. E. Egler. 1950. Vegetation of the Wequetequock-Pawcatuck tidal marshes, Connecticut. *Ecological Monographs* 20:143-172.
- Mitsch, W. J. and J. G. Gosselink. 1993. *Wetlands*. New York: Van Nostrand Reinhold.
- Moeller, I., T. Spencer, and J. R. French. 1996. Wind wave attenuation over saltmarsh surfaces: Preliminary results from Norfolk, England. *Journal of Coastal Research* 12:1009-1016.
- Morgan, P.A. 2000. Conservation and ecology of fringing salt marshes along the southern Maine/New Hampshire coast. Ph.D. Thesis, University of New Hampshire.
- Niering, W. A. and R. S. Warren. 1980. Vegetation patterns and processes in New England salt marshes. *Bioscience* 30:300-306.
- Nixon, S. W. 1982. The ecology of New England high salt marshes. Office of Biological Services, U.S. Fish and Wildlife Service, Report No. FWS/BS-81/55.
- Nixon, S. W. and C. A. Oviatt. 1973. Analysis of local variation in the standing crop of *Spartina alterniflora*. *Botanica Marina* 16:103-109.
- Osgood, D. T. and J. C. Zieman. 1993. Factors controlling aboveground *Spartina alterniflora* (smooth cordgrass) tissue element composition and production in different-age barrier island marshes. *Estuaries* 16:815-826.
- Patrick, W. H. and R. D. DeLaune. 1990. Subsidence, accretion and sea level rise in south San Francisco Bay marshes. *Limnology and Oceanography* 35(6):1389-1395.
- Redfield, A. C. 1972. Development of a New England salt marsh. *Ecological Monographs* 42:201-237.
- Reed, D. J. 1989. Patterns of sediment deposition in subsiding coastal salt marshes, Terrebonne Bay, Louisiana: The role of winter storms. *Estuaries* 12:222-227.
- Roman, C.T., N. Jaworski, F. T. Short, S. Findlay, and R. S. Warren. 2000. Estuaries of the northeastern United States: Habitat and land use signatures. *Estuaries* 23(6):743-764.

- Schmitt, C., N. Weston, and C. Hopkinson. 1998. Preliminary evaluation of sedimentation rates and species distribution in Plum Island Estuary, Massachusetts. *The Biological Bulletin* 232-233.
- Short F. T., D. M. Burdick, C. A. Short, R. C. Davis, and P. A. Morgan. 2000. Developing success criteria for restored eelgrass, salt marsh and mud flat habitats. *Ecological Engineering* 15(3):239-252.
- Stumpf, R. P. 1983. The process of sedimentation on the surface of a salt marsh. *Estuarine, Coastal and Shelf Science* 17:495-508.
- Teal, J. M. 1986. The ecology of regularly flooded salt marshes of New England: A community profile. U.S. Fish and Wildlife Service, Biological Report No. 85(7.4).
- Taylor, P.H. 2008. Salt marshes in the Gulf of Maine: Human impacts, habitat restoration and long-term change analysis. Gulf of Maine Council on the Marine Environment. www.gulfofmaine.org. IV+42 pp.
- Tobias, C. R., I. C. Anderson, E. A. Canuel, and S. A. Macko. 2001a. Nitrogen cycling through a fringing marsh-aquifer ecotone. *Marine Ecology Progress Series* 210:25-39.
- Tobias, C. R., S. A. Macko, I. C. Anderson, E. A. Canuel, and J. W. Harvey. 2001b. Tracking the fate of a high concentration groundwater nitrate plume through a fringing marsh: A combined groundwater tracer and in situ isotope enrichment study. *Limnology and Oceanography* 46(8):1977-1989.
- Warren, R. S. and W. A. Niering. 1993. Vegetation change on a Northeast tidal marsh: Interaction of sea-level rise and marsh accretion. *Ecology* 74:96-103.

Figure 1. Locations of fringing salt marsh and meadow salt marsh study sites. Complete site names and locations (latitude/longitude) are listed in Table 1. Sites ending in "F" are fringing marshes; sites ending in "M" are meadow marshes.

Figure 2. Relative sizes of five meadow and five fringing salt marsh sites (all shown at the same scale). Study sites are dark grey; surrounding salt marshes are medium gray.

Figure 3. Means of physical characteristics for each marsh type and for sample points at each site. Bars in each graph followed by the same letter are not significantly different from each other according to the pairwise comparison listed. Error bars are ± 1 SE.

- a. Mean porewater salinities of five fringing marsh and five meadow marsh sites. Means are significantly different from each other (ANOVA, $p < 0.05$).
- b. Mean elevations of five fringing marsh and five meadow marsh sites. Means are significantly different from each other (ANOVA, $p < 0.01$). Elevation units are meters above 0' tide.
- c. Mean areas of five fringing marsh and five meadow marsh sites. Means are significantly different from each other (ANOVA, $p < 0.001$, square root transformed data).
- d. Mean distance from water's edge to nine sample quadrats at five fringing marsh and five meadow marsh sites. Means are significantly different from each other (ANOVA, $p < 0.01$, log transformed data).
- e. Mean surface slope of five fringing marsh and five meadow marsh sites. Means are significantly different from each other (ANOVA, $p < 0.01$, log transformed data).

- f. Porewater salinities at each site. Means are significantly different from each other (ANOVA, $p < 0.001$). Pairwise comparisons made with Student-Newman-Keuls (SNK).
- g. Elevations at each site. Means are significantly different from each other (ANOVA, $p < 0.001$). Pairwise comparisons made with Scheffe's S. Elevation units are meters above 0' tide.
- h. Areas of fringing and meadow marsh sites.
- i. Mean distance from water's edge to nine sample quadrats at each site. Means are significantly different from each other (ANOVA, $p < 0.001$, log transformed data). Pairwise comparisons made with SNK.
- j. Percent surface slope at each site. Means are significantly different from each other (ANOVA, $p < 0.001$, log transformed data). Pairwise comparisons made with SNK.

Figure 4. (a) Aboveground and (b) belowground plant biomass of fringing and meadow salt marsh sites. Error bars are ± 1 SE from the mean. Neither aboveground ($p = 0.924$) nor belowground ($p = 0.195$, $1/x$ transformed) biomass was significantly different between fringing and meadow marshes.

Figure 5. (a) Percent organic matter content of fringing and meadow salt marsh soils. Error bars are ± 1 SE from the mean. Means are significantly different ($p < 0.05$, elevation covariate $p = 0.001$). (b) Relationship between marsh surface elevation and soil percent organic matter content for five fringing and five meadow salt marsh sites. Elevation units are meters above 0' tide.

Figure 6. Amount of sediment deposited on meadow marshes, fringing marshes, and areas where no marsh was present. Error bars are ± 1 SE from the mean. (a) Randomly distributed traps ($p = 0.374$, log transformed data) and (b) Traps placed 1m from the water's edge ($p = 0.120$, log transformed data).

Figure 7. Relationship between the suspended sediment concentration of tidal waters coming onto marsh sites and sediment deposition on (a) randomly distributed traps and (b) traps placed 1 m from the water's edge.

Figure 8. Wave profiles at MRM, a meadow marsh. Values for wave peaks and troughs were taken from videos simultaneously recording the passing waves at (a) 0 m and (b) 5 m along the transect.

Figure 9. Percent reduction in (a) maximum wave height and (b) 'three wave mean' height in fringing, meadow and no marsh areas. Error bars are ± 1 SE from the mean. At 7 m, the difference between marsh and 'no marsh' areas was statistically significant (ANOVA $p < 0.05$, square root transformed data; Student-Neuman-Keuls test, $p < 0.05$).