

University of New England
DUNE: DigitalUNE

Environmental Studies Faculty Publications

Environmental Studies Department

2009

Aggressive Response Of Adult Bobolinks To Neck Ligatures On Nestlings

Lynn P. Little
University of Vermont

Allan M. Strong
University of Vermont

Noah G. Perlut
University of New England, nperlut@une.edu

Follow this and additional works at: http://dune.une.edu/env_facpubs



Part of the [Ornithology Commons](#)

Recommended Citation

Little, Lynn P.; Strong, Allan M.; and Perlut, Noah G., "Aggressive Response Of Adult Bobolinks To Neck Ligatures On Nestlings" (2009). *Environmental Studies Faculty Publications*. Paper 20.
http://dune.une.edu/env_facpubs/20

This Article is brought to you for free and open access by the Environmental Studies Department at DUNE: DigitalUNE. It has been accepted for inclusion in Environmental Studies Faculty Publications by an authorized administrator of DUNE: DigitalUNE. For more information, please contact bkenyon@une.edu.

ACKNOWLEDGMENTS

We are grateful to those who helped rescuing owls and the staff of Wildlife Rehabilitation Center “*La Tahonilla*” (Cabildo Insular de Tenerife). Ricardo Mesa, Juaní Curbelo, and Rubén García provided valuable information about the genus *Setaria* and entanglement observations. G. R. Bortolotti, J. S. Marks, an anonymous reviewer, and C. E. Braun provided useful comments on an early draft of the manuscript.

LITERATURE CITED

- DEL HOYO, J., A. ELLIOTT, AND J. SARGATAL (Editors). 1999. Handbook of the birds of the world. Volume 5. Barn-owls to hummingbirds. Lynx Editions, Barcelona, Spain.
- GLAYRE, D. 1959. Une Choutte effraie capturée pas des graminées. *Nos Oiseaux* 266:122.
- IZQUIERDO, I., J. L. MARTÍN, N. ZURITA, AND M. ARECHAVALETA. 2004. Lista de Especies Silvestres de Canarias (Hongos, Plantas y Animales Terrestres). Consejería de Medio Ambiente y Ordenación Territorial, Gobierno de Canarias, Santa Cruz de Tenerife, Spain.
- MENDELSON, J. M. 1983. Causes of mortality in Black-shouldered Kites. *Bokmakierie* 35:11–13.
- MIKKOLA, H. 1983. Owls of Europe. T. & A.D. Poyser, London, United Kingdom.
- MOLNAR, G. 1996. Ragados muhar (*Setaria verticillata*) mint bagolycsapda. *Tuzok* 1:185.
- NEGRO, J. J., C. PERTOLDI, E. RANDI, J. J. FERRERO, J. M. LÓPEZ-CABALLERO, D. RIVERA, AND E. KORPI-MÄKI. 2006. Convergent evolution of *Elanus* kite and the owls. *Journal of Raptor Research* 40:222–225.
- NOZERAND, R. 1994. Hiboux Moyens-Ducs *Asio otus* emprisonnés dans des graminées. *Alauda* 62:116.
- TAYLOR, I. 1994. Barn Owls. Predator-prey relationships and conservation. Cambridge University Press, Cambridge, United Kingdom.

The Wilson Journal of Ornithology 121(2):441–444, 2009

Aggressive Response of Adult Bobolinks to Neck Ligatures on Nestlings

Lynn P. Little,^{1,2} Allan M. Strong,^{3,5} and Noah G. Perlut⁴

ABSTRACT.—We monitored provisioning behavior at 18 Bobolink (*Dolichonyx oryzivorus*) nests during 240.5 min of videotape data from June to July 2006, and observed 64 nest visits by adults while nestlings were fitted with neck ligatures. Adults pecked or pulled at the ligatures, often aggressively, at 72% of nests ($n = 18$) and 52% of visits ($n = 64$). These behavioral responses by adults indicate the neck ligature technique is more invasive than previously believed. We documented no mortality as a result of ligature placement, but researchers should minimize the time that ligatures are in place to reduce stress to both parents and nestlings. *Received 8 September 2008. Accepted 30 January 2009.*

Quantitative assessments of avian diets may be critical for evaluating habitat quality. However, methods used to quantify avian diets have associated biases and/or shortcomings as most studies require techniques tailored to specific studies and hypotheses (Rosenberg and Cooper 1990). Neck ligatures have been used to quantify diets of nestling birds as this technique allows collection of prey items prior to onset of digestion. Modifications have been suggested to improve ligature function and minimize negative effects on nestlings. For example, Johnson et al. (1980) described abnormal behavior of nestlings after leaving ligatures in place for 1 hr and suggested that collection of prey immediately after each parental visit would minimize biases. Further, Mellott and Woods (1993) found that cable ties simplified ligature placement compared to coated wire, especially when used by untrained personnel.

Neck ligatures are considered an invasive technique (Rosenberg and Cooper 1990, Poulsen and Aebischer 1995), but most studies addressing their effects have focused on the be-

¹ Department of Animal Science, University of Vermont, Burlington, VT 05405, USA.

² Current address: 5060 Bentley Lane, Columbus, OH 43220, USA.

³ Rubenstein School of Environment and Natural Resources, 81 Carrigan Drive, University of Vermont, Burlington, VT 05405, USA.

⁴ Vermont Cooperative Fish and Wildlife Research Unit, The Rubenstein School of Environment and Natural Resources, University of Vermont, Burlington, VT 05405, USA.

⁵ Corresponding author; e-mail: allan.strong@uvm.edu

havioral responses of nestlings. We used video cameras and neck ligatures simultaneously in a study of nestling Bobolink (*Dolichonyx oryzivorus*) diets. We report on the response of adult Bobolinks to placement of neck ligatures on nestlings.

METHODS

Study Site.—Our study was conducted during June–July 2006 in three hayfields in Hinesburg, Shelburne, and Charlotte, Chittenden County, Vermont, USA. Bobolinks breed in hayfields and pastures throughout this agricultural region (Shustack 2004, Perlut et al. 2006) and their ground nests are relatively easy to locate.

Video and Ligature Data Collection.—Nest observations, video monitoring, and ligature sampling occurred between 0400 and 1300 hrs EST during precipitation-free periods. A small, wide-angle “lipstick” lens (www.helmetcamera.com) was mounted 10 cm from the nest when nestlings reached 6 days of age. The lens was attached by cable to an 8-mm camcorder (Sony DCR-TRV460 Digital8 Handycam) placed 2 m from the nest and concealed by vegetation. Recording sessions commenced with an initial 45-min acclimation period during which nestlings were left undisturbed. Recording was paused while neck ligatures were placed on nestlings following Johnson et al. (1980) and subsequent modifications (Mellott and Woods 1993). We placed ligatures on a maximum of three nestlings per nest using plastic cable ties (10 cm length before cutting off excess, 2.5 mm width), temporarily removing any additional nestlings for the remainder of the videotaping period (generally 45 min). Removal of some of the nestlings reduced the number of nestlings that were not fed by parents during a feeding session (adults were not observed feeding all nestlings in a single feeding trip when a nest contained five or six nestlings; N. G. Perlut, unpubl. data) and reduced the number of nestlings that would be exposed to any potential stress during the ligature process. We attempted to remove ligatures and collect diet samples from the nestlings as soon as we observed the adults return to the nest with food. Prolonged use of ligatures can induce abnormal swallowing and gaping behavior in nestlings, which leads to the redistribution of prey

items among nestlings by adults (Johnson et al. 1980).

RESULTS

We monitored 18 Bobolink nests during 240.5 min of videotape data. We applied neck ligatures to 50 nestlings, and successfully collected 99 prey items from 28 of those nestlings. No nestling mortality occurred while nests were videotaped. Leaf hoppers (27.3% based on numerical abundance), holometabolous larvae (24.2%), grasshoppers (16.2%), mayflies (8.1%), spiders (7.1%), and moths (5.1%) were the most common prey types fed to nestlings. Mean \pm SD prey length was 14.5 \pm 10.2 mm with 17% of prey items delivered >25 mm. Sixteen of the 18 nests were successful with 56 nestlings fledged.

We observed 64 nest visits by adults while nestlings were fitted with ligatures. Seventy-two percent of visits were by females and 28% by males. On average, females and males visited nests 2.6 and 1.0 times, respectively while ligatures were in place (~45 min). Adults pecked at ligatures of nestlings in 13 of 18 nests (72%) and in 33 of 64 (52%) visits. They directed pecks at more than one nestling in 70% of the visits during which adults pecked at the ligatures. Females had a greater propensity to peck at ligatures with 59% of all female visits eliciting pecks to ligatures compared to 33% of all male visits. The first adult to return to the nest pecked at the ligatures in 12 of 13 nests in which adults pecked at ligatures. There was no indication that probability of pecking changed across the time period ligatures were in place (logistic regression, $\chi^2 = 1.68$, $df = 1$, $P = 0.20$).

Adults generally pecked at the thin band of the cable tie. In these cases, the adults appeared to be assessing whether or not the ligature could be easily removed from the chicks. Adults also grasped and pulled at the bulkier locking mechanism of the cable tie with the appearance that removal was the goal. Parents were notably aggressive in about half of the nests in their attempts to remove the ligatures, grasping the ligature and forcibly pulling the nestlings' heads upwards or sideways. In one instance, a female inspected, grasped, and pulled at a nestling's ligature for 30 sec. No aggressive actions toward nestlings were noted when ligatures were not present

on nestlings based on video data prior to ligature placement and a few tapes which we left running after the ligature sessions. One female visually inspected her nestling's throats upon her first visit to the nest ~20 min after removal of the ligatures.

DISCUSSION

Use of neck ligatures has been shown to provide quality dietary data for nestling birds (Orians 1966, Martin et al. 2000, Clotfelter et al. 2007). Our data support these results as we were able to collect a relatively large sample of prey items over a short period of time. Prey length data suggest that adults did not adjust the size of prey items brought to nestlings with ligatures. However, we do not have adequate control data (nestlings without ligatures) for a quantitative comparison.

Our data show that undesirable behavioral responses to ligatures are not restricted to nestlings. The strong and consistent response of adults to ligatures suggests this technique elicits stress that was not previously documented. Attempts by female Red-winged Blackbirds (*Agelaius phoeniceus*) to remove pipe cleaner ligatures have been documented, but no behavioral data were provided (Robertson 1966). Our results indicate that Bobolinks are strongly attuned to the appearance of their nestlings and, in most cases, neck ligatures trigger a response upon the first visit to the nest. Females in our sample were more likely to attempt to remove ligatures than males (but also made more nest visits). Approximately half of the attempts to remove the ligatures were aggressive with parents lifting and/or dragging nestlings by the ligature. These responses indicate the neck ligature technique is more invasive than previously believed. Comparable data from other species would be useful to better address the generality of our findings.

Gaunt and Oring (1999) noted the potential for changes in blood circulation, tracheal function, and food delivery rates with use of neck ligatures. Our data suggest there may be additional stress to nestlings beyond physical placement of the ligatures. The majority of nest visits by adults included pecking or pulling at neck ligatures and there was no indication that adults became habituated to the presence of ligatures. Neck ligatures remain a

relatively safe, inexpensive, and informative method, but do present welfare concerns. Investigation into the efficacy of less-invasive alternatives such as videography should be conducted, especially when working with rare and declining species. We documented no mortality as a result of ligature placement, but researchers should minimize the time that ligatures are in place to reduce stress to both parents and nestlings.

ACKNOWLEDGMENTS

This research was conducted under University of Vermont Institutional Animal Care and Use Committee protocol 05-039. The project was supported by the National Research Initiative of the USDA Cooperative State Research, Education and Extension Service (grant number 03-35101-13817) and the Natural Resource Conservation Service's Wildlife Habitat Management Institute. Supplies and summer support to LPL were provided by the University of Vermont's Hughes Endeavor for Life Science Excellence (HELIX) program. Additional funding was provided by University of Vermont's College of Agriculture and Life Science. We thank Shelburne Farms and Sam Dixon, and the Ross and Thibault families for allowing us to conduct research on their property. Field assistance was provided by Thomas Lawrence, Stacey Thompson, Roger Masse, Christopher Lang, and Zachary Rowe. T. B. McFadden, R. C. Hovey, F.-Q. Zhao, and the Strong Graduate Laboratory provided support and comments on earlier versions of the manuscript.

LITERATURE CITED

- CLOTFELTER, E. D., C. R. CHANDLER, V. NOLAN JR., AND E. D. KETTERSON. 2007. The influence of exogenous testosterone on the dynamics of nestling provisioning in Dark-eyed Juncos. *Ethology* 113: 18–25.
- GAUNT, A. S. AND L. W. ORING. 1999. Guidelines to the use of wild birds in research. Second Edition. The Ornithological Council, Washington, D.C., USA.
- JOHNSON, E. J., L. B. BEST, AND P. A. HEAGY. 1980. Food sampling biases associated with the "ligature method". *Condor* 82:186–192.
- MARTIN, P. A., D. L. JOHNSON, D. J. FORSYTH, AND B. D. HILL. 2000. Effects of two grasshopper control insecticides on food resources and reproductive success of two grassland songbirds. *Environmental Toxicology and Chemistry* 19:2987–2996.
- MELLOTT, R. S. AND P. E. WOODS. 1993. An improved ligature technique for dietary sampling in nestling birds. *Journal of Field Ornithology* 64:205–210.
- ORIAN, G. H. 1966. Food of nestling Yellow-headed Blackbirds, Cariboo Parklands, British Columbia. *Condor* 68:321–337.
- PERLUT, N. G., A. M. STRONG, T. M. DONOVAN, AND

- N. J. BUCKLEY. 2006. Grassland songbirds in a dynamic management landscape: behavioral responses and management strategies. *Ecological Applications* 16:2235–2247.
- POULSEN, J. G. AND N. J. AEBISCHER. 1995. Quantitative comparison of two methods of assessing diet of nestling Skylarks (*Alauda arvensis*). *Auk* 112: 1070–1073.
- ROBERTSON, R. J. 1973. Optimal niche space of the Red-winged Blackbird. III. Growth rate and food of nestlings in marsh and upland habitat. *Wilson Bulletin* 85:209–222.
- ROSENBERG, K. V. AND R. J. COOPER. 1990. Approaches to avian diet analysis. *Studies in Avian Biology* 13:80–90.
- SHUSTACK, D. P. 2004. Bobolink and Savannah Sparrow habitat selection in the Champlain Valley. Thesis. University of Vermont, Burlington, USA.