

2-12-2017

Utilization Of Postural Control Training To Improve Gait Symmetry And Walking Ability In A Patient Following A Lacunar Stroke: A Case Report

Hannah C. Wilder
University of New England

Follow this and additional works at: http://dune.une.edu/pt_studcrpaper

 Part of the [Physical Therapy Commons](#)

© 2017 Hannah Wilder

Recommended Citation

Wilder, Hannah C., "Utilization Of Postural Control Training To Improve Gait Symmetry And Walking Ability In A Patient Following A Lacunar Stroke: A Case Report" (2017). *Case Report Papers*. 69.
http://dune.une.edu/pt_studcrpaper/69

This Course Paper is brought to you for free and open access by the Physical Therapy Student Papers at DUNE: DigitalUNE. It has been accepted for inclusion in Case Report Papers by an authorized administrator of DUNE: DigitalUNE. For more information, please contact bkenyon@une.edu.

20 **ABSTRACT**

21 **Background and Purpose**

22 Stroke affects approximately 800,000 people annually and alterations in gait is one of the most
23 noted impairments following stroke. The purpose of this case report is to outline physical therapy (PT)
24 rehabilitation that utilized postural control training, task-oriented training, and visual feedback to address
25 walking ability and functional capacity in a patient following a stroke.

26 **Case Description**

27 The patient was a 67-year-old male apple orchard owner three months post a lacunar ischemic
28 stroke affecting the posterior limb of the internal capsule, the basal ganglia, and part of the cerebellum.
29 His initial examination revealed impaired strength, sensation, range of motion, balance, endurance, and
30 mobility. This case report describes his initial ten outpatient PT visits primarily focused on improving the
31 patient's functional mobility, ambulation in particular, through postural control training and task-oriented
32 training.

33 **Outcomes**

34 After ten outpatient visits, the patient demonstrated improvements in gait and postural symmetry
35 on observation. Improvements were shown in both gait speed (from 0.24 m/s to 0.30 m/s) and gait
36 endurance (from feet to 130 feet) and although the improvements did not meet established minimally
37 important clinical difference values, he did demonstrate trends toward improvement.

38 **Discussion**

39 Postural control training and task oriented training are common PT interventions utilized in
40 patients following stroke. Utilizing postural control training and task-oriented training, the patient showed
41 initial improvements in postural symmetry and gait mechanics, which translated to improved access to his
42 environment. Despite initial improvements, the patient's various comorbidities likely contributed to his
43 plateau in progress. Future research on lacunar stroke should address how comorbidities affect the
44 acquisition of PT goals and improvements in gait speed.

45 Manuscript word count: 3,547 words

46 **BACKGROUND AND PURPOSE**

47 Stroke affects approximately 800,000 people annually in the United States, and survivors of stroke
48 report persistent difficulties with daily tasks as a direct consequence of stroke.¹ Furthermore, alterations in
49 gait is one of the most noted impairments following stroke, and improving walking ability is one of the
50 most common goals amongst patients with stroke undergoing rehabilitation.² Of the common gait
51 deviations observed following stroke, decreased gait speed is a significant limitation as a result of stroke.³

52 A case report by Lewek³ addressed the effectiveness of visual and proprioceptive feedback to
53 improve gait speed in two patients with chronic stroke. Improving mobility was the goal for the
54 intervention, so intensive gait training was performed by either overground or treadmill training. During
55 gait training, visual and proprioceptive feedback was given throughout. Following six weeks of
56 intervention, both patients had improved gait speed as well as spatiotemporal symmetry with gait
57 mechanics.

58 A systematic review by Eng and Tang² examined 39 randomized controlled trials (RCTs) assessing
59 different rehabilitation strategies for improving walking ability in people following stroke. The
60 interventions analyzed included neurodevelopmental training, strength training, treadmill training, and
61 task-specific training. Among these interventions, task-oriented training showed highly compelling
62 evidence of effectiveness in improving walking ability. These task-oriented training programs focused not
63 only on walking, but also on a broad array of other functional mobility tasks. They were shown to be
64 effective in improving walking ability following stroke, as well as including further benefits of functional
65 strengthening, balance improvements, and cardiorespiratory benefits.

66 Salbach⁴ used a RCT to further evaluate the effectiveness of task-oriented training on improving
67 walking speed and distance in patients following stroke. In the experimental group (n=44), ten functional
68 tasks were utilized that were hypothesized to improve lower extremity (LE) strength and subjects'
69 walking speed, distance, and balance. The control group (n=47) performed only upper extremity
70 activities. Following intervention, significant differences were shown between the experimental and
71 control groups in both walking speed ($p < 0.05$) and distance ($p < 0.05$) in favor of task-oriented training.

72 Andersson and Franzen⁵ evaluated weight-shift training and its ability to improve walking ability in
73 patients with chronic stroke (n=10). They hypothesized that gait abnormalities were largely a result of
74 asymmetry due to hemiparesis following stroke. By training patients to shift weight toward their involved
75 sides, they believed patients would have greater postural control, more symmetrical gait mechanics, and
76 improved walking ability. Following three weeks of weight-shift training, greater spatial and temporal
77 symmetry of gait and increased gait speed (p=0.037) was noted.

78 Based on the findings in the literature, visual and proprioceptive feedback are effective in improving
79 gait speed and mechanics following stroke.³ Additionally, task-oriented training has been shown to be
80 effective in improving walking ability, strength, and balance.^{2,4} Finally, based on the positive results
81 demonstrated with weight-shift training, postural control training may similarly improve walking ability
82 following stroke. The purpose of this case report is to outline PT rehabilitation that utilized postural
83 control training, task-oriented training, and visual feedback to address walking ability and functional
84 capacity in a patient following a stroke.

85

86 **CASE DESCRIPTION**

87 **Patient History and Systems Review**

88 The patient (JT) gave written consent to participate in this case report. JT was a 67-year-old male
89 referred to outpatient PT following a lacunar ischemic stroke affecting the posterior limb of the internal
90 capsule, the basal ganglia, and part of the cerebellum approximately three months prior. This case report
91 is based on the patient's first ten visits which comprise his initial phase of outpatient PT. JT was the
92 owner of a hotel and apple orchard and was very involved in his community. He lived in a two story
93 home with approximately four stairs to enter, but had not been able to access the second floor of his home
94 following the stroke. He had a very supportive wife who was highly involved in his rehabilitation and was
95 present for nearly every PT session. JT was unable to drive so his family drove him to therapy.

96 His stroke was a result of a blood clot that formed due to atrial fibrillation, which caused ischemia in
97 the brain. Following his stroke, he had an acute stay in the hospital for one week, and was then transferred

98 to a subacute rehabilitation facility where he stayed for 30 days. Following discharge, he returned home
99 and received home PT services for five weeks prior to initiating outpatient PT.

100 As a result of his stroke, he had hemiparesis of his dominant left side upper and lower extremities.
101 Upon his initial evaluation in outpatient therapy, his chief complaints included decreased endurance, and
102 pain in the left shoulder and left lower extremity (LLE) with ambulation. JT's left shoulder pain was a
103 result of chronic subluxation of the glenohumeral joint due to flaccid paralysis of his left upper extremity
104 (LUE), and his LLE pain was a result of intermittent claudication. The patient used a manual wheelchair
105 with left arm trough, and a wide base quad cane for ambulation. He also wore an ankle foot orthosis
106 (AFO) on his LLE, and a sling on his left upper extremity (LUE) for ambulation. In addition to his stroke,
107 the patient also had a history of high cholesterol, hypertension, peripheral artery disease, and atrial
108 fibrillation. Please see Table 1 for results of his systems review and Appendix 1 for his medication list. In
109 addition to outpatient PT, the patient was also receiving outpatient occupational therapy (OT), which
110 focused primarily on his LUE function.

111 JT's primary goal for PT was to improve his mobility and endurance to be more independent and less
112 reliant on his wheelchair.

113

114 **Clinical Impression 1**

115 Following a review of JT's history, his primary problems included impairments in strength,
116 sensation, range of motion (ROM), balance, endurance, and mobility as a result of an ischemic stroke.
117 Abnormal reflexes, clonus, and spasticity were also suspected. There were no differential diagnoses to be
118 addressed upon initial evaluation. Planned tests and measures were: the Fugl-Meyer assessment for the
119 LE, the Activities Specific Balance Confidence (ABC) scale, the Stroke Impact Scale (SIS-16), the
120 Disabilities of the Arm, Shoulder, and Hand (DASH) questionnaire (performed by OT), the Six Minute
121 Walk Test (6MWT), the Ten Meter Walk Test (10MWT), and the Five Times Sit to Stand Test (FTSST
122 or 5XSST). The patient remained an appropriate candidate for a case report because he was highly
123 motivated to improve his walking ability to return to work and enhance his function.

124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149

Examination – Tests and Measures

Tests and measures performed during the PT initial evaluation focused primarily on gait, balance, and LE function. Tone, balance, sensation, and ROM were grossly assessed through observation and patient report. Since the patient was also receiving OT, the majority of his UE tests and measures were performed by the occupational therapist. While in the waiting area prior to the start of the PT evaluation, JT was given the ABC, the SIS-16, and the DASH. The ABC is a self-reported measure of balance confidence.⁶ The SIS-16 is a shortened version of the Stroke Impact Scale and is used as a subjective measure to assess stroke-related physical disability.⁷ The DASH is a self-reported questionnaire of disability related to UE function.⁸

From the waiting area to the treatment room, the patient self-propelled in his manual wheelchair using his right upper and lower extremities. While in his wheelchair, a gross postural assessment was performed in sitting. The patient performed a stand pivot transfer from the wheelchair to a chair with supervision to assess his functional mobility and independence. Following the patient history, further tests and measures were done to obtain objective measures of neuromuscular, cardiopulmonary, and musculoskeletal system functions (see Table 1).

The 5XSST was performed to assess JT’s LE strength and function. It was used as a clinical assessment tool representing the body function domain of the International Classification of Functioning, Disability, and Health (ICF) model.⁹ The Fugl-Meyer Assessment of Motor Recovery after Stroke has both upper and LE scales to assess the motor recovery in patients who have had strokes with hemiplegia and hemiparesis.⁹ The LE scale of the Fugl-Meyer was completed with JT to assess his motor function, reflex integrity, coordination, and his ability to perform movements out of synergy. The Fugl-Meyer also represents the body function domain of the ICF model.

The 10MWT and 6MWT were used to assess walking ability and can be associated with both the activity and participation domains of the ICF model.^{9,10} The 10MWT was specifically used to measure gait speed. Moreover, there is predictability in a patient’s ability to be a home vs. community ambulator

150 based on his self-paced gait speed as determined by the 10MWT.¹⁰ The 6MWT was used to measure gait
151 endurance. Cardiovascular deconditioning can occur following stroke due to increased energy demands
152 from an inefficient gait pattern, making performing activities of daily living (ADLs) increasingly taxing.¹¹
153 Additionally, one of the patient's major complaints following his stroke was his limited endurance with
154 ambulation, and it has been shown that gait endurance is one of the most challenging areas for patients
155 following stroke.¹² Psychometric properties for tests and measures can be seen in Table 3.

156

157 **Clinical Impression 2**

158 The initial clinical impression was both confirmed, and rejected based on the examination data. It
159 was originally hypothesized that JT would present with impaired sensation as a result of his stroke;
160 however, sensation was grossly intact. He did, however, present with weakness, limited ROM, impaired
161 balance and endurance, abnormal reflexes, clonus, and spasticity which were all consistent with the initial
162 clinical impression.

163 Based on the patient's performance with his affected LLE on the Fugl-Meyer, it was determined
164 that despite the extensor synergy pattern, he was able to coordinate movements of the left leg out of
165 synergy, such as performing hip extension with knee flexion. It was also determined with the Fugl-Meyer,
166 that JT's impairments were predominantly found in the distal joints compared to the proximal joints,
167 affecting coordination and terminal accuracy of movements.

168 With the 10MWT, it was determined that the patient's gait speed was 0.24 meters per second
169 (m/s). Eng and Tang² report that patients with stroke who can ambulate with a speed of 0.4 m/s are more
170 likely to be able to be community ambulators. Based on this data, it was reinforced that improving JT's
171 gait speed would be a major focus of PT in order to improve his function and increase his participation in
172 the community. With JT's performance on the 6MWT, it was also reinforced that his endurance was
173 impaired as a result of his stroke. He ambulated 85 feet (25.91 meters) in six minutes, which is below
174 healthy age-matched norms (see Table 2). The patient continued to be appropriate for PT and this case
175 study to further develop the body of literature on postural control training for patients with lacunar

176 strokes, so the decision was made to continue with therapy.

177 Based on JT's medical history and the results of tests and measures, two rehabilitation ICD-10
178 codes were chosen. The primary ICD-10 code was I69.952, *hemiplegia and hemiparesis following*
179 *unspecified cerebrovascular disease affecting left dominant side*, and the secondary code chosen was
180 R26.9, *unspecified abnormalities of gait and mobility*.

181 There were several factors noted during the initial evaluation that were predicted to positively
182 affect the patient's prognosis. Prior to his stroke, JT was very active and performed jobs that required
183 high levels of mobility. He was also highly motivated and committed to the process of his recovery
184 through PT, and his family was very supportive. Based on his performance on the Fugl-Meyer, it was
185 concluded that the patient had the ability to move out of patterns of synergy, which indicated a greater
186 ability to make functional improvements. It has been shown that improvements in scoring on the Fugl-
187 Meyer are moderately correlated with improvements in gait speed in patients with stroke.² Additionally,
188 his stroke had occurred approximately three months prior to the initial outpatient PT evaluation, placing
189 him in the three to six-month window where the greatest recovery following stroke has been shown to
190 occur.¹⁴

191 In addition to these positive prognostic indicators, there were also some negative indicators that
192 needed to be considered in regards to JT's prognosis. His history of atrial fibrillation placed him at risk
193 for a second stroke, as well as other cardiovascular and pulmonary complications. He also had complaints
194 of pain in his left shoulder and LLE, which had the potential to limit his tolerance for activity. Based on
195 these prognostic indicators, JT's prognosis was determined to be fair.

196 No additional referrals were needed, but coordination and consultation with the patient's OT were
197 planned to take place as needed. It was also decided that additional testing would be performed as deemed
198 necessary, and reassessments of outcome measures would take place monthly, or at every tenth visit to
199 meet Medicare guidelines.

200 An intervention plan was developed following the patient's initial PT evaluation with the primary
201 focus on improving postural control in order to increase functional mobility. It was also determined that

202 there should be an emphasis on motor control training, and gait training to improve endurance. Eng and
203 Tang² recognize that walking endurance is an important factor that can help to predict the ability for
204 community reintegration in patients with stroke. Since decreasing dependence on his wheelchair and
205 returning to work were JT's primary goals for PT, these were important factors that helped to drive
206 decision-making for JT's therapeutic interventions. Additional PT goals were established and can be seen
207 in Appendix 2.

208

209 **Intervention**

210 Coordination and communication of the patient's care occurred on a weekly basis with his
211 occupational therapist (OT). Progress notes were sent to JT's primary care physician every eighth to tenth
212 visit to give updates on any changes in his care and progress he had made. Sixty-minute PT sessions
213 occurred three times weekly for ten weeks. The initial four weeks, or ten visits of outpatient PT
214 interventions, are further described. Notes were documented in the electronic medical record system for
215 the patient's treatment sessions as well as progress notes when necessary.

216 Patient-related instruction included patient education on the PT plan of care (POC), the
217 importance of proper use of his manual wheelchair and arm trough, the typical disease process of stroke
218 and the recovery timeline, and current evidence for improvements in gait following stroke. The patient
219 and his wife were also updated on his improvements and the clinical significance of gains achieved. JT
220 attended all scheduled PT sessions and was very compliant with his home program. At the start of every
221 PT session he and his wife gave updates on his improved ability to perform ADL's.

222 PT interventions primarily focused on improving JT's functional mobility, ambulation in
223 particular, through postural control training and task-oriented training. Andersson and Franzen⁵ found that
224 subjects with chronic stroke who underwent training to shift weight toward their hemiparetic side, had
225 improved gait and ambulation. JT's therapists hypothesized that by increasing weight bearing on his
226 hemiparetic L side in various positions, including sitting, standing, kneeling, and half kneeling, he would
227 be able to perform a more efficient gait pattern. Additionally, it has been shown that visual feedback with

228 postural control training can improve gait speed in individuals with stroke.³ Therefore, postural control
229 training with JT also involved visual feedback with a mirror, in addition to verbal and tactile feedback
230 provided by therapists. Task-oriented training involved functional tasks, such as sit-to-stand, quadruped to
231 half-kneeling, and stair climbing. Task-oriented training has been shown to have many benefits in
232 improving walking ability and functional task performance following stroke.^{2,4}

233 In addition to postural control training, and task-oriented training, other PT interventions were
234 implemented to improve walking ability and functional capacity, including therapeutic exercise (ther ex)
235 and neuromuscular re-education (NMR). See Table 4 for interventions. Ther ex interventions involved
236 posterior shoulder and parascapular muscle strengthening to combat shoulder subluxation in the patient's
237 LUE, and to prevent overuse injury of his RUE. Ther ex and NMR interventions were both geared toward
238 inhibiting JT's LLE tone through positioning and activation of antagonist muscles, including the hip
239 extensors, hip abductors, knee flexors, and ankle dorsiflexors.¹⁵ With improved activation and
240 strengthening of these muscles, it was hypothesized that JT would have a greater ability to coordinate and
241 achieve L foot clearance during the swing phase of gait. NMR also included interventions aimed at
242 improving JT's balance to enhance walking ability and decrease risk for falls.

243 Gait training interventions progressed over time throughout JT's POC (see Table 4). As he
244 progressed, the ambulation distance increased, and the amount of rest time decreased. Additionally, the
245 amount of support from the therapist and assistive device (AD) decreased. Early gait training utilized a
246 hemi walker in the hopes JT's gait speed would increase using a more supportive AD. Gait training was
247 then progressed to a less supportive wide base quad cane (WBQC), and TheraBand (TheraBand, Akron,
248 OH) wrap to the ankle, knee, and hip for assistance with LLE mechanics and foot clearance (see Figure
249 1). As the patient's gait continued to progress, AD use was discontinued in order to train proper
250 unsupported gait mechanics. O'Sullivan¹⁵ notes that although AD's can improve stability during gait early
251 on, they should be discontinued in patients who may be able to walk without them as they can inhibit
252 balance reactions and proper gait mechanics.

253 Interventions continued both with and without AD use depending on the desired therapeutic

254 response to the intervention. When the intervention was geared toward increasing gait speed and
255 endurance, an AD was utilized; when the purpose was improving gait mechanics, the patient ambulated
256 with therapist assistance only. It was planned as JT progressed in his ability to ambulate, use of external
257 supports would be discontinued. It was also planned that gait and neuromuscular reeducation
258 interventions would be progressed according to Gentile's Taxonomy of Tasks (see Appendix 4).¹⁶ This
259 allowed progression of therapeutic interventions as JT's motor learning and motor control improved by
260 manipulating either the environmental conditions, or the desired outcomes of the task to increase
261 complexity and demand.¹⁶

262

263 **OUTCOME**

264 PT interventions primarily focused on improving the patient's functional mobility, ambulation in
265 particular, through postural control training and task-oriented training. After ten outpatient visits, JT
266 demonstrated improvements in gait and postural symmetry on therapist observation. He showed improved
267 ability to function out of synergistic movement patterns and isolate single-joint movements in the
268 hemiparetic LLE. This resulted in improved step clearance which allowed the patient greater access to his
269 yard, apple orchard, and to operate his tractor. Improvements were shown in both gait speed and gait
270 endurance as measured by the 10 meter walk test and six minute walk test respectively. Although the
271 improvements were not deemed clinically meaningful based on established minimally important clinical
272 difference values, he did demonstrate trends toward improvement.^{12,13} However, with PT intervention,
273 the patient's improvements in gait allowed him to safely access his yard and orchards at home, advancing
274 him from a limited household ambulator to an unlimited household ambulator.¹⁷ Results of outcome
275 measures taken during week four can be seen in Table 2, and status of short and long-term goals
276 following four weeks of PT can be seen in Appendix 2.

277

278 **DISCUSSION**

279 Following ten visits of outpatient PT, the patient demonstrated improvements in mobility and

280 function in tasks that were important to him. Based on his self-reports, he perceived a greater ability to
281 negotiate stairs and access his tractor to mow his apple orchard. However, this did not correlate with
282 achievement of predetermined PT established goals. Although the patient's gait speed improved, and he
283 achieved his personal goal of being able to ride his tractor again, the patient's various comorbidities likely
284 contributed to his plateau in progress. With complex patient cases such as this, it is important to recognize
285 functional achievements, as they are meaningful.

286 This POC may be beneficial when applied to other patients with a similar presentation; however,
287 further investigation is warranted. A limitation to this case report is the short time frame of ten visits, and
288 although initial progress was seen, it is unclear what the patient's potential for further progress may have
289 been. JT showed initial improvements in postural symmetry and gait mechanics, however, it is unknown
290 whether these improvements remained over time. A long-term follow-up study would provide further
291 information on maintenance of these achievements in this patient population and this type of POC's
292 overall effectiveness through measurement of gait speed, endurance, and functional benchmarks over time
293 on a larger scale.

294 A clear evaluation of the plan's effectiveness was also limited due to the patient's various
295 comorbidities throughout PT intervention. Pain, for instance, was a confounding variable to patient
296 progress. The patient had pain in his LUE due to chronic shoulder subluxation, pain in his RUE due to
297 impingement as a result of overuse, and pain in his LLE due to claudication. Limitations in PT visits, and
298 the need to show progress for insurance purposes, made striking a balance between PT interventions for
299 functional mobility and pain management challenging. Further research on lacunar stroke should address
300 how comorbidities such as shoulder subluxation, impingement syndrome, and claudication, may affect the
301 achievement of PT goals as well as opportunities to improve gait speed. Co-morbidities such as these also
302 affect a patient's ability to appropriately use an AD, further limiting potential gait speed improvements.
303 Additionally, greater frequency of visits would allow targeted therapy to address both pain modulation
304 and functional mobility goals in order to reveal the patient's true potential for recovery.

305

306 **REFERENCES**

- 307 1) Winstein CJ, Stein J, Arena R, Bates B, Cherney LR, Cramer SC. Guidelines for Adult Stroke
308 Rehabilitation and Recovery: A Guideline for Healthcare Professionals from the American Heart
309 Association/American Stroke Association. *Stroke*. 2016; 47(6):98-169. doi:
310 10.1161/STR.0000000000000098. Accessed September 22, 2016.
311
- 312 2) Eng JJ, Tang PF. Gait training strategies to optimize walking ability in people with stroke: a
313 synthesis of the evidence. *Expert Rev Neurother*. 2007;7(10):1417-1436.
314 doi:10.1586/14737175.7.10.1417. Accessed June 26, 2016.
315
- 316 3) Lewek MD, Feasel J, Wentz E, Brooks FP, Whitton MC. Use of visual and proprioceptive
317 feedback to improve gait speed and spatiotemporal symmetry following chronic stroke: a case
318 series. *Phys Ther*. 2012;92(5):748-756. doi:10.2522/ptj.20110206. Accessed June 15, 2016.
319
- 320 4) Salbach NM, Mayo NE, Wood-Dauphinee S, Hanley JA, Richards CL, Côté R. A task-oriented
321 intervention enhances walking distance and speed in the first year post stroke: a randomized
322 controlled trial. *Clin Rehabil*. 2004;18:509-519. doi:10.119/0269215504cr763oa. Accessed June
323 27, 2016.
324
- 325 5) Andersson P, Franzen E. Effects of weight-shift training on walking ability, ambulation, and
326 weight distribution in individuals with chronic stroke: a pilot study. *Top Stroke Rehabil*.
327 2015;22(6):437-43. doi: 10.1179/1074935715Z.00000000052. Accessed July 9, 2016.
328
- 329 6) Botner EM, Miller WC, Eng JJ. Measurement properties of the Activities-Specific Balance
330 Confidence Scale among individuals with stroke. *Disabil Rehabil*. 2005; 27(4):156-63. doi:
331 10.1080/09638280400008982. Accessed June 30, 2016.
332
- 333 7) Duncan PW, Lai SM, Bode RK, Perera S, DeRosa J. Stroke impact scale-16: A brief
334 assessment of physical function. *Neurology*. 2003;60(2):291-6. doi:
335 10.1097/01.NPT.0000284780.19341.c9
336
- 337 8) Beaton DE, Katz JN, Fossel AH, Wright JG, Tarasuk V, Bombardier C. Measuring the whole or
338 the parts? Validity, reliability, and responsiveness of the Disabilities of the Arm, Shoulder and
339 Hand outcome measure in different regions of the upper extremity. *J Hand Ther*. 2001;14(2):128-
340 46. <http://www.ncbi.nlm.nih.gov/une.idm.oclc.org/pubmed/11382253>. Accessed June 30, 2016.
341
- 342 9) Beninato M, Portney LG, Sullivan PE. Using the International Classification of Functioning,
343 Disability and Health as a framework to examine the association between falls and clinical
344 assessment tools in people with stroke. *Phys Ther*. 2009;89(8):816-825. doi:
345 10.2522/ptj.20080160. Accessed June 30, 2016.
346
- 347 10) Tyson S, Connell L. The psychometric properties and clinical utility of measures of walking and
348 mobility in neurologic conditions: a systematic review. *Clin Rehabil*. 2009;23(11):1018-33. doi:
349 10.1177/0269215509339004. Accessed June 30, 2016.
350
- 351 11) Ivey FM, Macko RF, Ryan AS, Hafer-Macko CE. Cardiovascular health and fitness after stroke.
352 *Top Stroke Rehabil*. 2005;12(1):1-16. doi: 10.1310/GEEU-YRUY-VJ72-LEAR. Accessed June
353 30, 2016.
354
355

356 12) Tang A, Eng JJ, Rand D. Relationship between perceived and measured changes in walking after
357 stroke. *J Neurol Phys Ther.* 2012;36(3):115-21. doi: 10.1097/NPT.0b013e318262dbd0. Accessed
358 June 30, 2016.
359

360 13) Rehab Measures: Fuel-Meyer Assessment, Activities-Specific Balance Confidence Scale,
361 Disabilities of the Arm, Shoulder, and Hand, Six Minute Walk Test, Ten Meter Walk Test, Five
362 Times Sit to Stand. Rehab Measures website. Available at:
363 <http://www.rehabmeasures.org/rehabweb/allmeasures.aspx?PageView=Shared>. Accessed July 24,
364 2016.
365

366 14) Dashe J. Stroke Prognosis in Adults. *UpToDate*, 2014:14086.
367 <http://www.uptodate.com/contents/ischemic-stroke-prognosis-in-adults>. Accessed on July
368 5, 2016.
369

370 15) O’Sullivan SB. Stroke. In: O’Sullivan SB, Schmitz TJ, Fulk GD, eds. *Physical Rehabilitation*.
371 Philadelphia, PA: F.A. Davis; 2010:645-720.
372

373 16) Cavanaugh J. Functional Task Analysis. Lecture presented: Physical Therapy Management of
374 Adults with Disorders of the Neuromuscular System at the University of New England; January
375 29, 2016; Portland, ME.
376

377 17) Joa KL, Kwon SY, Choi JW, Hong SE, Kim CH, Jung HY. Classification of walking ability of
378 household walkers versus community walkers based on K-BBS, gait velocity and upright motor
379 control. *Eur J Phys Rehabil Med.* 2015;51(5):619-25. Available at:
380 [http://www.minervamedica.it/en/journals/europa-](http://www.minervamedica.it/en/journals/europa-medicophysica/article.php?cod=R33Y2015N05A0619)
381 [medicophysica/article.php?cod=R33Y2015N05A0619](http://www.minervamedica.it/en/journals/europa-medicophysica/article.php?cod=R33Y2015N05A0619). Accessed September 18, 2016.
382
383

384
385
386
387
388
389
390
391
392
393
394

395 **TABLES and FIGURES**

396 Table 1. Systems Review upon initial evaluation

Systems Review	
Cardiovascular/Pulmonary	Impaired: diagnosed with atrial fibrillation, history of peripheral artery disease (PAD) and hypertension
Musculoskeletal	Impaired: left shoulder separation, impaired range of motion of left hip and ankle
Neuromuscular	Impaired: Tone: increased tone in left upper and lower extremities, flexor spastic pattern of left upper extremity Posture: weight shifted to the right in sitting and standing Sensation: grossly intact Balance: decreased due to right weight shift
Integumentary	Unimpaired: rigid AFO on left lower extremity, sling on left upper extremity
Communication	Unimpaired
Affect, Cognition, Language, Learning Style	Unimpaired: preferred explanations and demonstrations for learning

397

398 Table 2. Tests and Measures

Tests & Measures	Age-Matched Norms	Initial Evaluation Results	Results at Week 4
Fugl-Meyer Assessment- Lower Extremity Scale	Not established	Affected: 20/34 Unaffected: 34/34	Not tested
Activities-Specific Balance Confidence (ABC) Scale	79.89/100 ¹³	32.5/100	Not tested
Stroke Impact Scale-16 (SIS-16)	Not established	54/100	Not tested
Disabilities of the Arm, Shoulder, and Hand (DASH)	Not established	70.7/100	Not tested
Six Minute Walk Test (6MWT)	527 m ¹³ (1,729 ft)	85 ft in 4:52 with WBQC too fatigued to continue	130 ft with hemi walker
Ten Meter Walk Test (10MWT)	1.36 m/s ¹³	0.24 m/s	0.30 m/s
Five Times Sit to Stand (5xSTS)	8.1 ± 3.1 seconds ¹³	15.3 seconds with use of right upper extremity	Not tested

399 m=meters, ft=feet, WBQC=wide base quad cane, m/s=meters per second

400

401

402

403

404 Table 3. Psychometric Properties of Tests and Measures

Tests & Measures	Psychometric Properties
Fugl-Meyer Assessment	Excellent test-retest and inter-tester reliability. ⁹
Activities-Specific Balance Confidence (ABC) Scale	High internal consistency and test-retest reliability in individuals with stroke. ⁶
Stroke Impact Scale-16 (SIS-16)	Good instrument reliability and concurrent validity. ⁷
Disabilities of the Arm, Shoulder, and Hand (DASH)	Evidence in the stroke population is limited, but it has been shown to be reliable and responsive for impairments of the entire upper extremity in adults with UE musculoskeletal impairments. ⁸
Six Minute Walk Test (6MWT)	Tested in stroke populations and has a strong predictability for community-based outcomes. ¹⁰
Ten Meter Walk Test (10MWT)	High predictive validity and excellent correlation with level of dependence in instrumental activities of daily living (IADLs) in patients with stroke. ¹⁰
Five Times Sit to Stand (5xSTS)	Excellent validity in measuring affected and unaffected lower extremity strength in patients with stroke. ⁹

405

406

407 Table 4. Interventions

	Week 1 Interventions Treatments 1-2	Week 2 Interventions Treatments 3-4	Week 3 Interventions Treatments 5-7	Week 4 Interventions Treatments 8-10
Gait training	Step ups on 4" step B/L, 5x3	Ambulation with hemi walker (on R) to increase speed and step length, 3x25 ft, min assist x1 Stepping in parallel bars tapping Airex* mat behind and stepping to a piece of tape in front to increase stride length on left Ambulation with hemi walker and WBQC (on R) with LLE blue band wrap Step facilitation in parallel bars with LLE blue band wrap, CGA x1, mod assist x1 for improved knee	Step facilitation in parallel bars from Bosu ball+ posteriorly to piece of tape anteriorly, mod assist x1 for knee flexion with swing Ambulation with no AD, min assist x2, 2x25 ft Ambulation with no AD, min assist x2, 2x60 ft Stepping over WBQC with LLE for foot clearance, min assist x2 Ambulation with no AD, CG assist x1,	Stepping in parallel bars with LLE over Dynadisc^ to increase foot clearance and step length Reassessment of 10MWT and 6MWT

		flexion with swing (see Figure 1)	min assist x1, 150 ft	
Therapeutic exercise	Wheelchair pulls with LLE to facilitate hamstring strengthening, 12 ft	RUE rows in sitting with blue TheraBand ^o	Supine on table, single leg bridges with LLE on 8" stool for hip extension on L, use of gait belt to bring R hip into flexion to reduce compensation Supine on table with LLE over edge to bring L hip into extension, use of gait belt to bring R hip into flexion, hamstring curls on L with yellow Theraband ^o (see Figure 2) LUE shoulder extensions in sitting with yellow Theraband ^o	Side stepping in parallel bars, x5 L hip abduction in parallel bars, 5x3
Neuro-muscular Reeducation	Sitting and standing postural corrections in mirror Seated isometric hamstring activation with LLE on Dynadisc [^] Seated L knee flexion with foot on towel to decrease friction Standing L hip extension and abduction with L foot on towel to decrease friction	Sit to stand from chair without use of arms, min assist x1 L trunk rotations in sitting with L scapular stabilization	Weight bearing through LUE on parallel bar Half kneeling on Airex [^] mat with RLE forward to bring L hip into extension and reduce tone Half kneeling Airex [*] mat with L foot forward with concentric lifting with verbal and tactile cues for L hamstring activation L hip extension with knee flexion onto 6" step in parallel bars (see Appendix 2)	Lunging on Airex [*] with R foot forward to stretch L hip flexor and reduce tone Trunk rotations in half kneeling Lunging with L foot forward, concentric lifting hamstring activation
Manual Therapy	Soft tissue mobilization to L lateral leg			
Modalities				Iontophoresis with

				Dexamethasone to R biceps tendon for pain management
Patient/Family Education	<p>Instruction on importance of home activities that facilitate dorsiflexion, knee flexion, and hip extension</p> <p>Correction of L AFO height and recommendation of R heel lift for increase L foot clearance with gait</p>	<p>Clinical importance of increasing step length and gait speed and how his gait speed compares to age-matched norms</p> <p>Importance of using LUE trough on wheelchair</p> <p>Pushing from chair for transfers instead of pulling grab bar</p>		Correct postural alignment with scapular activation

408 B/L=bilaterally, LLE=left lower extremity, RLE=right lower extremity, L=left, R=right, AFO=ankle foot orthosis,
 409 WBQC=wide bade quad cane, CGA=contact guard assist, min=minimum, mod=moderate, RUE=right upper
 410 extremity, LUE=left upper extremity, AD=assistive device, ft=feet


411
 412 * Airex, Chattanooga, TN

413 †Bosu, Ashland, OH

414 °TheraBand, Akron, OH

415 ^SPRI, Libertyville, IL

416



417 A. B.
418 **Figure 1:** Use of Theraband as a walking and stepping aid to assist with control at the left ankle, knee,
419 and hip joints. **A:** Active hip extension to initiate LLE swing phase of gait. **B:** Utilization of tape for
420 visual cue of step width.



421
422
423
424
425
426
427
428
429
430
431

Figure 2: Left lower extremity (LLE) hamstring curls in L hip extension with right lower extremity (RLE) inhibition.

432 **APPENDICES**

433 **Appendix 1. Medications at Initial Evaluation**

Medications	Dosage	Purpose
Ropinerole	0.5 mg daily	For restless leg syndrome
Terazosin	10 mg daily	For hypertension
Ambien	5 mg as needed	For sleep
Amlodipine	5 mg daily	For hypertension
Eliquis	10 mg daily	For peripheral artery disease
Aspirin	Low dose	For preventing blood clots
Atenolol	100 mg daily	For hypertension
Atorvastatin-	40 mg daily	For high cholesterol

434

435 **Appendix 2. Physical Therapy Short and Long Term Goals**

Time frame	Goal	At discharge
Short term: 4 weeks	Patient will demonstrate 60 ft improvement on the 6MWT as a result of improved endurance	Not achieved
	Patient will report improved functional capacity based on the SIS-16 with improvements between 4.7-7.1 points	Not tested/ deferred
	Patient will demonstrate improved static and dynamic awareness of his center of gravity with increased weight bearing symmetry through his lower extremities	Achieved
Long term: 8 weeks	Patient will demonstrate 120 ft improvement on the 6MWT as evidence of clinically meaningful detectable improvement for chronic stroke patients	Not achieved
	Patient will demonstrate meaningful improvement on the 5xSTS test scoring between 11-12 seconds	Not tested/deferred
	Patient will report improved functional capacity based on the SIS-16 with improvement between 9.4-14.1 points	Not tested/deferred
	Patient will be able to get on and off the floor with use of right UE and minimal assistance	Deferred
	Patient will increase gait speed by 0.17 m/s as measured by the 10MWT in order to have a minimally clinically important improvement	Not achieved
	Patient will report 75% reduced right lower leg pain during ambulation	Not achieved

436 ft=feet, 6MWT=six minute walk test, SIS-16=stroke impact scale 16, 5xSTS=five time sit to stand,
 437 UE=upper extremity, m/s=meters per second, 10MWT= ten meter walk test

438
 439
 440
 441
 442
 443
 444
 445
 446


447 **Appendix 3.** Example of neuromuscular re-education



448
449 Posterior stepping, L hip extension with knee flexion onto 6” step

450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469

470 **Appendix 4.** Gentile's Taxonomy of Tasks¹²



471 A tool for classification of tasks based on the goal of the action, and the environment in which it is
 472 performed. The complexity of a task is based on where it fits into the taxonomy.
 473