

STAFNE BONE CYST IN THE ANTERIOR MANDIBLE: AN UNUSUAL LOCATION

e-Poster: 261

Congress: ESHNR 2017

Type: Educational Poster

Topic: ESHNR 2017

Authors: R. Bignone, R. Mauceri, F.P. Lombardo, P. Purpura, G. La Tona, A. Lo Casto;
Palermo/IT

Keywords: Stafne Bone Cyst, Anterior mandible, Sublingual gland, Conventional radiology, CT, MRI

Any information contained in this pdf file is automatically generated from digital material submitted to e-Poster by third parties in the form of scientific presentations. References to any names, marks, products, or services of third parties or hypertext links to third-party sites or information are provided solely as a convenience to you and do not in any way constitute or imply ESHNR's endorsement, sponsorship or recommendation of the third party, information, product, or service. ESHNR is not responsible for the content of these pages and does not make any representations regarding the content or accuracy of material in this file.

As per copyright regulations, any unauthorised use of the material or parts thereof as well as commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is strictly prohibited.

You agree to defend, indemnify, and hold ESHNR harmless from and against any and all claims, damages, costs, and expenses, including attorneys' fees, arising from or related to your use of these pages.

Please note: Links to movies, ppt slideshows and any other multimedia files are not available in the pdf version of presentations.

<http://www.eshnr.eu>

1. Learning Objectives

To describe clinical features of the anterior Stafne bone cyst.

To supply radiologist a guide for diagnosis and classification of Stafne bone cyst.

2. Background

Stafne bone cyst (SBC) is a rare mandibular defect described by Stafne in 1942. It is not a real cyst but a bone defect of unknown origin. It is often a solitary and asymptomatic, incidentally discovered on routine radiological examination. It is mostly seen in aged males (between 50 and 70 years old), in the posterior portion of the mandible. Its anterior variant is uncommon and located in the premolar region of the mandible. Pressure of major salivary gland, with failure of normal bone deposition, is thought to be responsible of the SBC, even if the defect may contain other soft tissues. According to this widely accepted concept, the submandibular salivary gland is responsible for the posterior variants whereas the sublingual gland causes the anterior variant. A variety of imaging modalities, including conventional radiology, computed tomography (CT) or cone beam computed tomography (CBCT), magnetic resonance imaging (MRI) and sometimes sialography are employed to make the diagnosis.

Imaging techniques usefulness

Conventional radiology, usually panoramic radiograph. Initial, often incidental, evaluation. Follow up.

CT. Better definition of cortical bone, location and relationship with surrounding structures, thanks to 3D representation. Also useful to define the content (i.e. fat, glandular tissue, etc.).

MRI. Better definition of content, due to higher contrast resolution for soft tissue. Non ionizing radiations.

Sialography. It allows to determine the presence of glandular tissue in the cavity, but is invasive and uncomfortable for patients.

Classifications

According to the location in the mandible

Anterior (incidence: 0.009-0.3%; Fig. 1A).

Posterior (incidence: 0.1%-0.48%; Fig. 1B).

Ramus (about 20 cases reported).

c.jpg




Figure 1. A) 3D reformatted CBCT image. Anterior Stafne cyst. B) 3D reformatted CT image. Posterior Stafne cyst. C) Schematic drawing of different location of Stafne cyst: a, anterior, p, posterior, r, ramus (modified from: Ogunsalu C, Pillai K, Barclay S. Radiological assessment of type II Stafne idiopathic bone cyst in a patient undergoing implant therapy: a case report. West Indian Medical J 2006; 55:447-450)

According to the extension

Type I. The concavity does not reach the buccal cortical plate.

Type II. The concavity reaches the buccal cortical plate.

Type III. The concavity is characterized by buccal cortical plate expansion.

a.jpg




Figure 2. Modified from: Arijji E, Fujiwara N, Tabata O, et al. Stafne's bone cavity. Classification based on outline and content determined by computed tomography. Oral Surg, Oral Med, Oral Pathol 1993; 76:375-380

According to the cortical defect margins

Broad type.

Narrow type.

Smooth margin.

Irregular margin.

d.jpg




Figure 3. Modified from Minowa K, Inoue N, Sawamura T, Matsuda A, Totsuka Y, Nakamura M. Evaluation of static bone cavities with CT and MRI. Dentomaxillofac Radiol 2003; 32:2-7

According to the content

Type F. It is filled only with fat density.

Type S. Soft tissue structure inside, as lymph node, vessel, connective tissue or other.

Type G. Salivary gland entrapped in or located close to the concavity.

b.jpg




Figure 4. Modified from: Aiji E, Fujiwara N, Tabata O, et al. Stafne's bone cavity. Classification based on outline and content determined by computed tomography. Oral Surg, Oral Med, Oral Pathol 1993; 76:375-380

Stafne bone defect might be misdiagnosed as other cystic lesions of the mandible, such as solitary (traumatic) bone cyst, lateral periodontal cyst, radicular and residual cyst, keratocyst, and early stage focal cemento-osseous dysplasia.

No surgical treatment is required, but only periodic follow up, although the cavity is static and benign. Surgical exploration or biopsy should be performed in atypical cases.

3. Imaging Findings or Procedure Details

CASE REPORT

A 40 year-old-woman was referred by her dentist for further evaluation of a bone defect in the anterior mandible. The patient was asymptomatic and had previously made elsewhere a panoramic radiograph and a CBCT. A MR study was performed on a 1.5 T unit, before and after i.v. administration of a paramagnetic contrast medium, with TSE T1, T2 3D, STIR, and FFE T1 sequences.

Panoramic radiograph

On the panoramic radiograph an unilocular, oval radiolucency with well-defined margins (white arrow) was appreciable as an incidental finding in the anterior left mandible, premolar region. Panoramic radiograph was repeated in 2017 for follow up with no changes.

e.jpg



Figure 5. Panoramic radiograph in 2016 (A) and 2017 (B).

Cone-beam computed tomography (CBCT)

The CBCT made after in 2016 showed a well-defined, unilocular, corticated, round shape hypodense lesion in the anterior mandible (white arrow). The mandibular bone defect involved the lingual cortical bone but not reached the vestibular one.

f.jpg



Figure 6. CBCT. A) panoramic reformation. B) sagittal, C) coronal and D) axial planes.

i.jpg



Figure 7. 3D reformatted CBCT.

Magnetic resonance imaging (MRI)

On the MRI study performed in 2017 the left sublingual gland, herniated within the anterior Stafne cyst, was observed,

g.jpg




Figure 8. MRI study performed on a 1.5 T unit with the following multiplanar sequences: TSE T1 without and with fat saturation after i.v. administration of a paramagnetic contrast medium, T2 3D, STIR, and FFE T1 3D with fat saturation after i.v. administration of a paramagnetic contrast medium.

Case report classification

According to the location	ANTERIOR
According to the extension	TYPE I
According to the cortical defect margins	NARROW TYPE - SMOOTH MARGIN
According to the content	TYPE G

Table 1. Classification of the anterior Stafne cyst reported, according to the main classifications in the literature.

4. Conclusion


Anterior Stafne cyst is a rare, usually incidental occurrence on a panoramic radiograph. CT is useful for differential diagnosis with other cystic lesions of the mandible and classification. MRI, demonstrating sublingual gland herniation in the mandibular defect, can avoid bioptic or surgical procedures in asymptomatic patients.

5. References


- Ariji E, Fujiwara N, Tabata O, et al. Stafne's bone cavity. Classification based on outline and content determined by computed tomography. *Oral Surg, Oral Med, Oral Pathol* 1993; 76:375-380
- Campos PS, Panella J, Crusoé-Rebello IM, Azevedo RA, Pena N, Cunha T. Mandibular ramus-related Stafne's bone cavity. *Dentomaxillofac Radiol* 2004; 33:63-66
- Graham RM, Duncan KA, Needham G. The appearance of Stafne's idiopathic bone cavity on magnetic resonance imaging. *Dentomaxillofac Radiol* 1997; 26:74-75
- Kopp S, Ihde S, Bienengraber V. Differential diagnosis of Stafne idiopathic bone cyst with digital volume tomography (DVT). *J Maxillofac Oral Surg* 2010; 9:80-81
- Lee KH, Thiruchelvam JK, McDermott P. An unusual presentation of Stafne bone cyst. *J Maxillofac Oral Surg* 2015; 14: 841-844
- Minowa K, Inoue N, Sawamura T, Matsuda A, Totsuka Y, Nakamura M. Evaluation of static bone cavities with CT and MRI. *Dentomaxillofac Radiol* 2003; 32:2-7
- Ogunsalu C, Pillai K, Barclay S. Radiological assessment of type II Stafne idiopathic bone cyst in a patient undergoing implant therapy: a case report. *West Indian Medical J* 2006; 55:447-450
- Probst FA, Probst M, Maisterli I-Z, Otto S, Troeltzch M. Imaging characteristics of a Stafne bone cavity - panoramic radiography, computed tomography and magnetic resonance imaging. *Oral Maxillofac Surg* 2014; 18:351-353
- Schneider T, Filo K, Locher MC, et al. Stafne bone cavities: systematic algorithm for diagnosis derived from retrospective data over a 5-year period. *Br J Oral Maxillofac Surg* 2014; 52:369-374
- Sisman Y, Etoz OA, Mavili E, Sahman H, Ertas ET. Anterior Stafne bone defect mimicking a residual cyst: a case report. *Dentomaxillofac Radiol* 2014; 39:124-126
- Stafne EC. Bone cavities situated near the angle of the mandible. *J Am Dent Assoc* 1942; 29:1969-1972
- Taysi M, Ozden C, Cankaya B, Olgac V, Yildirim S. Stafne bone defect in the anterior mandible. *Dentomaxillofac Radiol* 2014; 43:20140075

6. Mediafiles


a.jpg




b.jpg



c.jpg



d.jpg




e.jpg



f.jpg



g.jpg



i.jpg

