

1 **The key role of forests in meeting climate targets requires science**
2 **for credible mitigation**

3

4

5 **Authors**

6 **Giacomo Grassi***, European Commission, Joint Research Centre, Ispra (VA), Italy

7 **Jo House**, Cabot Institute, Bristol University, UK

8 **Frank Dentener**, European Commission, Joint Research Centre, Ispra (VA), Italy

9 **Sandro Federici**, FAO consultant, Italy

10 **Michel den Elzen**, PBL Netherlands Environmental Assessment Agency, The Hague, The
11 Netherlands

12 **Jim Penman**, University College London, UK

13

14

15 *Corresponding author: giacomo.grassi@ec.europa.eu

16 **ABSTRACT**

17

18 **Forest-based climate mitigation may occur through conserving and enhancing the carbon sink**
19 **and through reducing greenhouse gas emissions from deforestation. Yet the inclusion of forests**
20 **in international climate agreements has been complex, often considered a secondary mitigation**
21 **option. In the context of the Paris Climate Agreement, countries submitted their (Intended)**
22 **Nationally Determined Contributions ((I)NDCs), including climate mitigation targets. Assuming**
23 **full implementation of (I)NDCs, we show that land use, and forests in particular, emerge as a**
24 **key component of the Paris Agreement: turning globally from a net anthropogenic source**
25 **during 1990-2010 (1.3 ± 1.1 GtCO₂e/y) to a net sink of carbon by 2030 (up to -1.1 ± 0.5**
26 **GtCO₂e/y), and providing a quarter of emission reductions planned by countries. Realizing and**
27 **tracking this mitigation potential requires more transparency in countries' pledges and an**
28 **enhanced science-policy cooperation to increase confidence in numbers, including reconciling**
29 **the ≈ 3 GtCO₂e/y difference in current estimates between country reports and scientific studies.**
30 **This represents a challenge and an opportunity for the scientific community.**

31 MAIN TEXT

32

33 In December 2015, 195 countries adopted the Paris Climate Agreement¹ at the 21st Conference of
34 Parties (COP-21) of the United Nations Framework Convention on Climate Change (UNFCCC). As
35 part of the process, 187 countries, representing more than 96% of global net emissions in 2012²,
36 submitted their Intended National Determined Contributions³ (INDCs, which become NDCs with the
37 ratification of the Paris Agreement⁴). The NDCs are the basis for implementing actions under the
38 Agreement, and the vast majority include commitments in the land-use sector.

39 Land use, including agriculture and forests, accounts for about 10% of global greenhouse gas (GHG)
40 emissions as CO₂, and nearly quarter including CH₄ and N₂O⁵⁻⁹. Also, about one third of the current
41 anthropogenic CO₂ emissions are removed by terrestrial ecosystems, mainly forests. While
42 deforestation is estimated to be the main GHG source in many tropical countries, forest sinks are
43 important globally with net sinks dominating in temperate and boreal countries¹⁰.

44 Including land use in the UNFCCC process has been long and complex. For forests, uncertainties of
45 GHG estimates and methodological issues such as additionality (i.e. showing that proposed mitigation
46 efforts go beyond Business-as-Usual (BAU) and separation of non-anthropogenic effects) and leakage
47 (displacement of land-use activities to other areas) have often led to controversies and compromises¹¹.

48 The UNFCCC requires that all countries report GHG inventories of anthropogenic emissions and
49 removals using methodologies developed by the Intergovernmental Panel on Climate Change (IPCC)
50 and adopted by UNFCCC¹². Developed countries report annual GHG inventories¹³, using mandatory
51 and voluntary land-use activities towards meeting their emission reduction targets where applicable
52 under the Kyoto Protocol¹⁴. Developing countries' GHG inventories have historically been reported
53 less frequently¹⁵, though biennial updates are now required¹⁶, and may undertake voluntary mitigation
54 activities, notably through the REDD+ process (Reducing Emissions from Deforestation, forest
55 Degradation, and other forest activities).

56 The Paris agreement is a potential game changer for land use mitigation. It calls explicitly for all
57 countries to make use of a full-range of land-based mitigation options, and to take action on REDD+.
58 Based on country information, this analysis quantifies the expected GHG mitigation role of the land-
59 use sector in the (I)NDCs to the year 2030, including activities conditional on finance, technology and
60 capacity-building support. It does not assess specific country policies. It focuses on CO₂ emissions
61 and removals and non-CO₂ emissions from Land Use, Land-Use Change and Forestry (LULUCF,
62 primarily deforestation and forest management), encompassing most of the land-use sector identified
63 in (I)NDCs. Harvested wood products are included for most developed countries. Non-CO₂ emissions
64 from agriculture are not included.

65

66 **Country mitigation targets are expressed in different ways**

67 Countries express their (I)NDC targets with different combinations of the following elements¹⁷⁻¹⁹
68 (Supplementary Tables 1-2), reflecting different national circumstances, i.e.:

- 69 • Quantifier - targets are expressed as either an *absolute quantity* e.g. amount of GHG reduction in
70 tonnes of CO₂ equivalent (tCO₂e), or as a change in the *emission intensity*, e.g. China and India
71 express a reduction of emission intensity per unit of GDP.
- 72 • Reference point – Emissions in the target year (e.g. 2025 or 2030) are compared to either a historic
73 base year (e.g. 1990, 2005) or to the target year in a BAU scenario. The BAU scenario assumes
74 either no mitigation activity, or some existing mitigation activity.
- 75 • Conditionality - While developed country (I)NDC targets are unconditional, most developing
76 countries expressed at least part of their targets as conditional on finance, technology or capacity-
77 building support.

78 The (I)NDCs vary in the way they include LULUCF. It may be fully included as part of the overall
79 target like other sectors, partially included through accounting rules to reflect the additional impact of

80 human actions or considered separately with special mitigation actions. Consequently, evaluating the
81 expected effect of LULUCF on the (I)NDC mitigation targets is complex.

82 Our analysis is based on information provided on LULUCF in the (I)NDCs³, and also (in order of
83 priority) other country reports to UNFCCC^{13,15,16,20,21}, other official country documents, and FAO-
84 based datasets for forest^{8,22} and for other land uses²³ (Supplementary Tables 4-5). Given the Paris
85 Agreement context of our analysis, we prioritized (I)NDCs and those country reports that are formally
86 reviewed or technically assessed by UNFCCC, with FAO-based datasets used for gap filling,
87 allowing global estimates covering 195 countries (see Methods). We found sufficient information to
88 analyse the LULUCF mitigation contribution for 68 countries (or 41 (I)NDCs, with the EU's NDC
89 representing 28 countries), representing around 78% of global net emissions in 2012² and 83% of the
90 global forest area²². For the remaining countries, where LULUCF is not expected to offer a large
91 mitigation potential (Supplementary Section 1), the future LULUCF mitigation contribution was
92 assumed to be zero.

93

94 **Historical and projected forest emissions and removals**

95 Fig. 1 shows, for all 195 UNFCCC countries, historical and future anthropogenic LULUCF emissions
96 and removals from this analysis, based on official country data. The Supplementary Sections 2 and 3
97 provide, respectively, additional country-specific assessments and an analysis of uncertainties for the
98 absolute level of net emissions and their trend^{24,25}, based on information from countries' reports.

99 While country information on uncertainty up to 2030 is not available, we conservatively assumed that
100 the uncertainty estimated for historical net emissions would also hold for the future.

101 Historically, global LULUCF net emissions decreased from 1.54 ± 1.06 GtCO₂e/y (95% CI) in 1990
102 to 0.01 ± 0.86 GtCO₂e/y in 2010 (slope of linear trend: -0.08 GtCO₂e/y). The trend and the inter-
103 annual variability over this period are influenced by: (i) deforestation in Brazil, with peak years in
104 1995 and 2002-2004 followed by a steep reduction of emissions by about -1.3 GtCO₂e/y till 2010; (ii)
105 high deforestation rates (1997-1999) and peak years in peat fire emissions (e.g., 1997) in Indonesia;

106 (iii) an increasing sink in managed temperate and boreal forests, of about $-0.8 \text{ GtCO}_2\text{e/y}$ from 1990 to
107 2010. By splitting the 1990-2010 period (average emissions: $1.28 \pm 1.15 \text{ GtCO}_2\text{e/y}$) into four sub-
108 periods, we conclude that the historical trend is statistically significant after 2000 (Supplementary
109 Section 3).

110 The wide range of future LULUCF net emissions depends on policy scenarios (Fig. 1). The ‘country-
111 BAU’ scenario foresees a marked increase in global net emissions (Supplementary Table 6), reaching
112 $1.94 \pm 1.53 \text{ GtCO}_2\text{e/y}$ in 2030. This is because several developing countries assumed BAU to be a no-
113 measures scenario, e.g. ignoring the existing policies to reduce deforestation. Under the ‘pre-(I)NDC
114 scenario’, i.e. considering policies in place prior to COP-21 (including the earlier Copenhagen
115 pledges²¹), global net emissions increase moderately, up to $0.36 \pm 0.94 \text{ GtCO}_2\text{e/y}$ in 2030. For the
116 ‘unconditional (I)NDC scenario’ the global net emissions slightly decrease, reaching a sink of $-0.41 \pm$
117 $0.68 \text{ GtCO}_2\text{e/y}$ in 2030. An additional reduction of net emissions is estimated for the ‘conditional
118 (I)NDC’ scenario, leading to a sink of $-1.14 \pm 0.48 \text{ GtCO}_2\text{e/y}$ in 2030.

119 The analysis of the emission trend over the entire period shows that the difference between the 1990-
120 2010 average and the net emissions in 2030 is not significant for the pre-(I)NDC scenario, but is
121 significant (95% CI) for both the unconditional and the conditional (I)NDC scenarios (Supplementary
122 Figure 3b). This indicates that the reduction of net emissions assumed by the (I)NDCs relative to the
123 historical period, if achieved, is statistically robust.

124

125 **Comparison with global datasets**

126 Fig. 2 compares the historical LULUCF trend from our analysis to other three well-known global
127 LULUCF datasets: (i) latest country reports to UNFCCC (ref^{13,15,16,20}); (ii) FAOSTAT for all land
128 uses²³; and (iii) IPCC Fifth Assessment Report (AR5) Working Groups (WG) I⁵ and III⁶ data used for
129 the global carbon budget.

130 The difference between this analysis and the UNFCCC country reports is because several (I)NDCs
131 updated past datasets, and because we used FAO-based data for gap-filling, instead of pre-2010
132 National Communications.

133 Differences between this analysis and FAOSTAT include the definition of forest (UNFCCC vs.
134 FAO); coverage of areas and of carbon pools; and differing estimation methods by reporting
135 agencies⁸ (see Methods).

136 There is a large difference of about 3 GtCO₂e/y between this analysis, based on country reports
137 following the IPCC Guidelines for national GHG inventories^{25,26} (IPCC GL), and the scientific studies
138 summarized by the IPCC AR5^{5,6}. For the period 2000-2009, the level of net emissions is on average
139 0.90 ± 1.11 GtCO₂e/y (95 % CI) in our analysis and 4.03 ± 2.93 GtCO₂e/y (90 % CI, reflecting both
140 methodological and terminological choices²⁷⁻²⁹) in IPCC AR5 (Fig. 2). The above differences are
141 linked to different scopes of the two IPCC work streams³⁰: the GL focus on internationally agreed
142 methodologies for national anthropogenic GHG estimation, recognizing different countries'
143 definitions and technical capabilities, whilst the AR5 focuses on assessing the state of the science on
144 the global carbon budget using globally applied data, definitions and modeling methods.

145 Specifically, LULUCF in the IPCC GL includes estimates of GHG emissions and removals from all
146 land uses, reported under either a stable or changed land-use status (typically in the last 20 years), e.g.
147 “forest remaining forest” or “forest converted to cropland” (or vice versa). There is a large scientific
148 challenge of providing a practicable methodology to factor out direct human-induced mitigation
149 action from indirect human-induced and natural effects^{31,32}, such as the natural aging of forests,
150 natural disturbances and environmental change (e.g. climate change, extended growing seasons,
151 fertilizing effects of increased [CO₂] and nitrogen deposition). Therefore, the IPCC GL^{25,26} use the
152 category of “managed land” as a default first order approximation of “anthropogenic” emissions and
153 removals, based on the rationale that the preponderance of anthropogenic effects occurs on managed
154 land³². The GHG inventories should report all emissions and removals for managed land, while GHG

155 fluxes from unmanaged land are excluded. What is included in “managed land” varies from country
156 to country, although the countries’ definition must be applied consistently over time.

157 In contrast, global models such as those used in IPCC AR5 and the Global Carbon Project take a
158 different approach to separate anthropogenic from natural effects. Anthropogenic fluxes (referred as
159 “net land-use change”^{5,9}, or “Forestry and Other Land Uses”⁶), are estimated by a bookkeeping
160 model²⁷ or by dynamic global vegetation models⁹ based on changes in land cover (i.e. between forest
161 and agriculture), forest regrowth and, depending on the modeling capability, some forms of
162 management (wood harvest and shifting cultivation). The difference between this modeled
163 “anthropogenic” flux and the estimated total net flux of CO₂ between the land and atmosphere⁹ is the
164 “residual terrestrial sink”^{5,6,9}, which is generally assumed to be a natural response of primary or
165 mature regrowth forests to environmental change^{9,27}.

166 The above methodological differences are reflected in the net emissions from developed countries,
167 where most of the ≈ 3 GtCO₂e/y difference between our analysis and IPCC AR5 occurs for the period
168 2000-2009: while these countries report a substantial “anthropogenic” sink (-1.9 GtCO₂e/y in
169 “UNFCCC Annex 1” countries), the bookkeeping model (IPCC AR5) finds a small net source (0.1
170 GtCO₂e/y, “OECD” in Fig. 11.7 of ref.⁶). This difference lies essentially in whether the large sinks in
171 areas designated by countries as “managed forest” (following IPCC GL), well documented in forest
172 inventories¹⁰, are attributed to “anthropogenic” (in the GHG inventories) or to “natural” fluxes (in
173 IPCC AR5).

174 To explore, at least in part, the impact of these different attribution methods, Fig. 3a compares what is
175 considered undisputedly “anthropogenic” by both IPCC AR5 (land-use change) and the country
176 reports (land converted to other land uses). These estimates, both predominated by tropical
177 deforestation, are of similar magnitude, especially after 2000. The other fluxes, where the attribution
178 differs more between IPCC AR5 and the countries, are shown in Fig 3b. Thus much of the sink that
179 countries report under ‘forest remaining forest’, the global models consider part of the natural flux.
180 This disaggregation suggests that the residual sink is at least partly influenced by management

181 practices not captured by global carbon models³³, but also that countries consider anthropogenic what
182 is partly influenced by environmental change and by recovery from past disturbances.

183 There are many reasons for the lower sink reported by countries in Fig 3b compared to the residual
184 sink from IPCC AR5³⁰, including the fact that countries do not report sinks for unmanaged lands (e.g.,
185 a large sink in tropical and boreal unmanaged forests¹⁰) and their reporting for managed land may be
186 incomplete, i.e. ignoring fluxes (e.g. sink in grasslands, wetlands or forest regrowth) or carbon pools.
187 There would be other factors to consider, including treatment of legacy fluxes from past land-use and
188 other definitional and methodological differences. These would require a more detailed analysis,
189 which is outside the scope of this paper.

190 Finally, the projections from this analysis can be compared to RCP scenarios used in IPCC AR5 up to
191 2030 (Fig. 3, dashed lines). For the undoubtedly “anthropogenic” fluxes (Fig. 3a), our country data
192 analysis falls broadly within the IPCC AR5 scenarios, supporting previous qualitative findings³⁴.

193 Overall, our analysis shows 1) that various global LULUCF datasets may be more consistent than
194 apparent at first glance, 2) unless the scientific and GHG inventory community appreciate these
195 definitional and methodological issues, conflicting numbers and messages are likely to appear in the
196 coming years, and 3) that several reasons for the differences among datasets can be further reconciled
197 in collaboration between the two communities, which would be a very useful contribution to science
198 and policy.

199

200 **Different perspectives on mitigation contribution by forests**

201 To reflect the complexity of approaches to (I)NDCs, this analysis assesses three different perspectives
202 on LULUCF mitigation:

203 (A) 2030 (I)NDC vs. 2005, i.e., the expected impact of full (I)NDC implementation. The year 2005 is
204 chosen as historically reliable in terms of data. Fig. 1 shows that the global LULUCF net emissions to
205 the atmosphere would transition from an estimated net anthropogenic source of +0.8 GtCO₂e/y in

206 2005 to a net sink of -0.4 GtCO₂e/y (unconditional (I)NDCs) or -1.1 GtCO₂e/y (conditional (I)NDCs)
207 in 2030.

208 (B) 2030 (I)NDC vs. 2030 alternative scenarios: country-BAU or pre-(I)NDC, i.e., the additional
209 LULUCF contribution relative to alternative scenarios (Fig. 1). The magnitude of the difference
210 between country-BAU and pre-(I)NDC (1.6 GtCO₂/y) may raise concerns about the expected results-
211 based payments under REDD+, which should be based on credible baselines and not on a no-
212 measures scenario. Clarification of the role of REDD+ in (I)NDCs should therefore be seen as a
213 priority by countries. Compared to the estimated pre-(I)NDC scenario, net emissions in 2030 are
214 lower by 0.8 GtCO₂e/y or 1.5 GtCO₂e/y for unconditional and conditional (I)NDCs, respectively. For
215 the ‘conditional (I)NDC vs. 2030 pre-(I)NDC’ scenario (Fig. 4a), this LULUCF contribution of 1.5
216 GtCO₂e/y (Fig. 4a, last column) represents 26% of the total mitigation expected from all GHG sectors
217 (5.9 GtCO₂e/y³⁵, Fig. 4a, third column). The countries contributing most to LULUCF mitigation
218 under this perspective are Brazil and Indonesia, followed by other countries focusing either on
219 avoiding carbon emissions (e.g. Ethiopia, Gabon, Mexico, DRC, Guyana and Madagascar) or on
220 promoting the sink through large afforestation programs (e.g. China, India).

221 (C) Country perspective on emissions reduction in the (I)NDC, i.e. what each country might consider
222 its “LULUCF contribution to the overall (I)NDC”, as part of its mitigation package, e.g. if a country
223 commits to reduce its all-sectors emissions by $x\%$ relative to y (reference point: base year or BAU-
224 scenario), what fraction of x is attributable to LULUCF? This approach looks at the way countries
225 define their (I)NDCs (e.g. reference point) and the way LULUCF is included within the (I)NDC (as
226 any other sector or with special accounting rules). Globally, under this perspective the LULUCF
227 contribution is 3.1 GtCO₂e/y (unconditional) or 3.8 GtCO₂e/y (conditional). The latter case (Fig. 4b,
228 last column) corresponds to 24% of total all-sectors emission reduction relative to the reference point
229 (i.e. 15.8 GtCO₂e/y, Fig. 4b, third column).

230 The emission reductions from a country perspective (Fig. 4b) are greater than the deviation from the
231 pre-(I)NDC scenario (Fig. 4a), because countries’ choices of reference point in their (I)NDCs tend to

232 maximize the accounted mitigation, i.e. countries that already reduced emissions used a historical
233 base year, whereas countries expecting a future increase of emissions used a future BAU-scenario.
234 This is evident under perspective C, where nearly one third of the contribution comes from Brazil,
235 followed by Indonesia and Russia (Fig 4b, last column). In Brazil, where total emissions have
236 declined after 2004 due to successful implementation of policies to reduce deforestation³⁶, the NDC
237 target (-43%) is relative to 2005. Our analysis suggests that in Brazil the LULUCF contribution to
238 NDC is greater than the all-sectors NDC target for 2030, i.e. the NDC allows emissions from other
239 sectors to increase. In Indonesia the conditional NDC target (-41%) is relative to the BAU-scenario in
240 2030. LULUCF represents about 65% of current (2010) total emissions and is expected to contribute
241 nearly two-thirds of the NDC emission reduction (relative to BAU) foreseen in 2030. Brazil and
242 Indonesia are representative examples of GHG emission trends in developing countries: with an
243 expanding and industrializing economy, the currently high LULUCF emissions are expected to
244 decrease, and be superseded by growing emissions from the energy sector. The (I)NDC target of
245 Russia (-30%) is relative to 1990, with LULUCF contributing by about two-fifths to this emission
246 reduction. Russia is more important in perspective C than in B because its specific accounting method
247 for LULUCF gives prominence to the contribution of the current forest sink to climate mitigation.

248 The (I)NDCs of the three countries above may be assessed also in terms of clarity and trust of
249 information provided (see Supplementary Section 2). Overall, Brazil's NDC is transparent on the
250 land-use sector and the underlying GHG estimates are based on a well-developed monitoring system.
251 The recent relevant upward revision of historical deforestation emissions in Brazil opens new
252 questions on the implementation of the NDC target and on how and when data consistency between
253 NDC, REDD+ and National Communications will be ensured. The relative ambiguity of Indonesia's
254 NDC on how it would address land use emissions is improved by the information in more recent
255 documents. Furthermore, recent monitoring efforts have improved the GHG emission estimates,
256 especially from peatland drainage and from forest degradation, whereas emissions from peat fires
257 remain very uncertain. These improvements are mainly due to the REDD+ process, which in many
258 developing countries is triggering unprecedented monitoring efforts. The challenge is increasingly to

259 transfer these improvements into the NDC process, and to clarify the often uncertain relationship
260 between the financially-supported REDD+ activities and the NDCs. For Russia, transparency of
261 mitigation efforts will crucially depend on clarifying the accounting method chosen for LULUCF. In
262 addition, credible GHG estimates will require reconciling or explaining the currently large difference
263 in the forest sink between the reports submitted by Russia to UNFCCC and to FAO.

264 In summary, the full implementation of (I)NDCs would turn LULUCF globally from a net source
265 during 1990-2010 (1.3 ± 1.1 GtCO₂e/y) to a net sink by 2030 (up to -1.1 ± 0.5 GtCO₂e/y). The
266 accounted LULUCF mitigation contribution in 2030 is very different depending on the way that
267 mitigation is calculated, ranging from 0.8 to 3.1 GtCO₂e/y for unconditional (I)NDCs and from 1.5 to
268 3.8 GtCO₂e/y for conditional (I)NDCs (Supplementary Table 3). However, in relative terms,
269 LULUCF would provide about a quarter of total emission reductions planned in countries' (I)NDCs
270 irrespective of the approach to calculating mitigation.

271 Whereas a similar trend of decreasing LULUCF net emissions with full (I)NDCs implementation has
272 been suggested also by other analyses (ref^{34,37}), the absolute level of net emissions differs
273 significantly: e.g., ref³⁷ reports net emissions about 3 GtCO₂e/y higher than ours, due to the
274 'harmonization' of different datasets (country projections and (I)NDCs were aligned to historical
275 FAOSTAT data). By contrast, our study is the first so far showing a global picture of country-based
276 LULUCF net emissions that is consistent between historical and projected periods, including
277 discussing the differences with other global datasets and different mitigation perspectives.

278

279 **Science can help countries to keep the forest mitigation promise**

280 Several studies suggest a theoretical mitigation potential from land use^{6,35,38} higher than in this
281 analysis, others suggest limits posed by ecological and socio-economic constraints (including land
282 availability)^{39,40}. Irrespective of the potential, in the past UNFCCC negotiations the LULUCF sector
283 has often been treated separately and considered as a secondary mitigation option, largely due to its
284 complexity and limited trust in data.

285 Our analysis shows a wide range of future LULUCF net emissions, depending on policy scenarios.
286 Through the implementation of (I)NDCs countries (especially developing ones) expect a key
287 contribution from LULUCF in meeting their (I)NDC targets, with a clear focus on forests. Achieving
288 this will require increasing the credibility of LULUCF mitigation, through more transparency in
289 commitments and more confidence in estimates. To this regard, the Paris Agreement includes a
290 “Framework for transparency of actions”, key for its credibility⁴¹, aimed at providing clarity on GHG
291 estimates and tracking of progress toward achieving countries’ individual targets.

292 More transparent commitments means that future updates of the NDCs should provide more details
293 on how LULUCF mitigation is calculated towards meeting the target and how the financially-
294 supported REDD+ activities contribute to the pledges. More confidence in LULUCF estimates will
295 require improving the country GHG inventories in terms of transparency, accuracy (including
296 information on uncertainties), consistency, completeness and comparability⁴², especially in
297 developing countries.

298 This is a challenge and an opportunity for the scientific community. Supporting country GHG
299 estimation includes regular reviews of the latest science (e.g. ref⁴³), expanding the scope of the
300 operational methods in the IPCC guidance, as has been done for REDD+⁴⁴, and incorporating
301 opportunities offered by emerging satellite data^{45,46} available through highly accessible products⁴⁷.
302 More confidence also requires independent checks of the transparency and reliability of data, e.g. by
303 reproducing and verifying countries’ GHG estimates. According to IPCC guidance²⁵, verification of
304 GHG inventories is key to improve scientific understanding and to build confidence on GHG
305 estimates and their trends. This can be achieved by comparing GHG inventories with scientific
306 studies using partially or totally independent datasets and/or different methods (e.g. ref⁴⁸), including
307 greater integration of modeling and measurement systems of land use-related net emissions⁹.
308 Meaningful verification requires improving mutual understanding and cooperation between the
309 scientific community and the developers of national GHG inventories.

310 Finally, increasing trust in proposed LULUCF mitigation options will require reconciling the current
311 ≈ 3 GtCO₂e/y difference in global LULUCF net emissions between country reports and scientific
312 studies (as reflected in IPCC reports). Among the many possible reasons for these differences^{30,49}, we
313 suggest that what is considered “anthropogenic sink” is key and deserves further analyses. While
314 recognizing differences in scopes among these communities, reconciling differences in estimates is a
315 necessity, as the “Global stocktake”, i.e. the foreseen five-yearly assessment of the collective progress
316 toward achieving the long-term goals of the Paris Agreement, will be based on both country reports
317 and IPCC reports. Without speaking the same language, the “balance between anthropogenic GHG
318 emissions by sources and removals by sinks in the second half of this century”¹, needed to reach the
319 ambitious “well-below 2°C” target, cannot be properly assessed.

320

321

322 **Correspondence and requests for materials:** giacomo.grassi@ec.europa.eu

323 **Disclaimer:** The views expressed are purely those of the writers and may not in any circumstances be
324 regarded as stating an official position of the European Commission.

325 **Author Contributions** G.G. conceived the analysis on (I)NDCs, executed the calculations and
326 drafted the paper. J.H., F.D., M.d.E. and J.P. contributed to the analysis and to the writing of the paper.
327 S.F. provided data from FAO FRA-2015 and contributed to the analysis. J.H. was supported by
328 Leverhulme Foundation and EU FP7 through project LUC4C (GA603542).

329 **Competing financial interests.** The authors declare no competing financial interests.

330 **Acknowledgements.** Author Jim Penman passed away recently. Jim was the UK and EU
331 negotiator on LULUCF for many years, coordinator of key IPCC methodological reports and
332 credited as one of the key architects of the LULUCF process under the UNFCCC. He was
333 awarded an OBE (Order of the British Empire) for his work. He was an outstanding scientist
334 and negotiator, who strived always towards a better world.

335

336 **References**

337

338 1 UNFCCC. Adoption of the Paris Agreement. Report No. FCCC/CP/2015/L.9/Rev.1,

339 <http://unfccc.int/resource/docs/2015/cop21/eng/l09r01.pdf>. (UNFCCC, 2015).

340 2 European Commission, Joint Research Centre (JRC)/Netherlands Environmental Assessment

341 Agency (PBL). Emission Database for Global Atmospheric Research (EDGAR), release

342 version 4.3.1 <http://edgar.jrc.ec.europa.eu/overview.php?v=431>, 2016.

343 3 UNFCCC. INDCs as communicated by Parties,

344 <http://www4.unfccc.int/submissions/indc/Submission%20Pages/submissions.aspx>.

345 (UNFCCC, 2015).

346 4 UNFCCC. Paris Agreement - Status of Ratification.

347 http://unfccc.int/paris_agreement/items/9444.php (UNFCCC 2016)

348 5 Ciais, P. *et al.* in *Climate Change 2013: The Physical Science Basis. Contribution of Working*

349 *Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*

350 (eds T.F. Stocker *et al.*) Ch. 6, 465-522 (Cambridge University Press, 2013).

351 6 Smith, P. *et al.* in *Climate change 2014: mitigation of climate change. Contribution of*

352 *Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate*

353 *Change* (eds O. Edenhofer *et al.*) Ch. 11, 811-886 (Cambridge University Press, 2014).

354 7 Tubiello, F. N. *et al.* The Contribution of Agriculture, Forestry and other Land Use activities

355 to Global Warming, 1990-2012. *Global Change Biology* **21**, 2655-2660 (2015).

356 8 Federici, S., Tubiello, F. N., Salvatore, M., Jacobs, H. & Schmidhuber, J. New estimates of

357 CO₂ forest emissions and removals: 1990-2015. *Forest Ecology and Management* **352**, 89-98

358 (2015).

359 9 Le Quéré, C. *et al.* Global Carbon Budget 2015. *Earth System Science Data* **7**, 349-396

360 (2015).

361 10 Pan, Y. *et al.* A large and persistent carbon sink in the world's forests. *Science* **333**, 988-993

362 (2011).

363 11 Schlamadinger, B. *et al.* Options for including land use in a climate agreement post-2012:
364 improving the Kyoto Protocol approach. *Environmental Science and Policy* **10**, 295-305
365 (2007).

366 12 UNFCCC. Decision 2/CP.17. Outcome of the work of the Ad Hoc Working Group on Long-
367 term Cooperative Action under the Convention, FCCC/CP/2011/9/Add.1,
368 <http://unfccc.int/resource/docs/2011/cop17/eng/09a01.pdf> (UNFCCC, 2011).

369 13 UNFCCC. Greenhouse Gas Inventories,
370 [http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/](http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/8812.php)
371 [items/8812.php](http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/8812.php). (UNFCCC, 2015).

372 14 Grassi, G., den Elzen, M. G. J., Hof, A. F., Pilli, R. & Federici, S. The role of the land use,
373 land use change and forestry sector in achieving Annex I reduction pledges. *Climatic Change*,
374 1-9 (2012).

375 15 UNFCCC. National Communications Non-Annex 1, [http://unfccc.int/nationalreports/non-](http://unfccc.int/nationalreports/non-annexinatcom/submittednatcom/items/653.php)
376 [annexinatcom/submittednatcom/items/653.php](http://unfccc.int/nationalreports/non-annexinatcom/submittednatcom/items/653.php) (UNFCCC, 2015).

377 16 UNFCCC. Biennial Update Reports, [http://unfccc.int/national_reports/non-](http://unfccc.int/national_reports/non-annex_i_natcom/reporting_on_climate_change/items/8722.php)
378 [annex_i_natcom/reporting_on_climate_change/items/8722.php](http://unfccc.int/national_reports/non-annex_i_natcom/reporting_on_climate_change/items/8722.php) (UNFCCC, 2015).

379 17 WRI. CAIT Climate Data Explorer. Paris Contributions Map, <http://cait.wri.org/indc/>. (World
380 Resources Institute, 2015).

381 18 Admiraal, A. *et al.* Assessing Intended Nationally Determined Contributions to the Paris
382 climate agreement – what are the projected global and national emission levels for 2025–
383 2030? Report No. PBL 1879, [http://www.pbl.nl/en/publications/assessing-intended-](http://www.pbl.nl/en/publications/assessing-intended-nationally-determined-contributions-to-the-paris-climate-agreement)
384 [nationally-determined-contributions-to-the-paris-climate-agreement](http://www.pbl.nl/en/publications/assessing-intended-nationally-determined-contributions-to-the-paris-climate-agreement). (PBL, 2015).

385 19 Rogelj, J. *et al.* Paris Agreement climate proposals need a boost to keep warming well below
386 2 °C. *Nature* **534**, 631-639, doi:10.1038/nature18307 (2016).

387 20 UNFCCC. National Communications Annex 1,
388 <http://unfccc.int/nationalreports/annexinatcom/submittednatcom/items/7742.php>; (UNFCCC,
389 2015).

- 390 21 UNFCCC. Compilation of information on nationally appropriate mitigation actions to be
391 implemented by Parties not included in Annex I to the Convention,
392 FCCC/AWGLCA/2011/INF.1, <http://unfccc.int/resource/docs/2011/awglca14/eng/inf01.pdf>.
393 (UNFCCC, 2011).
- 394 22 FAO-FRA. Country reports (2015), [http://www.fao.org/forest-resources-assessment/current-](http://www.fao.org/forest-resources-assessment/current-assessment/country-reports/en/)
395 [assessment/country-reports/en/](http://www.fao.org/forest-resources-assessment/current-assessment/country-reports/en/). (Food and Agricultural Organization of the United Nations,
396 2015).
- 397 23 FAOSTAT. Land use emissions. (Food and Agricultural Organization of the United Nations
398 (FAO), Rome, Italy, http://faostat3.fao.org/download/G2/*/E, 2015).
- 399 24 Grassi, G., Monni, S., Federici, S., Achard, F., and Mollicone, D. Applying the
400 conservativeness principle to REDD to deal with the uncertainties of the estimates.
401 *Environmental Research Letters*, 3(3) (2008)
- 402 25 IPCC. in *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (eds H.S.
403 Eggleston *et al.*) (National Greenhouse Gas Inventories Programme, Institute for Global
404 Environmental Strategies, Hayama, Japan, 2006).
- 405 26 IPCC. in *Good Practice Guidance for Land Use, Land-Use Change and Forestry* (eds J.
406 Penman *et al.*) (Institute for Global Environmental Strategies, 2003).
- 407 27 Houghton, R. A. *et al.* Carbon emissions from land use and land-cover change.
408 *Biogeosciences* **9**, 5125-5142 (2012).
- 409 28 Hansis, E., Davis, S. J. & Pongratz, J. Relevance of methodological choices for accounting of
410 land use change carbon fluxes. *Global Biogeochemical Cycles* **29**, 1230-1246 (2015).
- 411 29 Pongratz, J., Reick, C. H., Houghton, R. A. & House, J. I. Terminology as a key uncertainty
412 in net land use and land cover change carbon flux estimates. *Earth System Dynamics* **5**, 177-
413 195 (2014).
- 414 30 Federici, S. *et al.* GHG fluxes from forests: An assessment of national reporting and
415 independent science in the context of the Paris Agreement,

416 <http://www.climateandlandusealliance.org/reports/ghg-fluxes-from-forests/> (Climate and
417 Land Use Alliance, 2016).

418 31 IPCC. in *IPCC Meeting on Current Scientific Understanding of the Processes Affecting*
419 *Terrestrial Carbon Stocks and Human Influences on them* (eds D. Schimel & M. Manning)
420 (National Oceanic & Atmospheric Administration, 2003).

421 32 IPCC. in *Revisiting the use of managed land as a proxy for estimating national*
422 *anthropogenic emissions and removals* (eds S Eggleston, N Srivastava, K Tanabe, & J
423 Baasansuren) (IGES, 2010).

424 33 Erb, K. H. *et al.* Bias in the attribution of forest carbon sinks. *Nature Climate Change* **3**, 854-
425 856 (2013).

426 34 UNFCCC. Synthesis report on the aggregate effect of the intended nationally determined
427 contributions, FCCC/CP/2015/7, <http://unfccc.int/resource/docs/2015/cop21/eng/07.pdf>
428 (UNFCCC, 2015).

429 35 UNEP. The Emissions Gap Report 2015,
430 http://uneplive.unep.org/media/docs/theme/13/EGR_2015_301115_lores.pdf. 1-98 (United
431 Nations Environment Programme, 2015).

432 36 Lapola, D. M. *et al.* Pervasive transition of the Brazilian land-use system. *Nature Clim.*
433 *Change* **4**, 27-35 (2014).

434 37 Forsell, N. *et al.* Assessing the INDCs' land use, land use change, and forest emission
435 projections. *Carbon Balance Manage* 11:26, doi: 10.1186/s13021-016-0068-3 (2016).

436 38 Houghton, R. A., Byers, B. & Nassikas, A. A. A role for tropical forests in stabilizing
437 atmospheric CO₂. *Nature Climate Change* **5**, 1022-1023 (2015).

438 39 Mackey, B. *et al.* Untangling the confusion around land carbon science and climate change
439 mitigation policy. *Nature Climate Change* **3**, 552-557 (2013).

440 40 Clarke, L. *et al.* in *Climate Change 2014: Mitigation of Climate Change. Contribution of*
441 *Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate*
442 *Change* (eds O. Edenhofer *et al.*) Ch. 6, 413-510 (Cambridge University Press, 2014).

443 41 Tollefson, J. Paris climate deal hinges on better carbon accountancy. *Nature* **529**, 450 (2016).

444 42 UNFCCC. Guidelines for the preparation of national communications from Parties not
445 included in Annex I to the Convention, Bonn, Decision 17/CP.8,
446 <http://unfccc.int/resource/docs/cop8/07a02.pdf> (UNFCCC, 2003).

447 43 GOFC-GOLD. A sourcebook of methods and procedures for monitoring and reporting
448 anthropogenic greenhouse gas emissions and removals associated with deforestation, gains
449 and losses of carbon stocks in forests remaining forests, and forestation, GOFC-GOLD
450 Report version COP21-1, University of Wageningen, the Netherlands,
451 <http://www.gofcgold.wur.nl/redd/>. (GOFC-GOLD, 2015).

452 44 GFOI. Integrating remote-sensing and ground-based observations for estimation of emissions
453 and removals of greenhouse gases in forests: methods and guidance from the Global Forest
454 Observations Initiative. (Global Forest Observations Initiative, 2014).

455 45 Hansen, M. C. *et al.* High-resolution global maps of 21st-century forest cover change.
456 *Science* **342**, 850-853 (2013).

457 46 Achard, F. & House, J. I. Reporting carbon losses from tropical deforestation with Pan-
458 tropical biomass maps. *Environmental Research Letters* **10**, doi:10.1088/1748-
459 9326/10/10/101002 (2015).

460 47 Global Forest Watch <http://www.globalforestwatch.org/>. (2016).

461 48 Pilli, R., Grassi, G., Kurz, W.A., Abad Viñas, R., Guerrero, N. Modelling forest carbon stock
462 changes as affected by harvest and natural disturbances. I. Comparison with countries'
463 estimates for forest management. *Carbon Balance Manage*, 11:5, doi: 10.1186/s13021-016-
464 0047-8 (2016)

465 49 Roman-Cuesta, R.M. *et al.* Multi-gas and multi-source comparisons of six land use emission
466 datasets and AFOLU estimates in the Fifth Assessment Report, for the tropics for 2000–2005.
467 *Biogeosciences*, **13**, 5799–5819 (2016).

468 50 Friedlingstein, P. *et al.* Update on CO₂ emissions. *Nature Geoscience* **3**, 811-812 (2010).

469 51 IPCC. in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group*
470 *I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds
471 T.F. Stocker *et al.*) Ch. SPM, 1-29 (Cambridge University Press, 2013).
472

473 **METHODS**

474

475 This analysis quantifies the mitigation role of Land Use, Land Use Change and Forestry (LULUCF,
476 mainly forests), based on the (I)NDCs^{3,4} submitted by Parties in the context of the Paris Climate
477 Agreement¹, complemented with information from other countries' official documents. This analysis
478 does not aim to assess specific country policies or the quality of country data in comparison with
479 independent sources.

480 Our analysis of LULUCF net emissions over time covered all 195 UNFCCC countries, with
481 assumptions necessary in some cases (i.e. using the latest historical data where no (I)NDC projection
482 was available, see below). However, due to constraints, our estimation of the LULUCF mitigation
483 contribution was possible only for 68 countries (41 (I)NDCs), covering 83% of global forest area
484 (based of FAO-FRA 2015²²). Other countries were not included either because LULUCF was not
485 clearly included in the target or because the LULUCF contribution was not entirely clear or directly
486 quantifiable (see Supplementary Section 1, Supplementary Information).

487 Our analysis is based on countries' documents submitted up to February 2016. However, the most
488 relevant recalculations made by countries after that date and before December 2016 (e.g. Brazil,
489 Indonesia and USA) are briefly discussed in the Supplementary Section 2.

490

491

492 **Information used in this analysis**

493 The methodological approach applied in this analysis required collecting information on:

- 494 (i) Countries' historical data and projections up to 2030 (for all 195 UNFCCC countries),
495 using countries' documents submitted up to end of February 2016, with the following
496 priority: (I)NDCs³; other country data submitted to UNFCCC (2015 GHG Inventories¹³)

497 (GHGI) for developed countries, and GHGIs included in recent National
498 Communications^{15,20} (NC) and in Biennial Update Reports¹⁶ (BUR) for developing
499 countries); other official countries' documents (e.g. ref.²¹); FAO-based datasets (for
500 forests^{8,22} and non-forest emissions²³). Despite gaps in country reports (especially for
501 non-forest land uses in developing countries), this priority is justified by the fact that
502 country reports to UNFCCC are formally reviewed or technically assessed by UNFCCC
503 (GHGIs of developed countries are formally reviewed annually, with biennial technical
504 assessment for developing country inventories), and are the means by which countries
505 assess their progress towards targets. Furthermore, FAO-FRA reports²² are not primarily
506 for reporting CO₂ emissions and removals, while UNFCCC country reports specifically
507 address emissions and removals. The range of historical country datasets (dotted line in
508 Fig. 1) reflects alternative selections of country sources, i.e. GHGIs for developed
509 countries (selected for both the lower and the upper range), plus FAO-based datasets
510 (upper range) or NCs (lower range) for developing countries. This alternative selection
511 assumes a high reliability of GHGIs for developed countries, while providing an idea of
512 the impact of choosing only NCs (including old NCs) vs. FAO-based datasets for
513 developing countries. See Supplementary Table 4 for an overview of historical datasets
514 used.

515 For **historical data**, GHGIs with a time series from 1990 to 2013 were available for all
516 developed countries, in most cases including Harvested Wood Products. For developing
517 countries, data are from BURs when available or from latest NCs, typically not including
518 Harvested Wood Products. When only few years were available (typically at least two
519 between 1990 and 2010), 5 or 10 years averages were used. Sometimes, especially for
520 older NCs, data from NCs contain ambiguities, or are not fully comparable across
521 countries (e.g. while most countries implicitly report only emissions and removals from
522 “managed forests”, in accordance with the recent IPCC guidance, a few countries include
523 the sink from apparently unmanaged forests). To reduce the risk of using old or

524 inappropriate data, the more recent FAO datasets were used instead of NCs prior to 2010.
525 Net emissions from forests (e.g., sink from forest management and emissions from
526 deforestation) usually dominate the LULUCF fluxes in country reports, although in some
527 case emissions from croplands and grasslands (rarely reported by developing countries)
528 are also relevant, especially from organic soils.

529 Based on available information from countries' reports to UNFCCC complemented by
530 expert judgment, we performed an analysis of the uncertainties for LULUCF absolute
531 GHG net emissions (level) and for the associated trends (see Supplementary Section 3,
532 Supplementary Table 7 and Supplementary Figures 2 and 3).

533 The FAO-based datasets include country data on forest carbon stock change from the
534 Forest Resource Assessment (FAO-FRA 2015²², as elaborated by ref⁸) and FAOSTAT²³
535 data on country-level non-forest land use emissions (CO₂, CH₄ and N₂O from biomass
536 fires, including peatlands fires, and from drainage of organic soils). The overall small
537 difference between the FAO-FRA 2015 forest carbon stock data used in our analysis
538 (based on ref.⁸) and the FRA-2015 forest carbon stock data included in FAOSTAT²³ is
539 that the gap-filling methods differ (although for the biomass pools such difference does
540 not impact the total net CO₂ emissions/removals across the time series), and that we
541 include both living biomass (above and below-ground) and dead organic matter if
542 reported by countries, while FAOSTAT only considers living biomass. Overall, for the
543 historical period we only used FAO-based datasets to fill the gaps for a relative large
544 number (60), but typically rather small developing countries (covering 11% of global
545 forest area). The significant difference between this analysis and FAOSTAT (Fig. 2 of the
546 paper) is due to several factors, including higher non-forest land use emissions in
547 FAOSTAT for developing countries (especially in Indonesia, Sudan, Zambia) and higher
548 forest land use emissions in FAOSTAT for both developing countries (e.g. Colombia,
549 Liberia, Madagascar, Myanmar, Nigeria, Philippines, Zimbabwe) and developed ones
550 (USA and Russia).

551 For **projections**, data from (I)NDCs (with some expert-judgment interpretation when
552 needed), or NCS²⁰ were available for almost all developed countries. For developing
553 countries, if no projection was available in the (I)NDCs, BURs or NCS, FAO-FRA 2015
554 country projections^{8,22} were used in few cases. Where no projection was available, the
555 latest historical country data available were used (i.e. continuing the recent estimates).

556 While almost no country provided formal information on uncertainties in their
557 projections, in the analysis of uncertainties (see Supplementary Section 3) we
558 conservatively assumed that the uncertainties estimated for the past will hold for the
559 future. In addition, the different scenarios that our analysis identified up to 2030 (Fig. 1)
560 may also give an order of magnitude of the uncertainties. The range “LULUCF
561 projections min-max” shown in Fig. 1 is slightly broader than the various scenarios (by
562 about 500 MtCO₂e/y, or 0.5 GtCO₂e/y, in 2030) because in few cases countries provide a
563 range of projections and not all the various projections can be associated with the four
564 scenarios analyzed. The overall difference of about 500 MtCO₂e/y is essentially due to
565 the range of projections from the US (the difference between the “high” and a “low”
566 sequestration scenario in their latest National Communication amounts to 370 MtCO₂e/y
567 in 2030), and due to Russia (the difference between the various sequestration scenario in
568 their latest National Communication amounts to about 150 MtCO₂e/y in 2030).

569 With regards to the GHGs considered, this paper focuses on CO₂ emissions and removals
570 and on available data on non-CO₂ emissions (CH₄ and N₂O), based on the information
571 included in countries’ documents. National GHGIs are required to report on all GHGs,
572 but in some developing countries the information on non-CO₂ gases is incomplete. Based
573 on available information, and excluding agricultural emissions, the importance of non-
574 CO₂ gases is typically small relative to the total GHG fluxes (see ref³⁰ for details),
575 representing about 2-3% of total CO₂-equivalent forest flux, with slightly higher values
576 found where forest fires are important in the overall GHG budget. This suggests that,
577 when comparing different datasets (Fig. 2), the possible different coverage in the

578 (I)NDCs and other documents of non-CO₂ gases does not represent a major reason for
579 discrepancy.

580 (ii) *Type of mitigation target elaborated in each countries' (I)NDC* (Supplementary Table 1),
581 i.e. change in absolute emissions or intensity, either relative to a base year or to a BAU
582 scenario (i.e. 2025 or 2030 scenario year); target 'unconditional' or 'conditional' (i.e.
583 related to the provision of finance, technology or capacity-building support). (I)NDCs
584 expressing only 'policies and measures' (without quantitative targets) were not taken into
585 account.

586 (iii) *Modality of inclusion of LULUCF within each countries' (I)NDC* (Supplementary Table
587 1), i.e. it may be treated in the same way as other sectors (fully included as part of the
588 overall target), or partially included (only forest activities), or considered separately with
589 special mitigation actions and/or accounting rules.

590 Some additional expert evaluation was included where necessary.

591

592 **(I)NDC cases**

593 The (I)NDCs were classified into four '(I)NDC cases' (Supplementary Table 2). Based on the
594 availability of country LULUCF information, enough information was found to assign 68 countries to
595 these different "(I)NDC cases", and to quantify directly the expected LULUCF mitigation. These 68
596 countries include all countries with a major forest coverage and correspond to 78% of global
597 emissions in 2012 (including LULUCF emissions and international aviation and marine emissions)².

598

599 **Different mitigation perspectives**

600 The quantification of the mitigation role of LULUCF has been undertaken using different approaches,
601 reflecting different perspectives, according to the questions addressed (Supplementary Table 3).

602

603 **Estimation of LULUCF mitigation**

604 Whereas estimates for perspective ‘A’ (LULUCF net emissions over time) could be made for all 195
605 UNFCCC countries, the information needed for the LULUCF mitigation contribution under
606 perspectives ‘B’ ((I)NDC compared to alternative future scenarios) and ‘C’ (country perspective on
607 calculating emissions reduction (I)NDC) was available only for the 68 countries (41 (I)NDCs)
608 included in Supplementary Table 1. For the remaining countries, the additional mitigation in
609 perspectives ‘B’ and ‘C’ were assumed to be zero relative to other sectors. This assumption is
610 probably conservative (see Supplementary Section 1).

611 Based on the four (I)NDC cases (Supplementary Table 2), and using the available country
612 information (generally with limited expert judgment), this analysis quantified the LULUCF mitigation
613 perspectives (Supplementary Table 3) following the method illustrated in Supplementary Fig. 1. In
614 the very few cases where the target is expressed for 2025, we assumed that the same target applies to
615 2030, allowing us to sum up all the countries’ contribution to 2030.

616

617 **Contribution of the land sector to mitigation activity across all sectors**

618 The LULUCF mitigation perspectives ‘B’ and ‘C’ were compared to the expected (I)NDC mitigation
619 efforts across all sectors, for each country and at a global level. The global-level all-sectors ‘pre-
620 (I)NDC’ and ‘(I)NDC unconditional + conditional’ are taken from UNEP³⁵. All-sector emissions at
621 the ‘reference point’ (i.e. base year or BAU scenario for target year 2025 or 2030) are from: (i)
622 countries or (ii) from ref¹⁸ (for the BAU estimates for China and India). These two sources of
623 information were sufficient for countries representing 87% of global GHG emission in 2012.
624 Emissions for the remaining countries were approximated by assuming the same ratio of emissions at
625 reference point (i.e. estimates from available sources were multiplied by 100/87).

626

627 **Comparison of this analysis with IPCC AR5**

628 In order to make a meaningful comparison of country data (this analysis) with IPCC AR5^{5,6}, we
629 disaggregated country data between “land converted to another land use” and “land remaining under
630 the same land use”. While this disaggregation was directly available in all developed country reports,
631 and was largely available for the most important developing countries (e.g. Brazil, Indonesia, India,
632 China, Mexico), for the remaining developing countries information was generally available only for
633 deforestation. In these cases, unless specified otherwise, the other emissions and removals were
634 assigned to “land remaining under the same land use”.

635

636 **Data availability**

637 This study is primarily based on countries’ (I)NDCs^{3,4} and other GHG reports submitted to
638 UNFCCC^{13,15,16,20,21}, complemented by FAO-based datasets^{8,22,23}. A large part of elaborated data used
639 to support our findings are available in the Supplementary Information, including:

640 (i) *Country-specific information* for 68 countries (41 (I)NDCs), in terms of general features
641 of the (I)NDCs (Supplementary Tables 1 and 2) and of data and sources of information of
642 LULUCF net emissions for the historical period 1990-2010 and for 2030, as expected for
643 unconditional and conditional (I)NDC targets (Supplementary Table 5).

644 (ii) *Aggregated information* on uncertainties (Supplementary Figures 2 and 3), on LULUCF
645 mitigation perspectives (Supplementary Table 3) and on LULUCF net emissions
646 (Supplementary Table 6).

647 Any other raw or elaborated data used in this study are available from the corresponding author upon
648 request.

649