

THE FLORA AND VEGETATION OF THE MUREŞ (MAROS) VALLEY

CONSTANTIN DRĂGULESCU

Introduction

Having its springs in the Oriental Carpathians, running across the Transylvanian Plateau, separating the Occidental Carpathians from the Meridional ones and finally flowing through the plain from western Romania and eastern Hungary, the Mureş and its valley include several various ecotopes with a rich flora and diverse vegetation.

Although some botanical investigations were done in the Mureş Valley by J. Lerchensfeld and P. Sigerus as early as the 18th century (*Herb. Sibiu*), the first pieces of botanical information were given by J. Chr. G. Baumgarten (1816), P. Kitaibel (1863), E. Schur (1866), E. Porcius (1878), A. Cserni (1879), L. Walz (1878), J. Csató (1896), A. Halász (1889) and particularly by M. Fuss (1866) and L. Simonkai (1886, 1893).

During the 20th century floristic and phytocenological research was conducted by E. I. Nyárády (1914, 1931), I. Prodan (1928), L. Tímár (1952, 1957), R. Soó (1938, 1940, 1943), St. Csürös (1956, 1970), E. Pop (1960), Al. Borza, V. Lupşa (1964-1965, 1968), I. Gergely (1964), Fl. Rațiu (1968, 1969, 1971, 1972), I. Pop (1978, 1979), E. Täuber (1973, 1986), R. Samu (1982) and others. "Flora R.P.R.", respectively "Flora R.S.R.", collating the results of these botanists, returned 915 species of cormophytes in the Mureş Valley.

This study presents an enumeration of the flora and vegetation which grows in the Mureş river valley, from its spring (*Izvorul Mureşului*) to its confluence with the Tisza River (Szeged).

Methods

In order to offer a floristic-phytocenologic and ecological characterization as detailed as possible of this river, and, also to reduce the dimensions of our study especially of the chorological informations (the river runs through more than 100 localities), we divided the whole water course of the river into five segments (sectors), pointed out in the study by roman numerals I through V: I. Izvorul Mureşului- Topliţa, respectively the Depression of Giurgeu (Gheorgheni); II. Topliţa-Deda, which corresponds to the Mureş Strait between the Mts. of Călimani and Gurghiu; III. Deda-Alba Iulia or the central-Transylvanian water course of the Mureş; IV Alba Iulia-Lipova or the Mureş Corridor between the Occidental Carpathians (Metalliferous and Zărand Mts.) and the Meridional Carpathians (Sureanu Mts., Poiana Rusă Mts.); V. Lipova-Szeged, the flatland course of the river.

These segments distinguish themselves not only by discharge and water speed of the river, but also by its level of pollution, by zooanthropic influences on woody and grassy

vegetation which grows on both banks of the waterway, by the geological structure, the geomorphology of the valley, etc.

Relying on the consulted bibliography and after our site investigations, we drew up a card index of species and plant associations. The index was set up according to the phylogenetic system used in "Flora R.S.R." from which we also adopted the names of the species with the underlined corrections in volume XIII. Each plant species is accompanied by its ecological indices (humidity, temperature and soil reaction), the life strategy (life form), the floristic element (geoclement) and the waterway segment where it grows. The value of the ecological indices is given according to R. Soó (1964-1980) and V. Sanda et al. (1983).

Classification of the life forms follows C. Raunkiaer's system, developed by J. Braun-Blanquet (1951). The determination of the floristic elements was carried out according to the works of R. Soó (1964-1980) and V. Sanda et al. (1983).

For vegetation analysis we adopted J. Braun-Blanquet's method, adjusted for conditions in Romania and Hungary. The classification of the plant associations is in accordance with the systems of E. Oberdorfer (1970, 1977) and R. Soó (1964-1980). By each plant association, the waterway segment is marked where they occur.

Abbreviations in the text

Life-forms: Ph- phanerophyte (MPh- megaphanerophyte, mPh- mezophanerophyte, nPh- nanophanerophyte); Ch- chamaephyte; H- helicryptophyte; G geophyte; T therophyte (Th- annual, TH- biennial); Hh- helohydrophyte; Ep- epiphyte.

Floristic elements: Cosm- cosmopolitan; Cp- circumpolar; Eua- Eurasian; E- European; Ec- Central-European; Atl-M- Atlantic-Mediterranean; M- Mediterranean; MP- Mediterranean-pontic; P- Pontic; Pn- Pannonic; B- Balcanic; D- Dacic; Carp- Carpathic; Alp- Alpin; Cauc- Caucasian; Anat- Anatholian; Adv- adventive.

Ecological indexes: H- humidity; T- temperature; R- soil reaction.

I, II, III, IV, V- waterway segments of Mureş.

The mark (!) indicates that plant (or association) was noticed by the author.

Results

The flora of the Mureş Valley

After volumes I-XIII of "Flora R.S.R." had been issued, 472 species were newly identified by us and also published by different botanists. Our research, correlated to the data published by other authors (see References), point out 1316 species of higher plants (cormophytes) plus 71 hybrids (altogether 1387 items), to which must be added 60 subspecies, 231 varieties and 168 forms. These taxa belong to 502 genera of 117 families. Among genera Carex (53 species), Hieracium (32 species), Euphorbia (25 species), Centaurea, Ranunculus (22-22 species), Veronica (21 species), Rumex, Trifolium (19-19 species), Silene (17 species), Chenopodium, Cirsium (14-14 species), Juncus, Salix, Viola (13-13 species), Potentilla, Senecio (12-12 species), Potamogeton (11 species), Rorippa and Vicia (11-11 species) are better represented. The families richest in species are Asteraceae

(206 species), Poaceae (100 species), Cyperaceae (68 species), Brassicaceae (67 species), Caryophyllaceae (65 species), Fabaceae (64 species), Rosaceae (54 species), Lamiaceae, Ranunculaceae (52-52 species), Scrophulariaceae (51 species), Apocynaceae (49 species), Liliaceae (37 species), Chenopodiaceae (32 species) and Polygonaceae (31 species).

From all the 1846 referred taxa, 220 spp. are considered to be rare or protected plants, such as *Botrychium multifidum*, *Marsilea quadrifolia*, *Taxus baccata*, *Betula humilis*, *Betula x warnstorffii*, *Silene x grecescui*, *Euphorbia x csatoi*, *Trollius europacus*, *Nymphaea alba*, *Nuphar luteum*, *Elatine alsinastrum*, *Sempervivum heuffelii*, *Ribes heteromorphum*, *Spiraea salicifolia*, *Trapa natans*, *Vitis silvestris*, *Pleurospermum austriacum*, *Cnidium dubium*, *Lysimachia thyrsiflora*, *Pedicularis sceptrum-carolinum*, *Plantago schwarzengergiana*, *Nymphoides peltata*, *Bryonia dioica*, *Achillea x girgioensis*, *Petasites x celakovskyi*, *Ligularia sibirica*, *Centaurea x nyaradiana*, *Hieracium x auriculoides*, *Typha laxmannii*, *Lilium martagon*, *Fritillaria meleagris*, *F. montana*, *Scilla autumnalis*, *Streptopus amplexifolius*, *Tamus communis*, *Iris nyaradyana*, *Traunsteinera globosa*, *Calla palustris*, *Arum maculatum*. The 1316 species from the following list were analysed by ecological preferences (humidity, temperature, soil reaction), life strategies (life forms) and the floristic element to which they belong in order to give a characterization of the flora of the Mureş Valley from spring to mouth, as accurate and detailed as possible.

The flora

LYCOPODIACEAE

1. *Lycopodium annotinum* L.: I-II (22, 23); Ch, Cp; H4T2.5R2

EQUISETACEAE

2. *Equisetum arvense* L.: I-V; G, Cosm; H3T3R0
- *f. obtusatum* Kluge: IV-V (43, !)
3. *E. fluviatile* L. (*E. limosum* L.): I-III (6, 23, 31, 22, 14, !); Hh, Cp; H5T3R0
4. *E. hiemale* L.: II (14); G, Cp; H3.5T2.5R4
5. *E. x littorale* Kuhl. (*arvense x fluviatile*): III (5, 14);
6. *E. palustre* L.: I-V; G, Cp; H5T2R0
7. *E. pratense* Ehrh.: II (14); G, Cosm; H3.5T2.5R4
8. *E. ramosissimum* Desf.: III-V (14, 5, 36, 40, 25, 18); G, Cosm; H2T0R0
- *f. simplex* (Doll) Milde: III (14)
9. *E. siliculosum* L.: I (14, 23, 30, !); G, H3.5T2R0
10. *E. telmateja* Ehrh. (*E. maximum* Lam.); II-IV (22, 14, !); G, Cp; H3.5T2R0

OPHIOGLOSSACEAE

11. *Ophioglossum vulgatum* L.: I (14), V (37, 40, 14); G, Cp; H4T3R0
12. *Botrychium lunaria* (L.) Sw.: I (14); G, Cosm; H2.5T2R0
13. *B. multifidum* (S.G. Gmel.) Rupr.: I (14); G, Cp; H2.5T2R0

HYPOLÉPIDACEAE

14. *Pteridium aquilinum* (L.)Kuhn: II-V (22,14,25,40,!); G, Cosm; H3T3R0

ASPLENIACEAE

15. *Asplenium adiantum-nigrum* L.: III-IV (14,29,25,43); H, E; H2.5T3R3
- *f. argutum* Heuil.: IV (14,25)
16. *A. ruta-muraria* L.: IV V (14,37,40,!); H, Cp; H1.5T3R5
17. *A. septentrionale* (L.)Hoffm.: II (14), IV (14,25); H, Cp; H1T3R2
18. *A. trichomanes* L.: II (22,!); IV (14,37,25,5,!); H, Cp; H3T0R4
19. *Phyllitis scolopendrium* (L.)Newman: II (14,!); IV (14); G, Cp; H3.5T3R5

ATHYRIACEAE

20. *Athyrium filix-femina* (L.)Roth: I-II (14,22,23,30,!); IV (14,37,25,!); H, Cosm;
H4T2.5R0
21. *Cystopteris fragilis* (L.)Bernh.: II (22,!); IV (14,37,25); H, Cosm; H3.5T0R0
- ssp. *fragilis* f. *anthriscifolia* (Hoffm.)Koch: IV (14) and f. *angustata* Koch: IV (14,37,25)
- ssp. *alpina* Hartm. (C. *regia* (L.)Presl.): IV (14)
22. *C. sudetica* A. Br. et Milde: II (22); H, Eua; H3.5T2R0
23. *Matteuccia struthiopteris* (L.)Tod.: II (22,!); IV (25); H,Cp; H4T2R0

ASPIDIACEAE

24. *Polystichum braunii* (Spenn.)Fee: II (14,22); H, Cp; H3.5T2.5R3
25. *P. setiferum* (Forsk.)Woynar: IV (14,25); H, Cosm; H3.5T0R4
26. *Dryopteris carthusiana* (Vill.)H.P Fuchs (*D. spinulosa* O.F. Mull.): I (23); H, Cp;
H4T3.5R0
27. *D. cristata* (L.)A. Gray: I (14,23,!); H, Cp; H3.5T2R3
28. *D. filix-mas* (L.)Schott: I-V; H, Cosm; H4T3R0

THELYPTERIDACEAE

29. *Thelypteris palustris* Schott (*Dryopteris thelypteris* (L.)A. Gray): I (23); Hh, Cp;
H4T0R3
30. *T. phegopteris* (L.)Slasson (*Phegopteris polypodioides* Fcc): I-II (22,23); G, Cp;
H3.5T2R2

POLYPODIACEAE

31. *Polypodium vulgare* L.: II (22,!); IV (25); G, Cp; H3.5T3R4

MARSILEACEAE

32. *Marsilea* (*Marselia*) *quadrifolia* L.: IV (14,25,38); Hh, Eua (M); H6T3R0

SALVINIACEAE

33. *Salvinia natans* (L.)All.: IV V (37,25,38,14,!); Hh, Eua; H6T3R3

TAXACEAE

34. *Taxus baccata* L.: II (14,22); mPh, E; H3T3.5R4

PINACEAE

35. *Abies alba* Mill.: II (22); MPh, Ec; H4T3R0

36. *Picea abies* (L.)Karst.: I-II (22,14,!); MPh, E; H0T0R0

37. *Pinus silvestris* L.: I-II (22,14); MPh, Eua; H0T0R0

CUPRESSACEAE

38. *Juniperus communis* L.: I (6,23,30,!); IV (25); mPh, Cp; H2T0R0

EPHEDRACEAE

39. *Ephedra distachya* L.: III (14,4,5); mPh, Eua (C); H2T4.5R4.5

BETULACEAE

40. *Betula humilis* Schrank: I (23,31,32); mPh, Eua; H5T2.5R1

41. B. x *hybrida* Bechst. (pubescens x pendula): I (14,23,!);

42. B. *pendula* Roth (B. *verrucosa* Ehrh.): I-II (6,23,22,!); III (29), IV (5,25); MPh-mPh, Eua; H3T2R2

43. B. *pubescens* Ehrh.: I-II (14,23,6,30,31,32,!); MPh-mPh, Eua; H0T0R0

- f. *glabra* (Fieck.)Georgescu: I (14)

44. B. x *warnstorffii* C.K.Schneid. (*humilis* x *pubescens*): I (23);

45. *Alnus glutinosa* (L.)Gaertn.: I-V; Mph-mPh, Eua; H5T3R3

46. A. *incana* (L.)Mnch.: I-IV (30,22,14,!); MPh-mPh, Eua; H4T2R4

CORYLACEAE

47. *Carpinus betulus* L.: II-IV (14,22,29,5,!); V (40); MPh-mPh, E; H3T3R3

48. *Corylus avellana* L.: I-IV; mPh, E; H3T3R3

FAGACEAE

49. *Fagus sylvatica* L.: II (22,!); MPh-mPh, E; H3T3R0

50. *Quercus cerris* L.: IV V (14,37,25,24,27,!); MPh-mPh, M; H2T3.5R3

- f. *austriaca* (Willd.)Hegi: V (40)

51. Q. *frainetto* Ten.: IV (14,37,25); MPh, B; H2T4R3

52. Q. *petraea* (Matt.)Liebl.ssp. *petraea*: II-IV (22,29,5,25,!); MPh-mPh, E; H2.5T3R0

53. Q. *pubescens* Willd.: IV (14,25), III (14); MPh, M; H1.5T4R5

- var. *undulata* (Kit.)Schwz.: IV (43)

54. Q. *robur* L.: II-V; MPh, E; H3.5T3R0

- var. *tardiflora* Cernaiov: V (14,40)

- var. *puberula* (Lasch)Schwz.: V (40)

55. Q. *virgiliiana* (Ten.) Ten.: I V (14,5); MPh, M; H2T4R4

SALICACEAE

56. *Populus alba* L.: III-V (14,24,18,!); MPh-mPh, Eua; H3.5T3R3
 57. *P. x canescens* Sm. (*alba* x *tremula*): V (18,40,!);
 58. *P. nigra* L.: III-V (14,25,27,40,!); MPh, Eua; H4T3R4
 59. *P. tremula* L.: I-V; MPh-mPh, Eua; H3T2R2
 60. *Salix alba* L.: I-V; Mph-mPh, Eua; H5T3R4
 61. *S. aurita* L.: I (14); mPh, H4.5T0R2
 62. *S. caprea* L.: III-V (23,27,5,!); mPh, Eua; H3T3R3
 63. *S. cinerea* L.: I-V; mPh, Eua; H3T3R3
 64. *S. daphnoides* Vill.: III (14); mPh, E; H4.5T2.5R4.5
 65. *S. fragilis* L.: I-V; mPh-MPh, Eua; H4.5T3R4
 66. *S. pentandra* L.: I-III (14,22,23,31,29,!); MPh, Eua; H4.5T0R3.5
 67. *S. purpurea* L.: I-V; mPh, Eua; H5T3R4.5
 - f. *angustifolia* Kern.: IV (25)
 68. *S. rosmarinifolia* L.: I (14,23,6,32,!); mPh, Eua; H4T0R0 69. *S. x rubens* Schrank (*alba* x *fragilis*): V (40);
 - var. *excelsior* (Host.) A. ct G.: V (40,27)
 - var. *palustris* (Host.) Seem: V (40)
 70. *S. triandra* L.: I-V; mPh, Eua; H5T3R0
 - f. *glaucophylla* Scr.: V (40)
 - f. *semperflorens* (Host.) Beldie: I (!), V (40)
 71. *S. x undulata* Ehrh. (*alba* x *triandra*): IV V (14,37,40);
 72. *S. viminalis* L.: I-V; mPh, Eua; H5T2R4.5
 - f. *tenuifolia* Kern.: I (23,!)

ULMACEAE

73. *Ulmus glabra* Huds. (*U. montana* Stokes): II (22). V (24,27,40); mPh-MPh, Eua; H4T3R3
 74. *U. levis* Pall.: V (14,25,40,27,!); Mph-mPh, E; H4T3R3
 75. *U. minor* Mill. (*U. foliacea* Gilib.): III-V (37,40,9,5,24,!); MPh, Eua; H3T3R4
 - var. *minor* f. *carpinifolia* (Bokh.) Beldie: IV (25) and f. *suberosa* (Henry) Beldie: IV V (25,43,40,!)
 - var. *stricta* (Lindl.) Rehd.: IV (25)
 - var. *asperrima* Simk.: IV (43)

CANNABINACEAE

76. *Humulus lupulus* L.: I-V; H, Eua; H3.5T3R4

URTICACEAE

77. *Urtica dioica* L.: I-V; H-G, Cosm; H3T3R4
 - f. *pubescens* (Ldb.) Trautv : V (14)
 78. *U. urens* L.: III-V (5,18,!); Th, Cosm; H3T3R4
 79. *Parietaria officinalis* L.: IV (14,25); H, M; H4T3.5R4

SANTALACEAE

80. *Thesium alpinum* L.: I (14); H, E; H2.5T2R2.5
 81. *T. arvense* Horvatovszky (*T. ramosum* Hayne): V (18,40); TH-H, Eua(C):
 H1.5T3.5R4.5
 82. *T. dollineri* Murb.: IV V (14,5,40); Th-H, P-B; H2T3R0
 83. *T. linophyllum* L.: IV (14,5,!); G-H, Ec; H2T4R4

LORANTHACEAE

84. *Loranthus europaeus* Jacq.: III-V (14,37,25,40,29); Ch-nPh, E:H3T3.5R0
 85. *Viscum album* L.: III-V (14,25,!); Ch-nPh,Eua; H3.5T3R0

POLYGONACEAE

86. *Polygonum amphibium* L.: I (31), III-V (29,5,25,40,!); G-Hh, Cosm; H6T3R0
 - f. *aquaticum* (Leyss)Grint.: IV-V (25,40,14,!)
 - f. *terrestrre* (Leyss)Grint.: I (31)
 87. *P. arenarium* W. et K.: IV (14,25); Th, P-Pn; H1T3.5R0
 88. *P. aviculare* L.: I-V; Th, Cosm; H2.5T0R3
 - var. *erectum* (Roth)Hayne: V (40)
 89. *P. bistorta* L.: I-III (14,30,23,6,22,29,!); H,Eua; H4T2.5R3
 - f. *puberulum* Beck: I (14)
 90. *P. hydropiper* L.: I-V; Th, Eua (M); H4.5T3R4
 91. *P. lapathifolium* L.: III-V (14,9,6,27,40,!); Th,Cosm; H4T0R4
 - var. *tomentosum* (Schrank)Beck: III-V (14,26,40)
 - f. *lanceolatum* (A.Br.)A. et G.: V (40)
 - var. *brittingeri* (Opiz)Beck: V (14,40)
 92. *P. minus* Huds.: III-V (14,5,25,40);Th, Eua; H5T3R4
 93. *P. mite* Schrank: III (14), V (14,25,40); Th, Eua : H5T3R4
 - f. *angustifolium* (A.Br.)Beck: III (14)
 94. *P. persicaria* L.: III-V (29,14,27,25,40,!); Th, Eua; H4.6T3R0
 95. *P. x subglandulosum* Borb. (*hydropiper x minus*): II (14);
 96. *Bilderdykia convolvulus* (L.)Dumort (*Fagopyrum convolvulus* (L.) H. Gross): III-V
 (14,5,25,18,47); Th, Eua; H2.5T3R3
 97. *B. dumetorum* (L.) Dumort (*Fagopyrum* (L.)Schreb.): IV V (14,37,40,25,43,!); Th, Eua
 (M); H2.5T3R3
 98. *Rumex acetosa* L.: I-V; H, Cosm; H3T0R0
 99. *R. acetosella* L.: I-V; H-G, Cosm; H2T3R2
 - f. *multifidus* (L.)Prod.: I (14), III (14)
 100. *R. x acutus* L. (*crispus x obtusifolius*): III (14,29), IV (14);
 101. *R. aquaticus* L.: I (14,30,23,31,!); V (14); Hh, Cp; H4.5T0R4
 102. *R. conglomeratus* Murray: III-V (14,40,27,5,!); H, Cp; H4T4R4
 103. *R. crispus* L.: I-V; H, Eua; H4T3R0
 104. *R. x erubescens* Simk. (*obtusifolius x patientia*). III (14);
 105. *R. x heteranthos* Borb. [(*maritimus x conglomeratus*)x *stenophyllus*]:V (14,40);
 106. *R. hydrolapathum* Huds.: III-V (14,29,5,40,!); H(G), E; H6T4R4
 107. *R. x intercedens* Rech (*crispus x stenophyllus*) ssp. *aradensis* Prod.:V (14)

108. *R. kerneri* Borb.: V (14,25,40); H, D-Pn-B; H3T5R3
 109. *R. maritimus* L.: III (14). V (14,40,!); Th, Eua: H5T3.5R4.5
 110. *R. obtusifolius* L.: III-V (14,18,40,!); H, E; H4T0R3
 - ssp. *silvestris* (Lam.)Rech: III-V (14,40)
 - ssp. *transiens* (Simk.)Rech f: III (14)
 - ssp. *subalpinus* (Schur)Simk.: III, V (14)
 111. *R. palustris* Sm. (*R. limosus* Thuill.): III-V (14,5,40,!); Th-TH,Eua;H5T3R4
 112. *R. patientia* L.: I (14), II-V (5,25,14,40,!); H, Eua (C); H3T4R0
 - ssp *recurvatus* (Rech)Rech f: I, III-IV (14)
 113. *R. pulcher* L.: IV V (14,25,40); Th-TH, Atl-M; H4T3R3
 114. *R. sanguineus* L.: III-V (14,37,25,!); H, E ; H4T3R4
 - var. *viridis* (Sibth.)Koch: V (18,40)
 115. *R. stenophyllum* Ldb.: III-V (14,34,18,45,!); H, Eua (C); H5T4R4
 116. *R. x stenophyloides* Simk. (*maritimus* x *stenophyllum*): V (14,37,40)

CHENOPODIACEAE

117. *Polyneum arvense* L.: IV V (14,25,47,40); Th, Eua (M); H2T3R3
 118. *P. majus* A. Br.: III-V (14,5,40); Th, Eua (M); H1.5T4.5R4
 119. *Chenopodium album* L.: I-V; Th, Cosm; H3T3R0
 - ssp. *spicatum* (Koch)Nyár. var. *pracacutum* (Murr)Prod.: IV V (14,25)
 - f. *laciniatum* (Murr)Prod.: V(14)
 - ssp. *viride* (L.)Murr var. *rhombeum* Peterm.: III (14)
 - ssp. *striatum* (Kras.)J. Murr (*C. strictum* Roth): V (47)
 - f. *krasani* Beck: V (14)
 120. *C. bonus-henricus* L.: I-V; H, E; H3.5T2R3
 121. *C. botrys* L.: III-V (14,9,5,40,25,43); Th, Cosm; H3.5T4R0
 122. *C. ficifolium* Sm.: III (14), V (14,40); Th, Eua (M); H3.5T4R0
 123. *C. foliosum* (Mxch.)Aschers.: II (14); Th, Eua (M); H0T0R0
 124. *C. glaucum* L.: III-V (29,5,14,45,!); Th,Eua: H3.5T4R0
 125. *C. hybridum* L.: III-V (14,29,25,47,40); Th, Eua (M);H3T3R0
 126. *C. murale* L.: III (14); Th, Adv; H2.5T4R0
 127. *C. opulifolium* Schrad.: III-V (14,42,40); Th, Eua: H2T3.5R4
 128. *C. polyspermum* L.: IV V (5,47,40,37); Th, Eua; H3T4R0
 129. *C. rubrum* L.: III (14,42), V (14,40); Th, Cp; H3.5T0R0
 - ssp. *blitoides* (Lej.)A. et G.: V (14,37,40)
 130. *C. schraderianum* Schult. (*C. foetidum* Schrad.): III (14); Th, Adv; H2T3.5R4
 131. *C. urbicum* L.: III-V (14,5,25,47); Th, Eua (M); H3T0R3
 132. *C. vulvaria* L.: III-V (14,5,40); Th, Eua (M); H3T4R4
 133. *Atriplex hastata* L.: III-V (14,37,40); Th,Cp; H3.5T0R0
 - var. *microtheca* C.F.Schumach: V (40,47)
 134. *A. littoralis* L.: III (14), V (14,45); Th, Eua (M); H0T0R0
 135. *A. nitena* Schuhr.: III-V (14,25,40,!); Th, Eua (C); H3T3R0
 136. *A. oblongifolia* W et K: III-IV (14,5,25); Th, Eua (C); H2T3.6R4
 137. *A. patula* L.: III, V (14,18); Th,Cp; H0T0R0

- var. *angustifolia* (Sm.)Lange: V (14)
- var. *erecta* (Huds.)Lange: V (14,40)
- 138. *A. rosea* L.: III-V (14,37,40); Th, Eua (C); H3.5T3R3
- 139. *A. tatarica* L.: III (14), V (18,47); Th, Eua (M); H2T4R0
- 140. *Camphorosma annua* Pall. (*C. ovata* W et K.): V (13,18); Th, P-Pn; H2T4R5
- 141. *Kochia laniflora* (Gmel.)Borb.: IV (14); Th, Eua(C); H1.5T4.5R4.5
- 142. *K. prostrata* (L.)Schrad.: III (14,4); Ch-nPh,Eua (C);H1.5T4R4.5
- 143. *K. scoparia* (L.)Schrad.: III (14), V (14,40); Th, Eua (Adv);H3T3.5R0
- 144. *Corispermum nitidum* Kit.: IV (14); Th, P-Pn; H2.5T4R0
- 145. *Salicornia europaea* L. (*S. herbacea* L.): III (14,29,!); Th,Cosm; H4T0R5
- 146. *Salsola kali* L. ssp. *ruthenica* (Iljin)Soó (*S. ruthenica* Iljin): V (14,40); Th, Eua (C); H0T4R4
- f. *tenuisolia* Tauschin: V (40)
- 147. *Suaeda maritima* (L.)Dum.: V (18,40); Th, Cosm; H4.5T3.5R5
- 148. *Petrosimonia triandra* (Pall.)Simk.: III (14,29); Th, Eua(C); H2T4R4.5

AMARANTHACEAE

- 149. *Amaranthus albus* L.: III-V (9,25,47); Th,Adv; H3T3R3
- 150. *A. blitoides* S. Watson: V (46,47); Th,Adv; H2T4R0
- 151. *A. crispus* (Lesp. et Thev)N. Terrac.: III-V (14,9,25,!);Th, Adv; H3T4R3
- f. *ruber* F. Zimm. et Thell.: III (14,!)
- f. *macrophyllus* Deg. et Thell.: IV V (14)
- 152. *A. deflexus* L.: V (14,40); H, Adv; H2.5T4R4
- f. *scandens* (L. f.)Thell.: V(14)
- 153. *A. hybridus* L. (*A. hypochondriacus* L.): III (14), V (47);Th,Adv;H3T3R0
- var. *chlorostachys* (Willd.)Thell.: V (47)
- 154. *A. lividus* L.: III (14), V (14,40); Th, Cosm; H3.5T4R4
- var. *ascendens* (Lois)Thell.: V (37,40,18)
- f. *procumbens* (Spenn.)Morariu: III (14)
- 155. *A. retroflexus* L.: III-V (14,5,47,!); Th, Adv; H3T3R0

PORTULACACEAE

- 156. *Portulaca oleracea* L.: III-V (9,5,40,18,!); Th, Cosm; H3T0R0

CARYOPHYLLACEAE

- 157. *Myosoton aquaticum* (L.)Mnch. (*Stellaria aquatica* (L.)Scop.): III-V (9,29,14,25,27,18); Th-TH, Eua (M);H4T3R0
- 158. *Stellaria graminca* L.: I-V; H, Eua (M); H2.5T2R3
- 159. *S. holostea* L.: III-V (29,5,25,37,40,!);H-Ch, Eua; H3T3R0
- 160. *S. longifolia* Muhl.: I (23,32); H, Cp; H4.5T2R2
- 161. *S. media* (L.)Cyr.: I-V; Th-Th, Cosm;H3T0R0
- 162. *S. nemorum* L.:I-V; H, E; H3.5T3R3
- 163. *S. palustris* Ehrh.: IV (14,5); H, Eua; H4.5T3R3
- 164. *Cerastium arvense* L.: I (14), III (14,29), IV (5); Ch, Cp, H2.5T0R3.5

165. *C. brachypetalum* Desp.: III-V (14,37,47,40,!); Th, M; H3T3R0
 - f. *strigosum* Fries: III (14)
 - f. *taurinum* (Spr.)Prod. : IV V (14,40)
166. *C. dubium* (Bast.)Guepin (*C. anomalum* W et K.): III-V (35,14,25);Th, P-M; H3T3R0
167. *C. fontanum* Baumg. ssp. *triviale* (Link.)Julas (*C. caespitosum* Gilib.): I-V; H-Ch,
 Cosm; H3T0R0
168. *C. glomeratum* Thuill.: IV V (5,37,40); Th, Cosm; H2.5T3R0
169. *C. pumilum* Curt.: IV V (5,25,37,40); Th, Eua (M);H2T3R0
170. *C. semidecandrum* L.: III-V (37,40,!); Th, E; H2T3.5R0
171. *Holostecum umbellatum* L.: III-V (15,14,29,5,25,18); Th, Eua (M);H2T3.5R0
172. *Sagina procumbens* L.: V (37,40,25,27,!); H(Ch), Cp; H4T3R3
173. *Arenaria procera* Spreng. ssp. *glabra* (F.N. Williams)Holub (*A. graminifolia* Schrad.):
 IV V (14,37,40); H,Eua (C); H2T3.5R4
174. *A. serpyllifolia* L.: III-V (5,25,37,40,18,!); Th,Cp; H2T2.5R0
175. *Moehringia muscosa* L.: I-II (22,14), IV (14,5,25); H, E; H4T2R4
176. *M. trinervia* (L.)Clairv : III-V (15,29,5,25,40,!); Th-TH, Eua (M); H2.5T3R3
177. *Spergula arvensis* L. III-V (15,5,40,!); Th, Cosm; H3T0R0
178. *Spergularia marina* (L.)Grisb. (*S. salina* J. et C. Presl.): III-V(14,34,40,!); Th-TH, Eua;
 H2T3R2
179. *S. media* (L.)C. Presl. (*S. marginata* Kitt.): V (40); Th-H, Eua(M);H4T0R0
180. *S. rubra* (L.) J. et C. Presl.: V (37,40,!); Th (H). Cp; H4T3R4
181. *Herniaria glabra* L.: III-IV (14); Th-H, Eua(M); H2.5T3.5R3
182. *H. incana* Lam.: III-IV (14,5); H (Ch), Eua(M); H2T3.5R4.5
183. *Scleranthus annuus* L. ssp. *annuus*: II-V (22,5,14,25,!); Th, Eua; H2T3R2
 - ssp. *polycarpus* (Torn.)Thell. (*S. polycarpus* Torn.): V (40); Th, E(M); H2T3R2
184. *S. perennis* L.: IV V (37,14,25,!); H-Ch. Eua; H3T0R3
 - var. *dichotomus* (Schur)Prod.: IV (5)
185. *S. uncinatus* Schur: II (14); Th TH, Carp-B-Anat-Cauc; H3T2R0
186. *Agrostemma githago* L. : III-V (29,5,25,!); Th, Eua(M); H2T4R0
187. *Lychnis coronaria* (L.)Desr.: IV (14,25); H.M; H2.5T4R3
188. *L. flos-cuculi* L.: I-V; H, Eua; H3.5T2.5R0
189. *L. viscaria* L. (*Viscaria vulgaris* Rohl.): I-IV (6,22,34,29,25,!); H, Eua; H3T4R0
190. *Silene alba* (Mill.)E.H.L. Krause (*Melandrium album* (Mill.)Gaeke): III-V (5,18,!);
 Th-TH,Eua; H3.5T2R3
191. *S. armeria* L.: I-II (14,22,!); IV (14); Th, M; H2.5T4R3
192. *S. bupleuroides* L. (*S. longiflora* Ehrh.): III-V (14,29,40); H, P-M; H1.5T4.5R4
193. *S. conica* L.: III-IV (14,29,25); Th, Eua(M); H1T3.5R4
194. *S. dioica* (L.)Clairv (*Melandrium rubrum* (Weig.)Gaecke): I-II (14); H, Eua;
 H3.5T0R4
195. *S. dubia* Herb.: II (48,22,!); H, Carp (End); H2T3R0
196. *S. flavescens* W et K.: IV (14,25); H, D-B-Pn; H2T3R4
197. *S. x grecescui* Gusul. (*nutans* x *viridiflora*): III (14):
198. *S. heuffelli* Soó (*Melandrium nemorale* (Heuff)A Br.): IV-V (35,14,25,24,!); Th-TH,
 Carp-B; H3.5T2R0

199. *S. italica* (L.)Pers.: I (!), IV (5,25); H, Eua (M); H3T0R3
 - var. *nemoralis* (W et K.)Heuff.: I (14)
200. *S. multiflora* (Ehrh.)Pers.: IV (14); H, Eua (C); H3.5T3R4
201. *S. noctiflora* L. (*Melandrium noctiflorum* (L.)Fries): IV-V (5,40); Th-TH, Eua;
 H2T3.5R0
202. *S. nutans* L.: I-IV (14.22,35,!); H, Eua; H2T3R4
203. *S. otites* (L.)Wib. ssp. *otites*: III-V (14,29,5,18,!); H, Eua (C); H1.5T4R4.5
 - var. *pseudotites* (Bess.)Borb.: V (40)
 - var. *effusa* (Otth.)Gusul.: III (14)
204. *S. viridiflora* L.: III-IV (14,5,25,!); H, M; H2T3.5R3
205. *S. viscosa* (L.)Pers. (*Melandrium viscosum* (L.)Celak.): III (14,29), V (40); Th-TH,
 Eua (C); H2.5T3R4
206. *S. vulgaris* (Mnch.)Gaerke (*Bchen vulgaris* Mnch.) ssp. *vulgaris*: III-V
 (9,29,5,25,40,!); H (Ch), Eua; H3T3R4
207. *Cucubalus baccifer* L.: III-V (29,24,25,40,!); H, Eua; H3.5T3R4
208. *Gypsophila muralis* L.: III-V (14,5,25,!); Th, Eua(C); H2T3R2
 - f. *ramossissima* (Schur)Prod.: III (14)
 - f. *capillaris* Fiek et Schuber: IV (25)
209. *G. paniculata* L.: IV (14); G (Ch), Eua (C); H2T4R4.5
210. *Petrorhagia prolifera* (L.)P.WBall. et Heywood (*Tunica prolifera* (L.)Scop.): III-V
 (14,29,25,37,40); Th.P-M; H1.5T4R3
211. *Vaccaria pyramidata* Medik.: V (18,40); Th, Eua (M); H3T3R0
 -ssp. *graminiflora* (Fisch)Prod.: III (14), V (14,37,40)
212. *Dianthus armeria* L. ssp. *armeria*: IV V (25,24,40); Th-TH, E:H2T3R3
 - f. *glaber* Scholz: IV (43)
 - ssp. *armeriastrum* (Wolfn.)Volen. (*D. armeriastrum* Wolfn.): IV (14,25); Th-TH,
 E:H2.5T4R0
213. *D. barbatus* L. ssp. *compactus* (Kit.)Tutin (*D. compactus* Kit.): II (14,!); IV (5); H,
 Alp-Carp-B; H2.5T3.5R4
 - f. *umbrosus* Nyár.: II (22,14,!)
214. *D. carthusianorum* L.: I-IV (6,22,29,5,!); H,E; H2T4.5R5
 - ssp. *latisolius* (Gris. et Schenk)Hegi var. *parviflorus* Celak.: I (14)
215. *D. giganteus* D'Urv : IV (14); H,B; H2.5T3R4
 - f. *luxurians* Prod.: IV (14)
216. *D. kitaibelli* Janka ssp. *spiculisoli* (Schur)Sanda (*D. spiculisoli* Schur): IV (5,14);
 H, Carp (End);H2T3.5R4
217. *D. pontederac* Kern.: V (37,14,25,40); H, B-Pn; H2T5R5
 - ssp. *giganteiformis* (Borb.)Soó (*D. giganteiformis* Borb.): I (14), V (14, 43,25); H, Pn;
 H3T4R0
218. *D. puberulus* (Simk.)Kern.: III-IV (14,5,25); H, B; H2T4R4
 - var. *laevigatus* (Simk.)Hand.-Mazz.: IV (14)
219. *D. superbus* L.: I (23,!); H, Eua; H3T0R0
220. *D. trifasciculatus* Kit.: III-IV (37,14,25); H, B; H3T3R3
221. *Saponaria officinalis* L.: II-V (29,5,37,14,25,!); H, Eua(M);H3T3R0

EUPHORBIACEAE

222. *Euphorbia amygdaloides* L.: IV (!); Ch, E(M); H3T3.5R4
223. *E. angulata* Jacq.: III-IV (14,35,5); H-G, P-M; H2.5T3R4.5
224. *E. x angustata* (Roch.) Borza (*salicifolia* x *virgata*): III, V (14);
225. *E. carniolica* Jacq.: IV (14.25); H, Ec; H3T4R4
226. *E. x csatoi* (Simk.) Borza (*agraria* x *paradoxa*): III (14);
227. *E. cyparissias* L.: III-V (29,5,33,18,!); H (G), Eua; H2T3R4
228. *E. dulcis* L.: III (29); H-G, Ec; H4T3R3
229. *E. epithymoides* L. (*E. polychroma* Kern.): III-IV (14,29,5,25); H, Pn-B; H2.5T4R3
230. *E. esula* L.: IV-V (37,14,25,40); H, Eua; H2T3R3
- var. *riparia* Schur: IV (14)
231. *E. exigua* L.: III (!); Th, E (M); H2.5T3.5R4.5
- f. *tricuspidata* (Lapeyr.) Borza: III (14.29)
232. *E. falcata* L.: III-V (29,14.5,40,!); Th, Eua (M); H2T3.5R4.5
233. *E. helioscopia* L.: III-V (35,29,5,25,18,!); Th, M; H3T3R0
234. *E. lucida* W et K.: IV V (14.18,47); H, E (C); H5T3R4
235. *E. maculata* L.: IV (14); Th, Adv; H2T3.5R4.5
236. *E. nicaeensis* All. (*E. pannonica* Host.): V (14,40); H, P-Pn-B; H1.5T5R5
237. *E. palustris* L.: IV V (37,14,5,40); H-Hh, E; H4.5T3.5R4.5
238. *E. x paradoxa* (Schur) Podp. (*esula* x *salicifolia*): III-V (14,25,40);
239. *E. platyphyllus* L.: IV V (25,40,!); Th, Ec(M); H3T3R3
240. *E. x pseudolucida* Schur (*lucida* x *virgata*): V (40);
241. *E. salicifolia* Host.: III-V (29,5,25,40,!); H, P-Pn; H2T3.5R3
242. *E. segetalis* L.: V (14,40); Th-H, M; H3T3.5R0
243. *E. seguieriana* Neck.: IV-V (14,5,25,40,!); H, Eua(M); H1T3.5R4
244. *E. stricata* L.: IV-V (25,40); Th, E(C); H4T3R5
245. *E. villosa* W et K.: III-V (14,5,37,25); H, P-M; H3T3.5R0
246. *E. virgata* W et K.: III-V (35,37,14,18,47,!); H, Eua(C); H2T4R3
- f. *angustissima* Schur: V (46)
247. *Mercurialis perennis* L.: II-V (22,29,25,!); H-G, E; H3.5T3R4

CALLITRICHACEAE

248. *Callitricha cophocarpa* Sendtn. (*C. polymorpha* Lomnr.): III-V (14,38,!); Hh, Eua;
H6T3R0
249. *C. palustris* L. em. Druce (*C. verna* L.): I (14,31,38); Hh, Cp; H6T3R0

RANUNCULACEAE

250. *Trollius europaeus* L.: I (30,6,23,14,!); III (14.29); H, E; H4T2R4
251. *Caltha palustris* L. ssp. *lactea* (Schott, Nyman et Kotschy) Hegi: I-V; H,E; H5T3R0
- var. *pseudocornuta* Zap.: I-II (14)
- var. *alpina* (Schur) Graebn.: I-IV (14,!)
252. *Helleborus purpurascens* W et K.: III-IV (14,29,25,!); H, Carp-B-Pn; H2.5T3R4
- f. *baumgartenii* (Kov.) Nyár.: III-IV (14.25)

253. *Nigella arvensis* L.: III-V (14,29,25,18); Th, E(M); H2T4R4
254. *Isopyrum thalictroides* L.: III-V (15,29,14,24,25); G, Ec; H3T3.5R3
255. *Actaea spicata* L.: II-III (14,29); H, Eua; H3.5T3R3
256. *Consolida orientalis* (J.Gray) Schroding. (*Delphinium orientale* Gray): V (14,18,47,!); Th, Eua(C); H2.5T4R4.5
 - f. *rhodochroa* (Soc)A.Ny r.: V (40)
257. *C. regalis* S.F.Gray (*Delphinium consolida* L.): III-V (15,29,5,25,18,!); Th, Eua; H2T4R4
258. *Aconitum anthora* L.: II-IV (37,14); H,E(C); H2T3R5
259. *A. firmum* Rchb. ssp. *multifidum* (Rchb.)Grint. (*A. multifidum* Rchb.): II (14); H, Carp-B-Sudet; H2.5T2.5R4.5
260. *A. moldavicum* Haenq.: I-II (22,14,!); H, Carp (End); H3T2R3
 - var. *haequetianum* G. Grint.: I-II (14)
 - f. *piliferum* G. Grint.: II (14)
 - var. *confusum* G. Grint.: II (14)
261. *A. toxicum* Rchb.: I (23,30,!); H,Carp-B; H4T2.5R4.5
262. *A. variegatum* L. ssp. *gracile* (Rchb.)Gay (*A. gracile* Rchb.): III (29,14); H, Alp-Carp; H4T2.5R4.5
263. *Anemone nemorosa* L.: I-V; G, E; H3.5T4R0
264. *A. ranunculoides* L.: II-V (22,29,40,!); G, E; H3.5T3R4
265. *A. silvestris* L.: III-IV (14,29); H, Eua (C); H, Eua(C); H2T3.5R4
266. *Hepatica nobilis* Mill.: II-IV (15,22,29,14,!); G, E; H3T3R4
267. *H. transsilvanica* Fuss: I-II (14); G, Carp (End); H3T2R4
268. *Clematis alpina* (L.)Mill. (*Atragene alpina* L.): I-II (14,30,!); H(nph),Eua; H3T2R2
269. *C. integrifolia* L.: IV V (14,43,40,!); H, Eua (C); H3T3.5R5
270. *C. recta* L.: III-V (37,14,40,5,!); H, P-M; H2.5T3R4
271. *C. vitalba* L.: II-V (27,29,5,40,!); nPh, Ec (M); H3T3R3
272. *Myosurus minimus* L.: III-V (14,!); Th, Cp; H4T4R3
273. *Ceratocephalus testiculatus* (Cr.)Roth (*C. orthoceras* DC): III-V(14,5,40); Th, P-PN; H2T4R4.5
274. *Ranunculus acris* L.: I-V; H, Eua (M); H3.5R0R0
 - f. *stipitatus* Ny r.: III (14)
 - ssp. *strigulosus* (Schur)Hyl. (*R. stevenii* Andrz.): I (14,!);III-IV (5,14, 25); H-G, P-M; H3.5T2R3
 - var. *friseanum* (Jord.)A.Ny r.: I (14)
275. *R. x alliariifolius* (Rchb.)A.Ny r. (*cassubicus* x *flabellifolius*): III (14)
276. *R. arvensis* L.: III-V (14,29,5,18,!); Th, Eua (M); H3T3R0
 - var. *tuberculatus* (DC)Mert. et Koch: IV (5)
277. *R. auricomus* L.: I-V; H, Eua; H3.5T3R3
278. *R. bulbosus* L.: III-V (42,40,!); H-G, E; H2T3R3
279. *R. carpaticus* Herb.: II (14); G, Carp (End); H3.5T2R4
280. *R. cassubicus* L. : IV-V (24,25); H, E(C); H3.5T3R0
281. *R. x fallax* (Wimm. et Grab.)Kern. (*auricomus* x *cassubicus*):III (14),V (!);
 - var. *incisifolius* (Rchb.)A.Nyár.: III, V (14)

282. *R. sicaria* L. (*Ficaria verna* Huds.) ssp. *sicaria*: I-V; H-G, Eua (M); H3.5T3R3
 - ssp. *calthifolia* (Rchb.) Vcl.: V (40); H-G, E: H3.5T3R3
283. *R. flammula* L.: I (6.23,!). II (!): H, Eua: H4.5T3R0
 - f. *serratus* (DC) Prod.: I (14)
284. *R. lateriflorus* DC.: II (22), V (40); Th, Eua (C); H5T3R5
285. *R. lingua* L.: I (23), III (14), V (40); Hh, Eua: H6T3R4
286. *R. pedatus* W et K.: V (18,40); H-G, Eua (C); H1.5T3R4
287. *R. peltatus* Schrank (*Batrachium triphyllum* Waller.) Dum.: IV-V (14,40); Hh, E; H6T3R0
288. *R. platanifolius* L.: I (14); H, E; H3.5T2.5R0
289. *R. polyanthemos* L.: I-V; H, Eua (c); H2.5T3R3
290. *R. repens* L.: I-V; H, Eua (M); H4T0R0
291. *R. orcopophilus* M.B.: I (14); H, Alp-Carp-B; H2.5T4R4
292. *R. rionii* Lagg. (*Batrachium rionii* (Lagg.) Nym.): III (14), V (40); Hh, Eua (M); H6T3R
293. *R. sardous* Cr.: IV-V (5.47); Th-H, Eua; H3T3R4
294. *R. sceleratus* L.: I (32), III-V (14,27,5,25,18,!); Th, Cp; H4.5T3R4
295. *R. trichophyllum* Chaix. (*Batrachium divaricatum* (Schrank) Schur.): III-V (14,5,25,34,40,18,!); Hh, E: H6T3R0
296. *Thalictrum aquilegifolium* L.: I-IV (6,23,22,6,29,!); H, E; H2.5T2.5R4
297. *T. lucidum* L.: I-V; H, Ec; H4.5T3R5
 - var. *lucidum* f. *angustissimum* (Cr.) Nyár.: V (40)
 - f. *peucedanifolium* (Gris. et Schenk) A. Nyár.: V (18,40)
 - var. *stenophyllum* (Wimm. et Grab.) Hay: I-II (14)
 - f. *subglabrum* (Simk.) Nyár.: V (37,14)
 - var. *heterophyllum* (Wimm. et Grab.) Hay: I-IV (14,!)
 - f. *scopolii-nigricans* Nyár.: I (23)
298. *T. flavum* L.: III-V (14,29,40); H, Eua: H4.5T0R4.5
299. *Adonis aestivalis* L.: III-V (15,29,5,25,18,47); Th, Eua (M); H3T4R3
300. *A. annua* L. (*A. autumnalis* L.): IV (14); Th, Adv; H2T3.5R3.5
301. *A. flammea* Jacq.: V (40); H, P-M; H2T3.5R3.5

ARISTOLOCHIACEAE

302. *Asarum europaeum* L.: III-IV (14,5,25); H-G, Eua; H3.5T3R4
303. *Aristolochia clematitis* L.: III-V (14,29,25,5,18,!); H-G, Ec (M); H2.5T3.5R5

BERBERIDACEAE

304. *Berberis vulgaris* L.: III-IV (15,29,14,5); mPh, E; H2T3R4

NYMPHAEACEAE

305. *Nuphar luteum* (L.) Sm.: III (14,29,!); Hh, Eua (M); H6T0R3.5
 - var. *sericum* (Lang) Kitt.: III (14)
 - var. *tenellum* (Rchb.) Richt.: III (14)
306. *Nymphaea alba* L.: III (14), V(14,40,!); Hh, E (M); H6T0R4
 - var. *minoriflora* (Borb.) A. et G.: V (40)

CERATOPHYLLACEAE

307. *Ceratophyllum demersum* L.: III-V (14,37,38,25,40,!); Hh, Cosm; H6T3R0
- f. leve Crepin: V (40,14)
308. *C. submersum* L.: III-V (37,14,40); Hh,Eua (M); H6T3,5R0
- var. *haynaldianum* (Borb.)Beck: V (14,40)

PAPAVERACEAE

309. *Glaucium corniculatum* (L.)J.H.Rudolph: III-V (14,29,25,40); Th-TH, M: H2T4R3
- var. *phoenicuum* (Cr.)DC: V (40)
310. *Chelidonium majus* L.: III-V (15,29,5,18,!); H, Eua; H3T3R4
311. *Papaver dubium* L.: IV-V (14,25,40); Th, M; H2T3,5R3
312. *P. rhoeas* L.: IV-V (5,14,18,!); Th, Eua (M); H3T3,5R4
- ssp. *strigosum* (Boenningh.)Simk. var. *verum* A. Ny r.: V (14)
- var. *trifidum* (O.Ktze.)Fedde: V (14)
- var. *agrivagum* (Jord.)Beck: IV-V (14)
- var. *magnogenuinum* Nyár.: IV (14,25)
313. *Corydalis bulbosa* (L.)Pers. (*C. cava* (L.)Schweigg. et Koerte):
- ssp. *cava*: III (14,29), V (40); G, Ec; H3T3R0
314. *C. capnoides* (L.)Pers.: I (14); Th-TH, Eua; H3T3R4
315. *C. solida* (L.)Sw : II-V (14,22,5,25,40); G, E; H3T3R0
316. *Fumaria officinalis* L.: III-V (29,14,5,40); Th, Eua (M); H3T0R,3,5
317. *F. rostellata* Knaf.: IV V (5,14,40); Th, Ec-B; H3T0R3,5
318. *F. schleicheri* Soyer-Willemet: I (14), III-V (14,5,18); TH,Eua(M);H2,5T4R4
319. *F. vaillantii* Lioscl.: III-V (5,14,18,4?,40); Th,Eua; H2,5T3,5R4,5

BRASSICACEAE (CRUCIFERAE)

320. *Sisymbrium altissimum* L.: III-IV (14); Th-TH, Eua(C); H2T3,5R0
321. *S. loeselii* Jusl.: III-V (14,5,25,37,40): Th-TH, Eua (C); H2,5T4R3
- f. *ciliatum* (Beck)Nyár.: III (14)
322. *S. officinale* (L.)Scop.: III-V (14,29,5,40,!); Th, Eua (M);H2,5T4R3
323. *S. orientale* Torn.: V (47,40); Th-TH, Eua (M); H2,5T4R3
- f. *hygrophilum* (Fourn.)Thell.: II (14)
324. *S. strictissimum* L.: III (14,29); H, Ec; H3,5T4R4,6
325. *Descurainia sophia* Webb. (*Sisymbrium sophia* L.): II-V (14,22,29,5,18,!); Th,Eua;
H2,5T4R4
326. *Alliaria petiolata* (M.B.)Cavara et Grande (*A. officinalis* Andr.):III-
V(14,15,29,5,25,24,40,!); Th-TH,Eua (M); H3T3R4
327. *Arabidopsis thaliana* (L.)Heynh.: III-V (29,5,37,14,25,40); Th-TH, Eua(M); H2T3R3
328. *Myagrum perfoliatum* L.: III-V (14,5,40); Th, M; H2T4R3
329. *Isatis tinctoria* L.: III-V (29,25,14,40); H, P-Pn; H1,5T3,5R4
330. *Bunias erucago* L.: IV (14); TH, Adv; H3T3,5R3
331. *B. orientalis* L.: II-V (22,14,5,40); TH-H, Eua (C); H3T3,5R3
332. *Erysimum cheiranthoides* L.: II-V (14,29,5,18); Th, Cp; H3T0R4

333. *E. odoratum* Ehrh. (*E. pannonicum* Cr.); IV V (5,14,40); H-Th,P; H2.5T3R4
 - var. *speciosum* Nyár.: I (14)
334. *E. repandum* Hojer: III-V (15.29,14.18,!); Th. Eua (C); H2.5T4R4.5
335. *Hesperis silvestris* Cr.: III-V (14,29); H, M-P; H4T2R3
 - var. *runcinata* (W et K.)Borb.: III (14), V (40)
336. *Euclidium syriacum* (L.)R.Br.: V (40); Th, Eua (C); H2T4R4
337. *Barbaraca vulgaris* R. Br : I (14), III-V (14,40,5,29,34); TH-H,Eua (M); H3.5T3R3
338. *Rorippa amphibia* (L.)Bess. : III-V (14,29,25,18);Hh,Eua (M); H6T3R4
339. *R. x armoracioides* (Tsch.)Fuss (austriaca x silvestris): III-V (14,40);
 - var *integrifolium* (Tsch.)Nyár.: III (14), V (40)
 - var. *semisilvestris* Borb.: III(14)
 - var. *pinnatifida* (Tsch.)Borb.: III-V (14,40)
340. *R. austriaca* (Cr.)Bess.: II-V (22,14,%18); H-G, Ec; H4T3.5R4
 - var. *angustifolia* (Schur)Nyár.: V (14)
341. *R. x barbaraеoides* (Tsch.)Cel. (islandica x silvestris): III (14)
 - var. *capillipes* Borb.: III-V (14)
 - var. *reichenbachii* Knaf III (14), V (40)
 - f. *pubescens* (Borb.)Ny á.: III (14)
 - f. *arenaria* (Knaf)Nyár.: III (14)
 - f. *macrostylis* (Tsch.)Nyár.: III (14)
342. *R. x hungarica* Borb. (amphibia x austriaca): V (40)
 - var. *riparia* (Simk.)Nyár.: V (37,14,40)
343. *R. islandica* (Oed.)Borb.: I (14,!); III-V (14,29,27,25,40,!); Th-TH. Cosm; H5T3R4
344. *R. x permixta* Borb. (barbaraеoides x silvestris): III (14)
345. *R. prolifera* (Heuff.)Neilr.: IV (14); TH, B; H4T4R4
346. *R. pyrenaica* (L.) Rchb.: II-IV (14,22,5,!); H, M; H2.5T3R3
347. *R. silvestris* (L.)Bess.: I-V; H-G, E; H4T3R4
 - f. *dentata* (Koch)Borb.: I,III (14)
 - f. *tenuifolia* (Tsch.)Beck: III (14)
 - f. *densiflora* Borb.: III (14)
348. *R. x stenophylla* Borb. (silvestris x pyrenaica): III (14)
349. *Armoracia macrocarpa* (W.et K.)Baumg.: V (18); H, Pn; H3T3.5R0
350. *A. rusticana* (Lam.)G.M.Sch.: III-V (14,40,!); G (H)Adv; H3T3.5R0
351. *Nasturtium officinale* R. Br.: III-IV (29,5); Hh, Cosm; H5T2.5R4
352. *Cardamine amara* L.: I (30), IV (14); H, Eua (M); H5T0R0
353. *C. glanduligera* Schw. (Dentaria glandulosa W et K.): I-II (14,22,23); G, Carp (End);
 H4T2.5R4
354. *C. impatiens* L. : II-V (22,14,25,27); Th-TH. Eua (M): H4T3R3
355. *C. pratensis* L. ssp. *pratensis*: I-V; H, Cp; H5T3R0
 - var. *grandiflora* Neilr.: IV (14)
 - var. *palustris* W et Gr.: V (14,40)
 - ssp. *matthiolii* (Moretti)Soc (C. matthiolii Moretti, C. hayneana Welw): II (14), IV (25), V (40)
356. *Cardaminopsis arenosa* (L.)Hay : II (22,140, IV V (37,14,40);TH-H.Ec; H2.6T3R4

- var. perturbata Nyár.: II (14)
 - var. segetalis (Schur)Nyár.: III (14)
357. *Arabis glabra* (L.) Bernh. (*Turritis glabra* L.): II-IV (22,14,29,5,25,!); TH, Cp; H2T3R3
358. *A. hirsuta* (L.) Scop. ssp. *hirsuta*: III-V (14); TH-H, Cp; H1.5T3R4
- f. *sagittata* (Betol.) Tuzs.: V (14,37,40)
359. *Alyssum alyssoides* (L.) L.: II (22), IV (5), V (40); Th-TH, E (C); H1T3R0
360. *A. desertorum* Stapf.: V (40,14); Th, Eua (C); H1.5T4R4
361. *Berteroia incana* (L.) DC.: III-IV (29,6,25,!); Th-TH, Eua (C); H2T3.5R0
362. *Draba muralis* L.: II (22,14,!); III (29); Th, E (M); H2.5T3.5R4.5
363. *D. nemorosa* L.: I (14), III-V (14,5,40); Th,Cp; H3T0R4.5
364. *Camelina alyssum* (Mill.) Thell.: I-II (14); Th, E; H3T3R3
- f. *integrisolia* (Fr.) K. Maly; I-II (14)
365. *C. sativa* (L.) Cr. ssp. *sativa*: III-V (14,29,40); Th, Eua; H3T3R3
- ssp. *microcarpa* (Andrz.) E. Schmid (*C. microcarpa* Andrz.): IV-V (14,18); Th, Eua; H3T3R0
366. *Neslia paniculata* (L.) Desv.: III-V (14,40); Th, Eua; H2.5T3R4.5
367. *Capsella bursa-pastoris* (L.) Medik.: I-V; Th, Adv (M); H3T0R0
368. *Thlaspi aliaicum* L.: IV (14,25); Th, Atl-M; H2T4R0
369. *T. arvensc* L.: III-V (29,5,25,18,!); Th, Eua (M); H2T3R4
370. *T. perfoliatum* L.: III-V (15,29,5,40,18); Th, Eua; H2.5T3.5R4.5
371. *Cardaria draba* (L.) Desv (*Lepidium draba* L.): III-V (5,25,18,!); H, Eua; H2T4R4
372. *Lepidium campestre* (L.) R.Br.: III-V (14,25,40); Th, E (M); H2.5T3R0
373. *L. cartilagineum* (L. May) Thell.: V (40); G,Pn; H3.5T4R4
374. *L. persoliatum* L.: III (14,18); V (14,18); Th, Eua(C); H2T4R3
375. *L. ruderale* L.: IV-V (5,18); Th, Eua; H2T4R3
376. *Coronopus squamatus* (Forsk.) Aschers. (*C. procumbene* Gilib.): V (47,40); Th, M; H3.5T4R4
377. *Conringia orientalis* (L.) Andrez.: III-V (14,5,40); Th, Eua (M); H2T3.5R5
378. *Diplotaxis muralis* (L.) DC.: III-V (5,14,25,47,40); TH(H), M; H2.5T3.5R4
- f. *caulescens* Kit.: V (40)
379. *D. viminea* (L.) DC.: IV (5,14); Th, M; H2.5T4R0
380. *Brassica nigra* (L.) Koch: III-V (14,29,5,40); Th,Eua (M);H3T4R0
381. *Sinapis alba* L.: III (14), IV (25); Th, Adv; H3T4R4
382. *S. arvensis* L.: III (14,29,!); V (14,18); Th, Cosm; H3T3R3
- var. *arvensis* f. *orientalis* (Jusl.) Godr.: V (37,14,18)
 - var. *schkuhriana* (Rehb.) Hagenb.: III (14), V (25,40)
383. *Erucastrum gallicum* (Willd.) O.E.Schulz: III-IV (14,5); Th-TH,Atl-M; H2.5T3.5R4.5
384. *Rapistrum perenne* (L.) All.: I (14), III-V (14,4,18,40); TH-H (G),E (M); H2T3.6R4
385. *Calepina irregularis* (Asso) Thell.: III-V (14,5)
386. *Raphanus raphanistrum* L.: III-V (15,29,5,40,!); Th, M; H2.5T3R0

RESEDACEAE

387. *Reseda lutea* L.: III-V (29,5,14,43,18); Th-H,Eua (M); H2T3.5R4.5

388. *R. luteola* L.: IV (14,5,43,26,40); TH, Eua (M); H2T3R0

CISTACEAE

389. *Helianthemum nummularium* (L.)Mill. ssp. *nummularium*: I (6), III (29), IV (5,14); Ch-H, Ec-M; H2T3R4
- ssp. *obscurum* (Pers.)Prod. (*H. hirsutum* (Thuill.)Merat): I-II(14,22), IV (43); Ch-H. Ec-M; H2,5T3R4
- f. *lanceolatum* Willk.: IV (43)

TAMARICACEAE

390. *Myricaria germanica* (L.)Desv : I, III (14); nPh, Eua; H0T0R4.5

ELATINACEAE

391. *Elatine alsinastrum* L.: V (14); Hh, Eua (M); H5T4R3

DROSERACEAE

392. *Drosera rotundifolia* L.: I (14,23,6,!); H, Cp; H5T2.5R1

VIOLACEAE

393. *Viola alba* Bess.: IV (5,14); H, Ec-M; H3T4.5R4
394. *V. arvensis* Murr.: III-V (14,5,25,18,47); Th, Eua; H3T3R0
395. *V. canina* L.: I-IV (15,29,14,22,6,5,!); H, Eua; H2,5T3R2
396. *V. clatior* Fries: V (40,18); H, Eua; H4T4R4.5
397. *V. epipsila* Ldb.: I (23,32,14,!); H, Eua (C); H5T0R1.5
398. *V. hirta* L.: III-V (29,5,25,40,!); H, Eua; H2T3R4
399. *V. luteola* (Schur)Gay : II (22); Th-H, Eua; H2,5T3R0
400. *V. montana* L.: I (23); H, Eua; H2T3R2
401. *V. odorata* L.: III-V (5,29,25,40,!); H, AtI-M; H2,5T3.5R4
- f. *simonkaiana* Gay : III (14)
402. *V. persicifolia* Schreb. (*V. stagnina* Kit.): I (32), IV-V (14,40); H, Eua; H4,5T3R3.5
403. *V. pumila* Chaix: I (14), IV-V (5,14,40); H, Eua; H3T3R4
404. *V. reichenbachiana* Jord. (*V. silvestris* Lam.): IV V (25,24,40);H, Eua; H3T3R3.5
405. *V. tricolor* L. ssp. *tricolor*: I-V; Th-H, Eua; H2,5T3R0
- ssp. *subalpina* Gaud. (*V. bielziana* Schur): I (14)
- var. *perrobusta* Borb.: I (14)

HYPERICACEAE

406. *Hypericum hirsutum* L.: IV-V (25,24); H, Eua; H3T3R3
407. *H. maculatum* Cr.: I (6,32,!); IV (25); H, Eua; H4T3R2
408. *H. montanum* L.: I-II (30,14); H, E; H3T3R4
409. *H. perforatum* L.: I-V; H, Eua; H3T3R0
- var. *latifolium* Koch: IV (14)
- var. *angustifolium* DC.: I,III (14)
410. *H. tetrapterum* Fr. (*H. acutum* Mnch.): IV-V (14,40); H, E; H4T3R4

CRASSULACEAE

- 411. *Sedum annuum* L.: I (22.); Th-TH, Eua; H3T2R0
- 412. *S. caespitosum* (Cav.) DC.: V (14,40); Th, M; H2T4R4
- 413. *S. cepaea* L.: IV (25); Th-TH, M; H3T4R0
- 414. *S. hispanicum* L.: II (22.!), IV (25); Th(Ch), M; H1T3.5R4
- 415. *S. maximum* (L.) Hoffm.: II-IV (5,14,29,22,25,!); H (G), Eua (M); H2T3R0
- 416. *S. reflexum* L.: I (14); Ch, E; H1.5T3R2.5
- 417. *Sempervivum heuffelii* Schott: II (!); Ch, Carp-B; H1.5T2R0
- 418. *S. marmoreum* Gris. (*S. schlehani* Schott) ssp. *blandum* (Schott) Hay: II (14,!); Ch, Carp-B; H1.5T2.5R2.5

SAXIFRAGACEAE

- 419. *Saxifraga tridactylites* L.: II (14); Th, E (M); H2T3.5R4
- 420. *Chrysosplenium alternifolium* L.: I-V; H.Cp; H4T2R4

PARNASSIACEAE

- 421. *Parnassia palustris* L.: I (23.31,!); H, Cp; H4.5T2R1.5

GROSSULARIACEAE

- 422. *Ribes grossularia* L.: I-II (30.22,14.); mPh, Eua; H0T3R0
- 423. *R. heteromorphum* Topa: I (23,14); mPh, Eua; H4T2.5R3.5
- 424. *R. nigrum* L.: I (23.30,!); H0T0R3
- 425. *R. rubrum* L.: I (23); mPh, Eua; H4.5T0R3.5

THYMELAEACEAE

- 426. *Thymelaea passerina* (L.) Coss. et Germ.: IV V (14,25,40); Th,Eua(C);H1T4R3
- 427. *Daphne mezereum* L.: I-III (30,22,29,!); nPh, Eua; H3.5T3R3

ROSACEAE

- 428. *Spiraea salicifolia* L.: T (23,32,14,!); mPh, Eua (C); H4.5T2.5R2
- 429. *S. ulmaria* Scop.: I-II (30,22,!); mPh, Eua; H3.5T2.5R0
- 430. *Aruncus dioicus* (Walter) Fernald. (*A. vulgaris* Raf.): II (22.); H, Eua; H4T2.5R3
- 431. *Pyrus pyraster* (L.) Burgsd.: I-V; MPh-mPh, E; H2T3R4
- 432. *Malus silvestris* (L.) Mill.: I (23), III-V (29,6,25,24.); mPh, E; H3.5T3R4
- 433. *Sorbus aucuparia* L.: I-II (22,23,!); IV (5,26,!); MPh-mPh, E; H3T2.5R2
 - var. *lanuginosa* (Kit.) Beck: I (14,23)
- 434. *S. torminalis* (L.) Cr.: IV (5,14,!); MPh, E(M); H2.5T3R4
- 435. *Crataegus x intermixta* (Wenzig) Beck (*laevigata* x *monogyna*): V (40)
- 436. *C. laevigata* (Poir.) DC. (*C. oxyacantha* Auct. non L.): IV (14,25), V (24,27); mPh, Ec; H3T3R3
 - var. *kyrtostyla* (Fingerh.) Beck: V (40,14)
- 437. *C. monogyna* Jacq.: II-V (22,6,25,24,!); mPh, E; H2.6T3R3
 - var. *kyrtostyla* (Fingerh.) Beck: V (40,14)
- 438. *Rubus candidans* Whc.: II (14), IV (14,25); nPh, M; H2T2.5R0

- var. menyhazensis (Simk.)Nyár.: II (14)
- 439. *R. caesius* L.: I-V; H (nPh), Eua (M); H4.5T3R4
- var. *arvalis* Rchb.: IV (14, !)
- 440. *R. hirtus* W et K.: IV (25); nPh, E; H3T2.5R3
- 441. *R. idaeus* L.: I-IV (22.29.25, !); nPh, Cp; H3T3R3
- 442. *R. x longiramulus* Sabr. (rivularis x serpens): II (14)
- 443. *R. plicatus* Whc. et Nees: III(29); nPh, Atl-Ec; H3.5T3.5R2
- 444. *R. rivularis* P.J.Mull. et Wirtg.: II(14); nPh, Atl-Ec:H2.5T2.5R2
- var. *incultus* (Wirtg.)Focke: II (14)
- 445. *R. saxatilis* L.: I (30); nPh, Eua; H3T1.5R0
- 446. *R. x scythicus* (Sabr.)Gay (rivularis x tereticaulis): II (14)
- 447. *R. suberectus* G. Anders.: II (14); H (nPh), Ec; H4T2.5R3
- 448. *Fragaria vesca* L.: I-V; H, Eua; H3T2.5R0
- 449. *F. viridis* Duch.: I (14), IV (5,14.25); H, E (C); H2T4R3
- 450. *Potentilla anserina* L.: I-V; H, Cosm; H4T3R4
- 451. *P. argentea* L.: I-V; H, Eua; H2T4R2
- var. *dissecta* Walr.: I (14)
- 452. *P. chrysanthia* Trev.: IV V (14,24,27); H, Eua (C); H2T0R3
- 453. *P. erecta* (L.)Rauschel: I-IV (6,31,29,25, !); H, Eua(M); H0T0R0
- 454. *P. inclinata* Vill. (*P. canescens* Bess.): I (14); H, Eua (C); H3T0R4
- var. *laciniosa* Beck: I (14)
- 455. *P. palustris* (L.)Scop. (*Comarum palustre* L.): I (23,31,14, !); Hh(Ch),Cp; H5T2R3
- 456. *P. patula* W et K.: II (22); H, P-Pn; H2.5T3.5R3
- 457. *P. recta* L.: II-IV (22.29.5, !); H, Eua (C); H1.5T3.5R4
- 458. *P. reptans* L.: I-V; H, Cosm; H3.5T0R4
- 459. *P. x semiargentea* Borb. (*argentea* x *inclinata*): III (14)
- 460. *P. supina* L.: III-V (5,25,40, !); Th-H, M; H4T3R0
- 461. *P. thuringiaca* Bernh.: II-IV (14,22,25, !); H, Ec; H2T3R3
- var. *nestleriana* (Tratt.)Schinz et Kell.: III-IV (14,25, !)
- var. *elongata* (Goldb.)Th. Wolf. II (14)
- 462. *Geum aleppicum* Jacq.: II (22,14, !); H, Cp; H3.5T2.5R4.5
- 463. *G. rivale* L.: I-II (6,23,14, !); H, Cp; H4.5T0R4.5
- 464. *G. urbanum* L.: II-V (22.29.5,24.25, !); H, Eua (M); H3T3R4
- 465. *Waldsteinia ternata* (Stephan)Fritsch: I (14); Ch, Eua; H4T2.5R4
- 466. *Filipendula ulmaria* (L.)Maxim: I-IV (30,12,22,31,29, !); H, Eua:H4.5T2R0
- ssp. *denudata* (J. et C Presl.)Hayek: I (14)
- 467. *F. vulgaris* Mnch. (*E. hexapetala* Gilib.): II-V (22,34,5,25,40, !); H, Eua; H2.5T3R0
- 468. *Alchemilla glabrescens* Wallr. (*A. hybrida* Acut. non Mill.): I (14); H.Ec; H2.5T2.5R2.5
- ssp. *plicata* (Buser)Palitz: I (14); H, E; H3T2.5R3
- 469. *A. vulgaris* L.: I-III (31,29, !); H, Ec; H3.5T2R2
- ssp. *acutangula* (Buser)Paalitz: II (22,14); H, E; H3T2R0
- ssp. *micans* (Buser)Palitz: I-II (14); H, E; H3T2R0
- 470. *Agrimonia eupatoria* L.: I-V; H, Eua; H2.5T3R4

- f. fallax (Fick.)Buia: I (14)
- 471. *Sanguisorba officinalis* L.: I (23,31,!); III-V (14,29,25,40); H.Eua; H3T3R0
- 472. *Rosa arvensis* Huds.: II-IV (14,29); nPh, Atl-M; H3T3R2.5
- 473. *R. canina* L.: II-V (22,5,14,29,40,!); nPh, E; H2T3R3
 - var. *lutetiana* (Lem.)R.Kell. f. *muscrolunata* (Desegl.)Borb.: IV (14)
 - var. *transitoria* R. Kell. f. *marisensis* Simk.: IV (14)
- 474. *R. corymbifera* Borkh. (*R. dumictorum* Thuill.): III-V (14,24,!); nPh,E; H2,5T3R3
- var. *lembachensis* J.B.Kell. f. *ciliata* (Borb.)Borza: IV (14,25)
- var. *thuillieri* Chr. f. *solstitialis* (Bess.)Borb.: III-IV (14)
- 475. *R. dumalis* Bechst.: III-V (14,40,46,!); nPh, E; H2,5T3R4
- var. *villosumcula* (Rip.)H.Br. f. *villosumcula*: IV-V (14)
- 476. *R. gallica* L.: III-V (14,25,18,40); nPh, M; H2T4R4
- 477. *R. pendulina* L.: I-II (23,22,14,!); nPh, Ec; H3T2,5R3
- 478. *R. spinosissima* L.: IV V (14); nPh, Eua; H2T3R4
- 479. *Prunus avium* L. (*Cerasus avium* L.): IV (25,!); Mph-MPh, E; H3T3R3
- 480. *P. padus* L. (*Padus racemosa* (Lam.)C.KSchneid.): I (23,30), III (14,29), V (14); MPh, Eua; H3,5T3,5R4
- 481. *P. spinosa* L.: III-V (29,5,25,24,18,!); mPh, Eua; H2T3R3

FABACEAE (LEGUMINOSAE)

- 482. *Genista tinctoria* L. ssp. *tinctoria*: I-V; Ch-nPh, Eua; H2,5T3R2
 - ssp. *ovata* (W et K.)Arc.: IV (5,14); Ch-nPh, Alp-Carp-B; H2,5T3R3
 - var. *transsilvanica* Lerch. f. *mayeri* (Jka.)Borza: IV (5)
- 483. *Cytisus banaticus* Gris. et Sch. (*C. albus* Hacq. var. *pallidus* Schrad.): V (14); Ch-nPh, B-Pn; H1,5T4R3
- 484. *C. hirsutus* L. ssp. *hirsutus* (*C. leucothricus* Schur): II-IV (22,29,14,!); nPh, Ec (M); H2T3,5R4
- 485. *C. nigricans* L.: II-IV (22,15,29,5,25,!); nPh, Ec; H2,5T3R0
- 486. *Ononis arvensis* L. (*O. bircina* Jacq.): I (6), II-V (14,29,34,5,25,!); Ch-H, Eua (C); H3T4R0
- 487. *O. pseudohircina* Schur: IV V (5,14,40,!); Ch-H, D-B-Cauc; H2,5T3R0
- 488. *Trigonella procumbens* (Bess.)Rchb.: III-IV (14,5), V (40); Th, P-M; H4T4R4,5
- 489. *Medicago falcata* L.: II-V (22,34,29,18,!); H, Eua (M); H2T3R5
- 490. *M. lupulina* L.: I-V; Th-TH, eua; H2,5T3R4
- 491. *M. sativa* L.: III-V (14,5,18,!); H, M; H2T3R5
- 492. *M. x varia* Martyn (*sativa* x *falcata*): IV-V (37,14,25,40)
 - f. *pseudofalcata* (Rouy)Nyár.: IV (25)
- 493. *Melilotus albus* Desr.: III-V (34,5,25,40,!); Th-TH, Eua; H2,5T3R0
- 494. *M. dentatus* (W et K.)Pers.: IV V (14,40); TH (Th), Eua (C); H4T3,5R4
- 495. *M. officinalis* (L.)Pall.: I (32,!); III-V (34,5,25,18,!); TH-TH, Eua; H2,5T3,5R0
- 496. *Trifolium alpestre* L.: I-IV (6,22,14,5,25,!); H, E (M); H2,5T3R4
- 497. *T. angulatum* W et K.: III (34), V (14,27); Th, B-Cauc; H0T4R4,5
- 498. *T. arvense* L.: III-V (14,34,29,25,40,!); Th, Eua (M); H1,5T3R4
- 499. *T. aureum* Pollich (*T. strepens* Cr.): I (14,32), IV (14,25); Th-TH, Eua (M); H3T3R0

500. *T. campestre* Schreb.: I (31), II (!), III-V (29,6,40,18,1); Th-TH, E; H3T3R0
 - f. *pseudoprocumbens* (Gmel.) A. Ny r: III, V (14)
501. *T. dubium* Sibth.: II-V (14,5,40,27,!); Th-TH, E(M); H3,5T2R0
502. *T. fragiferum* L.: III-V (15,29,%18,!); H, Eua; H3T3R5
503. *T. hybridum* L.: I-V; H, E(M); H3,5T3R4
 - var. *fistulosum* (Gilib.) Hegi: III (14)
 - var. *elegans* (Savi) Boiss.: V (14)
504. *T. medium* L.: I-V; H, Eua; H3T3R0
505. *T. montanum* L.: I-III (6,22,29,!); H, Eua (C); H2,5T2R4
506. *T. ochrolellum* Huds.: I (6), III-IV (14,25,!); H, Ec-M; H2,5T2R4
507. *T. ornithopodioides* Sm.: V (14); Th, Atl-M; H4T4R2
508. *T. pallidum* W et K: V (14,27,40); Th-TH, M; H2T4R4
509. *T. pannonicum* Jacq.: I-IV (6,14,22,5,!); H, P-M; H2T3R0
510. *T. pratense* L.: I-V; H-TH, Eua; H3T0R0
511. *T. repens* L.: I-V; H, Eua; H3,5T0R0
512. *T. retusum* Hojer (*T. parviflorum* Ehrh.): V (27,14); Th, P-M; H2T4R4,5
513. *T. spadicium* L.: I (14,32,23); Th, E; H4,5T3R2
514. *T. striatum* L.: IV-V (37,14,25); Th, Atl-M; H1,5T3R4
515. *Anthyllis vulneraria* L.: II-IV (22,29,5,33,25,!); H, E(M); H2T0R4
 - ssp. *kernerii* (Sag.) Domin: II-IV (14)
516. *Lotus angustissimus* L.: V (14,40); Th, P-M; H2T4R4
517. *L. corniculatus* L.: I-V; H, Eua; H2,5T0R0
518. *L. tenuis* Kit.: III (14,34,!); V (14,!); H, Eua(M); H3,5T3R4
519. *Amorpha fruticosa* L.: III-V (4,46,25,!); mPh, Adv; H3T4R0
520. *Galega officinalis* L.: III-V (15,6,18,!); H, P-M; H4,5T3R4
521. *Astragalus cicer* L.: IV V (5,37,40,!); H, E(C); H2,5T4R4
522. *A. glycyphyllos* L.: II-V (22,5,25,24,40,!); H, Eua; H3T3R4
523. *Glycyrrhiza echinata* L.: IV V (14,27,40,!); H, P-M; H4T4R0
524. *Coronilla varia* L.: II-V (22,29,6,26,18,!); H, Ec-M; H2T3R4
525. *Onobrychia viciifolia* Scop.: II-V (22,14,5,40,!); H,M; H2T4R4,5
526. *Vicia cracca* L.: I-V; H, Eua; H3T0R3
527. *V. dumetorum* L.: III (14), V(40); H, H3T3R4,5
528. *V. grandiflora* Scop.: III-IV (14,26); Th-TH, B-P-Cauc; H3T3R0
 - var. *sordida* (W et K.) Gris.: III-IV (14,42,5)
529. *V. hirsuta* (L.) S.F.Gray: IV-V (5,14,25,40); Th, Eua (M); H2,5T3,5R4
530. *V. lathyroides* L.: IV-V (14,18,40); Th, Atl-M; H2T4R2,5
531. *V. pannonica* Cr.: III-V (29,5,40); Th, P-M; H2,5T3,5R4
 - ssp. *striata* (M.B.) Nym. (*V. striata* M.B.): III (14), V(40)
532. *V. pisiformis* L.: V (14,40); H, Ec; H2T3R4,5
533. *V. sativa* L. ssp. *sativa*: IV-V (14,18,40); Th, Adv; H0T3R0
 - ssp. *nigra* (L.) Ehrh. (*V. angustifolia* L.): II (22), III-V (14,6,40,47,!); Th, Eua (M); H0T3R0
534. *V. sepium* L.: I-V; H, Eua; H3T3R3
535. *V. tetrasperma* (L.) Schred.: IV V (5,26,40,!); Th, Eua; H3,5T3R3

536. *V. villosa* Roth: I (6,30,31), III-V (5,14,29,18); Th-TH, M; H2.5T3.5R2.5
 537. *Lathyrus aphaca* L.: III-V (14,29,5,47,40); Th,M;H3T3R3
 538. *L. hirsutus* L.: III-V (15,29,5,37,40); TH,Eua: H3T3.5R4
 539. *L. latifolius* L.: III-V (14,29,5,25); H, M; H2T3.5R4
 - var. *rotundifolius* (Willd.)Rchb.: V (14)
 540. *L. paluster* L.: I (31), III, V (14,40); H, Cp; H5T0R4.5
 541. *L. pannonicus* (Jacq.)Gaege ssp. *aspodocloides* (Gouan)Bassler (*L. versicolor* Auct. non Beck): III-IV (14,6,4,42); H-G, Eua (C); H2T4R4.5
 542. *L. pratensis* L.: I-V; H, Eua; H3.5T3R4
 543. *L. silvestris* L.: II-V (14,40,!); H, E (M); H2.5T3R4
 - var. *oblongus* Scr.: II-III (14)
 - var. *platyphyllus* (Retz)Aschers.: V (40,14,43)
 544. *L. tuberosus* L.: III-V (29,5,25,18,!); H (G), Eua (M); H2T4R4
 545. *Pisum sativum* L. ssp. *arvense* (L.)Cel. (*P. arvense* L.): II-III (14); Th, E; H3T3R4

LYTHRACEAE

546. *Peplis portula* L.: I (14); Th, All-M; H4T3R0
 547. *Lythrum hyssopifolia* L.: III-V (14,40); Th,Cosm; H4T3R0
 548. *L. salicaria* L.: I-V; H-Hh, Cosm; H4T3R0
 - f. *glabrescens* (Neilr)Todor: III (14)
 549. *L. virgatum* L.: III-V (14,34,43,18,!); H-Hh, Eua (C); H4.5T3.5R4

ONAGRACEAE (OENOTHERACEAE)

550. *Epilobium adenocaulon* Hausskn.: II,IV (14); H, Adv; H4T3R3
 551. *E. angustifolium* L. (*Chamaenerion angustifolium* (L.)Scop.): I-IV (30,22,29,5,!); H, Cp; H4T1.5R0
 552. *E. hirsutum* L.: I (31,!); III-V (34,29,25,40,!); H (Hh). Eua (M); H4T3R3
 553. *E. montanum* L.: II (22,!); IV (25,!); V (27,40); H, Eua (m); H3T0R3.5
 554. *E. palustre* L.: I-II (23,12,!); IV (5); H, Cp; H5T0R2
 655. *E. parviflorum* (Schreb.)Wither: III-IV (29,5,25,!); H, Eua; H5T3R4.5
 556. *E. tetragonum* L. (*E. adnatum* Gris.) ssp. *tetragonum*: IV-V (14,25,40); H, Eua; H4.5T3R0
 - ssp. *lamyi* (F.Schultz)Nym. (*E. lamyi* F. Schultz): IV-V (14,25,40); H, E; H0T3.5R0
 557. *E. x wiessenburgense* E. Schultz (adnatum x parviflorum): V (40)
 558. *Oenothera biennis* L.: III-V (14,5,25,18,!); Th, Adv; H2T4R0
 559. *Circaeа lutetiana* L.: II-V (29,24,40,!); G, Eua; H3.5T3R4

TRAPACEAE

560. *Trapa natans* L.: IV V (14,25,!); Hh, Eua (M); H6T4R4
 - var. *lacvigata* (Nath.)Gluck: V (40)
 - var. *conocarpa* Aresch: V (38)

HALORAGACEAE

561. *Myrophillum spicatum* L.: III (14,38,29), V (38,40); Hh, Cp; H4T0R4.5

562. *M. verticillatum* L.: III (14,38,!); IV (5), V (38,40,!); Hh, Cp; H6T3.5R3.5
- f. *pinnatifidum* (Wallr.) Topa: V (40)

MALVACEAE

563. *Hibiscus trionum* L.: III-V (14,5,18,!); TH, Eua; H2.5T4R4
564. *Abutilon theophrasti* Medik.: III (42,14), V (14,40,!); Th, Eua; H3T3R3
565. *Althaea officinalis* L.: III-V (25,18,!); H, Eua (C); H3T4R4
566. *Lavatera thuringiaca* L.: III-V (29,5,25,40,!); H, Eua (C); H2.5T3R0
567. *Malva neglecta* Wallr.: II-V (29,5,18,40,!); Th-TH, Eua (M); H3T3R3
568. *M. pusilla* Sm.: III-V (9,5,18,!); Th-H, Eua (m); H3.5T3R3
- f. *glabrescens* Morariu: III (14)
569. *M. silvestris* L.: III-V (5,29,25,18,!); Th-H, Eua (Cosm); H3T3R0
- var. *hispidula* Beck: III (14)

TILIACEAE

570. *Tilia cordata* Mill.: I-IV (23,22,29,5,25,!); MPh, E; H3T3R3
571. *T. platyphyllos* Scop.: IV V (5,25,24,!); MPh, Ec; H2.5T3R4
572. *T. tomentosa* Mnch.: IV (14,25,!); MPh, B; H2.5T3.5R3

LINACEAE

573. *Linum catharticum* L.: I-IV (6,31,12,33,5,25,!); Th (TH), E(M); H3T2R4
574. *L. flavum* L.: III-IV (29,5); H, P-Pn-B; H2T4R4
- var. *angustifolium* J.v.: I (14)
575. *L. perenne* L.: IV (5,14); H, Eua (C); H0T3.5R4

OXALIDACEAE

576. *Oxalis acetosella* L.: II (22,!); H-G, Cp; H4T3R3
577. *O. europaea* Jord. (*O. stricta* Auct. non L.): II-III, V (14,!); Th-H, Adv; H3.5T0R0

GERANIACEAE

578. *Geranium columbinum* L.: IV (5,25); Th, Eua; H2T3.5R4
579. *G. divaricatum* Ehrh.: I, IV (14); Th, Eua (M); H2.5T3R4
580. *G. palustre* Torn.: I (23,30,!); III (29,14); H, Eua (C); H4T3R4.5
581. *G. phacum* L.: II-IV (22,15,5,!); H, Ec; H4T3R3
582. *G. pratense* L.: I-V; H, Eua; H3.5T3R5
583. *G. pusillum* Burm.: IV-V (5,18,!); Th, E (M); H2.5T3R0
584. *G. robertianum* L.: II (!), IV-V (5,25,40,!); Th, Cosm; H3.5T3R3
585. *G. rotundifolium* L.: IV (14,5,!); Th, Eua (M); H2T3.5R4
- var. *trichospermum* Sanio et Bob.: IV (14,25)
586. *Erodium cicutarium* (L.) L'Hérit: IV-V (5,18,!); Th, Cosm; H2.5T0R0

BALSAMINACEAE

587. *Impatiens glandulifera* Royle (*I. roylei* Walp.): I, IV (14); Th, Adv; H4T4R4
588. *I. noli-tangere* L.: II-IV (29,25,!); Th, Eua; H4T3R4

ZYGOPHYLLACEAE

589. *Tribulus terrestris* L.: IV V (14.5); Th. Cosm; H0T4R4
- var. *orientalis* (Kern.) Beck: V (14,40)

POLYGALACEAE

590. *Polygala amara* L.: I (14); H (Ch), E; H0T2R4.5
591. *P. comosa* Schkuhr.: I (6.32), II (34); H (Ch), Eua; H2T4R4
592. *P. vulgaris* L.: I (14.6,!); III-V (29.5,!); H (Ch), Eua; H3T3R3
- f. *rosulata* (Fries) Hegi: I (14)
- f. *major* (Koch) Hegi: I (14)

ACERACEAE

593. *Acer campestre* L.: III-V (15.29.6.25.24,!); MPh-mPh, E; H2.5T3R3
594. *A. platanoides* L.: II-III (14.22.29); MPh, Eua; H3T3R3
595. *A. tataricum* L.: III-V (14.5.40.25,!); mPh-MPh, E (C); H2.5T3.5R4

CELASTRACEAE

596. *Eonymus europaeus* L.: I (30.23), III-V (5.27.18,!); mPh, E; H3T3R3
- f. *angustifolia* (Schultz.) Borza: V (14)
597. *E. nana* M.B.: I (23.14); mPh, Eua (C); H0T3R3.5
598. *E. verrucosus* Scop.: II-IV (22.9.5,!); mPh, E; H2.5T3R4

STAPHYLEACEAE

599. *Staphylea pinnata* L.: III-V (14.5.25.24,!); mPh, E(M); H2.5T3.5R4

RHAMNACEAE

600. *Frangula alnus* Mill.: I (23.31,!); IV V (25.24.40); mPh, Eua; H4T3R3
601. *Rhamnus catharticus* L.: IV V (5.27.40); mPh, Eua; H2T3R4

VITACEAE

602. *Vitis silvestris* Gmel.: V (14.24.40,!); mPh-Ep, P-M; H3.5T4.5R4.5

CORNACEAE

603. *Cornus mas* L.: IV V (25.24.27,!); mPh, P-M; H2T3.5R4

604. *C. sanguinea* L.: III-V (29.5.24.40,!); mPh, Ec; H3T3R4

ARALIACEAE

605. *Hedera helix* L.: III-IV (29.5,!); nPh-Ep, A(l)-M; H3T3R3

APIACEAE (UMBELLIFERAEE)

606. *Sanicula europaea* L.: II (22,!); H, A(l)-M; H3.5T3R4
607. *Eryngium campestre* L.: III-IV (29.5.25,!); H.P; H1T5R4
608. *E. planum* L.: III-V (14.5.40,!); H, Eua (C); H2T3R4
- f. *subglobulosum* (Uechter.) Schubc: IV (43)

609. Chacrophyllum aromaticum L.: I-II (14,22,!); H, Ec (C); H3.5T3R3
 - var. longipilum Thell.: II (14)
 - f. cinerascens Borb.: I (14)
 - var. brevipilium Murbeck f. tomentellum Todor: I (14)
610. C. bulbosum L.: III-IV (29,5,!); TH-H, E (C); H4T3.5R4.5
 - f. tenuissimum (Vel.)Jav : III (14)
611. C. hirsutum L. (C.cicutaria Vill.): I-III (31,14,29,!); H, Ec; H4.5T2R0
612. C. temulum L.: III-V (29,14,25,24); Th-TH, E: H3T3R4
613. Anthriscus caucalis M.B. (A. scandicina (Web.)Mansf): IV-V (5,15,40); Th, P-M:
 H2T4R0
614. A. cerefolium (L.)Hoffm. ssp. trichosperma (Spreng.)Arc. (A. trichosperma Spreng.):
 IV-V (5,18,!); Th,M; H2.5T4R4
615. A. silvestris (L.)Hoffm.: II-V (22,29,14,5,24,!); H, Eua (M); H3T3R4
 - f. glabrescens (Schur)Todor: III-IV (14)
616. Turgenia latifolia (L.)Hoffm. (Caucalis latifolia L.): IV V(14,40); Th, Eua (M);
 H2.5T4R0
617. Caucalis platycarpos L. (C. lappula (Web.)Grande): III-V (29,6,40); Th, Ec-M;
 H2T4R5
 - var. muricata (Bisch.)Gren. et Godr.: IV-V (14,25,5,18)
618. Torilis arvensis (Huds.)Link: IV-V (14,40); Th, Ec-M; H2.5T3.5R4
 - var. aglochis (Simk.)Jav : V (14)
619. T. japonica (Houtt.)DC. (T. rubella Mnch.): III-V (29,5,40); Th-TH,Eua; H3T3.5R4.5
620. Daucus carota L.: I-V: TH-H, Eua (M); H2.5T3R0
621. Bifora radians M.B.: III-IV (14,29,!); Th, M; H3T4R0
622. Conium maculatum L.: III-V (29,5,25,18,!); Th-TH, M; H3T3R3
623. Pleurospermum austriacum (L.)Hoffm.: I (14); H, Ec; H3T2R4
624. Bupleurum falcatum L.: I-IV (14,22,5,!); H, Eua; H2T3.5R4
625. B. tenuissimum L.: IV V (14,27,18); Th, AtI-M; H0T3.5R4.5
626. Cicuta virosa L.: I-V; Hh,Eua; H5T0R3
627. Trinia ramosissima (Fisch et Trev)Koch (T. kitaibelli M.B.): IV V (5,27,40); H, B-P:
 H2.5T3.5R4
628. Carum carvi L.: I-V; TH, Eua; H3.5T3R3
629. Aegopodium podagraria L.: I-V; H (G), Eua; H3.5T3R3
630. Pimpinella major (L.)Huds.: I (14,!); H, E; H3.5T0R4
 - var. rubra (Hoppe et Schleich.)Fiori et Paol.: I (14)
631. P. saxifraga L.: I-V; H, Eua; H2.5T0R3
 - var. ovata Spreng. f. pubescens (Mert. et Koch)Ny r : III, V(14)
632. Berula erecta (Huds.)Caville (Sium erectum Huds.): III (!); Hh, Cp: H6T3.5R0
633. Sium latifolium L.: III-V (14,40,!); Hh, Eua; H6T0R4
634. S. sisaroides DC. (S. lancifolium M.B.): IV-V (14,43,25,40);Hh, Eua (C) H5T3R4
635. Seseli annuum L.: I (34), IV (5); Th (th,H), E (C); H2T3R3
636. S. libanotis (L.)Koch (Libanotis montana Cr.): II-IV (14,22,5,!); H, Eua (C); H3T0R4
 - var. sibirica (L.)Patz, Mey et Eik.: II (14)
637. Oenanthe aquatica (L.)Poir.: I-V; Hh,Eua; H6T3R0

638. *O. banatica* Heuff.: I (14); III-V (14,25,27,!); H, D-B-Pn; H4T3.5R0
 639. *O. fistulosa* L.: III-IV (14,29); Hh, E; H5T4R4
 640. *O. silaifolia* M.B.: I (6,31); IV-V (14,6,40); H, M; H5T3.5R0
 641. *Actusa cynapium* L.: III-V (14,29,5,40,!); Th-TH, E; H3.5T3R0
 - ssp. *cynapioides* (M.B.)Nym.: III (14)
 - ssp. *agrestis* (Wallr.)Dost.: V (14,40)
 642. *Cnidium dubium* (Schkuhr)Thell.: I (23,32,14,!); TH (H), Eua; H4.5T4R3.5
 643. *Selinum carvifolia* L.: I (6,23,14,!); IV (14,25); H, Eua; H3.5T3R3
 644. *Angelica palustris* (Bess.)Hoffm.: I (14); H, Eua (C); H5T2.5R0
 646. *A. silvestris* L.: I-V; H, Eua; H4T3R3
 - var. *vulgaris* Fisch f. *stipulazis* (Schur)Thell.: III (14)
 646. *Peucedanum alsaticum* L.: IV-V (14,6,40); H, Ec; H2T3.5R4
 - f. *angustifolium* Erdn.: V (14)
 647. *P. carvifolia* Vill. (*P. chabraci* (Jacq.)Rchb.): IV (14,25,!); H, Ec; H3T3R4
 648. *P. officinale* L.: III (29). V (14,27); H, E; H2T3R3
 649. *P. orcosclinum* (L.)Mnch.: II-IV (22,29,5,!); H, Ec-M; H2.5T3R0
 650. *P. palustre* (L.)Munch.: I (14,23); H, Eua; H5T3R0
 651. *Pastinaca sativa* L.: II-V (34,29,5,!); TH-H, Eua; H3T4R4
 652. *Heracleum sphondylium* L.: II-IV (22,14,5,25,!); H, Eua; H3T2.5R0
 - ssp. *sphondylium* var. *branca-ursina* (Cr.)Thell.: III-IV (14,25)
 653. *Laserpitium latifolium* L.: I-II (14,22,!); IV (5); H, E; H0T0R4
 - var. *asperum* (Cr.)Soy-Will.: I (14)
 654. *L. ruthenicum* L.: I (14,23); H, Ec; H4T3.5R4
 - var. *hirtum* Wallr.: I (14)

PLUMBAGINACEAE

655. *Limonium gmelini* (Willd.)O. Ktze. (*Statice gmelini* Willd.): III-V (14,29,34,18,!); H, Eua (C); H3.5T4R4

PRIMULACEAE

656. *Lysimachia nummularia* L.: I-V; Ch,E; H4T3R0
 657. *L. punctata* L.: III-V (29,25,40); H, P-M; H3.5T3.5R3
 658. *L. thrysiflora* L.: I-II (14,23,31); Hh, Cp; H5T0R0
 659. *L. vulgaris* L.: I-V; H-Hh, Eua; H5T0R0
 660. *Angallis arvensis* L.: III-V (5,18,25,!); Th.Cosm; H3T3R0
 661. *A. foemina* Mill. (*A. coerulea* Schreb.): IV V (5,18,40,!); Th, Cosm; H3T3.5R0
 662. *Centunculus minimus* L.: IV (14,5); Th, Eua; H4T3.5R3
 663. *Samolus valerandi* L.: III (14); H, Eua; H4.5T3.5R0
 664. *Soldanella montana* Willd.: II (14); H, Ec; H3.5T2R1.5
 665. *Androsace elongata* L.: V (14,40); Th, Eua; H2T3.5R4
 666. *Hottonia palustris* L.: II-III (14,38); Hh, E; H6T3.5R3
 667. *Primula acaulis* (L.)Gruib.: IV (14,25); H, Atl-M; H3T3R3
 668. *P. veris* L. (*P. officinalis* (L.)Hill.): II-IV (22,29,25,!); H, Eua; H3T2R5

PYROLACEAE

669. *Pyrola rotundifolia* L.: I (14,23,6), IV (14); H, Cp: H3T0R2.5
670. *Monotropa hypopitys* L.: III (14,29) V (14,40); G, Cp: H3T2R0

ERICACEAE

671. *Vaccinium myrtillus* L.: I (!): nPh (Ch). Cp: H0T2R1
672. *V. vitis-idaea* L.: I (6,!); Ch (nPh), Cp: H3T2R1

CONVOLVULACEAE

673. *Calystegia sepium* (L.) Br : II-V (12,29,5,25,18,!); H, Eua; H4T3R4
674. *Convolvulus arvensis* L.: II-V (29,5,18,!); H-G,Cosm: H0T0R0

CUSCUTACEAE

675. *Cuscuta campestris* Yuncker ssp. *pentagona* (Engelm.) (C. *pentagona* Engelm.): IV-V (14,40); Th,Adv ; H3T3R0
676. C. *epithymum* (L.) Murr : I-II (14), IV-V (5,40,!); Th, Eua; H0T3R0
677. C. *europaea* L.: IV V (14,5,40); Th,Eua: H4T0R0
- var. *nefrens* Fries: IV (14)
678. C. *epilinum* Weihc: III-IV (14); Th, Eua; H0T4R0
679. C. *lupuliformis* Krock: IV V (14,43,40,!); Th,Eua (C);H0T3R0
680. C. *trifolii* Babigt.: II-V (14,40,!); Th, Eua : H0T3R0
- var *angustissima* (Engelm.)Buia f. *longisquama* Buia: II-III (14)
- var. *muresensis* Buia: III (14)

POLEMONIACEAE

681. *Polemonium caeruleum* L.: I-II (14,23,32); H, Cp: H4T2.5R4.5

HELIOTROPIACEAE

682. *Heliotropium europaeum* L.: III-V (14,5,47,40); Th, Ec-M; H2T4R0
683. H. *supinum* L.: V (40); Th, M; H0T4R4.5

BORAGINACEAE

684. *Cerinthe minor* L.: III-V (29,5,33,46,18); TH, P-M; H3T3R0
685. *Lithospermum arvense* L.: III-V (15,5,18); Th-TH, Eua; H0T0R4
- var. *caerulescens* DC.: III (14)
686. L. *officinale* L.: IV V (5,18); H, Eua; H2T3.5R4
687. L. *purpureo-caeruleum* L.: IV (5,25,!); H-G, Ec (M): H2.5T4R4.5
688. *Onosma arenarium* W et K: IV V (14,40); H, E (C); H1.5T3.5R4
689. *Echium vulgare* L.: II-V (22,29,5,18,!); Th, Eua; H2T3R4
690. *Myosotis arvensis* (L.) Hill.: III-V (29,5,18,40); TH,Eua: H3T3R4
691. M. *caespitosa* K.F.Schultz: V (40); Th-TH (H), Cp; H4.5T0R0
692. M. *alpestris* F.W. Schmidt ssp. *stenophylla* (Knaf.)Grint. et Ny r. (M. *stenophylla* Knaf.): II (14); H, Ec; H1.5T1.5R3

693. *M. scorpioides* L. (*M. palustris* (L.) Nathh.); I-V; H, Hh, Eua; H5T3R0
 - var *memor* Kitt.: I (14), V (14,25)
694. *M. silvatica* (Ehrh.) Hoffm.; II-IV (22,29,5,25); H, Eua; H3,5T3R3
695. *M. sparsiflora* Mikan; V (18,40); Th, Eua (C); H3,5T3R4
696. *Pulmonaria mollis* Wulfen (*P. montana* Lej.); IV (25); H, Eua; H2,5T3R4
 - ssp. *mollissima* (Kern.) Nym.; IV (25), V (40)
697. *P. officinalis* L.; III-V (14,29,6,40,!); H, E; H3,5T3R3
698. *P. rubra* Schott; I-II (14,22,!); H, Carp-B; H3,5T2R3
699. *Symphytum cordatum* W et K.; II (14,22,!); H-G, Carp-B; H3,5T2R3
700. *S. officinale* L.; I-V; H, Eua; H4T3R0
 - ssp. *uliginosum* (Kern.) Nym.; IV (14)
701. *S. tuberosum* L.; IV (5,25); H-G, Ec:H3T3R3
702. *Anchusa officinalis* L.; II-V (22,29,5,25,18,!); TH-H, E(M); H2T3,5R0
 - var. *arvalis* Dum.; II, V(14)
703. *Cynoglossum officinale* L.; IV V (5,18,40); TH, Eua (C); H2T3R4

SOLANACEAE

704. *Atropa belladonna* L.; II (22); H, Atl-M; H3T3R3
705. *Scopolia carniolica* Jacq.; II (14,22); G, E; H4T3R5
706. *Hyoscyamus niger* L.; III-V (29,5,18,14,!); TH-H, Eua(M); H3T3,5R4
 - f. *pallidus* (W et K.) Rchb.; IV (14)
707. *Physalis alkekengi* L.; III-V (29,5,40,!); H, Ec-M; H3T3R4
708. *Solanum alatum* Mnch.; IV V (14,40); Th, Eua (M); H3T4R3
709. *S. dulcamara* L.; I-V; Ch (nPh), Eua (M); H4,5T3R4
710. *S. luteum* Mill.; IV (14); Th, Eua (M); H3T3,5R4
711. *S. nigrum* L.; III-V (29,5,25,18,!); Th, Cosm; H3T4R0
712. *Datura stramonium* L.; III-V (15,29,6,25,18,!); Th, Cosm; H3T4R4

SCROPHULARIACEAE

713. *Verbascum blattaria* L.; III-V (34,29,5,25,18,!); H, Eua (M); H2,5T3,5R4
714. *V. densiflorum* Bertnl. (*V. taphsiforme* Schrad.); IV V (14,5); TH, E (M); H2,5T3,5R4,5
715. *V. lychnitis* L.; II-V (22,29,40,18,!); Th, E; H1T3R5
 - ssp. *kanitzianum* (Simk. et Walz.) Soc; II (14)
716. *V. nigrum* L.; III-V (29,25,27,40,!); TH-H, Eua; H2T3R4
717. *V. phlomoides* L.; III-V (6,25,18,!); TH, E; H2,5T3,5R4
 - f. *australe* (Schrad.) Ghisa; III-V (14)
718. *V. phoeniceum* L.; III-V (4,29,5,25,18,!); H, Eua (C); H2T4R4
719. *V. x schottianum* Schrad. (*chaixii* ssp. *austriacum* x *speciosum*); I (14)
720. *Linaria angustissima* (Lois.) Borb.; IV-V (14,5,40); H,M; H1T3,5R5
721. *L. genistifolia* (L.) Mill. ssp. *genistifolia*; III-V (14,6,25,40,!); H, Eua (C); H1T3,5R5
 - var. *procera* Sims.; IV-V (14,40)
 - var. *angustata* Wierzb.; V (40)
722. *L. x kocianovichii* Aschers. (*genistifolia* x *vulgaris*); IV V (14,25,40)

723. *L. vulgaris* Mill.: II-V (22,34,29,5,18,!); H (TH), Eua; H2T3R4
 - var. *glabra* Peterm.: IV V (5.40)
724. *Kickxia elatine* (L.)Dumort: III (42), V(40); Th, Ec (M); H2.5T3.5R4
725. *Scrophularia laciniata* W et K. ssp. *lasiocaulis* (Schur)Borza: IV (14); H, Carp-B;
 H2T3.5R0
726. *S. nodosa* L.: I (30), III-V (29,5,24,!); H, Eua; H3.5T3R0
727. *S. scopolii* Hoppe: I (30,!); IV V (25,40,!); H, P-m; H4T3R0
728. *Gratiola officinalis* L.: IV V (25,18,40,!); H, Eua; H4.5T3R4
729. *Limosella aquatica* L.: V (14,40); Th, Cosm; H4.5T3R0
730. *Lindernia procumbens* (Krocker)Philcox (*L. pyxidaria* All.): V (14,40); Th, Eua (m);
 H4.5T4R0
731. *Veronica acinifolia* L.: IV V (14,40); Th,M; H4T3.5R0
732. *V. anagallis-aquatica* L.: I (31,!); III-V (5,40,18,!); H-Hh, Cp; H5T0R4
 - f. *limosa* Krosche: V (14)
 - f. *maxima* (Schur)J. Kell.: III(14)
 - f. *tencrima* (Schm.)Vhal.: V (14)
733. *V. anagalloides* Guss.: IV-V (14,40,47); H-Hh, Eua; H4.5T0R4
734. *V. agrestis* L.: III-IV (29.5); Th, E; H3.5T2.5R4
735. *V. arvensis* L.: III-V (15,29,5,18,47); Th, Eua; H2.5T3R3
736. *V. beccabunga* L.: II-V (29,14,40,!); Hh-H, Eua; H5T3R4
 - var. *limosa* (Lej.)Math.: IV-V (14,40);
737. *V. catenata* Pennel (*V. aquatica* Bernh.): III (14); H-Hh, Cp; H5T3R4.5
738. *V. chamaedrys* L.: I-V; H-Ch, Eua; H3T0R0
739. *V. hederifolia* L.: IV-V (5,40,47,!); Th, Eua; H2.5T3R4
740. *V. longifolia* L.: I (14,23,!); V(18,40,!); H, Eua; H4T3R4
 - f. *media* (Schrad.)Koch: I.III (14)
 - f. *salicifolia* (Wallr.)Hay : V (14)
741. *V. officinalis* L.: II-IV (22,29,6,25,!); Ch, Eua; H2T2R2
742. *V. opaca* Fries.: IV-V (5,40); Th, E; H2.5T3R4.5
743. *V. paniculata* L. (*V. spuria* L.): IV-V (43,18); H, Eua; H0T3R4
744. *V. persica* Poir.: IV-V (5,40,!); Th, Adv; H3T0R4
745. *V. polita* Fries (*V. didyma* Auct.): IV-V (47,40); Th, Eua (M); H2.5T3.5R4.5
746. *V. praecox* All.: IV-V (14.5,25,40); Th, EC-M; H1.5T3.5R3
747. *V. prostrata* L.: III (33), V (18,40,!); Ch, Eua; H2T4R3
748. *V. scutellata* L.: I (14,31,!); III (14), IV (14,5); H-Hh, Cp; H4T3R4
749. *V. serpyllifolia* L.: I-V; H, Cosm; H3T3R0
750. *V. spicata*: III (34,29); H-Ch, Eua; H1T4R4
 - var. *lancifolia* Koch: V (40)
 - ssp. *orchidea* (Cr.)Celak.: II-V (34,33,40,!); H, P-M; H1.5T5R4
 - f. *pseudocrassifolia* Borb.: IV (14,25)
751. *V. triphyllus* L.: IV V (25,50); Th, E; H2T3.5R2
752. *Digitalis grandiflora* Mill.: II-IV (22,29,25,!); H, E; H3T3R3
753. *Odontites scrotilina* (Lam.)Rchb.: I-V; Th, Eua; H3T3R0
754. *Euphrasia rostkoviana* Hayne: I (14,!); III (29,!); Th.Ec; H3T3R3

- var. *montana* (Jord.) Wetst.: I (14,42)
- 755. *E. stricta* Host.: I (6,23,!); IV (25); Th. Ec; H3T3R0
- 756. *Pedicularis palustris* L.: I (23,!); IV (14.5); H, Eua; H5T0R0
- 757. *P. sceptrum-carolinum* L.: I (23,31,!); H, Eua; H4.5T2.5R4.5
- 758. *Rhinanthus angustifolius* Gmel. (*R. glaber* Lam.): I-II (22,!); Th, Eua; H0T0R0
- var. *bosnensis* Behr. et Stern: I (14)
- 759. *R. minor* L.: I-IV (14,40); Th, D-B-Anat; H3T4R0
- 760. *R. rumelicus* Velen.: IV-V (14.40); Th, D-B-Anat; H3T4R0
- var. *simonkaianus* (Soó) Nyár.: IV-V (14,40)
- 761. *Melampyrum barbatum* W. et K.: III-V (14,29,40); Th, Pn-D; H2.5T3.5R4
- var. *filarszkyanum* (Soó) Pauca et Nyár.: V (14)
- 762. *M. bihariense* Kern.: I (32), III-IV (29.14); Th, D-B; H2.5T3R3
- var. *kummerlei* (Dahl.) Nyár.: I (14)
- 763. *M. cristatum* L.: I (6,30,!); III-V (29,14,5,25,!); Th, Eua; H2T3R4.5
- var. *ronnigeri* (Poev) Beauv.: I, IV-V (14,40)

LENTIBULARIACEAE

- 764. *Pinguicula vulgaris* L.: I (6,23); H, Cp; H3T0R4
- 765. *Utricularia australis* R.Br. (*U. neglecta* Lchm.): II,III (14); Hh, Atl-M; H6T3.5R3
- 766. *U. vulgaris* L.: I (31,14), II (14), III (14,!); IV (5,38), V (14,40,!); Hh, Cp; H6T0R3.5

OROBANCHACEAE

- 767. *Orobanche lutea* Baumg.: II-IV (14,5); G, Eua (M); H2T3R4
- var. *buekiana* Koch: IV (14,25)
- f. *collecta* Beck: IV (14,25)
- 768. *O. picridis* F.W. Schultz: II,IV (14); G, Ec-M; H2T3R0
- 769. *O. ramosa* L.: IV-V (5,40,!); Th (G), Ec-M; H3T4R0
- 770. *O. vulgaris* Poir. (*O. caryophyllacea* Sm.): II-V (14,40,!); G, Ec-M; H4T4R0
- var. *incurvata* Beck f. *brevispicata* Beck: IV (25)

VERBENACEAE

- 771. *Verbena officinalis* L.: III-V (29,5,18,!); Th-H, Cosm; H3T3R4
- 772. *V. supina* L.: V (40,!); Th, P-M; H4T4R0

LAMIACEAE (LABIATAE)

- 773. *Ajuga genevensis* L.: I-IV (32,22,29,5,25,!); H, Eua (C); H2.5T3R4
- 774. *A. reptans* L.: I (30,23,!); III-V (29,25,40,!); H-Ch, E; H3.5T0R0
- 775. *Teucrium chamaedrys* L.: II-IV (22,29,5,25,!); Ch, Ec-M; H2T3.5R4
- 776. *Scutellaria galericulata* L.: I (23,31,!); III-V (14,5,25,40,!); H, Cp; H4T3R4
- 777. *S. hastifolia* L.: III-V (29,14,5,18,!); H, Ec; H5T3R3
- 778. *Marrubium peregrinum* L.: IV-V (14,5,40); H, Eua (M); H2T4R0
- 779. *M. x remotum* Kit. (*peregrinum x vulgare*): IV-V (14,5,40)
- 780. *M. vulgare* L.: III-V (14,5,25,18,!); H (Ch), Eua (M); H1.5T4R4
- 781. *Nepeta cataria* L.: III-V (14,29,40); H (Ch), Eua (M); H3T3R4

782. *N. pannonica* L.: II-IV (22,29,14,!); H-Ch, Eua (C); H2T3R0
 783. *Glechoma hederacea* L.: I-V; Eua; H3.5T3R0
 - ssp. *hirsuta* (W. et K.) F. Hermann (*G. hirsuta* W et K.): IV-V(25,24,40,!); H-Ch, P-M;
 H2.5T3R4
 784. *Prunella grandiflora* (L.)Jacq.: I (6), IV (5,!); H, E (M); H3T3R4.5
 785. *P. vulgaris* L.: I-V; H, Cp; H3T3R0
 786. *Lamium amplexicaule* L.: IV-V (5,18,!); Th, Eua (M); H2.5T3.5R0
 787. *L. album* L.: I-V; H, Eua; H3T3R0
 788. *L. galeobdolon* (L.)Nathh. (*Galeobdolon luteum* Huds.): II-IV (29,5,!); H (Ch), Ec;
 H3T0R4
 789. *L. maculatum* L.: II-IV (22,5,25,!); H (Ch), E: H3.5T0R4
 790. *L. purpureum* L.: III-V (29,5,25,18,40,!); Th (H), Eua; H3T0R4
 791. *Galcopsis ladanum* L.: III-V (15,29,5,40,!); Th, Eua; H2T0R4.5
 792. *G. pubescens* Bess.: I (23), IV (14,25); Th, Ec; H3T3R0
 793. *G. speciosa* Mill.: I-V; Th, Eua (C); H3T2R0
 794. *G. tetrahit* L.: I (23,!), III-IV (29,5,25,!); Th, Eua; H3T3R0
 - var. *sylvestris* Schlecht: I (14)
 795. *Chaiturus marrubiastrum* (L.)Rechb.: IV-V (5,25,40,!); Th-TH, Eua (C); H4T3R0
 796. *Leonurus cardiaca* L.: III-V (14,5,25,18,!); H, Eua; H3T4R4.5
 797. *Ballota nigra* L.: I-V; H (Ch), Ec-M; H2T3.5R4
 798. *Stachys alpina* L.: I (14); H, Ec; H3T2R0
 799. *S. annua* (L.)L.: III-V (29,6,18,!); Th,M; H3T3.5R4.5
 800. *S. palustris* L.: I (23!). IV-V (5,43,18,!); H (G), Cp; H4T3R4
 801. *S. recta* L.: III-V (14,29,5,33,18,1); H, P-m; H2T4R4.5
 802. *S. sylvatica* L.: III-V (29,5,25,24,40,!); H, Eua; H3.5T0R0
 803. *Betonica officinalis* L.: I-V ; H, Eua (M); H3T3R0
 804. *Salvia austriaca* Jacq.: III-V (33,6,18,!); H, P-Pn; H2T3.5R4
 805. *S. glutinosa* L.: II (22,!); H, Eua; H3.5T4R3
 806. *S. nemorosa* L.: III-V (29,5,25,40,!); H, Ec; H2.5T4R3
 807. *S. pratensis* L.: III-V (15,5,33,40,!); H, E(M); H2.5T3R4.5
 808. *S. × sylvestris* L. (*nemorosa* × *pratensis*): III (14), V (14,40)
 809. *Melissa officinalis* L.: IV-V (14,40); H,M; H2T4R0
 810. *Calamintha acinos* (L.)Clairv.: II (22,!); IV (5,25,!); Th-TH, E(M); H1.5T3.5R4
 811. *C. clinopodium* Spenn. (*C. vulgaris* (L.)Druce): IV-V (25,5,24,!); H, Cp; H2T3R3
 812. *C. sylvatica* Bromf.: IV-V (14,27,25,!); H, Ec-M; H2.5T3.5R5
 813. *Origanum vulgare* L.: II-IV (29,25,!); H, Eua (M); H2.5T3R3
 814. *Thymus glabrescens* Willd.: II-IV (14,25); Ch,P-Pn; H2T4R0
 815. *T. pannonicus* All. (*T. marschallianus* Wild.): II-V (22,14,40,5,!); Ch, Eua (C);
 H1.5T3.5R4
 816. *T. pullegioides* L.: I-IV 932,6,25,!); Ch,Ec; H2.5T3R3
 817. *Lycopus europaeus* L.: I-V (23,31,12,34,5,24,!); Hh, Eua; H5T3R0
 818. *L. exaltatus* L. f.: V (24,40,!); Hh, Eua (C); H5T3R0
 819. *Mentha aquatica* L.: I (31,!); IV (6,43,!); Hh-H, Eua; H5T3R0
 - var. *riparia* (Schreb.)Gusul.: III (14), V (14,40)

820. *M. arvensis* L. ssp. *arvensis*: I (30,!); IV (5), V 940,!; H-G, Cp; H4T3R0
 - var. *soliicoma* (Op.)Top: III, V (14)
 - var. *varians* (Host)Top: I (14)
 - var. *pascuorum* Top: V (40)
 - ssp. *austriaca* (Jacq.)Top var. *austriaca*: V (14,40)
 - var. *hostii* (Bor.)Top: IV (14)
 - var. *politzensis* Top: V (14)
 - var. *cuncifolia* Lej. et Court.: V (40)
821. *M. x dalmatica* Tausch (arvensis x longifolia): V (40)
 - var. *skofitziana* (A.Kern.)Briq.: IV-V (14,40)
822. *M. longifolia* (L.)Nathh. ssp. *longifolia*: I-V; H (G), Eua (M); H4.5T3R0
 - var. *ensidens* Briq.: V (40)
 - var. *barthiana* (Borb.)Top: IV-V (14)
 - var. *balsamiflora* (H.Br.)Top: V (40)
 - ssp. *mollissima* (Borkh.)Dom.: V (40)
 - var. *brassoensis* Top: III (14)
 - var. *eclytanthea* Top: IV (14)
 - var. *hollosyana* (Borb.)Top: III (14), V (14,40)
 - var. *ischnostachya* Top: IV (14)
 - var. *leconeura* (Borb.)Top: III-V (14,40)
 - var. *marisensis* (Simk.)Top: V (40)
 - var. *paramcophyllum* Top: III (14), V (40)
 - ssp. *incana* (Willd.)Gusul. var. *macilenta* Briq.: V (14,40)
 - var. *planitiensis* Top: V (14,40)
 - var. *retinervis* (Borb.)Gusul.: IV (14)
 - var. *wierzbickiana* (Op.)Briq.: III (14), V (40)
 - var. *viridescens* (Borb.)Gusul.: III, IV (14)
823. *M. pulchrum* L.: I-V; H, Eua (M); H4T3R5
824. *M. verticillata* L.: I (32); H, E; H4.5T0R0
 - var. *calaminthaefolia* (Host.)Top: III (14)
 - var. *ovatifolia* Top: III (14), V (14,40)
 - var. *serotina* (Host.)Top: I (23)

PLANTAGINACEAE

825. *Plantago altissima* L.: III (14); H, B-Pn; H4T3R4
826. *P. argentea* Chaix.: III-IV (14,29,25); H, M; H1.5T4.5R4
827. *P. lanceolata* L.: I-V; H, Eua; H0T0R0
828. *P. major* L.: I-V; H, Eua; H3T0R0
829. *P. media* L.: I-V; H, Eua; H2.5T0R4.5
830. *P. maritima* L.: III (14), V (18); H, Eua; H4T0R5
831. *P. schwarzenbergiana* Schur: III (14); H, Pn-D; H3.5T4R5

GENTIANACEAE

832. *Menyanthes trifoliata* L.: I-III (14,23,32,!); Hh, Cp; H5T5R5

833. *Nymphoides peltata* (S.G. Gmel.) O.Ktze.: V (!); Hh, Eua (M); H6T3R4
834. *Centaurium erythraea* Rafn. (*C. umbellatum* Gilib.): III-V (34,29,5,40,!); Th, Eua; H3T3R2
835. *C. littoralis* (D. Turner) Gilmour ssp. *uliginosum* (W. et K.) Rothm. (*C. uliginosum* (W. et K.) Beck): III (14.); Th-TH, Eua; H4T3R4.5
836. *C. pulchellum* (Sw.) Druce: I (14), III-V (14,40,25,!); Th, Eua; H4T3.5R4
837. *Gentiana asclepiadcea* L.: I-IV (6,23,29,14,!); H, Ec; H4T2R4
- f. *cruciata* Wartm. et Schlatt.: I (32)
838. *G. pneumonanthe* L.: I (23,32,!); III (14,29), IV (14); H, Eua (M); H4T3R0

VINCACEAE

839. *Vinca minor* L.: III-V (14,5,29,25); Ch, Ec-M; H3T3R3

ASCLEPIADACEAE

840. *Cynanchum vincetoxicum* (L.) Pers.: II-V (22,29,5,25,24,40,!); H, E (M); H2T4R4

OLEACEAE

841. *Fraxinus angustifolia* Vahl.: V (24,!); MPh, P-Pn; H4,5T4R4.5
842. *F. excelsior* L.: II (22,!); III (29), V (24,27,!); MPh, E; H3T3R4
843. *F. ornus* L.: IV (14,5,25); mPh-MPh, M; H1,5T3.5R5
844. *Ligustrum vulgare* L.: III-V (29,5,25,24,!); mPh, E (M); H2,5T3R3

RUBIACEAE

845. *Sherardia arvensis* L.: IV-V (14,5,25,40); Th, Eua; H3T3R3
846. *Asperula arvensis* L.: III (14), IV (14,25); Th, M; H2T4R4
847. *A. campanulata* Klokov (*A. glauca* (L.) Bess.): II-IV (14,22,29,5,25,!); H, P-M; H2T4R4
- f. *hirsuta* (Wallr.) Borza: III (14)
848. *A. cynanchica* L.: II (!), IV-V (5,40,!); H, P-M; H2T3.5R4.5
849. *A. odorata* L.: II-IV (22,29,25,!); G, Eua; H3T3R3
850. *A. rivalis* Sibth. et Sm.: I (14,23), III-IV (14); H, Eua; H5T3R3
851. *Galium aparine* L.: I-V; H Eua; H3T3R3
852. *G. boreale* L.: I-III (14,6,29,!); H, Eua; H4T2R4
853. *G. mollugo* L. ssp. *mollugo*: III-V (29,5,18,!); H, Eua; H3T0R3
- ssp. *erectum* (Huds.) Briq. var. *erectum*: I (14), II (14,22,!); III, V (14)
- f. *deminutum* (Nyár.) Borza: III (14)
854. *G. palustre* L.: I-II (23,22,!); V (27,40,!); H, Cp; H5T3R0
855. *G. schultesii* Vest.: IV-V (5,25,24,!); G, Ec; H2,5T3R3
856. *G. rubioides* L.: III-V (14,5,40); H, Ec; H4T3R4
857. *G. tricornutum* Dandy: III-V (14,5,25,40); Th, Eua (M); H2,5T3.5R0
858. *G. uliginosum* L.: I (6,23,14,!); IV (14); H, Eua; H4,5T3R4
859. *G. verum* L.: I-IV (6,22,34,29,25,!); H, Eua; H2,5T2,5R0
860. *Cruciata glabra* (L.) Ehrend. (*Galium vernum* Auct.): I-V; H, Eua; H3T2R2

861. *C. laevipes* Opiz. (*Galium cruciata* (L.) Scop.); I-IV; H, Eua; H2.5T3R3

CAPRIFOLIACEAE

862. *Sambucus ebulus* L.: III-V (29,5,18,!); H, Eua (M); H3T3R4.5

863. *S. nigra* L.: III-V (15,5,25,24,!); mPh-MPh, E; H3T3R3

864. *S. racemosa* L.: I (!). II (22,!); mPh, Eua; H3T2R3

865. *Viburnum lantana* L.: II-IV (22,6,25,!); mPh, Ec-M; H2.5T3R4.5

866. *V. opulus* L.: I-V; mPh, Cp; H4T3R4

867. *Lonicera nigra* L.: I (23,30,!); mPh, E; H3T2R3

868. *L. xylosteum* L.: III-IV (29,5,!); mPh, Eua; H3T3R4

VALERIANACEAE

869. *Valerianella locusta* (L.) Betske: III-V (14,29,47); Th, Ec-M; H3T3.5R4

- f. *dasyarpa* (Rchb.) Borza: V (40)

871. *Valeriana officinalis* L.: I-V; H, Eua (M); H4T3R4

- f. *altissima*

872. *V. simplicifolia* (Rchb.) Kabath: I (6,32,!); H, Ec; H5T2R2

873. *V. tripteris* L.: II (22,!); H, Ec; H3T0R4.5

DIPSACACEAE

874. *Cephalaria pilosa* (L.) Gren. (*Dipsacus pilosus* L.); IV-V (14,25,40,!); TH, Atl-M; H4T3.5R4

875. *Dipsacus laciniatus* L.: III-V (29,40,!); Th, Eua (C); H4T3.5R4

876. *Succisa pratensis* Mnch.: I (6,32,!); III-IV (14,29); H, Eua; H4T3R0

877. *Succisella inflexa* (Kluk) G. Beck: IV V (14,40); H, Ec; H4T3R0

878. *Knautia arvensis* Coult. ssp. *arvensis*: I (32,!); III (29), IV (5,25,!); H, E; H2.5T3R0

- var. *polymorpha* (Schm.) Szabó f. *pratensis* (Schm.) Szabó: I (14).II (22).V (14)

- f. *jasionea* (Borb.) Szabó: V (14)

- var. *kitaibelii* (Svhult.) Szabó: V (40)

- ssp. *rosea* (Baumg.) Borb. (K. dumetorum Heuff.): IV (5,!); H,D-B; H2T4R3

- f. *heterotoma* (Borb.) Szabó: IV (14)

- f. *nitidula* (Simk.) Szabó: IV (14)

- f. *rosea* (Baumg.) Szabó: IV (14)

879. *Scabiosa banatica* W. et K.: I (14), III (14), IV (14,25); H, D-B; H2T2.5R4

880. *S. ochroleuca* L.: I (6), III-V (34,5,18,!); H, Eua (C); H2T4R4

CUCURBITACEAE

881. *Bryonia alba* L.: III-V (29,5,40,!); H-G, Eua (C); H3.5T4R0

882. *B. dioica* Jacq.: IV (14,5,25); H, E; H3.5T3R5

883. *Echinocystis lobata* (Mchx.) Torr. et Gray (E. *echinata* (Muhl.) Britt., Stern et Pogg.)

II-V (25,!); Th, Adv; H4T0R4

884. *Sicyos angulata* L.: IV (14); Th, Adv; H4.5T3R4

CAMPANULACEAE

885. *Campanula abietina* Gris. et Sch.: II (14); TH, Carp-B; H3.5T2R2
 886. *C. cervicaria* L.: I (6); H, Eua (C); H2.5T3R3
 887. *C. glomerata* L.: I (6), III-V (29.5,25,40); H, Eua; H2.5T3R4
 - f. *ramosa* Nyár.: III(14)
 888. *C. latifolia* L.: III-IV (29.14); H, Eua; H3T0R3
 889. *C. patula* L.: I-IV (6,22,29,5,25,40,!); TH, E; H3T2.5R3
 - f. *flaccida* (Wallr.)Nyár.: IV (43)
 890. *C. persicifolia* L.: II (!), III-IV (29.5,!); H, Eua (M); H3T3R0
 891. *C. rapunculoides* L.: I (!), II (22,!); IV (5,!); H, Eua; H3T2R0
 892. *C. serrata* (Kit.)Hendrych (*C. napuligera* Schur): I (14); H, C-B; H0T2.5R0
 893. *C. trachelium* L.: II (22), IV (5), V (24,!); H, Eua (M); H3T3R3
 894. *Adenophora liliifolia* (L.)Bess.: I (23,14); H, Eua (C); H0T4R4
 - f. *stylosa* (Lam.)Jav : I (42)
 895. *Phyteuma tetramerum* Schur: I (14); H, Carp (End); H3T3.5R3

ASTERACEAE

896. *Eupatorium cannabinum* L.: II-V (29.5,25,40,!); H, Eua (M); H4T3R0
 897. *Solidago canadensis* L.: III-V (14,!); H, Adv; H3.5T3R3
 898. *S. gigantea* Ait. (*S. scrotina* Ait.): IV-V (14,40,25); H, Adv; H3.5T3R3
 899. *S. virgaurea* L. ssp. *virgaurea*: II-IV (29,!); H, Cp; H2.5T3R3
 - var. *latifolia* Koch f. *cylindrica* Ny r.: IV (43)
 900. *Bellis perennis* L.: IV-V (5,25,40,!); H, E (M); H3T2.5R0
 901. *Aster amellus* L.: III-IV (14,25); H, Eua (C); H2T3R4
 902. *A. punctatus* W. et K. ssp. *punctatus* (*A. sedifolius* L.): III-V (29,14,27,25,40,!); H,
 Eua (C); H4T3R2
 - var. *squamosus* (Lallem.)Morariu et Nyár.: V (14,40)
 903. *A. tripolium* L. ssp. *tripolium*: III (29,34,!); H, Eua (M); H5T0R5
 - ssp. *pannonicus* (Jacq.)Soó: V (45,!);
 904. *Erigeron acris* L. ssp. *acris*: I-IV (14,6,22,5,!); Th-H, Cp; H2.5T3R0
 905. *E. annuus* (L.)Pers. (*Stenactis annua* (L.)Necs) ssp. *annuus*: II-V (14,5,18,!); Th, Adv;
 H4T0R4
 - ssp. *strigosus* (Muhlbg.)Soó (*Stenactis ramosa* (Wallr.)Don.): III-V (25,43,18,!); Th, Adv;
 H3T3R0
 906. *E. canadensis* L.: II-V (5,25,18,!); Th-TH, Adv; H2.5T0R0
 907. *Filago arvensis* L.: III-V (29.5,25,40); Th, Eua (M); H2T3.5R0
 908. *F. germanica* L.: III-V (14,25,40,!); Th, cua (M); H2T3R0
 909. *F. minima* (J.E.Sm.)Pers. (*F. montana* L.): IV-V (14.5); Th, Eua; H2T3R4
 910. *Antennaria dioica* (L.)Gaertn.: I-II (6,23,22,!); H-Ch, Cp; H3T1R2.5
 911. *Gnaphalium luteo-album* L.: IV-V (43,14,40); Th, Cosm; H4T3.5R3
 912. *G. uliginosum* L.: III (!), V (18,!); Th, Eua:H5T3R4
 - var. *nudum* (Hoffm.)Koch-Hall. f. *pilulare* (Wahlbg.)Ny r.: V (40)
 - var. *strictum* Nyár.: III (14)
 913. *Inula britannica* L.: III-V (14,34,25,4,27,!); TH-H, Eua (M); H3T3R0

914. *I. helenium* L.: III-V (14,29,5,!); H, Adv; H4T3R3
 915. *I. hirta* L.: I (6,!); III-IV (14,29,5,!); H, Eua (C); H2T4R5
 916. *I. salicina* L. ssp. *salicina*: I (6,!); III-V (14,29,25,!); H, Eua; H2.5T3R3
 - var. *glabra* Beck: V (14)
 - var. *subhirta* C.A.Mey: III (14), V (40)
 917. *Pulicaria dysenterica* (L.)Gaertn.: IV-V (5,14,40,!); H, E (M); H4T3.5R0
 918. *P. vulgaris* Gertn.: III-V (14,25,18,!); Th, Eua (M); H4T3R3
 919. *Carpesium cernuum* L.: IV-V (14,25,24); Th, Eua (M); H3.5T3.5R5
 920. *Telekia speciosa* (Schreb.)Baumg.: II (22,!); IV (25); H, C-B-Cauc; H4T2R0
 921. *Xanthium italicum* Moretti: IV-V (14,25,43,40,!); Th, Adv; H3.5T4R0
 922. *X. spinosum* L.: III-V (29,5,18,!); Th, Adv; H2.5T4R3
 923. *X. strumarium* L.: III-V (14,29,18,!); Th, Eua; H3.5T3.5R4
 924. *Helianthus decapetalus* L.: III-IV (14,!); H, Adv; H3.5T3R4
 925. *Rudbeckia laciniata* L.: III (14); H, Adv; H4.5T4R4
 926. *Bidens cernua* L.: III-V (29,40,!); Th, Eua; H5T0R0
 927. *B. tripartita* L.: II-V (22,5,27,18,!); Th, Eua; H4.5T3R0
 - f. *dumosa* Nyár.: V (14)
 928. *Galinsoga parviflora* Cav.: III-V (40,!); Th, Adv; H3.5T0R3
 929. *Anthemis arvensis* L.: I-V; Th, E (M); H3T3R0
 930. *A. austriaca* Jacq.: III (42), V (47,40); Th, Eua (C); H2T4R4.5
 931. *A. cotula* L.: V (18,40); Th, Cosm; H2.5T4R0
 932. *A. macrantha* Heuff: IV (14); H, Carp-B; H3.5T2R4
 933. *A. ruthenica* M.B.: III-IV (14,5); Th, E (C); H2T4R4
 934. *A. tinctoria* L.: II (22,!); IV (25); H, Eua; H1.5T3R3
 - ssp. *tinctoria* f. *monocephala* Jka.: IV (14)
 - ssp. *fussii* (Gris.)Beldic: II (14); H, Carp-B; H2T2R3
 935. *Achillea collina* Becker: II-V (22,33,5,25,47,!); H, Ec; H2T3R3
 936. *A. chrymisiolia* W. et K.: IV-V (5,14,25,27); H, B-Pn; H2.5T4R0
 937. *A. distans* W. et K. ssp. *distans*: I (6,32,!); II (22,!); H, Ec; H2.5T3R4
 - ssp. *stricta* (Schleich.)Janch. (*A. stricta* Schleich.): I-II (14); H, Alp-B; H2.5T2R3
 938. *A. x gergiocensis* Ny r. (*impatiens* x *ptarmica*): I (14,23,!);
 - f. *serrata* Nyár.: I (14)
 - f. *incisa* Nyár.: I (14)
 939. *A. impatiens* L.: I (14,6,23); H, Eua (C); H4.5T2.5R2.5
 940. *A. x maxima* (Heuff.)Prod. (*distans* x *millefolium*): I (14,42)
 941. *A. millefolium* L.: I-V; H, Eua; H3T0R0
 942. *A. ptarmica* L.: I (14,23,6,!); IV (14); H, Eua; H4.5T0R2.5
 943. *A. setacea* W. et K.: I (!); IV-V (25,40,!); H, Eua (C); H2T3R5
 944. *Matricaria chamomilla* L.: III-V (29,5,18,!); Th, Eua (M); H3T3R0
 945. *M. matricarioides* (Less.)Porter: I-III (14,!); Th, Adv; H3T0R0
 946. *Tripleurospermum inodorum* (L.)Schultz-Bip. (*Matricaria inodora* L.): III-V
 (6,18,47,!); Th-TH, Eua; H0T3R3.5
 947. *Chrysanthemum corymbosum* L.: II-V (14,22,29,5,25,!); H, Eua (M); H2.5T2.5R3
 - ssp. *clusii* (Fischer)Heywod: I (14)

948. *C. leucanthemum* L. ssp. *leucanthemum*: I-V; H, Eua; H3T0R0
 - var. *lanceolatum* (pers.)Beck f. *lanceolatum*: I (!), IV (43)
 - var. *auriculatum* (Peterm.)Nyár. f. *pinnatifidum* (Lec. et Lam.) Jav.: I (14)
 - f. *ramosum* Nyár.: I (14)
 - f. *silvestre* (Pers.)Nyár.: I (14)
 - f. *praestans* (Burm.)Nyár.: III (14)
 - f. *acquidentatum* Nyár.: V (14)
 - f. *ramulosum* Nyár.: II, IV (14)
949. *C. serotinum* L.: IV-V (14); H, P-Pn; H4T4R0
950. *Tanacetum vulgare* L. (*Chrysanthemum vulgare* (L.)Burm.): II-V (29,27,40,!); H, Eua; H3T3R0
951. *Artemisia absinthium* L.: II-V (22,5,25,40,!); Ch-H, Eua (M); H2T3.5R0
952. *A. annua* L.: III-V (25,43,!); Th, Eua; H3T4R4
953. *A. maritima* L. ssp. *salina* (Willd.)Gams: III (14), V (!); Ch (H), Eua (C); H2.5T3R0
 - ssp. *monogyna* (W. et K.)Gams: III (14)
954. *A. pontica* L.: V (27,40,!); H (Ch), Eua (C); H2.5T4R4.5
955. *A. scoparia* W. et K.: IV-V (14,25); H, Eua (C); H3T3R0
956. *A. vulgaris* L.: II-V (22,34,18,!); H-Ch, Cp; H3T3R4
957. *Tussilago farfara* L.: II-V (22,29,5,40,!); G-H, Eua; H3.5T0R4.5
958. *Petasites albus* (L.)Gartn.: II-III (22,29,!); G (H), Eua; H3.5T0R0
959. *P. x celakovskyi* Matuschek (*albus* x *kablikianus*): II (14.22)
960. *P. hybridus* (L.)G.M.Sch.: II-IV (22,29,5,!); G (H), Eua; H5T3R3
961. *P. x intercedens* Matuschek (*hybridus* x *kablikianus*): II (14)
962. *P. kablikianus* Tausch.: II (22,!); G (H), Carp-B; H4T0R0
963. *P. x reichingeri* Hay. (*albus* x *hybridus*): II (14)
964. *Homogyne alpina* (L.)Cass.: I (23); H, E; H3.5T2.5R2.5
965. *Doronicum austriacum* Jacq.: I-II (23,22,!); H, Ec; H3.5T2R3
966. *D. hungaricum* (Sadl.)Rchb. f.: III-IV (14,29); G (H), Ec-B; H2T3R4
967. *Senecio barbaracifolius* (Krock.)Wimm. et Grab.: III (14), IV V (14,40); H, Ec; H3.5T3.5R4.5
 - f. *foliosus* Nyár.: III (14), IV (43)
968. *S. doria* Nath.: III-V (14,18,!); H, Eua (C); H3T0R3.5
 - ssp. *schwetzovii* (Korsch)Nyár.: III-V (14,5,40)
969. *S. crucifolius* L.: IV-V (14); H, Eua; H3T3.5R4.5
 - var. *tenuifolius* (Jacq.)DC.: IV (14,5)
970. *S. fluvialis* Wallr.: I (23,!); III-IV (14); H, Eua (C); H5T4R4
971. *S. jacobea* L.: I (6,!); III-V (29,5,40,!); H, Eua; H2.5T3R3
972. *S. nemorensis* L. ssp. *nemorensis*: I (!); H, Eua; H3.5T3R3
 - var. *germanicus* (Wallr.)Beck f. *jacquinianus* (Rchb.)Weiss: I (14)
 - ssp. *fuchsii* (Gmel.)Cekak. (*S. fuchsii* Gmel.): I (30,!)
973. *S. paludosus* L.: I (14,31,23,!); V (14,18,!); H, Eua; H4.5T3.5R0
 - var. *glabratus* Koch: V (40)
 - var. *tomentosus* (Host.)Koch: V (40)
974. *S. papposus* (Rchb.)Less.: I (14,32), IV (14); H, Carp-B; H3T2R2.5

- var. sulphureus (Baumg.)Cuf.: I (14)
 - var. integerimus (Schur)Cuf.: I (14)
 - f. wolffii (Schur)Cuf.: I (42)
 - var. heuffelii (Jav.)Cuf.: IV (14)
975. *S. sylvaticus* L.: IV-V (14); Th, E; H3T3R3
976. *S. vernalis* W. et K.: III-V (14,40,!); Th-TH, Eu (C); H2.5T4R0
977. *S. viscosus* L.: IV (14); TH, E (M); H2T3.5R2.5
978. *S. vulgaris* L.: III-V (29,25,18,!); Th-TH, Eu: H3T0R0
979. *Ligularia sibirica* (L.)Cass.: I (14,23,32,!); H, Eu: H4T2R3.5
- f. arancosa (DC.)E.Pop: I (23,!);
980. *Echinops commutatus* Jur.: III-V (14,5,!); H, Alp-Carp-B; H2T0R4
981. *E. sphaerocephalus* L.: III-V (14,29,5,43,40,!); H, Eu: (C); H2T4R4.5
982. *Xeranthemum annuum* L.: III-V (29,5,25,40); Th, P-M; H2T4R3
983. *X. foetidum* Munch.: V (14,!); Th, P-M; H1.5T4R3
984. *Carlina vulgaris* L.: II-IV (14,29,5,!); TH-H, Eu: (M); H2.5T3R0
985. *Arctium x ambiguum* (Cel.)Nym. (*lappa x tomentosum*): IV (14)
986. *A. lappa* L.: II-V (5,18,!); TH, Eu: (M); H3T3R4
987. *A. minus* (Hill.)Bernh.: II (14), V (40); TH, E; H3T3R4.5
- var. *involucratum* Nyár.: III (14)
988. *A. x nothum* (Ruhm.)Weiss (*lappa x minus*) f. *subracemosum* (Simk.) M th.: III-IV (14)
989. *A. nemorosus* Lej. (*A. vulgare* (Hill.)A.H.Evans): IV-V (14,25); TH, E; H3.5T3R0
990. *A. tomentosum* Mill.: IV-V (5,40,!); TH, Eu: H3T0R5
991. *Carduus acanthoides* L.: II-V (22,29,5,25,18,!); TH, E (M); H2T3R0
- var. *albisfrons* Schur: IV (43)
992. *C. candicans* W. et K.: IV-V (14); H, B-Pn; H2T3R5
993. *C. crispus* L.: III-V (29,5,25,40,!); TH, E; H4T3R0
994. *C. hamulosus* Ehrh.: III-V (14,40); TH, P-Pn; H2T4R2
995. *C. nutans* L.: IV-V (5,18,!); TH-Th, Eu: (M); H1.5T0R4.5
996. *C. x orthocephalus* Wallr. (*acanthoides x nutans*): V (14,40)
997. *C. personata* (L.)Jacq.: II, V(14); H, Ec; H4.5T2.5R4.5
998. *Cirsium arvense* (L.)Scop.: I-V; G, Eu: (M); H0T0R0
999. *C. brachycephalum* Jur.: V (14,40); TH-H, Pn (End); H4T3R0
1000. *C. canum* (L.)All.: I-V; G, Eu: (c); H4.5T3R4.5
- f. *sagittatum* Ny r.: III (14)
1001. *C. erisithales* (Jacq.)Scop.: I-II (6,32,!); H.Ec; H3T3R4.5
1002. *C. x crucagincum* DC. (*oleraceum x rivulare*): I (14)
1003. *C. furiens* Gcis. et Sch.: III-IV (14,5,!); Th, Carp-Pn; H2T3.5R4
1004. *C. oleraceum* (L.)Scop.: I-II (30,22,!); H, Eu: H4T3R4
1005. *C. palustre* (L.)Scop.: I-II (30,22,6,23,!); TH, Eu: (M); H4.5T3R2.5
- f. *glomeratum* Nacg.: I (14)
1006. *C. x praecalpinum* Beck (*erisithales x rivulare*): II (14,42)
1007. *C. rivulare* (Jacq.)Link.: I-II (23,6,22,!); H, Ec; H4T3.5R0
1008. *C. x subalpinum* Gaud. (*palustre x rivulare*): I (14).

1009. *C. x tataricum* (L.)All. (*canum* x *oleraceum*): I (14)
1010. *C. vulgare* (Savi)Ten. (*C. lancolatum* (L.)Scop.): III-V (34.5,25,18,!); TH, Eua; H3T3R0
1011. *C. waldsteinii* Rouy: II (14); H, Aip-Carp-B; H4T2R2
1012. *Onopordon acanthium* L.: III-V (29.5,18,!); TH, M; H2.5T4R4
1013. *Serratula tinctoria* L.: I (6,!); III-V (34.29.5,18,!); H, Eua (M); H3.5T3R0
 - var. *variifrons* (Beck)Nyár.: I, III (14)
1014. *Centaurea x austriacoides* Woloszcz. (*phrygia* x *jacea*): I (14.42)
1015. *C. x beckeriana* J. Wagn. (*jacea* ssp. *banatica* x *indurata*): IV (14.25)
1016. *C. x baumgarteniana* J. Wagn. (*indurata* x *pseudophrygia*): I (14.42)
1017. *C. x brasoviensis* (*melanocalathia* x *pseudophrygia*): I (14)
1018. *C. calcitrapa* L.: IV-V (14.5,40); TH-Th, Eua (M); H1.5T4R0
1019. *C. cyanus* L.: I (!), III-V (5,25,18,!); Th, Cosm; H3T4R0
1020. *C. x erdneri* J. Wagn. (*phrygia* x *pseudophrygia*): I-II (14(!))
1021. *C. indurata* Janka: I, III (!), IV V (40,!); H, D-Pn; H3T3R3
1022. *C. jacea* L.: I-II (!), III (14,34), IV-V (33,14,!); H, Eua; H3T0R0
 - ssp. *banatica* (Roch.)Fay (*C. banatica* Roch.): IV-V (14.5,40)
 - f. *fastigiata* Prod.: V (14)
1023. *C. x markiana* J. Wagn. (*jacea* ssp. *banatica* x *pseudophrygia*): IV-V (14.25)
1024. *C. melanocalathia* Borb.: I-II (14,6); H, Ec; H3T2.5R0
1025. *C. micranthos* Gmel.: II-V (14.5,22,34,4,25,40,!); Th-H, Ec; H2T3.5R4
1026. *C. nigrescens* Willd.: III-V (14,25,40); H, Ec; H3.5T3R3
1027. *C. x nyaradyana* J. Wagn. (*jacea* x *melanocalathia*): I (14)
1028. *C. x orodensis* J. Wagn. (*jacea* ssp. *banatica* x *nigrescens*): V (14.40)
1029. *C. phrygia* L. (*C. austriaca* Willd.): I-V; H, Ec; H3T2.5R3
1030. *C. pseudophrygia* C.A.Mey.: I-II (14.6,32); H, Ec; H3T0R3
1031. *C. scabiosa* L. ssp. *scabiosa*: I (14), V (18); H, Eua (M); H2.5T0R4
 - f. *silesiaca* (Borb.)Soó: I (14)
 - ssp. *spinulosa* (Roch.)Hay (*C. spinulosa* Roch.): IV-V (14.25);
1032. *C. solstitialis* L.: III-V (14.5,40,25,!); TH-Th, Eua (M); H2T4R0
1033. *C. x spuria* Kern. (*jacea* ssp. *angustifolia* x *stenolepis*): V (14)
1034. *C. stoebe* L. (*C. rhenana* Boreau): II-IV (14,!); TH-H, Ec (M); H2T3.5R4.5
1035. *C. x szöllősi* J. Wagn. (*indurata* x *jacea* ssp. *angustifolia*): III-V (14.26)
1036. *Cichorium intybus* L.: II-V (34,29,5,25,18,!); H-TH, Eua; H2.5T3.5R4.5
1037. *Lapsana communis* L.: IV-V (5.25,24,40,!); Th-TH, Eua (M); H2.5T3R3
1038. *Hypochaeris maculata* L.: I-III (6,32,29,22,!); H, Eua (C); H0T3.5R3.5
 - f. *Iciliophylla* (Borb.)Nyár.: I (14)
 - f. *nitida* Nyár.: I (14)
1039. *H. radicata* L.: III-IV (25,!); H, E; H3T3R2.5
1040. *Leontodon autumnalis* L.: I-V; H, Eua; H3T0R0
1041. *L. hispidus* L. ssp. *hispidus*: III-IV (33.5,25,!); I (6,!); H, Eua; H2.5T0R0
 - f. *denticulatus* Csong.: III (14)
 - f. *lobatus* Ny r.: III (14)
 - f. *variifrons* Ny r.: III-IV (14)

- ssp. hastilis (L.)Rchb. (*L. danubialis* Jacq.): I (6,!); II (!); III(29); IV (25); H; Eua;
H3T3R0
- f. integrifolius Csong.: I (14)
- f. sinuato-dentatus Csong.: I-II (14)
- 1042. *L. saxatilis* Lam. (*L. nudicaulis* (L.)Banks.): III (14,34); TH-H; Atl-M; H3T3R4
- 1043. *Picris echioides* L. (*Helminthia echioides* (L.)Gaertn.): V (14); TH-Th; Adv;
H2.5T4.5R4
- 1044. *P hieracioides* L. ssp. *hieracioides*: III-IV (29.5.25,!); V (18); TH-H; Eua; H1.5T3R4
- f. *racemosa* Nyár.: III (14)
- f. *ruderalis* (Schin.)Beck: III (14)
- 1045. *Tragopogon dubius* Scop.: III-IV (14,!); TH; P-M; H2.5T3.5R0
- var. *campestris* Bess.: IV (5)
- 1046. *T. orientalis* L.: II-V (22,33,5,40,!); TH-H; Eua; H3T3R4
- var. *hayekii* (Soe)Nyár.: III (14)
- 1047. *Scorzonera hispanica* L.: III-IV (14.29,25); H; Eua (M); H2T5R4
- 1048. *S. parviflora* Jacq.: III (14,34,!); V (!); TH-H; Eua (C); H4T3.5R4
- 1049. *Podosperum canum* C.A.Mey.: IV-V (14,40); H (TH), P-M; H2T4R4.5
- 1050. *P. laciniatum* (L.)DC.: III-V (14.5,18,!); Th (TH); Eua; H2T0R4
- var. *muricatum* (Baalb.)Bisch.: III (14)
- 1051. *Chondrilla juncea* L.: IV-V (5.25,40,!); H; Eua (C); H1.5T3.5R4
- var. *latifolia* (M.B.)Koch: V (40)
- 1052. *Taraxacum bessarabicum* (Hornem.)Hand.-Mazz.: III (34,!); V (14,40,!); H; Eua (C);
H4T3R4
- 1053. *T. officinale* Weber: I-V; H; Eua (Cosm); H3T0R0
- 1054. *T. serotinum* (W. et K.)Poir.: IV-V (14,18); H; P-Pn; H2T4R4.5
- 1055. *Mycelis muralis* (L.)Dum.: III-V (29.24.25); H; E; H3T3R0
- 1056. *Lactuca x dichotoma* Simk. (*saligna x serriola*): V (14,40)
- 1057. *L. quercina* L. ssp. *quercina*: III-IV (14.29,25); TH; Ec; H2.5T3.5R4
- ssp. *sagittata* (W. et Kc)Celak. (*L. chaixii* Auct.): III-IV(14.5.25); TH; P-Pn; H2T3.5R4
- 1058. *L. saligna* L.: IV-V (5.25,18,!); Th-TH; M; H1.5T4R4
- f. *runcinata* (Gren. et Godr.)Beger: V (40)
- 1059. *L. serriola* Torner: III-V (25,18,!); Th-TH; Eua(M); H1.5T3.5R0
- var. *integrata* Gren. et Godr.: V (40)
- 1060. *Sonchus arvensis* L.: III-V (5,25,!); H; Eua (Cosm); H3T3R4
- 1061. *S. asper* (L.)Hill.:III-V (5.25,18); Th; Eua; H3.5T3R4
- f. *glandulosus* Beckh.: III (14)
- 1062. *S. oleraceus* (L.)Gou.: II-V (29.25,18,!); Th; Eua; H2.5T3R4.5
- var. *laceratus* (Willd.)Wallr.: V (14,40)
- 1063. *S. palustris* L.: I (14), III (14.29); H; Eua; H4.5T3.5R4
- 1064. *Crepis biennis* L.: I-V; TH; E; H3T3R4
- var. *lacera* Wimm. et Grab. f. *banatica* (Roch.)Ny r.: V (40)
- 1065. *C. foetida* L. ssp. *foetida*: IV (26); Th; Eua; H2T3.5R3
- ssp. *rhocadifolia* (M.B.)Fiori et Paol. (*C. rhocadifolia* M.B.): IV-V (5,40,18,!); Th; Eua;
H2.5T3.5R3

1066. *C. paludosa* (L.) Mnch.: I (31), II (12); H, E; H4.5T0R4.5
1067. *C. praemorsa* (L.) Tausch: I (14), III (14,29), IV (14,25); H, Eua; H2T3.5R5
1068. *C. setosa* Hall.: V (18); Th, Atl-M; H2T3R3
1069. *C. tectorum* L.: III-V (29,40,!); Th, Eua (C); H2.5T0R0
1070. *Hieracium x atramentarius* N.P. (*aurantiacum* x *piloselloides*)
 - var. *atramentarius*: I (14)
 - f. *csikense* Nyár.: I (14)
 - var. *hypochoerisfolius* Nyár.: I (14)
1071. *H. aurantiacum* L. ssp. *aurantiacum*: I-II (6,31,23,22,!); H, Eua; H2.5T2.5R2
 - var. *aurantiacum* f. *brevipilum* (N.P.) Nyár.: I (14)
 - f. *longipilum* (N.P.) Nyár.: II (14)
 - ssp. *carpathicola* Naeg. et Peter: II (14)
 - var. *subkajancense* (Zahn) Nyár.: I (14)
 - f. *longifolium* Nyár. et Zahn: I (14)
 - f. *ramosum* Nyár.: I (14)
1072. *H. auricula* Lam. et DC.: I-IV (22,34,29,25,!); H, E; H3T0R3
 - var. *acutisquamum* (N.P.) Nyár.: II (14)
 - var. *lampreilema* (N.P.) Nyár.: I, III (14)
 - f. *astolonum* Nyár.: I (14)
 - f. *perlongum* Nyár.: I (14)
 - var. *melaneilema* (N.P.) Nyár. f. *subpilosum* (N.P.) Nyár.: III (14)
 - var. *amaurcilema* (N.P.) Nyár.: I (14)
 - var. *magnauricula* (N.P.) Nyár.: I (14)
1073. *H. x auriculoides* L. ng (*bauhini* x *echioides*): V (18,40)
 - var. *paniculosum* Nyár.: III (14)
 - f. *flagellatum* Nyár.: III (14)
 - var. *gyvergoense* Nyár.: I (14)
 - var. *flexiramum* (N.P.) Nyár.: I (14)
 - var. *tanythrix* (N.P.) Nyár.: I (14)
 - var. *sarmentosum* (Froel.) Nyár.: III (14)
 - var. *parvicapitulum* (N.P.) Nyár.: III (14)
 - var. *semipracox* (Zahn) Nyár.: III (14)
 - var. *mirum* (N.P.) Nyár.: III (14)
1074. *H. bauhini* Besscr: I-V; H, Eua (C); H1.5T3R3.5
 - ssp. *cryptomastix* (N.P.) Nyár. var. *gemmaescrum* (N.P.) Nyár.: IV (14)
 - ssp. *aerostolonum* Zahn var. *bükkense* Nyár.: III (14)
 - var. *pseudosparsum* (Zahn) Nyár.: I (14)
 - ssp. *magyaricum* (N.P.) Nyár. var. *magyaricum*: III (14)
 - f. *pilosum* N.P.: III (14)
 - var. *besserianum* (Sprng.) Nyár.: I (14)
 - f. *calvius* N.P.: I (14)
 - var. *nigrisetum* (N.P.) Nyár.: I (14)
 - var. *pseudoauriculoides* (N.P.) Nyár.: III (14)
 - var. *filiferum* (Tausch) Nyár.: I, IV (14)

- var. heothinum (N.P.)Nyár.: I (14)
- ssp. bauhini var. adenocladum (Rehm.)Nyár.: I (14)
- var. ingricum (N.P.)Nyár.: I, IV (14)
- var. melachaetum (Tausch)Nyár.: I (14)
- 1075. *H. bifidum* Kit.: I (14); H, Ec; H2.5T2R4.5
- 1076. *H. x blyttianum* Fr. (aurantiacum x auricula): I (14)
- var. sandui (Prod.)Nyár.: I (14)
- var. detonaticum (N.P.)Nyár.: I (14)
- 1077. *H. x brachiatum* Bertol. (bauhini (vel piloselloides) x pilosella): II (14.22)
- 1078. *H. caespitosum* Dumort ssp. *caespitosum*: I (23), III (14), IV (5); H, Eua; H3T3R3
- var. *silvicolum* Nyár.: I, III (14)
- 1079. *H. x cochleatum* (N.P.)Zahn (auricula x caespitosum): I (14)
- 1080. *H. cymosum* L.: III-IV (14.5,25); H, Eua (C); H2T3R4
- ssp. *cymosum* var. *cymosum* f. *geotropum* Borb.: III (14)
- 1081. *H. x diaphanoides* Lbg. (lachenalii x sylvaticum): I (14)
- 1082. *H. echioides* Lumn.: IV (25); H, Eua (C); H2T3.5R4
- 1083. *H. x fuscum* Vill. (aurantiacum x auricula): I (14)
- 1084. *H. hoppeanum* Schult.: I (14), III (34), IV (25); H, Ec-M; H3T0R2
- var. *leucolepioides* (Deg. et Zahn)Nyár.: IV (25)
- 1085. *H. x koernickeanum* N.P. (auricula x bauhini): I, IV (14)
- var. *bauhiniforme* Nyár.: I (14)
- var. *kreuzenense* (Durm. et Oborny)Nyár.: I, III (14)
- var. *abortistolonum* Nyár.: III (14)
- var. *denigratum* (N.P.)Nyár.: I (14)
- f. *paucicapitatum* Nyár.: I (14)
- var. *auriculifolium* Nyár.: I (14)
- f. *acutum* Nyár.: I (14)
- 1086. *H. x levicaule* Jord. (bifidum x lachenalii): I (14)
- var. *psammogeton* (Zahn)Nyár.: I (14)
- ssp. *triviale* (Norrl.)Nyár. var. *serratilanceum* (Zahn)Nyár.: I-II (14)
- 1087. *H. lavigatum* Willd. ssp. *tridentatum* (Fr.)Zahn: I (14); H, Eua; H3T3R2
- 1088. *H. x leptophyton* N.P. (bauhini x pilosella) var. *leptophyton*: I (14)
- f. *stolonicaule* Zahn: I (14)
- 1089. *H. x longiscapum* Boiss. et Ky. (auricula x caespitosum): I (14)
- 1090. *H. x megatrichum* Borb. (auriculoides x cymosum): I (14)
- 1091. *H. x phaedrocheilon* Zahn (lachenalii x pseudobifidum): I (14)
- 1092. *H. pilosella* L.: I-IV (32,22,29,5,25,!); H, Eua (M); H2.5T0R0
- var. *tricholepium* (N.P.)Nyár.: III (14)
- var. *tomentisquamum* (N.P.)Nyár.: IV (14)
- 1093. *H. piloselloides* Vill.: IV (14,25); H, E (M); H2.5T3.5R4
- var. *devanum* (Zahn)Nyár.: IV (14)
- 1094. *H. x praecurrentis* Vukot (muronum x transsilvanicum): I (14)
- var. *pseudodiplodiophylloides* (Maly et Zahn)Nyár. f. *palotae* Nyár. et Zahn: I (14)

1095. *H. racemosum* W. et K.: I (6,!); IV (5,25); H, Ec-M; H2T3R3
 1096. *H. sabaudum* L.: IV (14,25); H, E; H2.5T3.5R2.5
 1097. *H. x sulphureum* Doll. (*auricula x piloselloides*): I (14)
 1098. *H. sylvaticum* (L.) Grub. (*H. murorum* L.): IV (25); H, Eua; H3T0R3
 1099. *H. x schultesii* F. Sch. (*auricula x pilosella*): II (14,22)
 1100. *H. transsilvanicum* Heuf.: II (14); H, Carp-B; H3T0R0
 1101. *H. umbellatum* L.: I (23), IV (25,!); H, Cp; H2.5T3R3

ALISMATACEAE

1102. *Alisma graminaceum* Gmel.: I (31), V (14); Hh, Cp; H6T0R4.5
 1103. *A. lanceolatum* Wither.: I (14), III (34); Hh, Eua; H6T0R4
 1104. *A. plantago-aquatica* L.: I (31,!); III (34,29,!); IV-V (5,18,!); Hh, Cosm; H6T0R0
 1105. *Sagittaria sagittifolia* L.: III-V (14,43,25,40,!); Hh, Eua (M); H6T3R4

BUTOMACEAE

1106. *Butomus umbellatus* L.: III-V (29,5,18,!); Hh, Eua (M); H6T3R0

HYDROCHARITACEAE

1107. *Hydrocharis morsus-ranae* L.: II-V (14,29,38,!); Hh, Eua; H6T3.5R.3.5

JUNCAGINACEAE

1108. *Triglochin maritima* L.: II-IV (14,12,34,!); H, Cosm; H4T0R4
 1109. *T. palustris* L.: I-IV (31,12,14,5,!); H, Cp; H5T0R0

POTAMOGETONACEAE

1110. *Potamogeton acutifolius* Link.: III (14,39); Hh, E; H6T3R4
 1111. *P. alpinus* Balb.: I (14,23); Hh, Cp; H6T2.5R2.5
 1112. *P. crispus* L.: II-V (38,29,5,40,!); Hh, Cosm; H6T3.5R4
 - f. *rotundifolius* (Fisch.) Topa: V (14,39)
 - f. *cornutus* Topa: IV (14,39)
 1113. *P. x cymatodes* A. et G. (*crispus x perfoliatus*): II (14)
 1114. *P. gramineus* L.: I, V (14,39,!); Hh, Cp; H6T2.5R4
 - var. *lacustris* Fr.: V (14,39)
 - var. *amphibius* Fr.: V (14,39)
 - var. *myriophilus* (A. et G.) Soó: V (40)
 - ssp. *heterophyllum* Fr.: V (14,39,40)
 - var. *riparius* Fr.: V (14,39)
 - var. *terrestris* Fr.: V (14,39)
 1115. *P. lucens* L.: III (14,38), V (14,40); Hh, Eua (M); H6T0R4
 1116. *P. natans* L.: I-V; Hh, Cosm; H6T2.5R4
 - var. *prolixus* Koch: III (38), V (40)
 1117. *P. nodosus* Poir. (*P. fluitans* Roth): V (14,40); Hh, Cp; H6T3.5R4
 1118. *P. perfoliatus* L.: II-III (14,22,39,!); Hh, Cosm; H6T0R4
 1119. *P. pusillus* L.: I (38), III-V (14,29,38); Hh, Cosm; H6T3R4

- var. major Mert. et Koch: I (38)
- var. *tenuissimus* Mert. et Koch: I, V (14,39,40)
- 1120. *P. trichoides* Cham. et Schlecht.: I (14,23); Hh, Eua (M); H6T3R4
- 1121. *Zannichellia palustris* L.: III (14,29,38), V (39); Hh, Cosm; H6T0R4
- var. *arcuata* (Schur)Topa: III (14)

NAJADACEAE

- 1122. *Najas minor* All.: III-V (14,38,40); Hh, Eua; H6T4,5R4,5
- f. *intermedia* (Balb.)Ces.: III (14)

TYPHACEAE

- 1123. *Typha angustifolia* L.: I (23,!); III-V (29,5,18,!); Hh, Cosm; H6T4R0
- f. *media* Kronf.: III (14)
- 1124. *T. latifolia* L.: I-V; Hh, Cosm; H6T3,5R0
- 1125. *T. laxmannii* Lep.: V (!); Hh, Eua; H5T4R0

SPARGANIACEAE

- 1126. *Sparganium emersum* Rehman (*S. simplex* Huds.): I (14), III (14,29), IV (25); Hh, Eua; H6T3R3,5
- 1127. *S. erectum* L. ssp. *erectum* (*S. ramosum* Huds.): I (!), V (40,18,!); Hh, Eua; H5,5T3,5R0
- 1128. *S. minimum* Hill.: IV (14); Hh, Cp; H6T3R3

LILIACEAE

- 1129. *Colchicum autumnale* L.: I (31), III-IV (14,29,43,!); G, E-M; H3,5T3R4
- 1130. *Veratrum album* L.: I-IV (6,30,22,29,25,!); G, Eua; H4T2,5R4
- ssp. *album* f. *semilobelianum* Nyár.: I (14)
- 1131. *Anthericum liliago* L.: III-IV (14,29,5); G, Ec-M; H1,5T3,5R3
- 1132. *A. ramosum* L.: I (14), III-IV (14,29,25); G, Ec-M; H2,5T3,5R4
- 1133. *Gagea lutea* (L.)Ker.-Gawl.: III-V (14,25,24,40); G, Eua; H3,5T0R3
- 1134. *G. minima* (L.)Ker-Gawl.: III (14), IV (5), V (40); G, Eua (C); H3,5T3R4
- 1135. *G. pratensis* (Pers.)Dumort.: III-V (14,25,18,40); G, Ec; H2T3R3
- 1136. *G. villosa* (M.B.)Duby (*G. arvensis* (pers.)Dumort.): III-V (14,5,25,40); G, M; H2,5T4R0
- 1137. *Allium angulosum* L.: III (14); G, Eua (C); H4,5T0R4,5
- 1138. *A. atropurpureum* W. et K.: III-V (14,29,18); G, Pn-B; H2,5T3R4
- 1139. *A. flavescens* Bess. var. *ammophilum* (Heuff.)Zahariadi: III (34), IV (14); G, P-B; H1,5T4R4,5
- 1140. *A. oleraceum* L.: IV-V (14,25,40); G, Eua; H3T3R0
- 1141. *A. scorodoprasum* L.: IV-V (14,25,18,40); G, Ec-M; H2T3R4
- 1142. *A. vincale* L.: IV-V (14,40); G, Ec-M; H2T3R4
- 1143. *Lilium bulbiferum* L.: III (14,29); G, E (M); H3T2,5R3
- 1144. *L. martagon* L.: II-V (22,14,29,43,24,!); G, Eua; H3T0R4
- 1145. *Fritillaria meleagris* L.: I (31), II (14); G, E (M); H4T3,5R4

1146. *F. montana* Hoppe: I (14); V (14); G, B-Cauc; H3T3R4
 1147. *Scilla bifolia* L.: III-V (29.5.40); G, E; H3.5T3R4
 1148. *S. autumnalis* L.: V(!); G, M; H4T3R4.5
 1149. *Ornithogalum boucheanum* (Kunth)Aschers.: III (14), V (40); G, P-B; H2.5T4R4
 1150. *O. kochii* Parl. ssp. *kochii*: V (14.40); G, P-M; H2T4R4
 1151. *O. pyramidale* L.: III (14), V (14,40); G, M; H2.5T4R4
 1152. *O. umbellatum* L.: III (34), V (47,40); G, Ec-M; H0T3.5R4
 1153. *Muscari botryoides* (L.)Mill.: I11-IV (14,29.25); G, Ec-M; H2.5T3.5R4
 1154. *M. comosus* (L.)Mill.: III-IV (14,29,18); G, Ec-M; H1.5T3.5R0
 1155. *M. racemosus* (L.)Mill.: III (14), V (14,18,40); G, Ec-M; H1.5T4R5
 1156. *M. tenuiflorum* Tausch: III-V (14,25.5); G, P-Pn; H2T4R4
 1157. *Asparagus officinalis* L.: III-V (14,29.5,18.!). G, Eu (M); H1.5T4.5R3
 1158. *Majanthemum bifolium* (L.)F.W. Schm.: I (32!), III-IV (14,29,5); G, Eu; H3T3R0
 1159. *Polygonatum latifolium* (Jacq.)Desf.: III-V (14,24.5,18.!). G, P-Pn-B; H3T3.5R4
 1160. *P. multiflorum* (L.)All.: III-V (29.5,25,24); G, E; H3T3R3
 1161. *P. odoratum* (Mill.)Druce: I (14,!); III-V (14,29,25,24,!); G, Eu (M); H2T3R4
 1162. *P. verticillatum* (L.)All.: I-II (14,30,!); G, E; H3T2.5R2.5
 1163. *Streptopus amplexifolius* (L.)DC.: I (23); G, Cp; H4T2R2
 1164. *Convallaria majalis* L.: IV-V (25.5,24); G, E; H2.5T3R3
 1165. *Paris quadrifolia* L.: I (23.30.!). IV-V (5.27); H, Eu; H3.5T0R4

AMARYLLIDACEAE

1166. *Galanthus nivalis* L.: IV-V (5.24); G, E (M); H3.5T3R4
 1167. *Leucojum aestivum* L.: V (40); G, Atl.-M; H4.5T4R4

DIOSCOREACEAE

1168. *Tamus communis* L.: IV (14.25); G, Atl-M; H3T3.5R4

IRIDACEAE

1169. *Gladiolus imbricatus* L.: I (6.!), III (29); G, Eu (C); H4T3R3
 1170. *Crocus banaticus* Gay.: II (14), IV (14,25); G, D-B; H3T3R0
 1171. *C. heuffelianus* Herb.: II (22), V (24); G, Carp-B; H3T1R2
 1172. *Iris nyaradyana* Prod.: II (14); G, E; H3T2.5R3
 1173. *I. pseudacorus* L.: III-V (14,29,25,18.!). G-Hh, E; H5.5T0R0
 1174. *I. sibirica* L.: III, V (14,40); G, Eu (C); H4.5T3R4.5
 1175. *I. spuria* L.: III (14); G, Pn-D; H4T3.5R5
 1176. *I. variegata* L.: II (22,!); III-V (14); G, P-Pn-B; H2T3.5R4

JUNCACEAE

1177. *Juncus acutifolius* Ehrh.: I (14); G, E; H4.5T3R2
 1178. *J. articulatus* L.: I (31,!); III-V (34.5.18); H, Cp; H5T2R0
 1179. *J. atratus* Krick.: I (14.32.6), III (14,34); H, Eu; H4T3R4
 1180. *J. bufonius* L.: I (23,!); III-V (47,!); Th, Cosm; H4.5T0R3
 1181. *J. bulbosus* L. var. *uliginosus* Fries: I (14); H, E; H4.5T2.5R0

1182. *J. compressus* Jacq.: I (23), IV-V (14,18); G, Eua; H4T3R4
 1183. *J. conglomeratus* L.: I-V; H, Eua; H4.5T3R3
 1184. *J. effusus* L.: I-V; H, Cosm; H4.5T3R3
 1185. *J. gerardi* Lois.: I (32), III (34,14), V (14); G, Cp; H4.5T3R5
 1186. *J. inflexus* L.: I-II (23,12!); V (18,40); H, Eua (M); H4T4R4
 1187. *J. subnodulosus* Schrank.: IV (14); Hh, E; H4.5T3.5R0
 1188. *J. tenuis* Willd.: I-III (14,23,22,!); H, Adv; H3.5T3R4
 1189. *J. thomassii* Ten.: I (14); H, D-B; H4T2.5R3
 1190. *Luzula campestris* (L.) Lam. et DC.: I (6,!); IV-V (14,5,25,!); H, E (M); H3T0R3
 1191. *L. luzuloides* (Lam.) Dandy et Willmott: I (23), II(22,!); IV (14,25,!); H, E;
 H2.5T2.5R2
 1192. *L. multiflora* (Ehrh.) Lej.: I (32); H, Cp; H3T2R2
 1193. *L. pallescens* (Wahlbg.) Bess.: I (23); H, Eua; H3T2.5R3
 1194. *L. pilosa* (L.) Willd.: II (14), IV (14,25); H, Eua; H2.5T2R0
 1195. *L. silvatica* (Huds.) Gaud.: II (14); H, Ec; H3.5T2.5R2

CYPERACEAE

1196. *Scirpus sylvaticus* L.: I-V; Hh-G, Cp; H4.5T3R0
 1197. *Eriophorum angustifolium* Honckeny: I (23,6,!); G, Cp; H4.5T3R3
 1198. *E. latifolium* Hoppe: I-II (6,23,22,!); H, Eua; H5T0R4.5
 1199. *Bolboschoenus maritimus* (L.) Palla: III-V (34,25,18,!); Hh, Cosm; H6T0R4.5
 - f. *digynus* (Godr.) Jav.: III-V (14,5)
 1200. *Schoenoplectus lacustris* (L.) Palla: II (22), IV (5,25,!); Hh-G, Cosm; H6T3R4
 1201. *S. tabernaemontani* (Gmel.) Palla: III-V (14,34,5,40,!); G (Hh), Eua; H5T3R4
 1202. *Eleocharis acicularis* (L.) R.Br.: V (14,40,!); Th, Cp; H5.5T0R0
 1203. *E. carniolica* Koch: I (31); Th, Alp-Carp-B; H5T0R5
 1204. *E. palustris* (L.) R.Br.: I-V; G-Hh, Cosm; H5T0R4
 1205. *Cyperus fuscus* L.: III-V (14,40,!); Th, Eua (M); H6T3R4
 - var. *virescens* (Hoffm.) Vahl.: III-V (14,40)
 1206. *Blysmus compressus* (L.) Panz.: I (14,23); G, Eua; H4.5T3R4.5
 1207. *Chlorocyperus glomeratus* (Torn.) Palla: IV-V (14,40); Hh, Eua (M); H5T3R4
 1208. *C. glaber* (L.) Palla: V (40); Th, Eua (M); H5T3R4.5
 1209. *Pycreus flavescens* (L.) Rchb.: III-V (14,40,!); Th, Cosm; H6T3R4
 1210. *Dichostylis micheliana* (L.) Nees: V (40); Th, Eua (M); H4.5T4R4
 1211. *Acorellus pannonicus* (Jacq.) Palla: IV (14); Th, Eua (C); H4.5T3R5
 1212. *Carex acutiformis* Ehrh.: I (31,!); IV (5,25); Hh, Eua (M); H6T3R4
 1213. *C. x alsatica* Zahn (flava x oederi): II (14)
 1214. *C. atrata* L.: I (6); H, Cp; H3T1.5R3.5
 1215. *C. bukbaumi* Wahlbg.: I, III(14,23,32); G, Cosm; H4.5T0R0
 1217. *C. brunneascens* (Pers.) Poir.: I (31); H, Cp; H4T1.5R2
 1218. *C. brizoidis* Jusl.: IV-V (25,24); H-G, Ec; H3.5T3R2
 1219. *C. appropinquata* Schumacher: I (31,14,23); Hh, Eua; H5T3R4
 1220. *C. caespitosa* L.: I (14,30,31), II (12), III (14); Hh, Eua; H5T3R3

1221. *C. canescens* L.: I-II (14,31,!); H, Cp; H5T0R2
 1222. *C. caryophyllea* Latour.: III-IV (29,5,33,25); G, Eua (M); H2T2.5R0
 1223. *C. davalliana* Sm.: I (14,32); H, Ec; H3.5T2.5R3
 1224. *C. diandra* Schrank.: I (14,31,32); G, Cp; H4T2R3
 1225. *C. digitata* L.: IV (14,5,25); H, E; H3T3R3
 1226. *C. dioica* L.: I (14,23); G, Cp; H4T2R0
 1227. *C. distans* L.: I (23), III-IV (34,26,!); H, E; H4T3R4
 1228. *C. disticha* Huds.: I (14,23); G-Hh, Eua; H5T3R4
 1229. *C. divisa* Huds.: IV-V (14,5,40); G, Eua; H4T3.5R5
 1230. *C. divulsa* Stokes.: IV-V (25,24); H, Eua; H2.5T3R0
 1231. *C. elongata* L.: I (14,23); H, Eua; H5T2.5R4
 1232. *C. elata* All.: I (14); Hh, E; H5T2.5R4
 1233. *C. flava* L.: I-II (6,31,12,!); IV (5); H, Cp; H4.5T3R0
 1234. *C. gracilis* Curtis: I-V; Hh-G, Eua; H5T3R0
 1235. *C. hirta* L.: I-V; G, E (M); H0T3R0
 1236. *C. hordeistichos* Vill.: III (14), V (40); H, Pn-P-M; H4T4R4
 1237. *C. lasiocarpa* Ehrh.: III (14); Hh, Cp; H5T2.5R2.5
 1238. *C. lepidocarpa* Tausch: I-II (14,23,12,6,!); H, E; H4.5T3R0
 1239. *C. leporina* L.: I (6,23,31,!); V (27,40,!); H, Eua; H4T2.5R3
 1240. *C. melanostachya* Willd.: V (18); Hh, Eua (C); H4T3R0
 1241. *C. michelii* Host.: III-IV (14,34,25); G, Ec-P; H2T3R4
 1242. *C. nigra* (L.)Reichard (*C. fusca* All.) ssp. *nigra*: I (31,23,6,!); G, Cp; H4T3R2
 - ssp. *juncella* Fries.: I (31)
 1243. *C. pairaei* F. Schultz: I, IV (14); H, Eua; H3T3R0
 1244. *C. pallescens* L.: I (6,23,!); IV (25); H, Cp; H3.5T3R3
 1245. *C. panicula* L.: I (14,6,32); H (G), Eua; H3.5T3R0
 1246. *C. paniculata* Jusl.: I (31,32); Hh, Ec; H5T3R5
 1247. *C. pendula* Huds.: I (31); H, Atl-M; H4T2R3
 1248. *C. pilosa* Scop.: IV (25); H, Eua; H2.5T3R3
 1249. *C. praecox* Schreb.: I-V-V (5,18); G-H, Eua; H2T3R3
 1250. *C. pseudocyperus* L.: I (31), III (14); Hh, Cp; H6T3.5R3.5
 1251. *C. remota* Grub.: V (14,40); H, E; H4.5T3R3
 1252. *C. riparia* Curt.: I-II (31,22,!); V (18,40,!); Hh, Eua (M); H5T4R4
 1253. *C. rostrata* Stokes.: I-II (14,23,23,!); Hh, CP; H5T2R0
 1254. *C. serotina* Merat (*C. oederi* Auct.): II (14); H, Eua (M); H4.5T0R0
 1255. *C. spicata* Huds.: II (40); H, Eua (M); H0T3R0
 - var. *nemorosa* (Lumn.)Serb. et Nyár.: V (14)
 1256. *C. stellulata* Good.: I (14,23,!); H, Cp; H5T2R1
 1257. *C. stenophylla* Wahlbg.: IV V (14); G, Pn; H3T0R4.5
 1258. *C. sylvatica* Huds.: IV V (5,25,40); H, E; H3.5T3R4
 1259. *C. x schatzii* Kneucker (*lepidocarpa* x *oederi*): I (14)
 1260. *C. tomentosa* L.: IV (25,40); G, Eua (M); H3T3R0
 1261. *C. x toczensis* Simk. (*melanostachya* x *riparia*): V (14,40)
 1262. *C. umbrosa* Host.: I (14); H, Ec; H3T3R3

1263. *C. vesicaria* L.: I (14,23,31,!); III-V (14,25,40,!); Hh, Cp; H6T3R4
 1264. *C. vulpina* L.: I-V; Hh H. Eua (M): H4T3R4

POACEAE (GRAMINEAE)

1265. *Botriochloa ischaemum* (L.)Keng.: III-V (29,5,25,40,!); H. Eua (M): H1.5T5R3
 1266. *Digitaria ischaemum* (Schreb.)Muhlbg.: V (14,40); Th, Cosm; H3T3.5R3
 1267. *D. sanguinalis* (L.)Scop.: III-V (14,5,25,47,!); Th, Cosm; H1.5T0R4
 1268. *Setaria glauca* (L.)P.Beauv.: III-V (18,47,!); Th, Cosm; H3T4R0
 1269. *S. verticillata* (L.)P.Beauv.: III-V (14,5,25,18,!); Th, M; H2T4R0
 1270. *S. viridis* (L.)P.Beauv.: III-V (5,25,18,47,!); Th, Eua; H2T3.5R0
 1271. *Echinochloa crus-galli* (L.)P.Beauv.: III-V (25,27,18,!); TH, Cosm; H4T0R3
 - f. *oryzoides* (Ard.)Fritsch: III (14), IV (43)
 1272. *Tragus racemosus* (L.)All.: IV-V (14,40); TH, M; H0T0R4
 1273. *Anthoxanthum odoratum* L.: I-V; H, Eua; H0T0R0
 1274. *Typhoides arundinacea* (L.)Mnch. (*Phalaris arundinacea* L.): I-V; Hh-H, Cp;
 H5T3R0
 1276. *Leersia oryzoides* (L.)Sw.: V (40); Hh, Cp; H5T3R0
 - f. *patens* Wiesb.: III (14)
 1276. *Alopecurus acqualis* Sobol.: IV (26); H, Cp; H5T3R5
 1277. *A. geniculatus* L.: I (14,31), IV-V (14,5,43,18); H, E: H5T0R4
 1278. *A. myosuroides* Huds.: III (14,29); Th, M; H3.5T3.5R4
 1279. *A. pratensis* L.: I-V; H, Eua; H4T3R0
 1280. *Phleum montanum* K. Koch: I (14), III-IV (14,6,25); H, Carp-B-Cauc-Anat;
 H1.5T4.5R4
 1281. *P. phleoides* (L.)Karsten: I (6,!); III-IV (34,5,!); H, Eua; H2T3R4
 1282. *P. pratense* L.: I-V; H, Eua (M): H3.5T0R0
 1283. *Crypsis aculeata* (L.)Ait.: V (18,40); Th, Eua; H3.5T4R4
 1284. *Helcochloa alopecuroides* (Piller et Mitterp.)Host.: V (!); Th, Eua; H0T4R4.5
 - f. *angustifolia* Beck: V (40)
 1285. *H. schoenoides* (L.)Host.: III (14), V (14,40); Th, Eua; H0T4R4.5
 1286. *Beckmannia cruciformis* (L.)Host.: III (34), V (14,!); H, Cp; H4.5T3R4
 1287. *Cynodon dactylon* (L.)Pers.: III-V (5,25,18,!); G (H), Cosm; H2T3.5R0
 1288. *Agrostis canina* L.: I (31,!); IV-V (25,40); H, Eua; H3.5T3R3
 - var. *pudica* Doll.: I (14)
 1289. *A. stolonifera* L. ssp. *stolonifera*: I-V; H, Cp; H4T0R0
 - ssp. *gigantea* (Roth)Beldie var. *silvatica* (Host.)Beldie: V (40)
 1290. *A. tenuis* Sibth.: I-V; H, Cp; H0T0R0
 1291. *Apera spica-venti* (L.)P. Beauv.: IV (5,25); TH, Eua; H3.5T0R2.5
 1292. *Calamagrostis arundinacea* (L.)Roth: I (30); H (G), Eua; H2.5T3R2
 1293. *C. canescens* (Web.)Druce: I (14,23,6); H, Eua; H5T3R3
 1294. *C. epigeios* (L.)Roth: III-IV (34,25,5); H (G), Eua (M); H2T3R0
 1295. *C. neglecta* (Ehrh.)Gaertn.: I (14,23,6,31); H, Cp; H4.5T2R3
 - var. *fallax* Bancr: I (14)
 1296. *C. pseudophragmites* (Haller f.)Koeler: II (22), IV-V (25,40); H, Eua (C); H5T3R5

1297. *C. villosa* (Chaix) J.F.Gmel. var. *graciliscescens* (Blytt.) A. et G.: I (14,23); H, Eua; H4T2.5R1.5
1298. *Milium effusum* L.: V (24); H, Cp; H3.5T3R3
1299. *Stipa capillata* L.: III-IV (4,5,25); H, Eua (C); H1T5R4
1300. *S. pulcherrima* K.Koch: III (4); H, Eua (M); H1T4R5
1301. *Phragmites australis* (Cav.)Trin. et Steud. (*Ph. communis* Trin.):I-V; Hh. Cosm; H5T0R4
 - var. *flavescens* Custer: III-IV (14,40)
1302. *Sesleria heufleriana* Schur: IV (6); H, Carp (End); H2T3.5R4.5
1303. *Koeleria cristata* (L.)Pers. (*K. macrantha* (Ldb.)J.A. et J.H. Schult): I-V; H, Cp; H2T4R5
1304. *Melica nutans* L.: I (30), IV (5); H-G, Eua (M); H3T0R4
1305. *Holcus lanatus* L.: I-V; H, Eua; H3.5T3R0
1306. *Deschampsia cespitosa* (L.)P.Beaup: I-V; H, Cosm: H4T0R0
1307. *D. flexuosa* (L.)Trin.: I-II (30,22,!); H, Cp; H2T0R1
1308. *Arrhenatherum elatius* (L.)J. et C. Presl: II-V (22,28,25,5,40,!); H, E (M); H3T3R4
1309. *Avena fatua* L. ssp. *fatua*: III-V (14,40,!); Th, Eua (M); H3.5T0R4
 - var. *glabrata* Peterm.: V (14)
1310. *Helictotrichon pubescens* (Huds.)Pilger: IV (14,5); H, Eua; H3.5T2.5R4
1311. *H. alpinum* (Smith)Heward: I (14), III-IV (14,5); H, Carp; H2T2R3
1312. *H. pratense* (L.)Pilger: I (14,6); H, Eua; H2.5T3R0
1313. *Sieglungia decumbens* (L.)Bernh.: I (6,!); III (34); H, E; H0T3R2
1314. *Bromus arvensis* L.: II-V (22,34,25,40,!); Th-TH, Eua (M); H2.5T3R0
1315. *B. commutatus* Schrad.: IV-V (5,25,18,40); Th, E; H0T3R0
1316. *B. incermis* Leyss.: III-V (34,5,25,40,!); H, Eua (C); H2.5T4R4
1317. *B. japonicus* Thunb.: I (!); Th, Eua (M); H1.5T3.5R4
 - var. *transsilvanicus* (Auersw.)Hay : I (14), V (40)
 - var. *vestitus* (Schrad.)Stapf.: V (40)
1318. *B. mollis* L.: II-V (22,5,18,47,!); Th, Eua; H0T3R0
 - var. *mollis* f. *nanus* (Weig.)Nyár.: III (14)
 - var. *ciliatus* Schur: III (14)
1319. *B. ramosus* Huds.: II (22); H, Ec; H3T3R3
1320. *B. secalinus* L.: I, III (14), V (40); Th, Eua (M); H0T0R0
 - f. *submuticus* (Rehb.)Nyár.: III (14)
1321. *B. sterilis* L.: IV-V (25,18,!); Th, Eua (M); H2T4R4
1322. *Cynosurus cristatus* L.: I-V; H, E; H3T3R3
1323. *Brachypodium silvaticum* (Huds.)P.Beaup: III-V (25,40,!); H, Eua (M); H3T3R4
1324. *Eragrostis chilianensis* (All.)Link.: IV-V (14,6,40); Th, Eua; H2T4R4.5
1325. *E. poiooides* P.Beaup.: V (18,47,!); Th, Cosm; H3T4R0
1326. *Molinia coerulea* (L.)Mnch. ssp. *coerulea*: I (6,31,!); H, Eua; H4T3R0
 - ssp. *arundinacea* (Schrank.)Paul: I (14)
1327. *Dactylis glomerata* L.: I-V; H, Eua (M); H3T0R4
 - var. *pendula* Dumort: III (14)
1328. *Poa annua* L.: I-V; Th-TH, Cosm; H3.5T0R0

1329. *P. bulbosa* L.: IV-V (25,18,47); G-H, Eua (C); H2T3.5R4
 1330. *P. compressa* L.: II (22,!); IV (5,25); H, E: H1.5T3R0
 1331. *P. nemoralis* L.: II-V (22,5,25,40,!); H, Eua; H3T3R0
 1332. *P. palustris* L.: I (31,6,!); IV-V (25,40); H, Cp; H5T3R4
 1333. *P. pratensis* L. ssp. *pratensis*: I-V; H, Cp; H3T0R0
 - ssp. *angustifolia* (L.)Hay (*P. angustifolia* L.): II-IV (22,33,25); H, Eua; H2T3R0
 - var. *sctacca* (Hoffm.)Doll: I (14)
 1334. *P. stiriaca* Fritsch et Hay: I (14); H, E: H1.5T2.5R4
 1335. *P. trivialis* L.: I-V; H, Eua; H4T0R0
 1336. *Briza media* L.: I-V; H, Eua; H0T3R0
 1337. *Catabrosa aquatica* (L.)P.Beaup.: I-II (22,!); H, Cp; H5T2.5R4
 1338. *Glyceria fluitans* (L.)R.Br.: I-V; Hh-H, Cosm; H5T3R0
 1339. *G. maxima* (Hartm.)Holmberg: I-V; Hh-H, Cp; H5T3R4
 1340. *G. nemoralis* (Uccehr.)Uccehr. et Koern.: IV (14); Hh, Ec-Sarm:H5T3R3
 1341. *G. plicata* Fries.: I-II (23,31,22,!); Hh, Eua; H6T3R4.5
 1342. *Puccinellia distans* (L.)Parl.: III-V (34,5,!); H, Eua (C); H3.5T0R5
 1343. *P. limosa* (Schur)Holmberg: III-IV (14); H, Eua (C); H3.5T0R5
 1344. *Vulpia myuros* (L.)Gmel.: IV-V (14,25,27); Th, Eua (M); H1T3.5R2
 1345. *Festuca arundinacea* Schreb.: II (22), V (40); H, Ec; H4T3R4
 - var. *mediteranca* (Hack.)A. et G. f. *baltica* A. et G.: V (14);
 - var. *orientalis* (Kern.)A. et G.: II (14)
 1346. *F. drymeia* Mert. et Koch: II (14,22,!); IV (14,25,43); G-H, Carp-B; H4T2R3
 1347. *F. gigantea* (L.)Vill.: II (22), V (24,40); H, Eua; H4T3R2.5
 - f. *nemoralis* A. et G.: IV (14,43)
 1348. *F. pratensis* Huds.: I-V; H, Eua; H3.5T0R0
 - var. *subspicata* (G.F.W. Meyer)A. et G.: II (14)
 1349. *F. pseudovina* Hack.: III-V (34,5,!); H, Eua (C); H2T4R4
 1350. *F. rubra* L.: I-V; H, Cp; H3T0R0
 1351. *F. rupicola* Heuff. ssp. *rupicola*: II-V (22,33,5,25,40,!); H, Eua (C); H1.5T4R4
 - var. *rupicola* f. *hirsuta* Host.: III (14), V (40)
 - var. *sulcataeformis* Mgß-Dbg.: II (14)
 1352. *Pholiurus pannonicus* (Host.)Trin.: V (14,40); Th, P-Pn-B; H0T4R4.5
 1353. *Lolium multiflorum* Lam.: III-V (14,25,40); Th-TH, Adv (M); H2.5T4R0
 1354. *L. perenne* L.: I-V; H, Eua (M); H2.5T4R4.5
 - var. *tenue* (L.)Sm.: III (14)
 1355. *L. remotum* Schrank.: I (14); Th, Adv; H2T3R4
 1356. *L. temulentum* L.: IV (14,5); Th-TH, Adv; H2.5T4R4.5
 1357. *Hordelymus europacus* (L.)Jessen: II (22); H, E; H3.5T3R3
 1358. *Hordeum marinum* Huds.: V (14,18); Th, Atl-M; H2T4R3
 - ssp. *gussonianum* (Parl.)A. et G.: III (14), IV (14,25), V (14,!); Th, Eua; H2T4R4.5
 - f. *hirtellum* Degen: V (40)
 1359. *H. murinum* L.: III-V (5,25,18,!); Th, Eua (M); H2.5T4R0
 1360. *Agropyron caninum* (L.)P.Beaup.: IV-V (25,40); H, Cp; H3.5T0R4
 - var. *pauciflorum* (Schur)Volkart: V (40)

1361. *A. intermedium* (Host.)P.Beauv.: III-V (5,18,!); G, Eua; H2T4R4
 1362. *A. pectiniforme* Roem. et Schult.: V (40,!); H, Eua; H2T4R4.5
 - f. *ciliatum* Degen: V (40)
 1363. *A. repens* (L.)P.Beauv.: I-V; G, Eua; H0T0R0
 - var. *glaucum* (Host.)Doll: V (40)
 - var. *aristatum* (Neilr.)Hay: V (40)
 1364. *Nardus stricta* L.: I (6,31,!); IV (5); H, E; H0T0R1.5

ORCHIDACEAE

1365. *Orchis cordigera* Fries.: I-II (14,!); G, Alp-Carp-B; H4.5T2R2
 - var. *cordigera* f. *macrobracteata* Schur: II (14)
 - var. *siculorum* (Soó)Pauca et Beldie: I-II (14)
 1366. *O. laxiflora* Lam.: III-V (5,!); G, Eua (M); H4T3R0
 - ssp. *elegans* (Heuff.)Soó: III-V (34.29.5,!);
 - var. *javorkae* Soó: III (14)
 1367. *O. incarnata* L.: I (14,32,!); III-IV (14,!); G, Eua (M); H4.5T0R4
 1368. *O. maculata* L. ssp. *maculata*: I (6,31,!); IV (25,!); G, Eua (M); H0T0R0
 - var. *transsilvanica* (Schur)Doin: I (14,!);
 - ssp. *elodes* (Gris.)Camus var. *schurii* (Kingo)Pauca et Beldie: I (14,42)
 - ssp. *fuchsii* (Druce)Christens f. *karpati* Borsos: I (14)
 1369. *O. militaris* L.: III-IV (14); G, Eua; H3T3R4
 1370. *O. morio* L.: III-IV (34.29.5,!); G, Ec; H2.5T3R4
 1371. *O. sambucina* L.: II (14.22), IV (14); G, Ec; H3T2R3
 1372. *Traunsteinera globosa* (L.)Spreng.: II (22); G, Ec; H3T2R4.5
 1373. *Gymnadenia conopsea* (L.)R.Br.: I (6,!); III-V (14.6); G, Eua; H4T0R4.5
 1374. *Platanthera bifolia* (L.)L.C.Rich.: I (6,23,!); III-V (29.5.25,!); G, Eua (M);
 H3.5T0R3
 1376. *P. chlorantha* (Cust.)Rchb.: I, V (14); G, Eua (M); H3.5T3R3
 1376. *Listera ovata* (L.)R.Br.: I (32), IV (5), V (27); G, Eua (M); H3.5T0R4
 - f. *multinervia* (Peterm.)Hegi: III (14)
 1377. *Neottia nidus-avis* (L.)L.C.Rich.: IV-V (25.5.40); G, Eua (M); H3.5T3R3
 1378. *Cephalanthera longifolia* (Huds.)Fritsch.: II (22.14); G, E; H2.5T3R4
 1379. *Epipactis helleborine* (L.)Cr.: III-V (14.25.24.40,!); G, Eua; H3T3R3
 1380. *E. palustris* (L.)Cr.: I-II (14.6.23,!); G, Eua; H4.5T3R4.5

ARACEAE

1381. *Acorus calamus* L.: IV-V (14); Hh (G), Adv; H6T3.5R4
 1382. *Calla palustris* L.: I-II (14.22); Hh, Cp; H5T2.5R3.5
 1383. *Arum maculatum* L.: IV-V (5.18); G, Ec; H3.5T3.5R4
 - var. *alpinum* (Schott et Kotschy)Topa et Beldie: IV-V (14.40)

LEMNACEAE

1384. *Lemna gibba* L.: III-V (14.25.40); Hh, Cosm; H6T3.5R4
 1385. *L. minor* L.: I-V; Hh, Cosm; H6T0R0

1386. *L. trisulca* L.: III-V (25,38,40,!); Hh, Cosm; H6T0R4

1387. *Spirodela polyrrhiza* (L.)Schleichen: I (31), V (38,!); Hh, Cosm; H6T3.5R0

Study of the ecological preferences

Analysing the humidity (H) needs of the plant species from the Mureş Valley, we see that most of the plants belong to the mesophytic category (H3-H3.5). Forming 32.8% of the total flora list, they are here favoured by the moderate humidity both of the soil and from the atmosphere. The considerable representation (29.3%) of the mesohygrophytes (H4-H4.5), hygrophytes (H5-H5.5) and hydrophytes (H6) may be explained by the existence of several marshes, lakes and oxbow lakes of the Mureş. There are also xeromesophilous species (H2-H2.5) and xerophilous (H1-H1.5) ones in the flora, comprising 28.7% and 4.9%, respectively, which inhabit some rocky regions and sunny slopes characterized by water deficit.

Concerning the temperature factor (T) the richness of the micromesothermic species (T3-T3.5) holding for 56.6% can be noticed as the longest part of the waterway of Mureş is situated in a moderate climate with an annual mean temperature of about 9-10 ° Celsius. The microthermic species (T2-T2.5) are also well represented (10.9%), being favoured by the mountainous relief of the straits and particularly by the specific climate of the Depression of Giurgeu (Gheorgheni) which is one of the coldest poles of Transylvania with an annual mean temperature of only 4-5 ° Celsius. Both mesothermophilous (T4-T4.5) and thermophilous (T5) species are present (16.0%): their occurrence is connected to south-facing slopes from the pass Toplija- Deda, the Transylvanian Plateau and the Mureş Corridor.

In soil reaction (R), we notice the abundance of the weaker acid-neutrophilous (R4-R4.5), curvionic (R0) and acid-neutrophilous (R3-R3.5) species, in percentages: 40.3%, 28.5% and 20.6%, respectively. The acidophilous species (R1-R1.5) occur with a 6.5 % frequency, while the neutro-basiphilous ones (R5) represent a portion of 3.9 %. The above distribution reflects sufficiently well the preponderance of alluvial deposits and soils from the river and also the existence of the following soil types: brown podsolic, clayey podsoled, acid brown, peat and salty soils.

Life form composition

The analysis of life forms shows some characteristics of the biotopes and the influences exerted on them by different factors. The very high percentage of hemicryptophytes (42.7 %) is closely connected to the large surfaces of lawn and the presence of a grass layer in riverside coppices and woods. The therophytes (30.7 %) are the manifestation of the warmer climate from the center of Transylvania and the plain of Mureş, plus a pronounced anthropogenic influence on these areas. The helohydrophytes (7.2%) point out the lakes and the wet meadows of the riverside. The ratio between the number of therophytes and that of hemicryptophytes gives the anthropisation (altitude) coefficient

($K_a = T/H + Ch * 100$) (Pop & Drăgușescu 1983), by means of which it is possible to determine the level of degradation for an area. For the Mureș Valley its value is 72, which corresponds to a territory with considerable landscape disturbance.

Analysis of the floristic elements

The flora of the area is characterized by the predominance of eurasian (45.2%), European (13.1%), circumpolar (9.1%) and Central-European (8.3%) elements, respectively. The main components of the mesophilous and mesohydrophilous lawns, and of the riverside coppices and woods, are the temperate-continental climate and the geographical position of the region. The meridional elements (Mediterranean, Submediterranean, Mediterranean-Pontic, Pontic, Ponto-Pannonic and Balkanic) altogether represent 10.1%. They indicate some biotopes with a warmer microclimate from the Transylvanian Plateau and the neighborhood of the lower basin of Mureș upon the Pannonian province. The Daco-Balkanic plus Carpatho-Balkanic (2.1%) and endemic or Carpathic (0.8%) elements provide a peculiar color to the valley.

The vegetation

Relying on our phytocoenologic relevés, we identified 174 associations with 40 subassociations and facies. Until our research in the area, only 81 have been previously mentioned.

Associations

ASPLENIETEA RUPESTRIS H. Meier et Br.-Bl. 1934

ASPLENIETALIA SEPTENTRIONALIS Oberd. et al. 1967

Asplenion septentrionalis Gams 1927

1. *Asplenio trichomanes-Poaetum nemoralis* Boșcaiu (1970)1971: II (!)

2. *Sempervivetum heuffelii* E.Schneider-B. 1969: II (!)

3. *Hypno-Polypodictum vulgare* Jko. et Pec. 1963: IV (43)

LEMNETEA W. Koch et Tx. 1954

LEMNETALIA W. Koch et Tx. 1954

Lemnion minoris W. Koch et Tx. 1954

4. *Lemnetum minoris* (Oberd. 1957) Muller et Gors 1960: I-V

5. *Spirodeletum polyyrrhizae* W. Koch 1954: V (!)

6. *Lemno-Salvinietum natantis* Miyawaki et Tx. 1960: V (!)

7. *Salvinio-Spirodeletum polyyrrhizae* Slavnic 1956: V (!)

Utricularion vulgaris Pass. 1964

8. *Lemno-Utricularietum* Soó 1928: III (!)

- *lemnitetum trisulcae* (Kárpáti 1963) Soó 1964: III (!)

- *ceratophylletosum demersi* Soó (1957) 1964: III (!)

HYDROCHARIETALIA Rubel 1933

Hydrocharition Rubel 1933

9. Ceratophyllico-Hydrocharicum I.Pop 1962: III, V (!)
Ceratophyllum Den Hartog et Segal 1964
10. Ceratophylletum demersi (Soó 1927) Hild 1956: III (!)
 POTAMETEA Tx. et Prsg. 1942
- POTAMETALIA W. Koch 1926
Ranunculion aquatilis Pass. 1964
11. Ranunculetum (Batrachictum) trichophylli Soó (1927) 1971: V (!)
Potamion W. Koch 1926 emend. Oberd. 1957
12. Myriophyllo-Potametum Soó 1934: V (!)
13. Potametum crispi Soó 1927: II-III (!), IV (32a)
 - potametosum pusilli Soó (1927) 1973: V (!)
Nymphaeion Oberd. 1957 emend. Neuhausl 1959
14. Nymphacetum albo-luteac Nowinski 1928: III, V (!)
 - nymphacetosum V. Kárpáti 1963: V (!)
 - nupharetosum Soó (1957) 1964: III (!)
15. Polygonetum natantis Soó 1927: III, V (!)
16. Potametum natantis Soó 1927, Eggler 1933: III, V (!)
17. Nymphoidetum peltatae (Allorge 1922) Oberd. et Muller 1960: V (!)
18. Trapetum natantis Muller et Gors 1960: IV (26,43), V (!)
- PHRAGMITETEA Tx. et Prsg. 1942
- PHRAGMITETALIA W. Koch 1926 emend. Pign. 1953
Phragmition australis (communis) W. Koch 1926 emend. Soó 1947
19. Scripo-Phragmitetum W. Koch 1926: I-V
 - phragmitetosum Soó 1957: IV V (25,43. !)
 - butomosum Paun (1964) 1967: III (!)
 - hydrocharitosum I. Pop 1962: III, V (!)
 - solanetosum dulcamarae Krausch 1965: V (I)
20. Typhaetum angustifoliac (All. 1922) Pign. 1943: I, III, V (I)
21. Typhaetum latifoliac Soó 1927: I-III, V (!)
22. Typhaetum laxmannii (Ubrizsy 1961) Nedelcu 1968: V (!)
23. Schoenoplectetum lacustris Eggler 1933: I, V (!)
24. Glycerietum maximae Hueck 1931: I-V
25. Oenanthesetum aquaticaе Soó 1927. Eggler 1933: III, V (!)
Bolboschoenion maritimi Soó (1945) 1947
26. Bolboschoenetum maritimi Soó (1927) 1957: IV (6a, 32a, 25, 43), V (!)
 - schoenoplectetosum tabernaemontani Soó 1957: V (!)
27. Schoenoplectetum tabernaemontani Soó (1927) 1949: V (!)
28. Elcocharietum palustris Schennikow 1919: Soó 1933: I, III (!)
- NASTURTIO-GLYCERIETALIA Pign. 1953
Glycerio Sparganion Br.-Bl. et Sising ex Boer 1942
29. Sparganio-Glycerietum fluitantis Br.-Bl. 1925: IV (25,43), V (I)
30. Glycerietum plicatae Oberd. (1952) 1957: I (!)
31. Alismato-Elcocharitetum Máthé et Kovács 1967: III, V (I)
Phalarido Glycerion Pass. 1964

32. *Equisetum fluviatilis* (limosi) Soó (1927)1947: I (31)
 33. *Leersietum oryzoidis* Krause 1955 em. Pass.1957: I (49)
MAGNOCARICETALIA Pign.1953
Magnocaricion clatae W. Koch 1926
 34. *Caricetum rostratae* Rubel 1912: I (31..)
 35. *Caricetum paniculatae* Wangerin 1926. Soó 1969: I (31..)
 36. *Caricetum appropinquatae* (W. Koch 1926) Tx.1947: I (31)
 37. *Carici-Menyanthetum* Soó (1938)1955: I (32). II (?)
 38. *Caricetum gracilis* Almquist 1929; Tx. 1937: I (31..), II (12)
 39. *Caricetum acutiformis* Suer 1937: I (31..), III (4), IV (6a)
 40. *Caricetum ripariae* Soó 1928: I (31..)
 41. *Caricetum vesicariae* Br.-Bl. et Denis 1926; Zólyomi 1931: I (31..). IV-V (25,43..)
 42. *Caricetum vulpinac* Soó 1927: I (31), IV (6a)
 43. *Poetum palustris* Resmerită et Rađiu 1974: I (31)
 44. *Typhoidetum arundinaceae* Eggler 1933: I-III (31.4..), IV (32a), V (?)
ISOETO-NANOJUNCETEA Br.-Bl. et Tx. 1943
NANOCYPERETALIA Klika 1935
Nanocyperion flavescentis W. Koch 1926
 45. *Pucrictum* (Cyperetum) flavescenti-fusci W. Koch 1926 em. Philippi 1968: V(!)
 46. *Cypero-Juncetum* Soó et Csűrös 1944: III (9)
 47. *Juncetum bufonii* Morariu 1956; Philippi 1968: I, III, V (?)
Verbenion supinac Slavino 1951
 48. *Lythreto-Pulicarictum vulgaris* Tímár 1954: V (?)
SCHEUCHZERIO-CARICETEA NIGRAE (FUSCAE) Nordh.1936
SCHEUCHZERIO-CARICETALIA NIGRAE (FUSCAE) (WKoch1926)Gors et Muller ex Oberd.1967
Caricion lasiocarpae Van den Bergen 1949
 49. *Caricetum diandrac* (Jon.1932)Oberd.1957: I (32)
Caricion canescens nigrare (fuscae) (W. Koch 1926)Nrdh. 1936
 50. *Carici-Agrostietum caninae* Tx. 1937: I (32)
 51. *Caricetum nigrac* (fuscac) Br.-Bl.1915; W. Koch 1928: I (32..)
 52. *Carici stellulatae* (echinatae)-*Sphagnetum* Soó (1934)1954: I (32..)
 53. *Caricetum stellulatae* (echinatae) Csűrös 1956: III (?)
TOFIELDIETALIA Prsg. apud Oberd. 1949
Eriophorion latifolii Br.-Bl. et Tx. 1943
 54. *Carici flavae-Eriophoretum* Soó 1944: I-II (32.12..)
 55. *Valeriano-Caricetum flavae* Pawl., Pawłowska et Zazycki 1960: I (32)
MOLINIO-ARRHENATHERETEA Tx. 1937
CARICETALIA DAVALLIANAE Br.-Bl.1949
Caricion davallianae Klika 1934
 56. *Caricetum davallianae* W. Koch 1928: I (49)
MOLINIETALIA W. Koch 1926
Agrostion stoloniferae (albac) Soó (1933)1971
 57. *Agrostidetum stoloniferae* (Ujvárosi 1941) Burduja et all.1956: I-V

- clecocharctosum Soó 1964: III (!)
- 58. Poctum pratensis Rav., Cazac. et Turenschi 1956: III (!), IV-V (25.43, !)
- lolietosum perennis Grigore 1971: V (!)
- 59. Alopecuretum pratensis Nowinski 1928: IV (6a)
- 60. Agrostideto-Festucetum pratensis Soó 1949: III (!), IV-V (6a, 11a, 32a, !)
- 61. Lolietum perennis Safta 1943: III, V (!)
- 62. Lythro-Calnagrostidetum epigci I. Pop 1968: IV (!)
- Molinion coeruleae* W. Koch 1926
- 63. Molinietum coeruleae (All. 1922) W. Koch 1926: I (6, !)
- caricetosum nigrac (fuscac) Borza et F. Rajiu 1970: I (6, !)
- 64. Junco-Molinietum Prsg. 1910: I (6, !)
- nardetosum strictae (Jon. 1933) Kovács 1956: I (6)
- Calthion palustris* Tx. 1937
- 65. Calthactum lactae Krajina 1933: I (!)
- 66. Scirpetum sylvatici Schwick. 1944: I-V
- 67. Cirsictum cani Tx. 1951: I, III (!)
- 68. Trollio-Cirsictum (Kuhn 1937) Oberd. 1957: I (!)
- Filipendulo-Petasition* Br.-Bl. 1947
- 69. Petasitum hybridi (Dost. 1933) Soó 1940: II (!)
- 70. Filipendulo-Geranietum palustris W. Koch 1926: I (!)
- 71. Filipenduletum ulmariae W. Koch 1926: III-IV (!)
- 72. Chacrophylo-Equisetum palustre Vicol et Stoicovici 1977: I-II (!)
- ARRHENATHERETALIA Pawl. 1928
- Arrhenatherion elatioris* (Br.-Bl. 1925) W. Koch 1926
- 73. Arrhenatheretum elatioris (Br.-Bl. 1919 s.l.) Scherrer 1925; Soó 1969: III (28, !), IV (6a), V (11a)
- holcetosum Csürös 1970: III (11, 28)
- trisetosum flavescentis Horvatic 1930: III (11, 28)
- festucetosum rupicolae (sulcatae) Egller 1958: III (28)
- pastinacetosum comb. nova (Pastinaco-Arrhenatheretum elatioris) (Knapp 1954) Pass. 1964: II-III (!)
- geranictosum pratensis subass. nova: II-III (!)
- 74. Poaeto-Festucetum pratensis Soó 1949: III (28, !)
- Cynosurion cristati* Br.-Bl. et Tx. 1943
- 75. Anthoxantho-Agrostictum tenuis Sillinger 1933: Jurko 1969: IV (!)
- 76. Agrostideto-Festucetum rupicolae (sulcatae) Cs.-Káptalan (1962) 1964: III-IV (28, 25, 43, !)
- 77. Agrostideto-Danthonietum Soó 1947: III (28)
- 78. Agrosti-Cynosuretum Resmerita 1963: II-III (28, I)
- 79. Agrosti-Festuccetum rubrac Horv. (1951) 1952: I-II (!)
- 80. Lolio-Cynosuretum Tx. 1937: IV (6a, 43)
- DESCHAMPSIETALIA CAESPITOSAE Horvatic 1956
- Alopecurion pratensis* Soó 1938; Pass. 1946
- 81. Festucetum pratensis Soó 1938: III-IV (28, !)

- Deschampsion caespitosae* (Horvatic 1930) Soó 1971
82. Agrostideto-Deschampsietum caespitosae Ujvárosi 1947: I-II (!)
 - caricosum nigrae (fuscae) Stoicovici 1977: I (!)
- PUCCINELLO-SALICORNIETEA Topa 1939
- SALICORNIETALIA Br.-Bl. (1928) 1933
- Thero-Salicornion* Br.-Bl. (1930) 1933, Pign. 1933
83. Salicornietum europaeum Wendelbg. 1953: III (!)
- PUCCINELLIETALIA Soó 1940
- Puccinellion limosae* (Klika 1937) Wendelbg. 1943, 1950
84. Staticeto-Artemisiectum monogynac Topa 1939: III, V (!)
85. Hordeetum hystricis (Soó 1933) Wendelbg. 1943: V (!)
- Puccinellion pisonis* (Wendelbg. 1943) Soó 1957
86. Puccinellietum distantis Soó 1937, Knapp 1948: III, V (!)
- Juncion gerardii* Wendelbg. 1943
87. Juncetum gerardii (Warming 1906) Nordh. 1923; Wenzl 1934: III (4!), V (!)
88. Agrostio-Caricetum distantis (Rapaics 1927) Soó 1930: III (!), IV (6a), V (11a)
- Festucion pseudovinae* Soó 1933
89. Artemisio-Festucetum pseudovinae (Magyar 1928) Soó (1933) 1945: V (!)
- puccinellietosum Soó 1964: V (!)
- limonietosum Bodrogk. 1962: V (!)
90. Achilleo-Festucetum pseudovinae (Magyar 1928) Soó (1933) 1945: V IV (6a), V (11a, !)
- cynodontetosum Borza 1959; Bodrogk. 1965: V (11a, !)
- Beckmannion cruciformis* Soó 1933
91. Agrostio-Beckmannietum (Rapaics 1916) Soó 1933: V (!)
- FESTUCO-BROMETEA Br.-Bl. et Tx. 1943
- BROMETALIA ERECTI (Koch 1926 n.n.) Br.-Bl. 1936
- Bromion erecti* (Koch 1926 n.n.) Br.-Bl. 1936 s.str.
92. Brometum erecti Pázmány 1963: III (33)
- Cirsio-Brachypodion* Hadac et Klika 1944 emend. Krausch 1961
93. Cariceto humilis-Brachypodium pinnatum Soó (1942) 1947: III (4)
- globularietosum Borza et Lupșa 1962: III (4)
- salvietosum transsilvanicae Borza 1959: III (4)
- Festucion rupicolae (suicatae)* Soó (1929 n.n.) 1940 corr. Soó 1964
94. Cynodontio-Poetum angustifoliae (Rapaics 1926) Soó 1957: III (28!), V (11a)
95. Festucetum rupicolae Burduja et al. 1956: II-III (28!), V (11a)
96. Festuceto-Botriochloctum ischacmii Resmeriță 1965: III-IV (28!), V (11a)
- SECALIETEA Br.-Bl. 1931
- APERETALIA R. et J. Tx. 1960
- Aphanion* J. et R. Tx. 1960
97. Echo-Rumicetum acetosellae Soran 1962: III (!)
- SECALIETALIA Br.-Bl. 1931 emend. J. et R. Tx. 1960
- Caucalidion platycarpus* Tx. 1950 corr. Soó 1971
98. Consolido-Polygonetum convolvulus (Morariu 1943) Morariu 1967: III(4!), V (47)

- Trifolio Medicaginon sativae* Balázs 1944 emend. Soó 1959
99. *Plantagini lancolatae-Medicaginectum* (Balázs 1944) Soó et Tímár 1964: V(47)
- *verbenetosum officinale* Tímár 1953: V (!)
- ERAGROSTETALIA** J. Tx. 1961 emend. Soó 1968
- Consolido Eragrostion poidis* Soó et Tímár 1957
100. *Amarantho-Chenopodietum albi* (Morariu 1943) Soó (1947)1953: II (!). III (4.!). V (47,!)
- Tribulo Eragrostion poidis* Soó et Tímár 1957
101. *Hibisco-Eragrostetum poaeoidis* Soó et Tímár (1951)1957: V (!)
102. *Digitario-Portulacetum* (Felföldy 1942) Tímár et Bodrogk. (1943)1955: V (!)
- CHENOPODIETEA** Br.-Bl. 1951 emend. Lohn., J. et R. Tx. 1961
- SISYMBRIETALIA** J. Tx. 1961
- Sisymbrium officinalis* Tx., Lohm. et Prsg. 1950
103. *Hordeetum murini* Libbert 1932 em. Pass. 1964: III (!)
104. *Atriplicetum nitentis* Knapp 1945: III, V (!)
105. *Atriplicetum tataricac* (Prodan 1923) Borza 1926: III (!)
106. *Polygono avicularis-Amaranthetum crispī* Vicol. Schneider-B. et Täuber 1971: III,IV (!)
107. *Malvetum neglectae* Aichinger 1933 em. ass. 1966: I, III-IV (!)
108. *Descurainetum sophiae* Krecz 1935 corr. Oberd. 1970: III (!)
109. *Malvetum pusillae* Morariu 1943: III (!)
110. *Xanthio spinosae-Amaranthetum* Morariu 1943: III (!)
- Convolvulo arvensi Agropyrion repens* Gors 1966
111. *Agorpyretum repens* Felföldy 1942: III-IV (!)
- *convolvuletosum arvensis* Grigore 1971: III (!)
- Veronica Euphorbion* Siss. 1942
112. *Galeopsidetum speciosae* Krusem. et Vlieg. 1939: III (!)
- ONOPORDETALIA** Br.-Bl. et Tx. 1943 emend. Gors 1966
- Dauco Meliloton* Gors 1966
113. *Echio-Melilotetum albi* Tx. 1942: III-IV (4.!).
- Onopordion acanthi* Br.-Bl. 1926 s.str.
114. *Onopordetum acanthi* Br.-Bl. (1923)1936: III (4.!).
115. *Carductum acanthoidis* (Allorge 1922) Morariu 1939: II-III (!)
- POLYGONO-CHENOPODIETALIA** (Tx. et Lohm. 1950) J. Tx. 1961
- Panico-Setarion* Siss. 1946
116. *Setario-Galinsogetum* Tx. 1950: III-V (!)
- ARTEMISIETEA** Lohm., Prsg. et Tx. 1950
- ARTEMISIETALIA** Lohm. et Tx. 1947
- Arction lappae* Tx. 1937 emend. Siss. 1946
117. *Artemisiectum annuum* Morariu 1943 emend. Dihoru 1970: III (!)
118. *Tanaceto-Artemisiectum vulgaris* Br.-Bl. (1931)1949: I, III,V (4.!).
- *pastinacetosum* Szabó 1971: III (!)
119. *Coniectum maculati* I. Pop 1968: III, V (!)

120. Sambucetum cebuli (Kaiser 1926) Felsöldy 1942: III-IV (4,25,!)
 121. Arctio-Ballotetum nigrac (Felsöldy 1942) Morariu 1943: I (!), III (4,!), V (!)
 122. Urticetum dioicae Steien 1931; Turenschi 1966: I-V
 123. Potentillo-Artemisiectum absinthii (Prodan 1948) Falinski 1964: II-IV (!)
 124. Lycictum barbarum Felsöldy 1942 corr. Soó 1971: III, V (!)
Tussilaginion Szabó 1971 n.n.
 125. Tussilaginetum sarsarac Oberd. 1949: II, IV (!)
CALYSTEGIETALIA (CONVOLVULETALIA) SEPIUM Tx. 1950
Calystegion sepium Tx. 1947 ex Oberd. 1949
 126. Rudbeckio-Brachypodium silvaticae Szabó 1970: III (?)
 127. Glycyrrhizetum echinatae (Soó 1940 n.n.; Timár 1947) Slavnic 1951: V (!)
 128. Polygonetum cuspidati Tx. et Raabe 1950 apud Oberd. 1967: III, V (?)
 129. Helianthetum decapetalii Morariu 1967 n.n.: III (?)
 130. Eupatorietum cannabini Tx. 1937: II, III (?)
BIDENTETEA TRIPARTITI Tx., Lohm. et Prsg. 1950
BIDENTETALIA TRIPARTITI Br.-Bl. et Tx. 1943
Bidention tripartiti Nordh. 1940
 131. Bidentetum tripartiti (W. Koch 1926) Libbert 1932: III, V (?)
 132. Ranunculetum scelerati Siss. 1946 em. Tx. 1950: III (?)
 133. Bidentetum cernui Slavnic 1951: III (?)
 134. Xanthio strumarii-Chenopodietum albi (Timár 1947) I. Pop 1968: V (?)
Chenopodium fluviale (rubri) Tx. 1960
 135. Echinochloo-Polygonetum lapathifolii (Ujvárosi 1940) Soó et Csürös (1944) 1947: III-
 V (4,47,!)
 - chenopodiotosum albi Soó 1961: III (4,!)
 136. Echinochloo-Setarietum lutescentis Felsöldy 1942 corr. Soó 1971: III (?)
 137. Chenopodietum glauci (Wenzl. 1934) Raabe 1950: I, III (?)
 138. Xanthictum italicici Timár 1950: III-IV (25,!)

- PLANTAGINETEA MAJORIS** Tx. et Prsg. 1950
PLANTAGINETALIA MAJORIS Tx. (1947) 1950
Polygonion avicularis Br.-Bl. 1931 emend. Tx. 1950
 139. Lolio-Plantaginetum majoris (Linkola 1921) Beger 1930: IV-V (43,!)
 140. Poëtum annuae Gams 1927: I (!), III-IV (4,!)
 141. Polygonetum avicularis Gams 1927: II-V (4,25,!)
 - matricarietosum discoideac Morariu 1967 n.n.: II (?)
 142. Dauco-Matricarietum inodoreae I. Pop (1966) 1968: III (?)
 143. Juncetum tenuis (Diemont, Siss. et Westhoff 1940) Schwick. 1944: I-II (?)
Agropyro Rumicion crispí Nordh. 1940
 144. Myosuretum minimi Diem., Siss. et Westh. 1940: V (?)
 145. Trifolio repenti-Lolictum Krippelova 1967: IV (25)
 146. Lolio-Potentilletum anserinac Knapp 1946: II (!), IV (43)

147. Rorippo silvestri-Agrostictum stoloniferae (Moor 1958) Oberd. et Th. Müller 1961: IV-V (25, !)
148. Rorippo austriaceae-Agropyretum repentis (Timár 1947) Tx. 1950: IV (6a.32a)
149. Rumicetum-Alopecuretum geniculati Tx. (1937) 1950, Simon 1957: V (11a)
150. Ranunculetum repentis Knapp 1946 emend. Oberd. 1957: II-III (!)
151. Juncetum effusi Soó (1931) 1949: I-V
- EPILOBIETEA ANGUSTIFOLII Tx. et Prsg. 1950
- EPILOBIETALIA ANGUSTIFOLII Tx. 1950 corr. Soó 1961
- Atropion bella-donac* (Br -Bl. 1930) Tx. 1931
152. Rubo-Epilobietum Hadac et all. 1969: I-II (!)
- PETASITETO-CHAEROPHYLLETALIA Morariu 1967
- Telkion speciosae* Morariu 1967 n.n.
153. Telkio speciosae-Petasitetum albae Beldic 1967: II (1)
- SALICETEA PURPUREAE Moor 1958
- SALICETALIA PURPUREAE Moor 1958
- Salicion albae* (Soó 1930 n.n.) Müller et Gors 1958
154. Salici-Populetum (Tx. 1931) Mejer Drees 1936: III-IV (4,11a.32a, !)
155. Salicetum albae-fragilis Issler 1926 em. Soó 1957: II-V (22,25, !)
- amorphosum fruticosae Morariu et Danciu 1970: III-V (25, !)
- echinocystosum nov. fac.: III-V (!)
- Salicion triandrae* Müller et Gors 1958
156. Salicetum triandrac Malcuit 1929: III-V (!)
- amorphosum fruticosae Borza 1954 n.n.: IV-V (!)
- salicetosum viminalis Soó 1958: III-V (!)
157. Salicetum purpureae (Soó 1934 n.n.) Wendelbg.-Zelinka 1952: II-III (9, !)
- ALNETEA GLUTINOSAE Br -Bl. et Tx. 1943 em. Müller et Gors 1958
- SALICETALIA AURITAE Doing 1962 em. Westh. 1969
- Salicion cinereae* Müller et Gors 1958
158. Calamagrosti-Salicetum cinereae Soó et Zólyomi 1955: I (30, !), II, IV (!)
- spiraeaetosum salicifoliae Fl. Rațiu 1968: I (30, !)
- QUERCETEA PUBESCENTI-PETRAEAE (Oberd. 1948) Jakucs 1960
- ORNO-COTINETALIA Jakucs 1960
- Quercion farnetto* I. Horvat. 1954 corr. Soó 1960
159. Quercetum farnetto-cerris Georgescu 1945. Rudski 1949: IV (25, !)
160. Quercetum cerris Georgescu 1941: IV (25)
- PRUNETALIA Tx. 1952
- Prunion spinosae* Soó (1930 n.n.) 1940
161. Pruno spinosae-Crategetum (Soó 1927) Hueck 1931: IV-V (4.25, !)
162. Coryletum avellanae Soó 1927: II (22, !), IV (25, !)
- QUERCO-FAGETEA Br.-Bl. et Vliger 1937 em. Soó 1964
- FAGETALIA SILVATICA (Pawl. 1928) Tx. et Diem. 1936
- Aino Padion* Knapp 1942 em. Medvecka-Kornas 1957
163. Fraxino-Ulmetum (Tx. 1952) Oberd. 1953: V (24, !)

- fraxinetosum angustifoliac I. Pop 1979: V (24, !)
- quercetosum robori I. Pop 1979: V (25)
- 164. Alnetum incanae (Brockman 1907) Aichinger et Siegrist 1930: I (30, !), II (!)
- matteuccetosum Soó 1962; Lungu 1971: II (!)
- spiracactosum salicifoliac Fl. Rădu 1968: I (30)
- 165. Alnetum glutinosae-incanae Br.-Bl. (1915) 1930: II (!)
- 166. Aegopodio-Alnetum J. Kárpáti et Jurko 1961: IV (25, !)
Moehringia muscosa Acerion (Soó 1964) Boșcaiu 1979
- 167. Staphyleo-Tiliatum platyphylli Täuber 1986: IV (44)
- 168. Carpino-Tiliatum platyphylli Täuber 1986: IV (44)
- aceretosum tatarici Täuber 1986: IV (44)
Carpinton Oberd. 1953
- 169. Melampyro bihariense-Carpinetum Soó 1964: III-IV (4, 25)
- 170. Carpino-Quercetum petracae Borza 1941; I. Pop et Hodisan 1960: IV (25, !)
- 171. Carpino-Fagetum Pauca 1941: IV (25)
Symphyto Fagion (Vida 1959) Soó 1964
- 172. Festuco drymeae-Fagetum morarii et al. 1968: II (!), IV (25)
- 173. Chrysanthemo rotundifolio-Piceo-Fagetum Soó 1962, Vida 1959: II (!)
VACCINIO-PICEETEA Br.-Bl. 1939
- VACCINIO-PICEETALIA Br.-Bl. 1939 em. Hadac 1962
Vaccinio myrtillo-Piccion abietis Brezina et Hadac 1962
- 174. Hieracio tarnsilvanico-Piceetum abietis (Zlatnik 1935) Pawl. 1939: I-II (!)

Discussion

Analysis of the flora and vegetation points out that the Mureş Valley is now moderately degraded by human activity. The rate of degradation increases from the springs to the river mouth, and it has repercussions not only on the water ways of the Mureş, but also on the landscape of its banks. The best preserved sectors are I and II while the most deteriorated ones are V, III and IV, respectively. This fact is revealed also by the small number of therophyte species in the former two sectors (54 in I, 66 in II) as compared to the latter ones (318 in V, 304 in IV and 253 in III). The therophytes are indicators of drought and a degraded environment, by growing especially near localities as ruderal weeds and in crops as vegetal ones. The cosmopolites and adventive elements our allegation by their number in which they occur in the five sectors. Thus, the cosmopolites are distributed as follows: 33 in I, 32 in II, 60 in III, 64 in IV and 68 in V; the adventive elements: 5 in I, 8 in II, 24 in III, 30 in IV and 27 in V.

The vegetation of the Mureş valley is only to a limited extent natural and primary. It is about the riverside coppices (especially on the inferior course), the peat bogs and the rocky regions of the superior flow. Most of the vegetation is secondary, being represented by lawns (mainly sectors I and IV) and farming areas (particularly sectors V and III).

- The following areas must be at least partly protected and preserved for their floristic-phytocoenologic and landscape values: swamps from sector I (e.g. Voşlobeni, Joseni, Remetea), rocky regions of the Topliţa-Deda strait, woods near Arad (Ciala, Bezdin) and salt marshes at Idiceiu, Nagylak, Makó and Szeged.

References

1. Ardelean, A. (1973): Probleme ale oecotirii mediului înconjurător pe Valea Mureșului (Problems of plant protection on the Mureş Valley). - Cunoașterea și acțiune. Culegere de studii. Arad, 22-26.
2. Ardelean, V. (1971): Contribuții la cunoașterea florei și vegetației din împrejurimile comunei Lunca-Mureș (jud. Alba) (Review of the flora and vegetation of the Lunca-Mureş environs).- Luer. diplomă. Univ. Babeş-Bolyai Cluj-Napoca. Fac. biol.-geogr.
3. Baumgarten, J.Chr.G. (1816): Enumeratio stirpum Magno Transalvaniae. - Principatui... Vindobonae, I-III.
4. Borza, Al. Lupşa, Viorica, L. (1964, 1965): Flora și vegetația din jînul Blajului. I-II (Flora and vegetation of Blaj environs). - Contr. bot. Cluj. 147-166., 197-206.
5. Borza, Al. Lupşa, Viorica, L. (1968): Flora și caracterul geobotanic al împrejurimilor orașului Alba Iulia (Geobotanical characteristics and flora of the Alba Iulia). - Com. de bot. București, VI, 7-36.
6. Borza, Al., Rajtu Flavia. (1970): Molinetele din Bazinul Giurgeului (Molinea on the Giurgeu Basin). - Contr. bot. Cluj, 111-122.
- 6a. Boșcaiu, N. et al. (1972): Studiu fitocenologic al pajiștilor din sectorul hunedorean al văii Mureșului (Study on the phytocenology of the pastures in Valley Mureş on Hunedoara reaches).- Sargofia, ser.se. nat. Deva, IX, 53-75.
7. Csató, J. (1896): Alsófehér vármegye növényei (Plants of Alsófehér County). In: Alsófehér vármegye monografiája. II, 201-277.
8. Cserni, A. (1879): Gyulafelhérvarr és környékének flórája (Flora of Gyulafelhérvarr and its environs).- Gyulafelhérvarr.
9. Csûrös-Káptalan, M., Csûrös, St. (1972): Cercetări de vegetație în lunca Mureșului între Decea și Aiud (Research of the vegetation in the Mureş Valley between Decea and Aiud).- Contr. bot. Cluj. 237-246.
10. Csûrös, St. (1956): Vegetația din lunca Mureșului între Izvorul Mureș și Tg. Mureș (Vegetation of the Mureş Valley between Izvorul Mureş and Tîrgu Mureş). - (mscr.)
11. Csûrös, St. (1970): Despre vegetația ierbosă a luncilor din Transilvania (About vegetation of the grassy Transylvanian valleys). - Contr. bot. Cluj. 123-144.
- 11a. Csûrös, St., Fizitea, M. (1990): Cercetări de vegetație în împrejurimile localității Periam-Port (jud. Timiș) (Botanical research on the Periam-Port environs).- Contr. bot. Cluj. 3-16.
12. Diaconescu, B., Suteu, St. (1980): Analiză palinologică a lînajelor turboase de pe raza comunei Deda (Jud. Mureş) (Palinological analysis of the grassy peats on Deda environs). - Contr. bot. Cluj. 57-61.
13. Fölszegi, E. (1936): A Szegedi Fehértó növényzete (Vegetation of lake Fehértó near Szeged). - Debreceni Szemle, IX, 129-133.
14. Flora Republicii Populare Române (Flora Republicii Socialiste România) (1952-1976): Edit. Acad. București, I-XIII.
15. Fuss, M. (1866): Flora Transilvaniae excursioii Cibinii.
16. Gergely, I. (1964): Flora și vegetația regimului cuprinse între Mureș și masivul Bodeleu (Flora and vegetation between Mureş and Mt. Bodeleu). - (Dissertation), Univ. București.
17. Gyôrffy, I. (1929): Harasztok Csanád és Csongrád vármegyéból (Ferns from Csanád and Csongrád counties). - Acta Biol. Szeged. I,3.

18. Halász, A. (1889): *Enumeratio plantarum vesiculosarum oppidi Makó*. - Makói államilag segélyezett közs. polg. leányiskola évkönyve 1888-1889, Makó.
19. Jávorka, S. (1924-1925): *Magyar Flóra (The Hungarian Flora)*. Budapest.
20. Kitaibel, P. (1863): *Additamenta ad Floram Hungaricam*.
21. Nyárády, E.I. (1914): *Marosvásárhely és környékén élő tavaszi és nyáreleji növények határozója (A guide for the identification of the spring and summer time plants from Marosvásárhely and its environs)*.
22. Nyárády, E.I. (1931): *Die Vegetation des andesitischen Muresdurchbruchtales zwischen Toplița und Deda*. - Guide Six. Exs. internat. Roum., Cluj.
23. Pop, E. (1960): *Mlaștinile de turbă din R.P.Română (Peat bogs of Romania)*.- București.
24. Pop, I. (1979): *Considerații fitocenologice asupra pădurii Cială (jud.Arad) (Phytocenological studies of the environs of Cială forest)*. - Contr. bot. Cluj, 119-124.
25. Pop, I. et al. (1978): *Flora și vegetația Munților Zarand (Flora and vegetation of Mt. Zarand)*. - Contr. bot. Cluj, 3-216.
26. Porcius, Fl. (1878): *Enumeratio plantarum phanerogamicarum Districtus quondam Naszodiensis*.- Cluj.
27. Prodan, I. (1928): *Lesul dintre Dunăre și Tisa și cel dintre Tisa și ramificațiile Carpaților (Plain areas between Dunăre-Tisa and Tisa-Carpathian Mountains)*.- București.
28. Pușcaru-Sorceanu, Evd. et al. (1963): *Pășunile și finețele din Republica Populară Română (Pastures and hayfields of Romania)*.- Edit. Acad. R.P.R., București.
29. Răez, G., Voik, W. (1982): *Flora Reghinului în literatura botanică din secolul XIX (Flora of Reghin botanical literature of 19th century)*. - St. și com. Soc. st. biol. Reghin, 191-214.
30. Rațiu, Fl. (1968): *Vegetația lemnoasă din complexul mlașinilor eutrof Voglobeni (Jud. Harghita) (Woody vegetation of eutrophe marshy systems of Voșlobeni)* - Contr. bot. Cluj, 327-336.
31. Rațiu, Fl. (1971): *Asociații de rogozuri înalte din mlașinile eutrofe ale Depresiunii Giurgeului (Tall sedge associations of eutrophe marshes in the Giurgeu Basin)*. - Contr. bot. Cluj, 263-294.
32. Rațiu, Fl. (1972): *Asociații de rogozuri scunde din mlașinile eutrofe ale Depresiunii Giurgeului (Short sedge assotiations of eutrophe marshes in the Giurgeu Basin)*.- Contr. bot. Cluj, 161-186.
- 32a. Resmeriță, I. et al. (1972): *Vegetația din jurul Deva (Vegetation of the environs of Deva)*. - Sargetia, sc. nat.. Deva, IX, 23-52.
33. Resmeriță, I. Spârchez, Z. (1984): *Cercetări privind asociația Bromelium erecti Resmeriță et Spârchez 1963 (Research Bromelium erecti association Resmeriță et Spârchez 1963)*.-Contr.bot. Cluj, 169-174.
34. Samu, P. (1982): *Contribuții la cunoașterea florei săraturilor din zona Ideeiul Băi (Study of associations of the saline soils at Ideiul Băi)*. - St. și com. Soc. st. biol. Reghin, 191-204.
35. Schur, F. (1866): *Enumeratio plantarum Transsilvaniae*. Wien.
36. Simonkai, I. (1886): *Enumeratio Flora Transsilvaniae vasculosae critica*. Budapest.
37. Simonkai, I. (1893): *Arad vármegye és Arad város flórája (Flora of Arad County and Arad town)*.- Arad.
- 38a. Soó, R. (1928, 1938, 1946, 1947): *Zur Systematik und Soziologie der Phanerogamen Vegetation der Ungarischen Binnengewässer*. - Archivum Balatonicum. II, 45-79. - A magyar biológiai kutatóintézet munkáiból, Tihany. N., 174-194. - Annales Historico-Naturales Musci Nationalis Hungaria. Budapest. 44, 167-184. - Archiva Biologica Hungarica, Ser. II, 17, Tihany, 90-100.
39. Soó, R. (1938): *Die Arten und Formen der Gattung Potamogetum in der Flora des historischen Ungarn I, II.. Feddes Repertorium 45, 65-78.. 244-266.*
40. Soó, R. (1938): *A Tiszántúl flórája (Flora planitiei Hungariae Transtibicensis. Magyar flóraművek, II.. Debrecen.*
41. Soó, R. (1940, 1943): *A Székelyföld flórájának előmunkálatai (Preliminary studies of the flora of Szekler Land)*.- Kolozsvár. Supplementum I.
42. Soó, R. (1942): *Kritikai megjegyzések és újabb adatok a Székelyföld flórájához (Critical notes and recent data on the flora of Szekler Land)* Scripta Botanica Musei Transilvanici I. Kolozsvár, 38-52.
43. Tăuber, F. (1973): *Contribuții la flora și vegetația împrejurimilor orașului Lipova, Cunoaștere și aștiune (Contribution of the vegetation of environs of Lippa town)*. - Culegere de studii. Arad. 9-21.

44. Täuber, F. (1986): Sommerlinden-Waldgesellschaften vom unterlauf des Mures Flusses - Contr. bot. Cluj. 171-182.
45. Tímár, L. (1952): Egyéves növénytársulások a Szeged környéki szikesek iszapján I. (One yearly phytoecoses on the mud of saline lakes at Szeged environs) - Ann. Biol. Univ. Hung. tom. II., 311-321.
46. Timár, L. (1952): Adatok a Tisztántúl (Crisicum) flórájához (Data to the flora of Crisicum). - Ann. Biol. Univ. Hung., tom. II, 491-499.
47. Timár, L. (1957): Zönologische untersuchungen in den Ackerland Ungars.- Acta Botanica Acad. Sci. Hung. III., 1-2. Budapest.
48. Pop, I., Drăgușescu, C. (1983): Distribuția altitudinală a cormositelor pe cuprinsul Văii Sadului (Altitudinal distribution of the cormophytes on the Sad Valley) (munjii Cindrel-Stefoști).- Studia Univ. Babeș-Bolyai. Cluj-Napoca. 28, 3-8.
49. Sanda, V., Popescu, A., Doltu, M.I. (1980): Cenotaxonomia și corologia grupărilor vegetale din România (Cenotaxonomic and ecorologie of the plant groups in Romania). - St. și com. Muz. Brukenthal Sibiu, st. nat., 24. Suppl.
50. Sanda, V., Popescu, A., Doltu, M.I., Donita, N. (1983): Caracterizare ecologică și fitocenologică a speciilor spontane din flora României (Characterization of the ecology and phytocenology of spontaneous flora in Romania). - St. și com. Muz. Brukenthal Sibiu, St. nat., 25, Suppl.
51. Soó, R. (1964-1980): Synopsis systematico-geobotanica florae vegetationisquæ Hungariae. I-VI., Akad. Kiadó, Budapest.
52. Walz, L. (1878): A Görgény hegységen, a Maros mentén és Borszék vidékén 1858 nyarán gyűjtött növények jegyzéke (List of the plants collected on environs of Mt.Görgény, Maros and Borszék, summer 1858). - M.N.L., III., 65-72.

Constantin Drăgușescu, Sibiu University, Bul.Victoriei Nr. 39., 2400 Sibiu, Romania