

ZOOPLANKTON INVESTIGATIONS IN A LONGITUDINAL SECTION OF THE MAROS (MUREŞ) RIVER

KATALIN ZSUGA

Introduction

A detailed examination of a longitudinal section of the Maros River was conducted in August 1991. I performed the determination of the zooplankton from the biological examinations. The composition of the zooplankton stock, the large-scale presence or lack of certain organisms provides important information for evaluating the quality of a given waterway. I examined the groups of Rotatoria, Cladocera and Copepoda from the zooplankton elements in detail. In the course of the investigation of samples I addressed the following main questions:

- What sort of qualitative and quantitative changes characterize the zooplankton fauna of the Maros River?
- What sort of species describe the river in the given period?
- What sort of riparian categories are found along the longitudinal section? Are they separable, and, if so, what kinds of reaches are they?
- How can we describe the water quality of the Maros by the composition of the zooplankton fauna during the period of the examination?
- How can we evaluate the results of a single examination?

In Romania Rudescu (1960), Damian-Georgescu (1963,1970), Negrea (1983) refer to faunal, taxonomic research which mainly refers to the Danube, to the delta of the Danube, to the sea, to the high mountains, etc. I did not find any Romanian literature referring to the Maros. In the Hungarian reaches of this river Megyeri (1955,1970,1971,1972), Banesi (1981), Zsuga-Nagy (1989), Zsuga (1981,1990) performed examinations in the area around Makó and Szeged. I would like to contribute with this research to the disclosure of the Maros zooplankton fauna, to a more exact determination of changes in its water quality.

Material and methods

Time and location of the examination:

A zooplankton examination from the river Maros was performed in August 1991. The samples were taken from source to mouth in 15 segments (see Figure at p. 6).

Collecting method:

50-litre samples of water were filtered through plankton net, which is made of silk bolting cloth. The size of its mesh was 45 μm . The condensed samples were approx. 15-20 mls each, conserved on site with a 4-5% formaldehyde solution.

Processing method:

In the course of microscopic examinations I performed all the quantitative and qualitative processing of 15 samples. I used an Irgaval microscope and I did the counting in a box sized 80x35x6 mm and cubby-hole numbered with a graticule of 5x5 mm. For the preparation of mastax of Rotatoria I used hypoklorid (NaOCl). I gave the quantitative data in 100 i/l unit of measure. For identification of the species I used the taxonomic books from Banesi (1986,1988), Damian-Georgescu (1983,1970), Dévai (1977), Donner (1965), Carlin (1943), Gulyás (1974), Negrea (1983), Rudescu (1960), Ruttner-Kolisko (1974) and Voigt (1956).

Results

The development of the zooplankton of the rivers is influenced in great measure beside the known ecological factors (weather, nutrient state, temperature, etc.) by the hydrographical fundamentals of the area, the quality of the riverbed, the rise, the water speed, the quantity of the suspended load, etc. These effects are all observable in the development of the zooplankton of the Maros.

Rotatoria


The samples were taken in a period following a small flood. The numbers of the Rotatoria were rather few, in the 15 segments between 72 and 9120 i/100 l individual density was measured (Fig. 1). This great difference relates to the subsequent changes from the source to the mouth, to the differences between biotopes. During the examinations there were 62 species found altogether (Table 1). Around the source (Izvorul Mure^o) and downwards to it (Senetea, Suseni) a few species numbers were found beside the few individual numbers. The Maros River has a low water output here, with mountainous characteristics. In its Rotatoria fauna the organisms typical of low water, sources and streams (e.g. *Eencentrum orthodaetylum*, *Lecane arcuata*, *Trichocerca myersi* etc. Fig. 3) are present. Also there are a great number of representatives of the benthic and crust-dwelling creatures (e.g. *Cephalodella forficata* v. *macruca*, *Eencentrum grande*, *Lophocharis oxystemon*, *Notommata tripus*, *Pleurotrocha hyalina* etc. Fig. 5), as the planktonic and benthic living spaces do not separate definitely from each other as a consequence of the state of riverbeds, and the littoral region plays a great role as well.

In Sârma^o area greater individual numbers (640 i/100 l) and a higher number of species (19) were found than in the upper reaches (Figs. 1-2).

The Rotatoria fauna was the least at Răstolița and the species number was very few too, it was 4 altogether (Figs. 1-2). These organisms have a wide limit of tolerance

(*Cephalodella stercora*, *Lecane elosteroceera*, *Lepadella patella*, *Rotaria* sp.) and are even well adaptable to the great water-velocity too.

Fig. 1. Number of the Rotatoria (ind./100 l) in the River Maros



The most varied Rotatoria fauna (23 species) was developed in the neighborhood of Tîrgu Mureş and to Sîntimbru the characteristic organisms of both upper and lower courses can be found. The species of shallow waters, littoral region and euplanktonic elements were found equally (Fig. 2). The individual number grew as an effect of swelling and where this effect is not yet appreciable, decreased again.

Fig. 2. Number of Rotatoria species in the River Maros





Table 1. (continued)

<i>Keratella valga</i> f. <i>monospina</i>																		64
<i>Lecane arcuata</i>			4															
<i>Lecane bulla</i>						4	248				8							
<i>Lecane closteroceera</i>	20	4	4	48	12	12	16	64		80	552	240	64	24	12			
<i>Lecane lunaris</i>			8															
<i>Lepadella acuminata</i>						4												
<i>Lepadella ovalis</i>						4												
<i>Lepadella patella</i>	80	36	20	16	12	16	32		24	8								12
<i>Lepadella patella</i> v. <i>similis</i>			4	32								96						
<i>Lindia torulosa</i>									48									
<i>Lophocharis oxystemon</i>	20	4	4			8	8											12
<i>Lophocharis salpina</i>				16				32										
<i>Notommata tripus</i>		4																
<i>Platyias quadricornis</i>						8												
<i>Pleurotrocha hyalina</i>				16														
<i>Pleurotrocha petromyzon</i>						4												
<i>Polyarthra dolichoptera</i>																		
<i>Pompholyx sulcata</i>												24	96	32	24	48		
<i>Proales</i> sp.				16								48		12				
<i>Resticula melandocus</i>									24									
<i>Rotaria</i> sp.	80	44	32	144	36	192	496	784	408	264	408	336	320	96	72			
<i>Synchaeta pectinata</i>			4								48	96	64	12				
<i>Synchaeta tremula</i>			28					16					128	12	84			
<i>Testudinella mucronata</i>				16														
<i>Testudinella patina</i>						4		16		24								
<i>Trichocerca myersi</i>				16														
<i>Trichocerca pusilla</i>								64				192	32					
<i>Trichocerca</i> sp.			8															
Σ ROTATORIA	f100 l	260	120	128	640	72	336	952	2304	960	488	2904	9120	6656	624	2928		
CLADOCERA																		
<i>Alona guttata</i>			4															
<i>Alona rectangula</i>			4															
Σ CLADOCERA	f100 l	0	0	8	0	0	0	0	0	0	0	0	0	0	0	0	0	0
COPEPODA																		
nauplius			36	16	48		4	40	32			48		64	12	12		
copepodit		20						8	32				24					
Σ COPEPODA	f100 l	20	36	16	48	0	4	48	64	0	0	48	24	64	12	12		

The number of individuals grew at Ludu^o-Gheja too. The composition of the zooplankton stock relates to pollution in this area; the Rotatoria spp, which consume organic debris, dominated.

Fig. 3. Percentage composition of the Rotatoria in the River Maros



The following section of the Maros can be marked off from Alba Iulia, where the euplanktonic Rotatoria are found in the highest proportion (Fig. 3) (e.g. Anuraeopsis fissa, Brachionus angularis, Brachionus calyciflorus, Keratella cochlearis v. tecta, Filinia longiseta, Polyarthra dolichoptera) and the individual numbers multiplied proportionately to the upper areas (Fig. 1).

From the middle section downwards the trophic and saprobic degree grow on the flowing tributaries and pollution, the river becomes richer in nutrients and moderate pollution. This change was indicated by e.g. Lindia torulosa, Reticula melandocus, Pleurotrocha petromyzon, Brachionus spp., Epiphanes macrourus, etc. (Fig. 3).

Independent of the different section characteristics, Lecane closterocera and Rotatoria spp. were found at almost every sampling location. This relates to the wide range of tolerance of these organisms.

It is not typical in the course of the actual examination, but earlier examinations showed that very high individual densities can develop from time to time on the lower reaches of the river with the multitudinous swarming of 1-2 species (e.g. Brachionus spp., Anuraeopsis fissa etc.). In this case the influence of the Maros for the Tisza can grow considerably too. (Megyeri 1972, Zsuga-Nagy 1989.).


Crustacea

The results of the examinations showed the hydroecological conditions of the Maros were not favourable to Cladocera. They were only found in one area (Suseni), in small individual numbers (8 i/100 l). Both identified species (Alona guttata, Alona rectangulara) are curytop organisms, they can live in completely different waters. It is documented by the

earlier examinations performed in the neighborhood of Makó and Szeged that the Cladocera species are not frequent even in the lower reaches (Zsuga-Nagy 1989, Zsuga 1990).

The representatives of Copepoda are found at almost every sampling location though in few numbers (Fig. 4). The dominance of juvenile forms are identifiable by generation, thus nauplius and copepodite forms were found in different developmental phases and no adult species were found in the samples during the period of examination.

Fig. 4. Number of the Crustacea (ind/100 l) in the River Maros



Summary

The following conclusions may be drawn from this examination of the zooplankton in a longitudinal section of the Maros River.

-The quantity of zooplankton was generally low during the given time.

-By quantitative composition the proportion of Rotatoria dominated, the hydroecological conditions of the Maros were not favourable to Cladocera and in the Copepoda group the predominance of juvenile forms was characteristic in contradiction to adults.

-We could separate the Maros into three sections by the qualitative composition of Rotatoria, and by the presence of indicator species (Fig. 3).

1. Between Izvorul Mure^o and Răstolița the section has an upper course character, the oligotrophic, oligosaprobic water category was typical with low numbers of both species and individuals.

2. Between Tîrgu Mureş and Sîntimbru the most varied species-composition developed; the benthic, planktonic and littoral elements were mixed. The trophic and saprobic degree rose, the nutrient state and the pollution of the river grew.

3. Between Alba Iulia and Szeged the composition of the Rotatoria euplanktonic elements dominated. The number of the Rotatoria multiplied in proportion to conditions in the upper areas.

-In the different sections of the river, aside from the typical indicator species colouring elements were found which have good adaptability and a wide range of tolerance.

-This single examination gave only a few appreciable results for the characterization of Crustacea fauna of the Maros River. Repeated examinations are needed to know this group in greater detail.

-This results of this single examination have a disclosing character and indicating value. Because data in the scientific literature is limited, concerning zooplankton of the Maros, further examinations would be expedient for more detailed knowledge of the river. These present data may be considered as a basis for comparison.

References

- Bancsi L. 1986: A kerekcsférgek Rotatoria kishatározója I. (A guide for the identification of Rotatoria I). VIII.15. (Hungarian)
- Bancsi L. 1988: A kerekcsférgek Rotatoria kishatározója II. (A guide for the identification of Rotatoria II). VIII.17. (Hungarian)
- Bancsi L., Hamar J. 1981: Data on the Rotatoria and Zooflagellata fauna of the sediment in the Tisza and its tributaries. Tiseia XVI. 155-159.
- Bancsi L., Hamar J. 1981: Data on Rotatoria and Zooflagellata fauna of the sediment in the Tisza and its tributaries. Tiseia XVI. 155-159.
- Carlin B. 1943: Die Planktonrotatorien des Motalaström Meddelanden Fran Lunds Universitets Limnologiska Institution
- Damian-Georgescu A. 1963: Fauna Republicii Populare Romine, Crustacea Vol. IV. Fasc.6. Copepoda Cyclopidae, Ed. Acad. R.P.R., Bucureşti.
- Damian-Georgescu A. 1970: Fauna Republicii Socialiste România ,Crustacea Vol. IV. Fasc.11. Copepoda Harpacticoida, Ed. Acad. R.P.R., Bucureşti.
- Dévai I. 1977: Az evezőlábú rákok (Calanoida és Cyclopoida) alrendjeinek kishatározója (A guide for the identification of Calanoida and Cyclopoida). VIII.5. (Hungarian)
- Donner J. 1965: Ordnung Bdelloidea (Rotatoria, Radertiere)
- Gulyás P. 1974: Az ágasesápú rákok Cladocera kishatározója (A guide for the identification of Cladocera). VIII.2. (Hungarian)
- Megyeri J. 1955: Planktonvizsgálatok a Tisza szegedi szakaszán (Study of the plankton in reaches of the Tisza River at Szeged). Hidrológiai Közöny 35.évf. 7-8.sz. 280-292. (Hungarian)
- Megyeri J. 1970: A Tisza mesozooplanktonja I. Rotatoria (Mesozooplankton of the Tisza River I. Rotatoria). Szegedi Tanárképző Főiskola Tud. Közleményei 115-130. (Hungarian)
- Megyeri J. 1971: A Tisza mesozooplanktonja II. Entomostraca (Mesozooplankton of the Tisza River II. Entomostraca). Szegedi Tanárképző Főiskola Tud. Közleményei 99-110. (Hungarian)
- Megyeri J. 1972: Zooplankton vizsgálatok a Tisza mellékfolyóin (Investigations of the zooplankton in tributaries of the Tisza River). Szegedi Tanárképző Főiskola Tud. Közleményei 63-73. (Hungarian)

- Negrea S. 1983: Fauna Republicii Socialiste România, Crustacea Vol.4.Fasc.12.Cladocera. Ed. Acad. R.P.R., Bucureşti.
- Rudescu I. 1960: Fauna Republicii Populare România, Trochelminthes Vol.II.Fasc.II. Rotatoria. Ed. Acad. R.P.R., Bucureşti.
- Ruttner-Kolisko A. 1974: Plankton Rotifers Biology and Taxonomy. Die Binnengewasser XXVI:1.
- Zsuga K. 1981: Benthic Entomostraca fauna of the Tisza and its tributaries Tiscia Vol.XVI.183-190.
- Zsuga K.-Nagy M. 1989: Zooplankton vizsgálatok a Tisza vízgyűjtő területén (Investigation of the zooplankton on catchment area of the Tisza river). Természetvédelem 89-pályázat anyaga (Hungarian manuscript)
- Zsuga K. et al. 1990: A Tisza és mellékfolyóinak hidrológiai vizsgálata (Hydroecological study of the Tisza river and its tributaries). Kötivizig Zárójelentés 23-29. (Hungarian manuscript)

Katalin Zsuga, Water Management Directorate, H 5000 Szolnok, Ságvári krt. 4. Hungary