

USE OF WOOD ASH FOR FERTILIZING THE PLANTATION OF FAST-GROWING WOODY PLANTS – BASKET WILLOW (SALIX VIMINALIS L.)

Otepka Pavol, Ďuďák Ivan, Tóthová Slavka

Department of Sustainable Development, Faculty of European Studies and Regional Development,
Slovak University of Agriculture in Nitra, Tr. A. Hlinku 2, 949 76 Nitra, Slovak Republic
e-mail: Pavol.Otepka@uniag.sk

ABSTRACT

Energetic utilization of wood biomass obtained from fast-growing woody plants nowadays records more important capability. The article discusses the issue of research using wood ash as solid waste from the boiler while burning biomass and the intensification of short-rotation coppice basket willow – common osier (*Salix viminalis* L.) for energy use as a renewable energy source. Purpose of the article is to show the importance of basket willow (*Salix viminalis* L.) as short-rotation coppice with the emphasis of its production and remedial effects.

Key words: renewable energy resources (carriers), biomass, short-rotation trees, basket willow, common osier (*Salix viminalis* L.), wood ash

1. INTRODUCTION

The issue of renewable energy resources is still current, especially in times when oil crisis and negative consequences of global climate change are appearing in the human society. One of these sources is the biomass. It fulfills majority of benefits from all spectrum of renewable energy resources (Húska et al., 2000; Niňaj – Húska, 2005). In the spring of 1994, experiments to prove the possibility for cultivation of fast-growing varieties of basket willow – common osier (*Salix viminalis* L.) were carried out in Krivá research centre, located in the Orava region of the Slovak Republic. For this purpose, three varieties of willow bred at the Research Institute of Agriculture in Swale (Sweden), were used: ULV, ORM and RAPP (Habovštiak – Daniel, 2001). The same varieties are tested in the conditions of warm agro-climatic region in Dolná Malanta, Nitra, Slovakia. They have been grown here continuously since 1996. Experiments are established on Experimental Base of the Slovak University of Agriculture in Nitra, Slovak Republic (Otepka – Habán, 2006).

By the cultivation of fast growing coppice with short rotation period, the nutrients for growing plants are taking off intensively, reducing their storages in the soil. Partial compensation might be mineral multicomponent fertilization, for which wood ash with the content of wide spectrum of macro and microelements can fit well. Specific problem is that this type of solid waste generated in the boilers does not contain nitrogen, which had been released into the air after the biomass was burned. The project, supported by the Slovak Research and Development Agency is carried out by cooperation between Intech Slovakia, Ltd. and Slovak University of Agriculture in Nitra.

2. MATERIAL AND METHODS

Methodology of the experiment with wood species basket willow – common osier (*Salix viminalis* L.) has a long-term design (25 years). Stand was established by planting in the spring of 1996 (19th April). Planting material was originated from Swedish breeding (Swale); source was in Research Centre Krivá in the Orava region, the Slovak Republic (VÚTPaHP). Planted cuttings were done in length 0.20 m; thickness from 10 to 20 mm. Forecrop was *Amaranthus* sp. Soil preparation was done by plowing to the depth of 30 cm, dragging, harrowing. Planting method: manually by hand. Treatment during the vegetation period is aimed to regulation of harmful factors (weeds, pests and diseases) and to maintain the status and condition of the crop stand.

Total cultivation area is approximately 648 m²; area of individual plots (variants) is 108 m²; total number of tested plots is 6; identification of plots: ORM/A, ORM/B, ULV/A, ULV/B, RAPP/A RAPP/B. Cultivated varieties: ORM, ULV, RAPP. Design of experiment: A – Double row planting with row distance = 0.5 m, alternated 1.25 m and 0.75 m between rows, B – plant spacing 0.5 x 0.75 m. The article includes results from a third four-year harvesting period of willow in the presented locality (2004 – 2007). In this plantation, the crop biomass forecasting of willow is based on mathematical-statistical model, published by Špánik and Čimo (2009). Wood ash was applied in the autumn period (October 2009) and in the spring period (March 2010) in doses of 1.0, 3.0 and 5.0 t.ha⁻¹, while the control area was left without application of wood ash. Size of each area is: 2 x 162 m², i.e. 324 m².

3. RESULTS AND DISCUSSION

Number of individual willow plants (*Salix viminalis* L.) per unit area depends on the plant spacing of 2.00 individual plants per m² (planting space A) or 2.67 individual plants per m² (planting space B). According to Šimala – Habovštiak (1995), well-managed willow grows in the first year up to the height of 2.0 – 3.0 m and after the fourth year cultivation; the crop is 7.0 m high. Individual crops with height of 7.0 m can be harvested. From this finding it is difficult to conclude whether the authors considered these data also in assessing of the third harvest period, but for the comparison of results we indicate, that one year after the second harvest, the indicated height of all three varieties was not reached: ORM (1.70 m), ULV (1.77 m) and also variety RAPP (1.45 m, see Table 1) was below the height of 2.00 m. Better average height of rods was always reached at the plant spacing B, i.e. the one where denser planting was applied. In subsequent years, the height of the vegetation ranged from 1.95 m (variety, RAPP/A, in the 10th year after planting) to 4.27 m (variety ULV/B, in the 12th year after planting), and as it can be seen in Table 1 in neither of cases, the height reached 7.0 m. This we attribute to the impact of harmful agents, soil characteristics of the experimental unit as well as to the weather conditions during the period of observation. When observing an average rod thickness, the values were measured according to the forestry parameters i.e. at a level of 1.3 m above the soil surface. Based on observations in the second harvest period, in this parameter we reached relatively low values, since in neither of case the value got over 20 mm. For ULV variety grown in the plant spacing B, the highest average rod thickness observed was 18.2 mm.

Habovštiak – Daniel (2001) indicate that in the production year an average yield of 12.0 to 15.0 t.ha⁻¹ dry matter is probable. With this statement we can only agree with ULV variety, since its yield was higher than 12.0 t of dry matter.ha⁻¹.yr⁻¹ (plant spacing A: 12.46 t.ha⁻¹.yr⁻¹ and plant spacing B: 13.80 t.ha⁻¹.yr⁻¹).

Table 1. Measured parameters of basket willow – common osier (*Salix viminalis L.*) during four-year period (2004 – 2007)

Variety / Arrangement		ORM/A	ORM/B	ULV/A	ULV/B	RAPP/A	RAPP/B
Parameters							
Number of plants.m ⁻²		2.00	2.67	2.00	2.67	2.00	2.67
Average branch length (m)	9 th year	1.45	1.95	1.67	1.87	1.42	1.47
	10 th year	2.58	2.58	2.74	3.04	1.95	1.89
	11 th year	3.18	3.25	3.24	3.68	2.58	2.83
	12 th year	4.12	3.90	4.18	4.27	3.62	3.34
Average branch width (mm)	9 th year	8.5	10.8	9.5	9.3	8.1	7.6
	10 th year	10.7	13.4	11.0	12.4	10.2	9.8
	11 th year	12.8	15.7	13.8	15.4	12.5	12.1
	12 th year	13.7	17.6	16.6	18.2	14.0	13.2
Average weight of one branch after drying up (g)	9 th year	286	350	450	420	134	85
	10 th year	1050	960	1120	1090	460	385
	11 th year	2002	1680	1972	1851	820	765
	12 th year	2480	1940	2680	2490	1038	990
Weight of sample in dry matter (g)	9 th year	2860	3500	4500	4200	1340	850
	10 th year	10500	9600	11200	10900	4600	3850
	11 th year	20020	16800	19720	18510	8200	7650
	12 th year	24800	19400	26800	24900	10380	9900
Dry matter yield (t.ha ⁻¹)	9 th year	6.20	8.50	5.90	7.60	2.90	5.52
	10 th year	19.35	22.80	18.55	20.30	5.20	12.40
	11 th year	36.20	32.10	38.50	34.40	10.15	22.65
	12 th year	44.23	38.30	49.85	55.20	16.70	31.80

Wood ash obtained from a bottom area of the boiler after burning of biomass originated from the burning of clean and chemically untreated wood from the thermal CHP power plant in Hriňová. Content of hazardous elements in the applied wood ash (As, Cd, Cr, Hg, Pb) was lower than limit values of risk mineral elements of calcium and magnesium-calcium fertilizer under the Ministry of Agriculture Regulation from 15.12.2000, stipulating the types of fertilizers, the content of risk elements, conditions of sampling and methods for testing fertilizers, Coll. of Laws 26/2001, Supplement No. 2., content and other elements considered to be risky in the raw material for compost (Cu, Mo, Ni, Zn) was lower than the limit values of risk elements in raw materials for composting, according to STN 46 5735 "Industrial composts". Wood ash contained 11.5% calcium, 4% potassium and 1.4% magnesium.

4. CONCLUSION

Fast growing basket willow – common osier (*Salix viminalis* L.) has been grown on experimental base in Dolná Malanta, Nitra, Slovakia since 1996. In 12 years of its cultivation, the crop has been retained in a condition suitable for energetic, but also for therapeutic purposes and creates conditions for further observation and assessment in future. By experiment, the influence of factors on above-ground biomass production was studied: variety (ORM, RAPP, ULV), design of crop organization (method A – double rows = 0.5 m × 0.75 m or 1.25 m; method B = 0.5 m × 0.75 m) and the impact of crop years (2004 – 2007). From dried and weighted samples of plant rods following conclusions can be drawn. In the 12th year of cultivation, the height of vegetation ranged from 3.34 m (RAPP/B) to 4.27 m (ULV/B). The highest average height of vegetation was in variety ULV 4.23 m, followed by ORM 4.01 m and the lowest height was recorded in a variety RAPP 3.48 m. The average thickness of plant rods ranged from 13.2 mm RAPP/B to 18.2 mm ULV/B. Yield of the third harvest period measured in dry matter varied in significantly wide range: from 16.7 (RAPP/A) to 55.2 t.ha⁻¹ (ULV/B). Based on measured and calculated data the varieties of willow can be ordered according to their suitability for the conditions of southern Slovakia as follows: 1. ULV (52.5 t.ha⁻¹); 2. ORM (41.3 t.ha⁻¹); 3. RAPP (24.3 t.ha⁻¹). In terms of planting organization, better results were achieved in planting space B (41.8 t.ha⁻¹) than in planting space A (36.9 t.ha⁻¹).

The result of the project "Efficiency of fertilization potential of wood-ash for energy plants cultivation as renewable energy resources (Ash as fertilizer)" will be the determination of an appropriate procedure in application of wood ash for the intensification of energetic plants cultivation. Since, in accordance with the Slovak Republic legislation, it is allowed to use wood ash only for the scientific purposes, we assume that the results obtained in this project will be applied in the modification of legislative framework to enable the use of wood ash for applying nutrients in energetic plants cultivation.

ACKNOWLEDGEMENT

The work is carried out through funding by Applied Research and Development Project No.: VMSP-P-0035-09 under the "Research and development support for small and medium size enterprises" as was declared by the Slovak Research and Development Agency in 2009.

REFERENCES

1. ČIMO, J. – ŠPÁNIK, F. 2009. The agro climatic analysis phytomass production of *Salix viminalis*. In: Sustainable development and bioclimate, Stará Lesná. Bratislava : Slovenská bioklimatologická spoločnosť, 2009. pp. 142-143. ISBN 978-80-900450-1-9.
2. HABOVŠTIAK, J. – DANIEL, J.: Pestovanie odrôd vřby (*Salix viminalis*) na energetické účely. In: Naše pole, 5/2001, pp. 10. ISSN 1335-2466.
3. HÚSKA, J. et al.: Experimentálne výsledky z pestovania vřby košíkárskej (*Salix viminalis*). In: Nová energetická politika SR: obnoviteľné zdroje energie, aproximácia k politike EÚ (zborník z medzinárodnej konferencie). Bratislava: Dadaexpress, 2000. pp. 124-128. ISBN 80-968421-0-2.
4. JAMRIŠKA, P.: Rastliny na nepotravinárske využívanie. Nitra : SPU, 2001. 69 p. ISBN 80-7137-877-1.
5. MAGA, J. – NOZDROVICKÝ, L. – PEPICH, Š. – MARHAVÝ, Ľ. – HAJDU, Š. 2008. Komplexný model využitia biomasy na energetické účely. Nitra : SPU, 2008. 183 p. ISBN 978-80-552-0029-3.
6. NIŇAJ, M. – HÚSKA, J.: Vřba – zdroj energie. In: Úroda, 6/2005, pp. 38-39. ISSN 0139-6013.
7. OTEPKA, P. – HABÁN, M. 2006. Biomass yield of basket willow (*Salix viminalis* L.) cultivated as energy plant in a long-term experiment. In: Acta fytotechnica et zootechnica, Vol. 9, 2006, No. 3, pp. 70 -74. ISSN 1335-258X.