

ARTICLE

On the mechanism of stimulation by low-concentration stressors in barley seedlings

Erika Kovács^{1,2}, Péter Nyitrai^{1*}, Áron Keresztes²¹Department of Plant Physiology and Molecular Plant Biology, Eötvös Loránd University, Budapest, Hungary, ²Department of Plant Anatomy, Eötvös Loránd University, Budapest, Hungary

ABSTRACT Beneficial effect of low-concentration chemical stressors like Cd salt and a herbicide DCMU have been investigated previously in different plant model systems. The symptoms of stimulation are known from earlier studies, but information about the mechanism is limited. In this work, the mechanism of stimulation of low-concentration Cd ($5 \cdot 10^{-8}$ M) and 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU, 10^{-7} M) was investigated in barley seedlings. In treated plants, the amount of cytokinins increased in roots and, after being transported to the leaves, they caused stimulation of chlorophyll accumulation, photosynthetic activity, and delayed senescence. To identify the signal transduction pathway(s) involved in the primary stimulation of cytokinin synthesis in roots, specific phosphatidylinositol-4,5-bisphosphate – inositol-1,4,5-triphosphate/diacylglycerol (PIP₂-IP₃/DAG) and mitogen activated protein kinase (MAPK) signaling pathway inhibitors were added to the nutrient solutions. Both of them proved to be effective, eliminating the stimulation of the stressors

Acta Biol Szeged 52(1):179-180 (2008)

KEY WORDScadmium
chemical stressors
cytokinin
DCMU
MAPK pathway
PIP₂-IP₃/DAG pathway

Harmful effects of chemical stressors like heavy metals and herbicides have been widely investigated in the presence of their high concentrations (Krupa and Baszyński 1995; Sanitá di Toppi and Gabrielli 1998). Chemical stressors applied in low-concentration may have a beneficial effect (Zatykó 1973, Karavaev et al. 2001). These stimulative effects include increase of chlorophyll (Chl) content, changes in the composition of Chl-protein complexes, elevation of CO₂ incorporation, and changes in the ultrastructure of chloroplasts (Nyitrai et al. 2003, 2004, 2007).

In the present work, our aim was to explore the mechanism of stimulation. The role of cytokinins (CKs) could be expected on the basis of our earlier results with a rooting, detached bean leaf model (Nyitrai et al. 2004). Specific inhibitors for PIP₂-IP₃/DAG pathway and also for the mitogen activated protein kinase (MAPK) pathway were applied to examine whether these signaling pathway(s) were involved in the stimulation of plants.

Materials and Methods

Barley (*Hordeum vulgare* L. cv. Omega) were grown in ¼ strength Hoagland solution under standardized conditions: 14/10 h light/dark periods, at 24/18°C, photosynthetically active radiation of 100 μmol m⁻²s⁻¹ (Fodor et al. 1998). After 10 days of germination, hydroponic treatments were started under the same conditions with $5 \cdot 10^{-8}$ M Cd(NO₃)₂ or 10^{-7} M

DCMU. Barley seedlings were also treated with signalling pathway inhibitors (10^{-5} M LiCl, 10^{-5} M stearyl carnitine chloride – SCC, or $5 \cdot 10^{-5}$ M 2-(2'-amino-3'-methoxyphenyl)-oxanaphthalen-4-one – PD) added to control or together with the above-mentioned stressors in the nutrient solutions. Samples were taken after 0, 1 and 2 weeks of treatment from roots, and the first leaves.

Chl content were determined according to Porra et al. (1989).

Amount of active CKs were measured according to Biddington and Thomas (1973) with a modified *Amaranthus* betacyanin bioassay.

Results and Discussion

The Cd content of roots raised considerably, but its amount increased very little in leaves during the treatment (not shown). One week treatment with Cd already lead to the increase of the CK level in roots and it remained at a high level throughout the experiment (Fig. 1 A) compared to the control. During the second week, its amount increased almost three times in leaves (Fig. 1 B). After one and two weeks of DCMU treatment, the amount of CKs were considerably higher than the control level in roots and similarly in leaves (Fig. 1 A and B).

As a consequence of higher the level of CK in Cd and DCMU-treated leaves, Chl accumulation was facilitated slightly after one week of treatment, and more pronounced up to the end of treatment (Fig. 2). The signalling pathway inhibitors (Li, SCC, and PD) applied together with the stres-

*Corresponding author. E-mail: pnityi@ludens.elte.hu




Figure 1. Amount of CKs in roots (A) and the first leaves (B) of control (Ctrl), stressor (Cd, DCMU) treated or stressor plus inhibitor (Li, SCC, PD) treated barley seedlings. SDs are within 10%.

sors caused a decrease in the amount of of active CKs in their roots and leaves compared to the stressor-treated plants (Fig. 1 A and B). Similarly, the stressor plus inhibitor treated plants contained lower amount of Chl compared to the stressor-treated plants (Fig. 2).

Electron microscopy of control and treated chloroplasts supported the rejuvenating effect of stressors, mainly on the basis of the difference in total amount of plastoglobuli.

In summary, the signal from the low-concentration stressors is transmitted through PIP_2 - IP_3 /DAG and protein kinase C to MAPK pathway, and it causes the synthesis (and/or activation) of CKs in roots. CKs are then transported to the leaves and facilitate Chl accumulation via cytokinin transporters and the cytokinin signaling pathway extending the CK signal to the leaves.




Figure 2. Chl content in the first leaves of control (Ctrl), stressor (Cd, DCMU) treated or stressor plus inhibitor (Li, SCC, PD) treated barley seedlings. SDs are within 5%.

Acknowledgement

This work was supported by grant T-047243 from OTKA.

References

- Biddington NL, Thomas TH (1973) A modified *Amaranthus* betacyanin bioassay for the rapid determination of cytokinins in plant extracts. *Planta* 111:183-186.
- Fodor F, Cseh E, Varga A, Zárny Gy (1998) Lead uptake, distribution, and remobilization in cucumber. *J Plant Nutr* 21:1363-1373.
- Karavaev VA, Baulin AM, Gordienko TV, Dovyd'kov SA, Tikhonov AN (2001) Changes in the photosynthetic apparatus of broad bean leaves as dependent on the content of heavy metals in the growth medium. *Russ J Plant Physiol* 48:38-44.
- Krupa Z, Baszyński T (1995) Some aspects of heavy metal toxicity towards photosynthetic apparatus – direct and indirect effects on light and dark reactions. *Acta Physiol Plant* 17:177-190.
- Nyitrai P, Bóka K, Gáspár L, Sárvári É, Lenti K, Keresztes Á (2003) Characterisation of the stimulating effect of low-dose stressors in maize and bean seedlings. *J Plant Physiol* 160:1175-1183.
- Nyitrai P, Bóka K, Gáspár L, Sárvári É, Keresztes Á (2004) Rejuvenation of ageing bean leaves under the effect of low-dose stressors. *Plant Biol* 6:708-714.
- Nyitrai P, Mayer M, Óvári M, Keresztes Á (2007) Involvement of phosphoinositide signaling pathway in the anti-senescence effect of low-concentration stressors on detached barley leaves. *Plant Biol* 9:420-426.
- Porra RJ, Thomson WA, Kiedemann PE (1989) Determination of accurate extinction coefficient and simultaneous equations for assaying chlorophyll a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. *Biochim Biophys Acta* 975:384-394.
- Sanità di Toppi L, Gabrielli R (1998) Response to cadmium in higher plants. *Environ Exp Bot* 41:105-130.
- Simard S, Greiner G, Beaumont G (1990) Morphometric analysis of ultrastructural changes induced by a sublethal concentration of atrazine on young *Lemna* minor chloroplasts. *Plant Physiol Biochem* 28:49-55.
- Zatykó J (1973) Effect of low concentration colchicine on the cytokinin content of pinto bean leaves. *Die Naturwissenschaften* 60:105-106.