

## ARTICLE

# Changes of photosynthetic parameters in wheat/barley introgression lines during salt stress

Sándor Dulai<sup>1\*</sup>, István Molnár<sup>2</sup>, Márta Molnár-Láng<sup>2</sup><sup>1</sup>Department of Plant Physiology, Eszterházy College, Eger, Hungary, <sup>2</sup>Agricultural Research Institute, Hungarian Academy of Sciences, Martonvásár, Hungary

**ABSTRACT** Salt stress induced photosynthetic responses were investigated in the wheat/barley introgression lines developed in the Agricultural Research Institute, Martonvásár, Hungary, and in different parental wheat and barley *Hordeum vulgare* L. genotypes. An increase in NaCl concentration of the nutrient solution to 200 mmol L<sup>-1</sup> resulted in a considerable stomatal closure and a decreased net CO<sub>2</sub> assimilation rate (*A*) for wheat genotypes, barley cv. Betzes and 4H Asakaze komugi-Manas addition line while changes of these parameters were less significant for barley cv. Manas and the 7H Asakaze komugi-Manas and 6H Mv9 kr1-Betzes-Seneca addition lines. These preliminary results suggest that the 7H Akom-Manas addition line may be a good candidate for improving the salt tolerance of wheat in the future.

**Acta Biol Szeged 55(1):73-75 (2011)****KEY WORDS**salt stress  
wheat  
barley  
photosynthesis  
wheat/barley introgression lines

Salt stress is an important biomass-limiting stress factor in the field causing the suppression of cultivated plants in growth and in crop production (Belkhodja et al. 1999; Dulai et al. 2011). In crop plants several life processes are particularly sensitive to soil and water salinity, since high salt concentration has a three-fold effect: it limits water availability to the roots (osmotic stress); it can cause ionic stress when the salt is taken up by plants; and it also influences nutrient uptake and translocation (Bageci 2003; Siler et al. 2007). The reduction in photosynthesis during salt stress can be attributed to the reduced activity of primary photochemical processes (Kaiser et al. 1983) and the inhibition of carbon-dioxide fixation and assimilation (Siler et al. 2007; Dulai et al. 2011) related to metabolic processes and the diffusion of CO<sub>2</sub> to chloroplasts. Furthermore, the estimation of net assimilation rate might allow for a good discrimination between salt tolerant and non-tolerant cultivars of barley (Belkhodja et al. 1999).

The development of salt tolerant crops bears significant agricultural and economical importance, as soil salinity decreases the yield of cultivated plants. Barley is a potential gene source for wheat improvement, because of its generally good stress tolerance (Molnár et al. 2007). The introgressive hybridization of barley to wheat makes it possible to transfer some useful characteristics of salt tolerance as barley is regarded as being more salt tolerant than bread wheat (Colmer et al. 2005; 2006).

In this paper the photosynthetic responses to salt stress in the parental winter wheat (Mv9kr1, Asakaze komugi) and

barley (Manas, Betzes) cultivars were compared with those of 4H, 7H Asekazi komugi-Manas and 6H Mv9kr1-Betzes-Seneca addition lines, in order to indicate whether the introgression lines had better tolerance to salinity than wheat.

## Materials and Methods

The salt stress was applied on the wheat/barley introgression lines produced in Martonvásár (Molnár-Láng et al. 2000), together with the parental lines (Table 1). After germination, plants were grown in pots containing 1500 mL volume in half-strength modified Hoagland nutrient solution in growth chambers under normal CO<sub>2</sub> concentration at 20/25°C. Growth light intensity was 200 μmol m<sup>-2</sup> s<sup>-1</sup> with a 12/12 photoperiod. The water status of plants was traced by determining the relative water content (RWC).

CO<sub>2</sub> assimilation of intact leaves was measured with an infrared gas analyser (GFS-3000-FL gas exchange system, Walz, Effeltrich, Germany). The net CO<sub>2</sub> assimilation rate (*A*), stomatal conductance (*g<sub>s</sub>*), and intercellular CO<sub>2</sub> concentration (*C<sub>i</sub>*) were calculated in the light saturated state of photosynthesis by using the equations of von Caemmerer and Farquhar (1981).

## Results and discussion

In this study the photosynthetic performance of the newly developed wheat-barley introgression lines was examined and compared with those of parental genotypes under 200 mM salt concentration. These examination were encouraged by earlier findings suggesting that barley is regarded as being more salt tolerant than bread wheat (Colmer et al. 2005, 2006).

Accepted July 11, 2011

\*Corresponding author. E-mail: ds@ektf.hu

**Table 1.** Values of relative water content (RWC), stomatal conductance ( $g_s$ ) and net CO<sub>2</sub> assimilation rate (A) for NaCl treated (200 mmol L<sup>-1</sup>, Stress) and for control plants of similar age grown in nutrient solution without NaCl (Control). Mv9kr1, winter wheat originating from a Martonvásári wheat cultivar; Akom, winter wheat cv. Asekaze komugi; Manas, barley cv. Manas; Betzes, barley cv. Betzes; 4H add, 4H Asakaze komugi-Manas addition line; 7H add, 7H Asakaze komugi-Manas addition line; 6H add, 6H Mv9 kr1-Betzes-Seneca addition line.

Genotypes	Relative water content (RWC)		Stomatal conductance ( $g_s$ , mmol m <sup>-2</sup> s <sup>-1</sup> )		Net CO <sub>2</sub> assimilation rate (A, $\mu$ mol m <sup>-2</sup> s <sup>-1</sup> )	
	Control	Stress	Control	Stress	Control	Stress
Mv kr1	94.93±0.24	89.21±1.65	226.56±1.16	117.76±12.04	19.7±1.16	10.17±1.24
Akom	90.09±2.88	84.08±2.93	276.18±9.82	93.26±8.26	16.82±0.62	9.6±1.03
Manas	86.46±3.9	79.15±4.6	187.39±24.18	159.27±6.75	15.8±1.42	13.98±0.85
Betzes	94.79±0.8	81.68±4.18	234.85±5.98	112.71±3.26	20.63±0.29	9.9±0.97
4H add	84.25±1.64	83.45±2.96	299.92±51.86	88.24±8.7	16.99±1.61	11.18±0.43
7H add	93.6±1.38	82.52±0.73	189.81±7.12	132.87±13.35	15.79±1.23	14.35±1.71
6H add	93.68±0.83	86.13±3.62	208.47±9.86	144.67±15.6	15.01±1.51	13.75±0.87

As salt stress not only has an ionic effect but also lead osmotic stress the water balance of plants is changed during salt treatment, as a result of which the RWC of leaves decreases. Except the 4H add salt stress causes a reduction in RWC all of the examined genotypes (Table 1). Parallel with the decrease in water content the initially high stomatal conductance ( $g_s$ ) decreased significantly in wheat lines, Betzes and 4H add. Although the decrease in  $g_s$  was moderate in 6H and 7H add the water content remained at a level similar to the Akom.

As the stomata play an important role in the regulation of transpirational water loss during salt stress, the primary physiological effect of the salinity is the inhibition of photosynthetic CO<sub>2</sub> fixation partly by means of stomatal closure (Centritto et al. 2003). A decreased substantially as  $g_s$  fell in Akom, Mv9kr1, Betzes and 4H add at 200 mmol L<sup>-1</sup> NaCl (Table 1). At this level of salinity, however, A and  $g_s$  only decreased slightly in Manas, 6H add and 7 H add. These results indicate that 7 H add like the parental barley cv. Manas, was able to retain its CO<sub>2</sub> fixation rate during salt stress with relatively high  $g_s$ . Barley is regarded as being more salt tolerant than bread wheat and the role of 7H addition in modifying the Na<sup>+</sup> and K<sup>+</sup> contents in leaves during salt stress has also been described (Colmer et al. 2006). In connection with this, a previous study has reported that photosynthetic rate was unaffected in a salt resistant common centauray in this salinity range (Siler et al. 2007). Moreover, it is thought that the estimation of net assimilation rate is could be a good tool for discriminating between salt tolerant and non-tolerant cultivars (Belkhodja et al. 1999). It is interesting that 6H add seems also tolerant to the salt stress in spite of the salt sensitivity of its parental barley genotypes (Betzes).

Our results seem to suggest that the newly-developed 7H Akom-Manas addition line is able to maintain at a satisfactory level its photosynthetic activity with a low level of water loss under moderate salt stress. Consequently, the better tolerance of photosynthetic parameters to moderate salinity of parental barley cv. Manas might be manifested in this plant.

## Acknowledgment

We thank J. Prónay and B. Háló for technical assistance. This work was supported by the OTKA A grants K 75466, K 75381 and the Generation Challenge Program SP3 G4007.23. A part of this study was presented on the 10<sup>th</sup> Congress of the Hungarian Society for Plant Biology, August 31 - September 2, 2011, Szeged, Hungary.

## Abbreviations used

A, net CO<sub>2</sub> assimilation rate; Akom, Asekaze komugi; 4H add, 4H Asakaze komugi-Manas addition line; 6H add, 6H Mv9kr1-Betzes-Seneca addition line 7H add, 7H Asakaze komugi-Manas addition line;  $g_s$ , stomatal conductance; RWC, relative water content.

## References

- Bagci SA (2003) Determination of the salt tolerance of some barley genotypes and the characteristics affecting tolerance. *Turk J Agric For* 27:253-260.
- Belkhodja R, Morales F, Abadía A, Medrano H, Abadía J (1999) Effects of salinity on chlorophyll fluorescence and photosynthesis of barley (*Hordeum vulgare* L.) grown under a triple-line-source sprinkler system in the field. *Photosynthetica* 36:375-387.
- Centritto M, Loreto F, Chartzoulakis K (2003) The use of low [CO<sub>2</sub>] to estimate diffusional and non-diffusional limitations of Photosynthetic capacity of salt stressed olive saplings. *Plant Cell Environ* 26:585-594.
- Colmer TD, Flowers TJ, Munns R (2006) Use of wild relatives to improve salt tolerance in wheat. *J Exp Bot Salinity Special Issue*:1-20.
- Colmer TD, Munns R, Flowers TJ (2005) Improving salt tolerance of wheat and barley: future prospects. *Aust J Exp Agr* 45:1425-1443.
- Dulai S, Molnár I, Háló B, Molnár-Láng M (2010) Photosynthesis in 7h wheat/barley 'Asakaze komugi'/'Manas' addition line during salt stress. *Acta Agr Hung* 58:367-376.
- Kaiser WM, Weber H, Sauer M (1983) Photosynthetic capacity, osmotic response and solute content of leaves and chloroplasts from *Spinacia oleracea* under salt stress. *Z Pflanzenphysiol* 113:15-27.
- Molnár I, Linc G, Dulai S, Nagy ED, Molnár-Láng M (2007):The compensation ability of chromosome 4H for 4D in response to drought stress investigated in newly developed wheat-barley 4H(4D) disomic substitution line *Plant Breeding* 126:369-374.

*Photosynthesis and salt stress in wheat/barley lines*

- Molnár-Láng M, Linc G, Logojan A, Sutka J (2000) Production and meiotic pairing behaviour of new hybrids of winter wheat (*Triticum aestivum*) × winter barley (*Hordeum vulgare*). *Genome* 43:1045-1054.
- Siler B, Misić D, Filipović D, Popović Z, Cvetic T, Mijović A (2007) Effects of salinity on in vitro growth and photosynthesis of common centaury (*Centaurium erythraea* Rafn.). *Arch Biol Sci* 59:129-134.
- von Caemmerer S, Farquhar GD (1981) Some relationships between the biochemistry of photosynthesis and the gas exchange of leaves. *Planta* 153:376-387.