

New Descriptions of the Lovász Number, and the Weak Sandwich Theorem

Miklós Ujvári*

Abstract

In 1979, L. Lovász introduced the concept of an orthonormal representation of a graph, and also a related value, now popularly known as the Lovász number of the graph. One of the remarkable properties of the Lovász number is that it lies sandwiched between the stability number of the graph and the chromatic number of the complementary graph. This fact is called the sandwich theorem.

In this paper, using new descriptions of the Lovász number and linear algebraic lemmas we give three proofs for a weaker version of the sandwich theorem.

Keywords: Lovász number, weak sandwich theorem

1 Introduction

From the several remarkable properties of the Lovász number of a graph we mention here only the sandwich theorem: the Lovász number lies ‘sandwiched’ between the stability number, and the chromatic number of the complementary graph. A weaker form of this sandwich theorem will be derived here using new descriptions of the Lovász number. This weak sandwich theorem is the immediate consequence of the sandwich theorem, Brooks’ theorem (concerning an upper bound on the chromatic number), and the counterpart of Brooks’ theorem (concerning a lower bound on the stability number). In this paper our aim is to give more direct proofs.

We begin this paper with stating the above-mentioned results. First we fix some notation. Let $n \in \mathcal{N}$, and let $G = (V(G), E(G))$ be an undirected graph, with vertex set $V(G) = \{1, \dots, n\}$, and with edge set $E(G) \subseteq \{\{i, j\} : i \neq j\}$. The complementary graph will be denoted by \overline{G} . Thus $\overline{G} = (V(\overline{G}), E(\overline{G}))$ where $V(\overline{G}) = V(G)$ and $E(\overline{G}) = \{\{i, j\} \subseteq V(G) : i \neq j, \{i, j\} \notin E(G)\}$.

Let us define an *orthonormal representation* of the graph G (shortly, o.r. of G) as a system of vectors $a_1, \dots, a_n \in \mathcal{R}^m$ for some $m \in \mathcal{N}$, satisfying

$$a_i^T a_i = 1 \quad (i = 1, \dots, n), \quad a_i^T a_j = 0 \quad (\{i, j\} \in E(\overline{G})).$$

*H-2600 Vác, Szent János utca 1., Hungary. E-mail: ujvarim@cs.elte.hu

In the seminal work [8] L. Lovász introduced the following number, $\vartheta(G)$, now popularly known as the *Lovász number* of the graph G ([7]):

$$\vartheta(G) := \inf \left\{ \max_{1 \leq i \leq n} \frac{1}{(a_i a_i^T)_{11}} : a_1, \dots, a_n \text{ o.r. of } G \right\}.$$

(Here $(a_i a_i^T)_{11}$ denotes the upper left corner element of the matrix $a_i a_i^T$, that is the square of the first element of the vector a_i , and though not emphasized in the definition of $\vartheta(G)$, we suppose that $(a_i a_i^T)_{11} \neq 0$ for all $i \in V(G)$.)

By Lemma 3 in [8], the Lovász number $\vartheta(G)$ is an upper bound for the stability number $\alpha(G)$, the maximum cardinality of the (so-called stable) sets $S \subseteq V(G)$ such that $\{i, j\} \subseteq S$ implies $\{i, j\} \notin E(G)$. Moreover, by Theorem 11 in [8] if there exists an orthonormal representation of the graph G with vectors a_i in \mathcal{R}^m then $\vartheta(G) \leq m$. Specially, $\vartheta(G)$ is at most the chromatic number of the complementary graph, $\chi(\overline{G})$, where the chromatic number of a graph is the minimal number of stable sets covering the vertex set of the graph. Hence (see [8])

$$\alpha(G) \leq \vartheta(G) \leq \chi(\overline{G}),$$

a fact known as the sandwich theorem (see [7]).

The Lovász number can also be defined via orthonormal representations of the complementary graph: it is shown in [8] that $\vartheta(G) = \vartheta'(G)$ where the number $\vartheta'(G)$ is defined as

$$\vartheta'(G) := \sup \left\{ \sum_{i=1}^n (b_i b_i^T)_{11} : b_1, \dots, b_n \text{ o.r. of } \overline{G} \right\}.$$

(We remark that here the values $(b_i b_i^T)_{11}$ are allowed to be zero.) The proof of the equality $\vartheta(G) = \vartheta'(G)$ relies on strong duality between Slater-regular primal-dual semidefinite programs equivalent with the programs defining $\vartheta(G)$ and $\vartheta'(G)$, respectively. (See [8], [10] or [15] for the equivalency results; and, for example, [16], [17] for the duality results.) As a consequence of the sandwich theorem and the equality between the values $\vartheta(G)$ and $\vartheta'(G)$ we have

$$\alpha(G) \leq \vartheta'(G) \leq \chi(\overline{G}),$$

a fact that can also be derived easily from the definition of $\vartheta'(G)$.

For $i \in V(G)$ let $N(i)$ denote the set of vertices $j \in V(G)$ such that $\{i, j\} \in E(G)$. Let us denote by d_i the cardinality of the set $N(i)$, and let d_{\max} denote the maximum of the values d_i ($i \in V(G)$). We define similarly $\overline{N}(i)$, \overline{d}_i and \overline{d}_{\max} for the complementary graph \overline{G} instead of G .

The following theorem is well-known (see for example [9]):

Theorem 1.1. (*Brooks*) *For any graph G , the chromatic number $\chi(G)$ is at most $d_{\max} + 1$, with equality for a connected graph G if and only if the graph is a clique or an odd cycle.*

As a corollary of Theorem 1.1 and the sandwich theorem we obtain

Corollary 1.1. *The value $\vartheta(G)$ ($\vartheta'(G)$ also) is at most $\bar{d}_{\max} + 1$.*

The counterpart of the Brooks' theorem can be found for example in [1]. For further lower bounds on the stability number, see [3], [18].

Theorem 1.2. *(Caro-Wei) For any graph G , the stability number $\alpha(G)$ is at least $\sum_{i \in V(G)} 1/(d_i + 1)$, with equality if and only if the graph G is the disjoint union of cliques.*

Similarly as in the case of Theorem 1.1 we have the following corollary:

Corollary 1.2. *The value $\vartheta'(G)$ ($\vartheta(G)$ also) is at least $\sum_{i \in V(G)} 1/(d_i + 1)$.*

We will call the results described in Corollaries 1.1 and 1.2 together the weak sandwich theorem. In Sections 2 and 3 we give two proofs for this theorem using linear algebraic lemmas and new descriptions of the Lovász number. In Section 4 we present a new, elementary graph theoretical proof for Theorem 1.2 (which is a derandomization of the original proof) thus obtaining a third proof for the weak sandwich theorem.

2 First proof for the weak sandwich theorem

In the first proof of the weak sandwich theorem we will need the following lemma, implicit in the proof of Theorem 3 in [8]:

Lemma 2.1. *Let PSD denote the set of n by n real symmetric positive semidefinite matrices. Let P denote the following set of matrices:*

$$P := \left\{ \left(\frac{a_i^T a_j}{e_1^T a_i \cdot e_1^T a_j} - 1 \right) \mid \begin{array}{l} m \in \mathcal{N}; a_i \in \mathcal{R}^m \ (1 \leq i \leq n); \\ a_i^T a_i = 1 \ (1 \leq i \leq n) \end{array} \right\}.$$

Then $\text{PSD} = P$. (Here e_1 denotes the first column vector of the identity matrix I . Though not emphasized in the definition of the set P , we suppose that the vectors a_i have nonzero first coordinates, that is $e_1^T a_i \neq 0$ for $i = 1, \dots, n$.)

Proof. First we will prove the inclusion $P \subseteq \text{PSD}$. Let a_1, \dots, a_n be unit vectors. Then the vectors $a_i \cdot (e_1^T a_i)^{-1}$ can be written as $(1, x_i^T)^T$ with appropriate vectors x_i . We have

$$\left(\frac{a_i^T a_j}{e_1^T a_i \cdot e_1^T a_j} - 1 \right) = (x_i^T x_j) \in \text{PSD}.$$

Thus the elements of the set P are positive semidefinite.

To prove the reverse inclusion $\text{PSD} \subseteq P$, let X be a positive semidefinite matrix. Then, there exist vectors x_i such that $X = (x_i^T x_j)$. Let $a_i := \lambda_i (1, x_i^T)^T$ where

the constants λ_i are chosen appropriately so that $a_i^T a_i = 1$ holds. With these definitions we have

$$X = (x_i^T x_j) = ((1, x_i^T)(1, x_j^T)^T - 1) = \left(\frac{a_i^T a_j}{e_1^T a_i \cdot e_1^T a_j} - 1 \right).$$

Thus $X \in P$, which was to be shown. □

From Lemma 2.1 follows immediately that the program defining $\vartheta(G)$ and the following program are equivalent:

$$\inf \max_{1 \leq i \leq n} x_{ii} + 1, x_{ij} = -1 \ (\{i, j\} \in E(\overline{G})), X \in \text{PSD}. \tag{1}$$

(We remark that program (1) in an equivalent form was studied previously by Meurdesoif, see program $(\mathcal{P}_{\mathcal{L}})$ in [11].) We can see

Theorem 2.1. *The optimal value of program (1) is equal to $\vartheta(G)$, and it is attained.* □

Now, let X be the following matrix:

$$X := (x_{ij}), \text{ where } x_{ij} := \begin{cases} \overline{d}_i, & \text{if } i = j, \\ 0, & \text{if } \{i, j\} \in E(G), \\ -1, & \text{if } \{i, j\} \in E(\overline{G}). \end{cases}$$

Then $x_{ii} \geq \sum_{i \neq j} |x_{ij}|$ holds for $1 \leq i \leq n$, so the matrix X is positive semidefinite by the Gerschgorin's disc theorem, see [14]. (We can also use the fact that X is the Laplacian matrix corresponding to the adjacency matrix of \overline{G} , see [16] for another application of the Laplacian matrix.) Moreover, the matrix X is a feasible solution of program (1), with corresponding value $\overline{d}_{\max} + 1$. Thus we have $\vartheta(G) \leq \overline{d}_{\max} + 1$, and Corollary 1.1 is proved. □

Similarly on the dual side we can apply the variable transformation described in Lemma 2.1 to the program defining $\vartheta'(G)$. This way we obtain the following program:

$$\sup \sum_{i=1}^n \frac{1}{y_{ii} + 1}, y_{ij} = -1 \ (\{i, j\} \in E(G)), Y \in \text{PSD}. \tag{2}$$

The optimal value of program (2) is a lower bound of $\vartheta'(G)$, as when writing program (2) we considered only the representations (b_i) where the vectors b_i had nonzero first coordinates. From these considerations Corollary 1.2 follows similarly as in the case above Corollary 1.1. □

We remark that the program defining $\vartheta'(G)$, and the program (2) are not equivalent generally. In fact, let G_0 be the cherry graph:

$$G_0 := (\{1, 2, 3\}, \{\{1, 2\}, \{1, 3\}\}).$$

Then $\vartheta'(G_0) = 2$ by the sandwich theorem, but the program (2) has no feasible solution with corresponding value 2. Otherwise there would exist

$$Y = \begin{pmatrix} x & -1 & -1 \\ -1 & y & w \\ -1 & w & z \end{pmatrix} \in \text{PSD}$$

such that

$$\frac{1}{x+1} + \frac{1}{y+1} + \frac{1}{z+1} = 2.$$

All the principal submatrices of the positive semidefinite matrix Y have nonnegative determinants, see [14]. Hence, $xy, xz \geq 1$, $x, y, z > 0$, and

$$x = \frac{1 - yz}{2yz + y + z}$$

would hold. From these relations $(1 - yz)y \geq 2yz + y + z$, that is $0 \geq z(y + 1)^2$ would follow, which is a contradiction. This contradiction shows that there exist graphs such that in every optimal orthonormal representation (b_i) there exist at least one vector b_i with zero first coordinate.

In the next two propositions we describe two lower bounds for the optimal value of program (2).

Proposition 2.1. *The optimal value of program (2) is at least $n/\vartheta(\overline{G})$.*

Proof. Let $\varepsilon \geq 0$, and let $X = X(\varepsilon) \in \mathcal{R}^{n \times n}$ be a feasible solution of program (1) with \overline{G} instead of G , such that

$$\max_{1 \leq i \leq n} x_{ii} + 1 \leq \vartheta(\overline{G}) + \varepsilon.$$

Then the matrix X is a feasible solution of program (2), and

$$n = \sum_{i=1}^n \frac{1}{x_{ii} + 1} \cdot (x_{ii} + 1) \leq \max_{1 \leq i \leq n} (x_{ii} + 1) \cdot \sum_{i=1}^n \frac{1}{x_{ii} + 1}.$$

We can see that $n/(\vartheta(\overline{G}) + \varepsilon)$ is a lower bound for the optimal value of program (2), and $\varepsilon \rightarrow 0$ (or $\varepsilon = 0$) gives the statement. \square

Proposition 2.2. *The optimal value of program (2) is at least $\alpha(G)$.*

Proof. Let $S \subseteq V(G)$ be a stable set with cardinality $\alpha(G)$, and let $\varepsilon > 0$. Let us define the matrix $Y = Y(\varepsilon) \in \mathcal{R}^{n \times n}$ the following way:

$$Y := (y_{ij}), \text{ where } y_{ij} := \begin{cases} \varepsilon, & \text{if } i = j \in S, \\ 0, & \text{if } i, j \in S, i \neq j, \\ \lambda, & \text{if } i = j \notin S, \\ -1 & \text{otherwise.} \end{cases}$$

Here let $\lambda = \lambda(\varepsilon) \in \mathcal{R}$ be the minimum number such that Y is positive semidefinite, that is

$$\lambda := \left(1 + \frac{\alpha(G)}{\varepsilon}\right) \cdot (n - \alpha(G)) - 1.$$

(Schur complements [12] can be used to determine λ .) Then Y is a feasible solution of program (2). It can be easily seen that the corresponding value increases to $\alpha(G)$ while $\varepsilon > 0$ decreases to 0. \square

From Propositions 2.1 and 2.2 equality between the optimal value of program (2) and $\vartheta'(G)$ follows:

- for vertex-transitive graphs where the lower bound $n/\vartheta(\overline{G})$ (Proposition 2.1) and the upper bound $\vartheta(G)$ are equal, see Theorem 8 in [8];
- for perfect graphs where the lower bound $\alpha(G)$ (Proposition 2.2) and the upper bound $\vartheta(G)$ are equal by the sandwich theorem and the perfect graph theorem [9].

Note that in the case of vertex-transitive graphs the optimal value of program (2) is attained, while in the case of perfect graphs non-attainment is possible.

Equality holds in the general case as well:

Theorem 2.2. *The optimal value of program (2) and $\vartheta'(G)$ are equal.*

Proof. Let us denote by $\text{TH}(G)$ the set of vectors $x = (x_i) \in \mathcal{R}^n$ for which there exists a matrix $Z = (z_{ij}) \in \mathcal{R}^{n \times n}$ satisfying both

$$\begin{pmatrix} 1 & x^T \\ x & Z \end{pmatrix} \in \text{PSD}$$

and

$$z_{ii} = x_i \quad (i = 1, \dots, n), \quad z_{ij} = 0 \quad (\{i, j\} \in E(G)).$$

It can be shown (see Corollary 29 in [7]) that $\text{TH}(G)$ can be described alternatively as the set of vectors $x = (x_i) \in \mathcal{R}^n$ such that

$$x_i = (e_1^T b_i)^2 \quad (i = 1, \dots, n)$$

for some (b_i) orthonormal representation of the complementary graph \overline{G} .

Let $\text{TH}_+(G)$ denote the set of positive vectors of $\text{TH}(G)$. Then $\text{TH}_+(G)$ is a convex set (as $\text{TH}(G)$ is a convex set), and it is nonempty (as every graph can be represented by vectors with nonzero first elements). From this observation easily follows that

$$\text{TH}_+(G) \subseteq \text{TH}(G) \subseteq \text{cl TH}_+(G),$$

where cl denotes closure. Consequently we obtain the same value optimizing any linear function over $\text{TH}(G)$ and $\text{TH}_+(G)$; which is exactly the statement, for the linear function $(x_1, \dots, x_n) \mapsto \sum_i x_i$. \square

3 Second proof for the weak sandwich theorem

In this section we give an alternative proof for the weak sandwich theorem using a completely different technique than the one used in the previous section.

Let $\sigma(n)$ denote the number of integers s in the range $0 < s < n$ such that $s \equiv 0, 1, 2$ or $4 \pmod{8}$. For small values of n , the value $\sigma(n)$ can be read out from the following table:

Table 1: The values $\sigma(n)$ for $n = 1, \dots, 16$.

n	1	2	3,4	5,6,7,8
$\sigma(n)$	0	1	2	3
n	9	10	11,12	13,14,15,16
$\sigma(n)$	4	5	6	7

The table can be continued in a similar manner for larger values of n . With this notation the following combinatorial lemma holds:

Lemma 3.1. *If $n \geq 2$ then there exist n of $\sigma(n)$ -letter words made up from the letters a, b, c, d such that the number of letter-pairs (a, b) and (c, d) on the same position in any two of the words is altogether odd. (For example in the words “aa” and “cb” there is only one such letter-pair: (a, b) , on the second position.)*

Proof. For the values $2 \leq n \leq 9$ the following word-sets have the desired property:

$$\begin{aligned}
 n = 2, \sigma(n) = 1 : & \quad a, b \\
 n = 3 \text{ or } 4, \sigma(n) = 2 : & \quad \text{any } n \text{ words from the word-set } aa, cb, ba, db \\
 n = 5, 6, 7 \text{ or } 8, \sigma(n) = 3 : & \quad \text{any } n \text{ words from the word-set} \\
 & \quad aaa, ccb, cba, cdb, baa, dab, dbc, dbd \\
 n = 9, \sigma(n) = 4 : & \quad aaaa, accb, acba, acdb, abaa \\
 & \quad adab, adbc, cdbd, ddbd.
 \end{aligned}$$

For larger values of n we can use the following induction argument. Let us denote by S_1, \dots, S_9 the words defined above in the case $n = 9$. Suppose that for some n we have appropriate $\sigma(n)$ -letter words T_1, \dots, T_n . Then the word-set

$$S_1 \& T_1, \dots, S_9 \& T_1, bdbd \& T_2, \dots, bdbd \& T_n,$$

where $\&$ denotes concatenation, is made up of $n + 8$ of $(\sigma(n) + 4)$ -letter words, and it has the desired property, too. Thus the statement in the lemma is dealt with for all the values of n . \square

Now, let

$$A := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, B := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, C := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, D := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

These matrices are orthogonal, furthermore from the matrix set

$$A^T B, A^T C, A^T D, B^T C, B^T D, C^T D$$

the matrices $A^T B$ and $C^T D$ are skew-symmetric, the others are symmetric. Given a word made up of the letters $a, b, c,$ and d we can define a matrix by Kronecker-multiplying the corresponding matrices: for example the word “dbc” is transformed into the 8 by 8 matrix $D \otimes B \otimes C$ where \otimes denotes Kronecker product. (The *Kronecker product* of two matrices $X = (x_{ij}), Y$ is the block matrix $X \otimes Y := (x_{ij} \cdot Y)$, see for example [12].) The matrices obtained this way are orthogonal, as they are the Kronecker products of orthogonal matrices.

Using this construction, from Lemma 3.1 immediately follows

Lemma 3.2. *If $m = 2^{\sigma(n)}$ then there exist m by m orthogonal matrices C_1, \dots, C_n such that for each $i \neq j$, the matrix $C_i^T C_j$ is skew-symmetric.*

Proof. Transform a word-set with the properties described in Lemma 3.1 into a matrix-set using the construction described before Lemma 3.2. We claim that this matrix-set meets the requirements. For example consider the matrix-set

$$A \otimes A, C \otimes B, B \otimes A, D \otimes B.$$

As we have noted already, these m by m matrices are orthogonal. On the other hand,

$$\begin{aligned} (A \otimes A)^T \cdot (C \otimes B) &= (A^T \otimes A^T) \cdot (C \otimes B) = (A^T C) \otimes (A^T B) = \\ &= (C^T A) \otimes (-B^T A) = -(C^T \otimes B^T) \cdot (A \otimes A) = -(C \otimes B)^T \cdot (A \otimes A), \end{aligned}$$

and similarly for the other matrix-pairs:

$$(A \otimes A)^T \cdot (B \otimes A) = -(B \otimes A)^T \cdot (A \otimes A) \dots \text{etc.}$$

In the general case similar argument can be applied, so the lemma is proved. \square

We remark that in [13] Radon proved that there exist m by m orthogonal matrices $\tilde{C}_1, \dots, \tilde{C}_n$ such that for each $i \neq j$ the matrix $\tilde{C}_i^T \tilde{C}_j$ is skew-symmetric if and only if $m \equiv 0 \pmod{2^{\sigma(n)}}$ (see also [6], [12]). The “if” part is an easy consequence of Lemma 3.2: just Kronecker-premultiply the C_i matrices with an identity matrix of appropriate dimension. For a similar proof of this part of Radon’s theorem, see [4].

We will need one further lemma, concerning new descriptions of the Lovász number. The idea is to represent the graph G with matrices instead of vectors. Let us define a *matrix orthonormal representation* of the graph G (shortly, m.o.r. of G) as a block matrix $(A_1, \dots, A_n) \in (\mathcal{R}^{\ell \times s})^n$ for some $\ell, s \in \mathcal{N}$, satisfying

$$A_i^T A_i = I \ (i = 1, \dots, n), \ A_i^T A_j = 0 \ (\{i, j\} \in E(\overline{G})).$$

Then, let us define

$$\hat{\vartheta}(G) := \inf \left\{ \max_{1 \leq i \leq n} \frac{1}{(A_i A_i^T)_{11}} : (A_1, \dots, A_n) \text{ m.o.r. of } G \right\}$$

and

$$\check{\vartheta}(G) := \sup \left\{ \sum_{i=1}^n (B_i B_i^T)_{11} : (B_1, \dots, B_n) \text{ m.o.r. of } \overline{G} \right\}.$$

It is obvious that $\hat{\vartheta}(G) \leq \vartheta(G)$ and $\vartheta'(G) \leq \check{\vartheta}(G)$. Equalities here follow from Lemma 3.3.

Lemma 3.3. *With the above definitions the inequality $\check{\vartheta}(G) \leq \hat{\vartheta}(G)$ holds.*

Proof. We adapt the proof of Lemma 4 in [8].

Let (A_1, \dots, A_n) and (B_1, \dots, B_n) be matrix orthonormal representations of G and \overline{G} , respectively. Then,

$$(A_i \otimes B_i)^T \cdot (A_j \otimes B_j) = (A_i^T A_j) \otimes (B_i^T B_j) = 0 \quad (1 \leq i, j \leq n; i \neq j).$$

Thus the column vectors of the matrices $A_i \otimes B_i$ ($1 \leq i \leq n$) altogether form an orthonormal system. Hence,

$$\sum_{i=1}^n ((A_i \otimes B_i)(A_i \otimes B_i)^T)_{11} \leq 1,$$

which can be rewritten as

$$\sum_{i=1}^n (A_i A_i^T)_{11} \cdot (B_i B_i^T)_{11} \leq 1.$$

From this inequality

$$\min_{1 \leq i \leq n} (A_i A_i^T)_{11} \cdot \sum_{i=1}^n (B_i B_i^T)_{11} \leq 1$$

follows, and so

$$\sum_{i=1}^n (B_i B_i^T)_{11} \leq \max_{1 \leq i \leq n} \frac{1}{(A_i A_i^T)_{11}}$$

holds. We can see that $\check{\vartheta}(G) \leq \hat{\vartheta}(G)$, and the proof of the lemma is finished. \square

Lemma 3.3, together with the equality $\vartheta(G) = \vartheta'(G)$, gives

Theorem 3.1. *The values $\hat{\vartheta}(G)$ and $\check{\vartheta}(G)$ are equal to $\vartheta(G)$ ($\vartheta'(G)$ also), and are attained. \square*

The weak sandwich theorem is an easy consequence of Lemmas 3.2 and 3.3. Let C_1, \dots, C_n be m by m orthogonal matrices with the property described in Lemma 3.2. Let us define the matrices A_1, \dots, A_n the following way: the matrix will be $(1 + \bar{e})m$ by m where \bar{e} denotes the cardinality of $E(\overline{G})$. The first m by m block in A_i is $\alpha_i C_i$ where

$$\alpha_i := \frac{1}{\sqrt{\bar{e} + 1}}.$$

The further m by m blocks correspond to the edges of the complementary graph \overline{G} : let the block corresponding to the edge $\{i, j\}$ be $\alpha_i C_j$ in A_i , $\alpha_j C_i$ in A_j , and the zero matrix otherwise. Then, (A_1, \dots, A_n) is a matrix orthonormal representation of G , so

$$\max_{1 \leq i \leq n} \frac{1}{(A_i A_i^T)_{11}} \geq \hat{\vartheta}(G).$$

On the other hand,

$$\max_{1 \leq i \leq n} \frac{1}{(A_i A_i^T)_{11}} = \max_{1 \leq i \leq n} \frac{\overline{d}_i + 1}{(C_i C_i^T)_{11}} = \overline{d}_{\max} + 1$$

(note that the matrices C_i are orthogonal so the matrix $C_i C_i^T$ is the identity matrix). We obtained $\overline{d}_{\max} + 1 \geq \hat{\vartheta}(G)$. Similar construction on the dual side shows that $\sum_{i \in V(G)} 1/(d_i + 1) \leq \check{\vartheta}(G)$. The weak sandwich theorem now follows from Lemma 3.3 and the obvious inequalities $\check{\vartheta}(G) \geq \vartheta'(G)$, $\hat{\vartheta}(G) \leq \vartheta(G)$. \square

Note that instead of the matrices C_i in the above construction we can also use matrices D_i with the following properties: the matrices D_i are orthogonal; the matrices $D_i^T D_j$ are symmetric and have zero trace ($i \neq j$). (The only change is that the block corresponding to the edge $\{i, j\}$ is $\alpha_i D_j$ in A_i and $-\alpha_j D_i$ in A_j .)

It is an open problem to characterize the numbers m such that there exist m by m matrices D_1, \dots, D_n with the properties described above; but any power of 2, greater than or equal to n meets the requirements: n words of the same length $\log_2 m$, and made up from the letters a and d (or a and c) translate into appropriate matrices (see the proof of Lemma 3.2).

Using simultaneous diagonalization (see [12]), the open problem described above can be cast also in the following form: characterize the numbers (m, n) such that there exists a matrix $M \in \{\pm 1\}^{m \times n}$ such that $M^T M = mI$. This is a subproblem of the Hadamard's Maximum Determinant Problem (see [5]); the Hadamard conjecture in an equivalent form states that the (m, n) pairs satisfying the requirements are:

- $(m, 1)$ such that $m \geq 1$;
- $(m, 2)$ such that $m \geq 2$ is even;
- (m, n) such that $m \geq n$ and $m \equiv 0 \pmod{4}$.

4 New proof for the Caro-Wei theorem

In this section we will prove the Caro-Wei theorem (Theorem 1.2), the counterpart of Brooks' theorem (Theorem 1.1). We also describe the counterpart of Turán's theorem.

First we will show that

$$\alpha(G) \geq \sum_{i=1}^n \frac{1}{d_i + 1} \tag{3}$$

holds. We apply induction on the cardinality n of $V(G)$. In the case when $n = 1$, the statement is trivial; in what follows we will suppose that the number of vertices is $n > 1$ and that for graphs with smaller number of vertices the inequality (3) already holds. Note that we can suppose also that the graph G is α -critical (that is leaving out any edge the stability number becomes larger). In fact, otherwise delete edges from the graph while this operation does not change the stability number. In the end we get an α -critical graph, and the value on the right hand side of (3) became larger, while the value on the left hand side of (3) stayed the same. We can suppose also that the graph G is connected: if it has more than one components, then by induction the inequality (3) holds true for its components, and this implies the validity of (3) for the whole graph. Hence, it is enough to consider the case when the graph G is α -critical and connected.

Let v be a vertex of G such that $d_v = d_{\max}$. It is easy to prove that there exists a stable set of the size $\alpha(G)$ such that it does not contain the vertex v (see Exercise 8.12 in [9]). In fact, otherwise every maximum stable set in G contains the vertex v . As G is α -critical, so leaving out an edge $\{v, v'\}$, in the resulting graph there exists a stable set S' of the size $\alpha(G) + 1$ containing both v and v' . Then, $S' \setminus \{v\}$ is a maximum stable set in G ; contradicting the indirect assumption.

Let us denote by $G - v$ the graph with vertex-set $\{1, \dots, n\} \setminus \{v\}$, and with edge-set $\{\{i, j\} \in E(G) : i, j \neq v\}$. Then $\alpha(G - v) = \alpha(G)$. By induction, for the graph $G - v$ (3) holds, that is

$$\alpha(G - v) \geq \sum_{i \in N(v)} \frac{1}{d_i} + \sum_{i \notin N(v), i \neq v} \frac{1}{d_i + 1}. \tag{4}$$

As $d_v \geq d_i$ for all $i \in V(G)$, we have

$$\frac{1}{d_i} \geq \frac{1}{d_i + 1} + \frac{1}{d_v(d_v + 1)} \quad (i \in N(v)) \tag{5}$$

Writing this bound into (4) we obtain the following inequality:

$$\alpha(G - v) \geq \sum_{i \in N(v)} \frac{1}{d_i + 1} + \sum_{i \notin N(v), i \neq v} \frac{1}{d_i + 1} + \frac{1}{d_v + 1}.$$

As $\alpha(G - v) = \alpha(G)$, this inequality is in fact (3), and the first half of Theorem 1.2 is proved.

To prove the second half of the theorem we will show that if

$$\alpha(G) = \sum_{i=1}^n \frac{1}{d_i + 1} \tag{6}$$

holds then the graph G is the disjoint union of cliques (the other direction is obvious). Again we apply induction on n . Note that if (6) holds then the graph G is α -critical (otherwise G would have an edge such that after deleting this edge the stability number stays unchanged, while the value on the right hand side of (6)

becomes larger, contradicting (3)). We can suppose also that G is connected (if (6) holds then it holds for the components also). Thus it suffices to prove that if the graph G is α -critical and connected, furthermore (6) holds then G is a clique.

Let v be the same point as in the first half of the proof, and again consider the graph $G - v$. Let us denote by $\delta(G)$ the sum on the right hand side of (6). As we have seen it in the first half of the proof,

$$\alpha(G) = \alpha(G - v) \geq \delta(G - v) \geq \delta(G).$$

As now $\alpha(G) = \delta(G)$, we have equalities instead of inequalities, that is

$$\alpha(G) = \alpha(G - v) = \delta(G - v) = \delta(G).$$

It follows from the $\delta(G - v) = \delta(G)$ equality that $d_i = d_v$ ($i \in N(v)$) (as otherwise (5) would hold with strict inequality). Moreover by the $\alpha(G - v) = \delta(G - v)$ equality and by induction the graph $G - v$ is the disjoint union of cliques. As the graph G is connected, the set $N(v)$ intersects with all of these cliques. Let us choose one of the cliques, and a vertex $i \in N(v)$ from this clique. Then d_i equals d_v as well as the cardinality of the clique. Hence the components of $G - v$ all have the same cardinality d_v . Then $\alpha(G - v) = (n - 1)/d_v$. If the graph $G - v$ would have more than one components then we could chose from each component a vertex from $\overline{N}(v)$. These vertices together with the vertex v would constitute a stable set in G with cardinality larger than $\alpha(G - v)$. This would contradict the fact that $\alpha(G - v) = \alpha(G)$, so $G - v$ is a clique with cardinality d_v with vertices in $N(v)$. Thus the graph G is a clique, and the proof of the second half of Theorem 1.2 is finished also. \square

Note that the bound in Brooks' theorem, $\chi(G) \leq d_{\max} + 1$, is obvious (it can be proved using a simple greedy coloring algorithm), and together with (3) and the sandwich theorem gives

$$\sum_{i=1}^n \frac{1}{d_i + 1} \leq \alpha(G) \leq \vartheta(G), \vartheta'(G) \leq \chi(\overline{G}) \leq \overline{d}_{\max} + 1;$$

we obtained the third proof of the weak sandwich theorem.

We remark that Turán's theorem can be derived as a consequence of the Caro-Wei theorem, see [1]. Here the graph $T_{n,m}$ is defined as follows: Divide the vertex set $V(T_{n,m}) := \{1, \dots, n\}$ into m disjoint subsets $\overline{S}_1, \dots, \overline{S}_m$ such that the cardinality of \overline{S}_i and \overline{S}_j differ by at most one for each $i \neq j$. Then the edge set of the graph $T_{n,m}$ is

$$E(T_{n,m}) := \cup_{\ell=1}^m \{\{i, j\} \subseteq \overline{S}_\ell : i \neq j\}.$$

Corollary 4.1. (*Turán*) *Let G be a simple graph on n vertices with stability number $\alpha(G) \leq m$. Minimizing the number of the edges of G under these assumptions, the unique extremal graph is $T_{n,m}$.*

The following corollary describes the counterpart of Turán's theorem which in turn is a simple consequence of Brooks' theorem.

Corollary 4.2. *Let G be a simple graph on n vertices with chromatic number $\chi(G) \geq m$. Then the number of the edges of G is at least $m(m - 1)/2$. Equality holds if and only if G is the disjoint union of a clique and a stable set on m and $n - m$ vertices, respectively.*

Proof. It is well-known that the number of the edges of any simple graph G on n points is at least $\chi(G)(\chi(G) - 1)/2$, so it is enough to prove that if the number of the edges is $m(m - 1)/2$ and the chromatic number is m then G is isomorphic with the graph described in the statement.

Hence, we can suppose that the vertex set is the disjoint union of m stable sets with exactly one edge going between each two of them. Let us choose a connected component G' of the graph G such that its chromatic number is $\chi(G') = m$. By Brooks' theorem, then

$$m = \chi(G') \leq d_{\max}(G') + 1 \leq d_{\max}(G) + 1 \leq m,$$

and we can see that G' is a clique on m vertices; the statement is proved. \square

Finally, we mention an open problem. Wilf proved the following result (see [2]): the chromatic number $\chi(G)$ is at most $\alpha_{\max} + 1$ (where α_{\max} denotes the maximum eigenvalue of the adjacency matrix of G), with equality for a connected graph G if and only if the graph is a clique or an odd cycle. As $\alpha_{\max} \leq d_{\max}$ always holds (with equality for a connected graph if and only if the graph is regular), Wilf's theorem is stronger than Brooks' theorem. It would be interesting to see how Theorem 1.2 could be strengthened using spectral information. (The bound $n/(\alpha_{\max} + 1)$ [18] is not a strengthening of the Caro-Wei bound, as — using the convexity of the function

$$x \mapsto \frac{1}{d_G^T x + 1} \quad (0 \leq x \in \mathcal{R}^n), \quad d_G = (d_1, \dots, d_n)^T,$$

and Rayleigh's theorem [14] — it can be easily shown that

$$\frac{n}{\alpha_{\max} + 1} \leq \sum_{i=1}^n \frac{1}{d_i + 1}$$

holds, with equality if and only if the graph is regular.)

5 Conclusion

In this paper we presented a new proof for the counterpart of Brooks' theorem (the Caro-Wei theorem) concerning a simple lower bound on the stability number. As a consequence of the sandwich theorem, Brooks' theorem, and the Caro-Wei theorem we derived a weaker version of the sandwich theorem. For this weak sandwich theorem we gave another two, more direct proofs also, which are based on linear algebraic lemmas and new descriptions of the Lovász number.

Acknowledgements. I thank the two anonymous referees for their remarks that helped me to improve the presentation of the paper.

References

- [1] Alon, N. and Spencer, J. H. *The Probabilistic Method*. John Wiley & Sons, New York, 1992.
- [2] Cvetković, D. M., Doob, M., and Sachs, H. *Spectra of Graphs*. Academic Press, New York, 1979.
- [3] Edwards, C. S. and Elphick, C. H. Lower bounds for the clique and the chromatic numbers of a graph. *Discrete Applied Mathematics*, 5:51–64, 1983.
- [4] Geramita, A. V. and Pullman, N. J. A theorem of Hurwitz and Radon and orthogonal projective modules. *Proceedings of the American Mathematical Society*, 42(1):51–56, 1974.
- [5] Jablonszkij, SZ. V. and Lupanov, O. B., editors. *Diszkrét Matematika a Számítástudományban*. Műszaki Könyvkiadó, Budapest, 1980.
- [6] James, I. M. *The Topology of Stiefel Manifolds*. Cambridge University Press, Cambridge, 1976.
- [7] Knuth, D. The sandwich theorem. *Electronic Journal of Combinatorics*, 1:1–48, 1994.
- [8] Lovász, L. On the Shannon capacity of a graph. *IEEE Transactions on Information Theory*, IT-25:1–7, 1979.
- [9] Lovász, L. *Combinatorial Problems and Exercises*. Akadémiai Kiadó, Budapest, 1979.
- [10] Lovász, L. Semidefinite programs and combinatorial optimization. In Reed, B. A. and Sales, C. L., editors, *Recent Advances in Algorithms and Combinatorics*, CMS Books in Mathematics, Springer, pages 137–194, 2003.
- [11] Meurdesoif, P. Strengthening the Lovász $\theta(\overline{G})$ bound for graph coloring. *Mathematical Programming A*, 102:577–588, 2005.
- [12] Praszolov, V. V. *Lineáris Algebra*. Typotex Kiadó, Budapest, 2005.
- [13] Radon, J. Lineare scharen orthogonaler matrizen. *Abhandlungen aus dem Mathematischen Seminar der Hamburgischen Universität*, 1:1–14, 1923.
- [14] Strang, G. *Linear Algebra and its Applications*. Academic Press, New York, 1980.
- [15] Ujvári, M. *A Szemidefinit Programozás Alkalmazásai a Kombinatorikus Optimalizálásban*. ELTE Eötvös Kiadó, Budapest, 2001.
- [16] Ujvári, M. A note on the graph-bisection problem. *Pure Mathematics and Applications*, 12(1):119–130, 2002.

- [17] Ujvári, M. On a closedness theorem. *Pure Mathematics and Applications*, 15(4):469–486, 2006.
- [18] Wilf, H. S. Spectral bounds for the clique and independence numbers of graphs. *Journal of Combinatorial Theory B*, 40:113–117, 1986.

Received 29th November 2011