

## PARAGENESIS OF TYPOMORPHIC ACCESSORY MINERALS VS. TYPOLOGY OF GRANITIC ROCKS: EXAMPLES FROM WESTERN CARPATHIANS, SLOVAKIA

BROSKA, I.<sup>1\*</sup>, PETRÍK, I.<sup>1</sup> & UHER, P.<sup>2</sup>

<sup>1</sup> Geological Institute, Slovak Academy of Sciences, Dúbravská cesta 9, 840 05 Bratislava, Slovakia

<sup>2</sup> Department of Mineralogy and Petrology, Comenius University, Mlynská dolina G, 842 15 Bratislava, Slovakia

\* E-mail: [geolbros@savba.sk](mailto:geolbros@savba.sk)

Assemblages of accessory minerals are important criterion for the typological division of granitic rocks. On the example of the West-Carpathian granite suites have been shown that the most critical assemblages for granite classifications are the magmatic accessory paragenesis reflecting the primary character of former melts. The accessory minerals determining the character of primary melts are the typomorphic. Detailed study of accessory minerals in Variscan I- and S-type and post-Variscan Permian specialized S- and A-types granites in Tatric, Veporic and Gemic Units of the Western Carpathians enable us to characterize their differences and petrogenetic impact to origin of the suites (BROSKA *et al.*, 2011). Primarily relationship between *monazite* and *allanite* is important for the recognition of I- and S-type granitic rocks: *monazite*-(Ce) dominates in the S-type granites, on the other hand higher water and Ca activities stabilises *allanite*-(Ce) in the I-type granites. Exceptions represent more fractionated I-type granites where *monazite*-(Ce) is common. Although *monazite*-(Ce) in the hypersolvus A-type granites almost absent, in subsolvus granites occurs locally. Except *xenotime*-(Y), which may locally be abundant in S-type granites, A-type granitic rocks contain further Y-B-silicate phases (*gadolinite* and *hingganite*).

*Magnetite* as a typical mineral of the I-type granitoid paragenesis indicate higher oxidation level. In such rocks Ti-rich *magnetite* occurs first, which is in late-magmatic stage replaced by nearly pure *magnetite* in association with *titanite*. This is interpreted as result of late- to post-magmatic oxidation due to separation of fluid phase and following water dissociation. Another important basis for division is composition of *apatites* (hydroxylapatite to fluorapatite): low contents of Fe and Mn are typical of *apatites* from I-type granites, in contrast to S-type granite *apatites*, which are enriched in these elements. Similarly, *apatite* from A-type granites is commonly rich in Fe. The highest Mn contents accompanied by Sr are found in *apatites* from specialised S-type granites from the Gemic unit

*Zircon* composition and its morphology are the important markers for granite typology. Restite *zircon* holds many features of former granitic magma. A morphological boundary may be derived from comparison of *monazite/allanite* antagonistic relationship and *zircon* morphology based on I.T parameter equal 350; I.T < 350 indicates S-type granites, while I.T > 350 is characteristic of I-type granites, I.A parameter is close to 300

for both granitic types. A higher I.A parameter close to 400 indicates specialised S-type granites. The A-type granites have values in the range 650–700. Hypersolvus granites contain commonly *zircon* subtypes D and P5 with high I.T parameter around 700, whereas subsolvus A-type granites show lower I.T parameter, close to 300. Orthomagmatic *zircon*s show Zr/Hf<sub>wt</sub> ratio in S- and I-type granites roughly 35–45, late-magmatic *zircon*s in leucogranites have a lower ratio due to increase of Hf with differentiations. High Zr/Hf<sub>wt</sub> ratio (> 50), but low Y, REE, U, Th concentrations in early magmatic *zircon*s from hypersolvus A-type granites are in contrast to lower temperature subsolvus members. Similarly, highly fractionated S-type granites show Zr/Hf<sub>wt</sub> ratio under 30, and contents of P, Y, REE U, and Th are commonly ≥ 0.5 wt%.

*Tourmaline* supergroup minerals indicate increased boron and other volatile elements in the primary melt. *Schorl* to *foitite* occur in Permian, post-orogenic specialised S-type Gemic granites, locally are present also in some Veporic Permian S-type granites. Highly fractionated members of the specialised S-type granites contain *Nb-Ta oxide* minerals (mainly *columbite*-group minerals), *Nb-Ta rutile*, *cassiterite* and *ferberite*. Such mineralization typically occurs in greisenised granites. However, scarce *Nb-Ta rutile*, Ti-rich *ixiolite*, Fe-rich *columbite-tantalite*, and *ferrotapiolite* occur also in fractionated S-type leucogranites in the Tatric Unit. Moreover, some granitic *pegmatites* derived from S- and rarely I-type granitic magmas contain *beryl* and accessory minerals of *columbite*, rarely *tapiolite* and *wodginite* groups (Tatric Unit). This Nb-Ta-Sn-(Ti) suite is typical of granites-*pegmatites* of S- and I-type, in contrast to Y-REE-Ti-Nb-(Ta) suite [*fergusonite*-(beta)/*samaraskite*-(Y), *aeschynite/polycrase*-(Y), Nb-rich *rutile*?] in the hypersolvus A-type granites (Turčok, Gemic Unit). A special group of rare phosphates was found in Li-F-P topaz- and Li-mica-bearing granite from the Hnilec area (Gemic Unit) comprising *la-croixite*, *arrojadite*, *viitaniemiite*, *gorceixite* and *goyazite*.

### Reference

- BROSKA, I. PETRÍK, I. & UHER, P. (2011): Accessory minerals in the granitic rocks of the Western Carpathians. Veda Pub. (Bratislava), 250 pp, in press. (In Slovak with English summary)