



Energy decay of solutions for a wave equation with a constant weak delay and a weak internal feedback

Abbes Benaissa¹, Aissa Benguessoum¹ and Salim A. Messaoudi²

¹Laboratory of Analysis and Control of PDE's, Djillali Liabes University,
P. O. Box 89, Sidi Bel Abbes 22000, Algeria

²Department of Mathematics and Statistics, KFUPM, Dhahran 31261, Saudi Arabia

Received 2 September 2013, appeared 21 March 2014

Communicated by Paul Eloe

Abstract. In this paper, we consider the wave equation with a weak internal constant delay term:

$$u''(x, t) - \Delta_x u(x, t) + \mu_1(t) u'(x, t) + \mu_2(t) u'(x, t - \tau) = 0$$

in a bounded domain. Under appropriate conditions on μ_1 and μ_2 , we prove global existence of solutions by the Faedo–Galerkin method and establish a decay rate estimate for the energy using the multiplier method.

Keywords: nonlinear wave equation, delay term, decay rate, multiplier method.

2010 Mathematics Subject Classification: 35B40, 35L70.

1 Introduction

In this paper we investigate the decay properties of solutions for the initial boundary value problem for the linear wave equation of the form

$$\begin{cases} u''(x, t) - \Delta_x u(x, t) + \mu_1(t) u'(x, t) + \mu_2(t) u'(x, t - \tau) = 0 & \text{in } \Omega \times]0, +\infty[, \\ u(x, t) = 0 & \text{on } \Gamma \times]0, +\infty[, \\ u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x) & \text{on } \Omega, \\ u_t(x, t - \tau) = f_0(x, t - \tau) & \text{on } \Omega \times]0, \tau[, \end{cases} \quad (\text{P})$$

where Ω is a bounded domain in \mathbb{R}^n , $n \in \mathbb{N}^*$, with a smooth boundary $\partial\Omega = \Gamma$, $\tau > 0$ is a time delay and the initial data (u_0, u_1, f_0) belong to a suitable function space.

In absence of delay ($\mu_2 = 0$), the energy of problem (P) is exponentially decaying to zero provided that μ_1 is constant, see, for instance, [3, 4, 7, 8] and [12]. On the contrary, if $\mu_1 = 0$ and $\mu_2 > 0$ (a constant weight), that is, there exists only the internal delay, the system (P)

¹Corresponding author. Email: benaissa_abbes@yahoo.com

becomes unstable (see, for instance, [5]). In recent years, the PDEs with time delay effects have become an active area of research since they arise in many practical problems (see, for example, [1, 19]). In [5], it was shown that a small delay in a boundary control could turn a well-behaved hyperbolic system into a wild one and, therefore, delay becomes a source of instability. To stabilize a hyperbolic system involving input delay terms, additional control terms will be necessary (see [13, 15, 20]). For instance, the authors of [13] studied the wave equation with a linear internal damping term with constant delay ($\tau = \text{const}$ in the problem (P) and determined suitable relations between μ_1 and μ_2 , for which the stability or alternatively instability takes place. More precisely, they showed that the energy is exponentially stable if $\mu_2 < \mu_1$ and they also found a sequence of delays for which the corresponding solution of (P) will be unstable if $\mu_2 \geq \mu_1$. The main approach used in [13] is an observability inequality obtained with a Carleman estimate. The same results were obtained if both the damping and the delay are acting on the boundary. We also recall the result by Xu, Yung and Li [20], where the authors proved a result similar to the one in [13] for the one-space dimension by adopting the spectral analysis approach.

In [17], Nicaise, Pignotti and Valein extended the above result to higher space dimensions and established an exponential decay.

Our purpose in this paper is to give an energy decay estimate of the solution of problem (P) in the presence of a delay term with a weight depending on time. We use the Galerkin approximation scheme and the multiplier technique to prove our results.

2 Preliminaries and main results

First assume the following hypotheses:

(H1) $\mu_1: \mathbb{R}_+ \rightarrow]0, +\infty[$ is a non-increasing function of class $C^1(\mathbb{R}_+)$ satisfying

$$\left| \frac{\mu_1'(t)}{\mu_1(t)} \right| \leq M, \quad (2.1)$$

(H2) $\mu_2: \mathbb{R}_+ \rightarrow \mathbb{R}$ is a function of class $C^1(\mathbb{R}_+)$, which is not necessarily positive or monotone, such that

$$|\mu_2(t)| \leq \beta \mu_1(t), \quad (2.2)$$

$$|\mu_2'(t)| \leq \tilde{M} \mu_1(t), \quad (2.3)$$

for some $0 < \beta < 1$ and $\tilde{M} > 0$.

We now state a Lemma needed later.

Lemma 2.1 (Martinez [10]). *Let $E: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a non increasing function and $\phi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ an increasing C^1 function such that*

$$\phi(0) = 0 \quad \text{and} \quad \phi(t) \rightarrow +\infty \quad \text{as} \quad t \rightarrow +\infty.$$

Assume that there exist $\sigma > -1$ and $\omega > 0$ such that

$$\int_S^{+\infty} E^{1+\sigma}(t) \phi'(t) dt \leq \frac{1}{\omega} E^\sigma(0) E(S), \quad 0 \leq S < +\infty. \quad (2.4)$$

Then

$$E(t) = 0 \quad \forall t \geq \frac{E(0)^\sigma}{\omega|\sigma|}, \quad \text{if } -1 < \sigma < 0, \quad (2.5)$$

$$E(t) \leq E(0) \left(\frac{1 + \sigma}{1 + \omega\sigma\phi(t)} \right)^{\frac{1}{\sigma}} \forall t \geq 0, \quad \text{if } \sigma > 0, \quad (2.6)$$

$$E(t) \leq E(0)e^{1-\omega\phi(t)} \forall t \geq 0, \quad \text{if } \sigma = 0. \quad (2.7)$$

We introduce, as in [13], the new variable

$$z(x, \rho, t) = u_t(x, t - \tau\rho), \quad x \in \Omega, \rho \in (0, 1), \quad t > 0. \quad (2.8)$$

Then, we have

$$\tau z_t(x, \rho, t) + z_\rho(x, \rho, t) = 0, \quad \text{in } \Omega \times (0, 1) \times (0, +\infty). \quad (2.9)$$

Therefore, problem (P) takes the form:

$$\begin{cases} u''(x, t) - \Delta_x u(x, t) + \mu_1(t)u'(x, t) + \mu_2(t)z(x, 1, t) = 0, & x \in \Omega, t > 0, \\ \tau z_t(x, \rho, t) + z_\rho(x, \rho, t) = 0, & x \in \Omega, \rho \in (0, 1), t > 0, \\ u(x, t) = 0, & x \in \partial\Omega, t > 0, \\ z(x, 0, t) = u'(x, t), & x \in \Omega, t > 0, \\ u(x, 0) = u_0(x), u_t(x, 0) = u_1(x), & x \in \Omega, \\ z(x, \rho, 0) = f_0(x, -\tau\rho), & x \in \Omega, \rho \in (0, 1). \end{cases} \quad (2.10)$$

Let $\bar{\xi}$ be a positive constant such that

$$\tau\beta < \bar{\xi} < \tau(2 - \beta). \quad (2.11)$$

We define the energy of the solution by:

$$E(t) = \frac{1}{2} \|u'(t)\|_2^2 + \frac{1}{2} \|\nabla_x u(t)\|_2^2 + \frac{\bar{\xi}(t)}{2} \int_\Omega \int_0^1 z^2(x, \rho, t) d\rho dx, \quad (2.12)$$

where

$$\bar{\xi}(t) = \bar{\xi}\mu_1(t).$$

We have the following theorem.

Theorem 2.2. *Let $(u_0, u_1, f_0) \in (H^2(\Omega) \cap H_0^1(\Omega)) \times H_0^1(\Omega) \times H_0^1(\Omega; H^1(0, 1))$ satisfy the compatibility condition*

$$f_0(\cdot, 0) = u_1.$$

Assume that (H1) and (H2) hold. Then problem (P) admits a unique global weak solution

$$u \in L_{\text{loc}}^\infty((-\tau, \infty); H^2(\Omega) \cap H_0^1(\Omega)), u' \in L_{\text{loc}}^\infty((-\tau, \infty); H_0^1(\Omega)), u'' \in L_{\text{loc}}^\infty((-\tau, \infty); L^2(\Omega)).$$

Moreover, for some positive constants c, ω , we obtain the following decay property:

$$E(t) \leq cE(0)e^{-\omega \int_0^t \mu_1(s) ds}, \quad \forall t \geq 0. \quad (2.13)$$

Lemma 2.3. *Let (u, z) be a solution to the problem (2.10). Then, the energy functional defined by (2.12) satisfies*

$$\begin{aligned} E'(t) &\leq - \left(\mu_1(t) - \frac{\xi(t)}{2\tau} - \frac{|\mu_2(t)|}{2} \right) \|u_t(x, t)\|_2^2 - \left(\frac{\xi(t)}{2\tau} - \frac{|\mu_2(t)|}{2} \right) \|z(x, 1, t)\|_2^2 \\ &\leq 0. \end{aligned} \quad (2.14)$$

Proof. Multiplying the first equation in (2.10) by $u_t(x, t)$, integrating over Ω and using Green's identity, we obtain:

$$\begin{aligned} \frac{1}{2} \frac{\partial}{\partial t} \|u_t(x, t)\|_2^2 + \frac{1}{2} \frac{\partial}{\partial t} \|\nabla u(x, t)\|_2^2 \\ + \mu_1(t) \|u_t(x, t)\|_2^2 + \mu_2(t) \int_{\Omega} u_t(x, t - \tau) u_t(x, t) dx = 0. \end{aligned} \quad (2.15)$$

We multiply the second equation in (2.10) by $\xi(t)z$ and integrate over $\Omega \times (0, 1)$ to obtain:

$$\xi(t)\tau \int_{\Omega} \int_0^1 z_t(x, \rho, t) z(x, \rho, t) d\rho dx + \xi(t) \int_{\Omega} \int_0^1 z_{\rho}(x, \rho, t) z(x, \rho, t) d\rho dx = 0. \quad (2.16)$$

This yields

$$\frac{\xi(t)\tau}{2} \frac{d}{dt} \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx + \frac{\xi(t)}{2} \int_{\Omega} \int_0^1 \frac{\partial}{\partial \rho} z^2(x, \rho, t) d\rho dx = 0,$$

which gives

$$\begin{aligned} \frac{\tau}{2} \left[\frac{d}{dt} \left(\xi(t) \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx \right) - \xi'(t) \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx \right] \\ + \frac{\xi(t)}{2} \int_{\Omega} z^2(x, 1, t) dx - \frac{\xi(t)}{2} \int_{\Omega} u_t^2(x, t) dx = 0. \end{aligned}$$

Consequently,

$$\begin{aligned} \frac{\tau}{2} \frac{d}{dt} \left(\xi(t) \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx \right) \\ = \frac{\tau}{2} \xi'(t) \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx - \frac{\xi(t)}{2} \int_{\Omega} z^2(x, 1, t) dx + \frac{\xi(t)}{2} \int_{\Omega} u_t^2(x, t) dx. \end{aligned} \quad (2.17)$$

Combination of (2.15) and (2.17) leads to

$$\begin{aligned} \frac{1}{2} \frac{\partial}{\partial t} \left[\|u_t(x, t)\|_2^2 + \|\nabla u(x, t)\|_2^2 + \xi(t) \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx \right] \\ = -\mu_1(t) \|u_t(x, t)\|_2^2 - \mu_2(t) \int_{\Omega} z(x, 1, t) u_t(x, t) dx \\ + \frac{1}{2} \xi'(t) \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx - \frac{\xi(t)}{2\tau} \int_{\Omega} z^2(x, 1, t) dx + \frac{\xi(t)}{2\tau} \|u_t(x, t)\|_2^2. \end{aligned}$$

Recalling the definition of $E(t)$ in (2.12), we arrive at

$$\begin{aligned}
E'(t) &= - \left(\mu_1(t) - \frac{\bar{\xi}(t)}{2\tau} \right) \|u_t(x, t)\|_2^2 - \mu_2(t) \int_{\Omega} z(x, 1, t) u_t(x, t) dx \\
&\quad + \frac{1}{2} \bar{\xi}'(t) \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx - \frac{\bar{\xi}(t)}{2\tau} \int_{\Omega} z^2(x, 1, t) dx. \\
&\leq - \left(\mu_1(t) - \frac{\bar{\xi}(t)}{2\tau} \right) \|u_t(x, t)\|_2^2 - \mu_2(t) \int_{\Omega} z(x, 1, t) u_t(x, t) dx \\
&\quad - \frac{\bar{\xi}(t)}{2\tau} \int_{\Omega} z^2(x, 1, t) dx. \tag{2.18}
\end{aligned}$$

Due to Young's inequality, we have

$$\int_{\Omega} z(x, 1, t) u_t(x, t) dx \leq \frac{1}{2} \|u_t(x, t)\|_2^2 + \frac{1}{2} \|z(x, 1, t)\|_2^2. \tag{2.19}$$

Inserting (2.19) into (2.18), we obtain

$$\begin{aligned}
E'(t) &\leq - \left(\mu_1(t) - \frac{\bar{\xi}(t)}{2\tau} - \frac{|\mu_2(t)|}{2} \right) \|u_t(x, t)\|_2^2 - \left(\frac{\bar{\xi}(t)}{2\tau} - \frac{|\mu_2(t)|}{2} \right) \|z(x, 1, t)\|_2^2 \\
&\leq -\mu_1(t) \left(1 - \frac{\bar{\xi}}{2\tau} - \frac{\beta}{2} \right) \|u_t(x, t)\|_2^2 - \mu_1(t) \left(\frac{\bar{\xi}}{2\tau} - \frac{\beta}{2} \right) \|z(x, 1, t)\|_2^2 \leq 0. \tag{2.20}
\end{aligned}$$

This completes the proof of the lemma. \square

3 Global existence

Throughout this section we assume $u_0 \in H^2(\Omega) \cap H_0^1(\Omega)$ and $u_1 \in H_0^1(\Omega)$, $f_0 \in L^2(\Omega; H^1(0, 1))$.

We employ the Galerkin method to construct a global solution. Let $T > 0$ be fixed and denote by V_k the space generated by $\{w_1, w_2, \dots, w_k\}$ where the set $\{w_k, k \in \mathbb{N}\}$ is a basis of $H^2(\Omega) \cap H_0^1(\Omega)$.

Now, we define for $1 \leq j \leq k$ the sequence $\phi_j(x, \rho)$ as follows:

$$\phi_j(x, 0) = w_j.$$

Then, we may extend $\phi_j(x, 0)$ by $\phi_j(x, \rho)$ over $L^2(\Omega \times (0, 1))$ such that $(\phi_j)_j$ form a basis of $L^2(\Omega; H^1(0, 1))$ and denote by Z_k the space generated by $\{\phi_1, \phi_2, \dots, \phi_k\}$.

We construct approximate solutions (u_k, z_k) , $k = 1, 2, 3, \dots$, in the form

$$u_k(t) = \sum_{j=1}^k g_{jk}(t) w_j, \quad z_k(t) = \sum_{j=1}^k h_{jk}(t) \phi_j,$$

where g_{jk} and h_{jk} ($j = 1, 2, \dots, k$) are determined by the following system of ordinary differential equations:

$$\begin{cases} (u_k''(t), w_j) + (\nabla_x u_k(t), \nabla_x w_j) + \mu_1(t) (u_k'(t), w_j) + \mu_2(t) (z_k(\cdot, 1), w_j) = 0, \\ 1 \leq j \leq k, \\ z_k(x, 0, t) = u_k'(x, t), \end{cases} \tag{3.1}$$

associated with the initial conditions

$$u_k(0) = u_{0k} = \sum_{j=1}^k (u_0, w_j) w_j \rightarrow u_0 \text{ in } H^2(\Omega) \cap H_0^1(\Omega) \text{ as } k \rightarrow +\infty, \quad (3.2)$$

$$u'_k(0) = u_{1k} = \sum_{j=1}^k (u_1, w_j) w_j \rightarrow u_1 \text{ in } H_0^1(\Omega) \text{ as } k \rightarrow +\infty, \quad (3.3)$$

and

$$\begin{cases} (\tau z_{kt} + z_{k\rho}, \phi_j) = 0, \\ 1 \leq j \leq k, \end{cases} \quad (3.4)$$

$$z_k(\rho, 0) = z_{0k} = \sum_{j=1}^k (f_0, \phi_j) \phi_j \rightarrow f_0 \text{ in } L^2(\Omega; H^1(0, 1)) \text{ as } k \rightarrow +\infty. \quad (3.5)$$

By virtue of the theory of ordinary differential equations, the system (3.1)–(3.5) has a unique local solution which is extended to a maximal interval $[0, T_k[$ (with $0 < T_k \leq +\infty$) by Zorn lemma. Note that $u_k(t)$ is of class C^2 .

In the next step, we obtain a priori estimates for the solution of the system (3.1)–(3.5), so that it can be extended beyond $[0, T_k[$ to obtain a solution defined for all $t > 0$. Then, we utilize a standard compactness argument for the limiting procedure.

The first estimate. Since the sequences u_{0k} , u_{1k} and z_{0k} converge, then from (2.14) we can find a positive constant C independent of k such that

$$E_k(t) + \int_0^t a_1(s) \|u'_k(s)\|_2^2 ds + \int_0^t a_2(s) \|z_k(x, 1, s)\|_2^2 ds \leq E_k(0) \leq C, \quad (3.6)$$

where

$$E_k(t) = \frac{1}{2} \|u'_k(t)\|_2^2 + \frac{1}{2} \|\nabla_x u_k(t)\|_2^2 + \frac{\bar{\xi}(t)}{2} \int_{\Omega} \int_0^1 z_k^2(x, \rho, t) d\rho dx,$$

$$a_1(t) = \mu_1(t) \left(1 - \frac{\bar{\xi}}{2\tau} - \frac{\beta}{2} \right) \text{ and } a_2(t) = \mu_1(t) \left(\frac{\bar{\xi}}{2\tau} - \frac{\beta}{2} \right).$$

These estimates imply that the solution (u_k, z_k) exists globally in $[0, +\infty[$.

Estimate (3.6) yields

$$(u_k) \text{ is bounded in } L_{\text{loc}}^{\infty}(0, \infty; H_0^1(\Omega)), \quad (3.7)$$

$$(u'_k) \text{ is bounded in } L_{\text{loc}}^{\infty}(0, \infty; L^2(\Omega)), \quad (3.8)$$

$$\mu_1(t) (u'_k)^2(t) \text{ is bounded in } L^1(\Omega \times (0, T)), \quad (3.9)$$

$$\mu_1(t) (z_k^2(x, \rho, t)) \text{ is bounded in } L_{\text{loc}}^{\infty}(0, \infty; L^1(\Omega \times (0, 1))), \quad (3.10)$$

$$\mu_1(t) (z_k^2(x, 1, t)) \text{ is bounded in } L^1(\Omega \times (0, T)). \quad (3.11)$$

The second estimate. We first estimate $u''_k(0)$. Replacing w_j by $u''_k(t)$ in (3.1) and taking $t = 0$, we obtain:

$$\begin{aligned} \|u''_k(0)\|_2 &\leq \|\Delta_x u_{0k}\|_2 + \mu_1(0) \|u_{1k}\|_2 + |\mu_2(0)| \|z_{0k}\|_2 \\ &\leq \|\Delta_x u_0\|_2 + \mu_1(0) \|u_1\|_2 + |\mu_2(0)| \|z_0\|_2 \\ &\leq C. \end{aligned}$$

Differentiating (3.1) with respect to t , we get

$$(u_k''''(t) + \Delta_x u_k'(t) + \mu_1(t)u_k''(t) + \mu_1'(t)u_k'(t) + \mu_2(t)z_k'(1, t) + \mu_2'(t)z_k(1, t), w_j) = 0.$$

Multiplying by $g_{jk}''(t)$, summing over j from 1 to k , it follows that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} (\|u_k''(t)\|_2^2 + \|\nabla_x u_k'(t)\|_2^2) + \mu_1(t) \int_{\Omega} u_k''^2(t) dx + \mu_1'(t) \int_{\Omega} u_k''(t)u_k'(t) dx \\ + \mu_2(t) \int_{\Omega} u_k''(t)z_k'(x, 1, t) dx + \mu_2'(t) \int_{\Omega} u_k''(t)z_k(x, 1, t) dx = 0. \end{aligned} \quad (3.12)$$

Differentiating (3.4) with respect to t , we get

$$\left(\tau z_k''(t) + \frac{\partial}{\partial \rho} z_k', \phi_j \right) = 0.$$

Multiplying by $h_{jk}'(t)$, summing over j from 1 to k , it follows that

$$\frac{\tau}{2} \frac{d}{dt} \|z_k'(t)\|_2^2 + \frac{1}{2} \frac{d}{d\rho} \|z_k'(t)\|_2^2 = 0. \quad (3.13)$$

Taking the sum of (3.12) and (3.13), we obtain that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \left(\|u_k''(t)\|_2^2 + \|\nabla_x u_k'(t)\|_2^2 + \tau \int_0^1 \|z_k'(x, \rho, t)\|_{L^2(\Omega)}^2 d\rho \right) \\ + \mu_1(t) \int_{\Omega} u_k''^2(t) dx + \frac{1}{2} \int_{\Omega} |z_k'(x, 1, t)|^2 dx \\ = -\mu_2(t) \int_{\Omega} u_k''(t)z_k'(x, 1, t) dx - \mu_1'(t) \int_{\Omega} u_k''(t)u_k'(t) dx \\ - \mu_2'(t) \int_{\Omega} u_k''(t)z_k(x, 1, t) dx + \frac{1}{2} \|u_k''(t)\|_2^2 \end{aligned}$$

Using (H1), (H2), Cauchy–Schwarz and Young’s inequalities, we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \left(\|u_k''(t)\|_2^2 + \|\nabla_x u_k'(t)\|_2^2 + \int_0^1 \tau \|z_k'(x, \rho, t)\|_{L^2(\Omega)}^2 d\rho \right) \\ + \mu_1(t) \int_{\Omega} u_k''^2(t) dx + \frac{1}{2} \int_{\Omega} |z_k'(x, 1, t)|^2 dx \\ \leq |\mu_2(t)| \|u_k''(t)\|_2 \|z_k'(x, 1, t)\|_2 + |\mu_1'(t)| \|u_k''(t)\|_2 \|u_k'(t)\|_2 \\ + |\mu_2'(t)| \|u_k''(t)\|_2 \|z_k(x, 1, t)\|_2 + \frac{1}{2} \|u_k''(t)\|_2^2 \\ \leq \frac{|\mu_2(t)|^2}{2} \|u_k''(t)\|_2^2 + \frac{1}{2} \|z_k'(x, 1, t)\|_2^2 + \frac{|\mu_1'(t)|}{4} \|u_k''(t)\|_2^2 + |\mu_1'(t)| \|u_k'(t)\|_2^2 \\ + \frac{|\mu_2'(t)|}{4} \|u_k''(t)\|_2^2 + |\mu_2'(t)| \|z_k(x, 1, t)\|_2^2 + \frac{1}{2} \|u_k''(t)\|_2^2 \\ \leq c' \|u_k''(t)\|_2^2 + |\mu_1'(t)| \|u_k'(t)\|_2^2 + |\mu_2'(t)| \|z_k(x, 1, t)\|_2^2 + \frac{1}{2} \|z_k'(x, 1, t)\|_2^2 \\ \leq c' \|u_k''(t)\|_2^2 + M\mu_1(t) \|u_k'(t)\|_2^2 + \tilde{M}\mu_1(t) \|z_k(x, 1, t)\|_2^2 + \frac{1}{2} \|z_k'(x, 1, t)\|_2^2. \end{aligned}$$

Integrating the last inequality over $(0, t)$ and using (3.6), we get

$$\begin{aligned} & \left(\|u_k''(t)\|_2^2 + \|\nabla_x u_k'(t)\|_2^2 + \tau \int_0^1 \|z_k'(x, \rho, t)\|_{L^2(\Omega)}^2 d\rho \right) \\ & \leq \left(\|u_k''(0)\|_2^2 + \|\nabla_x u_k'(0)\|_2^2 + \tau \int_0^1 \|z_k'(x, \rho, 0)\|_{L^2(\Omega)}^2 d\rho \right) + 2M \int_0^t \mu_1(s) \|u_k'(s)\|_2^2 ds \\ & \quad + 2\tilde{M} \int_0^t \mu_1(s) \|z_k(x, 1, s)\|_2^2 ds + 2c' \int_0^t \|u_k''(s)\|_2^2 ds \\ & \leq C + C' \int_0^t \left(\|u_k''(s)\|_2^2 + \|\nabla_x u_k'(s)\|_2^2 + \tau \int_0^1 \|z_k'(x, \rho, s)\|_{L^2(\Omega)}^2 d\rho \right) ds. \end{aligned}$$

Using Gronwall's lemma, we deduce that

$$\|u_k''(t)\|_2^2 + \|\nabla_x u_k'(t)\|_2^2 + \tau \int_0^1 \|z_k'(x, \rho, t)\|_{L^2(\Omega)}^2 d\rho \leq Ce^{C'T}$$

for all $t \in \mathbb{R}^+$, therefore, we conclude that

$$(u_k'') \text{ is bounded in } L_{\text{loc}}^\infty(0, \infty; L^2(\Omega)), \quad (3.14)$$

$$(u_k') \text{ is bounded in } L_{\text{loc}}^\infty(0, \infty; H_0^1(\Omega)), \quad (3.15)$$

$$(\tau z_k') \text{ is bounded in } L_{\text{loc}}^\infty(0, \infty; L^2(\Omega \times (0, 1))). \quad (3.16)$$

Applying Dunford–Pettis' theorem, we deduce from (3.7), (3.8), (3.9), (3.10), (3.11), (3.14), (3.15) and (3.16), replacing the sequence u_k with a subsequence if necessary, that

$$u_k \rightarrow u \text{ weak-star in } L_{\text{loc}}^\infty(0, \infty; H^2(\Omega) \cap H_0^1(\Omega)), \quad (3.17)$$

$$u_k' \rightarrow u' \text{ weak-star in } L_{\text{loc}}^\infty(0, \infty; H_0^1(\Omega)),$$

$$u_k'' \rightarrow u'' \text{ weak-star in } L_{\text{loc}}^\infty(0, \infty; L^2(\Omega)), \quad (3.18)$$

$$u_k' \rightarrow \chi \text{ weak in } L^2(\Omega \times (0, T); \mu_1(t)),$$

$$z_k \rightarrow z \text{ weak-star in } L_{\text{loc}}^\infty(0, \infty; H_0^1(\Omega; L^2(0, 1))),$$

$$z_k' \rightarrow z' \text{ weak-star in } L_{\text{loc}}^\infty(0, \infty; L^2(\Omega \times (0, 1))), \quad (3.19)$$

$$z_k(x, 1, t) \rightarrow \psi \text{ weak in } L^2(\Omega \times (0, T), \mu_1(t))$$

for suitable functions

$$\begin{aligned} u & \in L^\infty(0, T; H^2(\Omega) \cap H_0^1(\Omega)), \quad z \in L^\infty(0, T; L^2(\Omega \times (0, 1))), \\ \chi & \in L^2(\Omega \times (0, T); \mu_1(t)), \quad \psi \in L^2(\Omega \times (0, T); \mu_1(t)), \end{aligned}$$

for all $T \geq 0$. We have to show that u is a solution of (P).

From (3.15) we have that (u_k') is bounded in $L^\infty(0, T; H_0^1(\Omega))$. Then (u_k') is bounded in $L^2(0, T; H_0^1(\Omega))$. Since (u_k'') is bounded in $L^\infty(0, T; L^2(\Omega))$, then it is bounded in $L^2(0, T; L^2(\Omega))$, too. Consequently, (u_k') is bounded in $H^1(Q)$.

Since the embedding $H^1(Q) \hookrightarrow L^2(Q)$ is compact, using the Aubin–Lions theorem [9], we can extract a subsequence (u_ζ) of (u_k) such that

$$u_\zeta' \rightarrow u' \text{ strongly in } L^2(Q). \quad (3.20)$$

Therefore

$$u_\zeta' \rightarrow u' \text{ strongly and a.e. in } Q. \quad (3.21)$$

Similarly we obtain

$$z_\zeta \rightarrow z \text{ strongly in } L^2(\Omega \times (0, 1) \times (0, T)) \quad (3.22)$$

and

$$z_\zeta \rightarrow z \text{ strongly and a.e. in } \Omega \times (0, 1) \times (0, T). \quad (3.23)$$

It follows at once from (3.17), (3.18), (3.19), (3.20) and (3.22) that for each fixed $v \in L^2(0, T; L^2(\Omega))$ and $w \in L^2(0, T; L^2(\Omega) \times (0, 1))$

$$\begin{aligned} & \int_0^T \int_\Omega (u_\zeta'' - \Delta_x u_\zeta + \mu_1(t)u_\zeta' + \mu_2(t)z_\zeta)v \, dx \, dt \\ & \rightarrow \int_0^T \int_\Omega (u'' - \Delta_x u + \mu_1(t)u' + \mu_2(t)z)v \, dx \, dt, \end{aligned} \quad (3.24)$$

$$\int_0^T \int_0^1 \int_\Omega \left(\tau z_\zeta' + \frac{\partial}{\partial \rho} z_\zeta \right) w \, dx \, d\rho \, dt \rightarrow \int_0^T \int_0^1 \int_\Omega \left(\tau z' + \frac{\partial}{\partial \rho} z \right) w \, dx \, d\rho \, dt \quad (3.25)$$

as $\zeta \rightarrow +\infty$. Thus the problem (P) admits a global weak solution u .

Uniqueness. Let (u_1, z_1) and (u_2, z_2) be two solutions of problem (2.10). Then $(w, \tilde{w}) = (u_1, z_1) - (u_2, z_2)$ satisfies

$$\begin{cases} w''(x, t) - \Delta_x w(x, t) + \mu_1(t)w'(x, t) + \mu_2(t)\tilde{w}(x, 1, t) = 0, & \text{in } \Omega \times]0, +\infty[, \\ \tau \tilde{w}'(x, \rho, t) + \tilde{w}_\rho(x, \rho, t) = 0, & \text{in } \Omega \times]0, 1[\times]0, +\infty[, \\ w(x, t) = 0, & \text{on } \partial\Omega \times]0, +\infty[, \\ \tilde{w}(x, 0, t) = w'(x, t), & \text{on } \Omega \times]0, +\infty[, \\ w(x, 0) = 0, \quad w'(x, 0) = 0, & \text{in } \Omega \\ \tilde{w}(x, \rho, 0) = 0, & \text{in } \Omega \times]0, 1[\end{cases} \quad (3.26)$$

Multiplying the first equation in (3.26) by w' , integrating over Ω and using integration by parts, we get

$$\frac{1}{2} \frac{d}{dt} (\|w'\|_2^2 + \|\nabla_x w\|_2^2) + \mu_1(t)\|w'\|_2^2 + \mu_2(t)(\tilde{w}(x, 1, t), w') = 0. \quad (3.27)$$

Multiplying the second equation in (3.26) by \tilde{w} , integrating over $\Omega \times (0, 1)$ and using integration by parts, we get

$$\frac{\tau}{2} \frac{d}{dt} \|\tilde{w}\|_2^2 + \frac{1}{2} \frac{d}{d\rho} \|\tilde{w}\|_2^2 = 0. \quad (3.28)$$

Then

$$\frac{\tau}{2} \frac{d}{dt} \int_0^1 \|\tilde{w}\|_2^2 \, d\rho + \frac{1}{2} (\|\tilde{w}(x, 1, t)\|_2^2 - \|w'\|_2^2) = 0. \quad (3.29)$$

From (3.27), (3.29), using the Cauchy–Schwarz inequality we get

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(\|w'\|_2^2 + \|\nabla_x w\|_2^2 + \tau \int_0^1 \|\tilde{w}\|_2^2 \, d\rho \right) + \mu_1(t)\|w'\|_2^2 + \frac{1}{2} \|\tilde{w}(x, 1, t)\|_2^2 \\ & = -\mu_2(t)(\tilde{w}(x, 1, t), w') + \frac{1}{2} \|w'\|_2^2 \\ & \leq |\mu_2(t)| \|\tilde{w}(x, 1, t)\|_2 \|w'\|_2 + \frac{1}{2} \|w'\|_2^2. \end{aligned}$$

Using Young's inequality, we obtain

$$\frac{1}{2} \frac{d}{dt} \left(\|w'\|_2^2 + \|\nabla_x w\|_2^2 + \tau \int_0^1 \|\tilde{w}\|_2^2 \, d\rho \right) \leq c \|w'\|_2^2,$$

where c is a positive constant. Then integrating over $(0, t)$, using Gronwall's lemma, we conclude that

$$\|w'\|_2^2 + \|\nabla_x w\|_2^2 + \tau \int_0^1 \|\tilde{w}\|_2^2 d\rho = 0.$$

Hence, uniqueness follows.

4 Asymptotic behavior

From now on, we denote by c various positive constants which may be different at different occurrences. We multiply the first equation of (2.10) by $\phi' E^q u$, where ϕ is a bounded function satisfying all the hypotheses of Lemma 2.1. We obtain

$$\begin{aligned} 0 &= \int_S^T E^q \phi' \int_{\Omega} u \left(u'' - \Delta u + \mu_1(t) u' + \mu_2(t) z(x, 1, t) \right) dx dt \\ &= \left[E^q \phi' \int_{\Omega} u u' dx \right]_S^T - \int_S^T (q E' E^{q-1} \phi' + E^q \phi'') \int_{\Omega} u u' dx dt \\ &\quad - 2 \int_S^T E^q \phi' \int_{\Omega} u'^2 dx dt + \int_S^T E^q \phi' \int_{\Omega} (u'^2 + |\nabla u|^2) dx dt \\ &\quad + \int_S^T E^q \phi' \mu_1(t) \int_{\Omega} u u' dx dt + \int_S^T E^q \phi' \mu_2(t) \int_{\Omega} u z(x, 1, t) dx dt. \end{aligned}$$

Similarly, we multiply the second equation of (2.10) by $E^q \phi' \zeta(t) e^{-2\tau\rho} z(x, \rho, t)$ and get

$$\begin{aligned} 0 &= \int_S^T E^q \phi' \int_{\Omega} \int_0^1 e^{-2\tau\rho} \zeta(t) z \left(\tau z_t + z_{\rho} \right) dx d\rho dt \\ &= \left[\frac{1}{2} E^q \phi' \zeta(t) \tau \int_{\Omega} \int_0^1 e^{-2\tau\rho} z^2 dx d\rho \right]_S^T - \frac{1}{2} \int_S^T \int_{\Omega} \int_0^1 (E^q \phi' \zeta(t) \tau e^{-2\tau\rho})' z^2 dx d\rho dt \\ &\quad + \int_S^T E^q \phi' \int_{\Omega} \int_0^1 \zeta(t) \left(\frac{1}{2} \frac{\partial}{\partial \rho} (e^{-2\tau\rho} z^2) + \tau e^{-2\tau\rho} z^2 \right) dx d\rho dt \\ &= \left[\frac{1}{2} E^q \phi' \zeta(t) \tau \int_{\Omega} \int_0^1 e^{-2\tau(t)\rho} z^2 dx d\rho \right]_S^T - \frac{\tau}{2} \int_S^T (E^q \phi' \zeta(t))' \int_{\Omega} \int_0^1 e^{-2\tau\rho} z^2 dx d\rho dt \\ &\quad + \frac{1}{2} \int_S^T E^q \phi' \zeta(t) \int_{\Omega} \left(e^{-2\tau} z^2(x, 1, t) - z^2(x, 0, t) \right) dx dt + \int_S^T E^q \phi' \zeta(t) \tau \int_0^1 \int_{\Omega} e^{-2\tau\rho} z^2 dx d\rho dt. \end{aligned}$$

Taking their sum, we obtain

$$\begin{aligned} A \int_S^T E^{q+1} \phi' dt &\leq - \left[E^q \phi' \int_{\Omega} u u' dx \right]_S^T + \int_S^T (q E' E^{q-1} \phi' + E^q \phi'') \int_{\Omega} u u' dx dt \\ &\quad + 2 \int_S^T E^q \phi' \int_{\Omega} u'^2 dx dt - \int_S^T \mu_1(t) E^q \phi' \int_{\Omega} u u' dx dt \\ &\quad - \int_S^T \mu_2(t) E^q \phi' \int_{\Omega} u z(x, 1, t) dx dt - \left[\frac{1}{2} E^q \phi' \zeta(t) \tau \int_{\Omega} \int_0^1 e^{-2\tau\rho} z^2 dx d\rho \right]_S^T \\ &\quad + \frac{1}{2} \int_S^T (E^q \phi' \zeta(t))' \tau \int_{\Omega} \int_0^1 e^{-2\tau\rho} z^2 dx d\rho dt \\ &\quad - \frac{1}{2} \int_S^T E^q \phi' \zeta(t) \int_{\Omega} \left(e^{-2\tau} z^2(x, 1, t) - z^2(x, 0, t) \right) dx dt, \end{aligned} \tag{4.1}$$

where $A = 2 \min\{1, e^{-2\tau_1}\}$. Using the Cauchy–Schwarz and Poincaré’s inequalities and the definition of E and assuming that ϕ' is a bounded non-negative function on \mathbb{R}^+ , we get

$$\left| E^q(t) \phi' \int_{\Omega} uu' dx \right| \leq cE(t)^{q+1}.$$

By recalling (2.14), we have

$$\begin{aligned} \int_S^T \left| qE'E^{q-1}\phi' \int_{\Omega} uu' dx \right| dt &\leq c \int_S^T E^q(t) |E'(t)| dt \leq c \int_S^T E^q(t) (-E'(t)) dt \\ &\leq cE^{q+1}(S), \end{aligned}$$

$$\int_S^T E^q \phi'' \int_{\Omega} uu' dx dt \leq c \int_S^T E^{q+1}(t) (-\phi'') \leq cE^{q+1}(S) \int_S^T (-\phi'') dt \leq cE^{q+1}(S),$$

and

$$\begin{aligned} \int_S^T E^q \phi' \int_{\Omega} u^2 dx dt &\leq c \int_S^T E^q \phi' \frac{1}{\mu_1(t)} \int_{\Omega} \mu_1(t) u^2 dx dt \\ &\leq \int_S^T E^q \frac{\phi'}{\mu_1(t)} (-E') dt. \end{aligned} \quad (4.2)$$

Define

$$\phi(t) = \int_0^t \mu_1(s) ds. \quad (4.3)$$

It is clear that ϕ is a non-decreasing function of class C^1 on \mathbb{R}^+ , ϕ is bounded and

$$\phi(t) \rightarrow +\infty \text{ as } t \rightarrow +\infty. \quad (4.4)$$

So, we deduce, from (4.2), that

$$\int_S^T E^q \phi' \int_{\Omega} u^2 dx dt \leq c \int_S^T E^q (-E') dt \leq cE^{q+1}(S), \quad (4.5)$$

By the hypothesis (H1), Young’s and Poincaré’s inequality and (2.14), we have

$$\begin{aligned} \left| \int_S^T E^q \phi' \int_{\Omega} uu' dx dt \right| &\leq c \int_S^T E^q \phi' \|u\|_2 \|u'\|_2 dt \\ &\leq c\varepsilon' \int_S^T E^q \phi' \|u\|_2^2 dt + c(\varepsilon') \int_S^T E^q \phi' \|u'\|_2^2 dt \\ &\leq \varepsilon' c_* \int_S^T E^q \phi' \|\nabla_x u\|_2^2 dt + c(\varepsilon') \int_S^T E^q \phi' \|u'\|_2^2 dt \\ &\leq \varepsilon' c_* \int_S^T E^{q+1} \phi' dt + cE^{q+1}(S). \end{aligned}$$

Recalling that $\zeta' \leq 0$ and the definition of E , we have

$$\begin{aligned} \int_S^T (E^q \zeta(t))' \tau \int_{\Omega} \int_0^1 e^{-2\tau\rho z^2} dx d\rho dt &\leq \int_S^T (E^q)' \zeta(t) \tau \int_{\Omega} \int_0^1 e^{-2\tau\rho z^2} dx d\rho dt \\ &\leq c \int_S^T E^q |E'| dt \\ &\leq c \int_S^T E^q (-E'(t)) dt \\ &\leq cE^{q+1}(S), \end{aligned}$$

$$\begin{aligned} \int_S^T E^q \xi(t) \int_{\Omega} e^{-2\tau} z^2(x, 1, t) dx dt &\leq c \int_S^T E^q \xi(t) \int_{\Omega} z^2(x, 1, t) dx dt \\ &\leq c \int_S^T E^q (-E') dt \\ &\leq cE^{q+1}(S), \end{aligned}$$

$$\begin{aligned} \int_S^T E^q \xi(t) \int_{\Omega} z^2(x, 0, t) dx dt &= \int_S^T E^q \xi(t) \int_{\Omega} u'^2(x, t) dx dt \\ &\leq cE^{q+1}(S), \end{aligned}$$

where we have also used the Cauchy–Schwarz inequality. Combining these estimates and choosing ε' sufficiently small, we conclude from (4.1) that

$$\int_S^T E^{q+1} \phi' dt \leq CE^{q+1}(S).$$

This ends the proof of Theorem 2.2.

Acknowledgements

This work is partially supported by KFUPM under Grant #IN121029.

References

- [1] C. ABDALLAH, P. DORATO, J. BENITEZ-READ, R. BYRNE, Delayed positive feedback can stabilize oscillatory system, *American Control Conference*, San Francisco, CA, USA, 1993, 3106–3107.
- [2] M. M. CAVALCANTI, N. A. LAR'KIN, J. A. SORIANO, On solvability and stability of solutions of nonlinear degenerate hyperbolic equations with boundary damping, *Funkcial. Ekvac.* **41**(1998), 271–289. [MR1662356](#)
- [3] G. CHEN, Control and stabilization for the wave equation in a bounded domain, Part I, *SIAM J. Control Optim.* **17**(1979), 66–81. [MR516857](#); [url](#)
- [4] G. CHEN, Control and stabilization for the wave equation in a bounded domain, Part II, *SIAM J. Control Optim.* **19**(1981), 114–122. [MR603084](#); [url](#)
- [5] R. DATKO, J. LAGNESE, M. P. POLIS, An example on the effect of time delays in boundary feedback stabilization of wave equations, *SIAM J. Control Optim.* **24**(1986), 152–156. [MR818942](#); [url](#)
- [6] A. HARAUX, Two remarks on hyperbolic dissipative problems, In: *Nonlinear partial differential equations and their applications. Collège de France seminar, Vol. VII (Paris, 1983–1984), Research Notes in Mathematics*, Vol. 122, 161–179, Pitman, Boston, MA, 1985. [MR879461](#)
- [7] V. KOMORNIK, *Exact controllability and stabilization. The multiplier method*, Masson-John Wiley, Paris, 1994. [MR1359765](#)

- [8] I. LASIECKA, R. TRIGGIANI, Uniform exponential energy decay of wave equations in a bounded region with $L^2(0, \infty; L^2(\Gamma))$ -feedback control in the Dirichlet boundary conditions, *J. Differential Equations* **66**(1987), 340–390. [MR876804](#); [url](#)
- [9] J. L. LIONS, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, Paris 1969.
- [10] P. MARTINEZ, A new method to obtain decay rate estimates for dissipative systems, *ESAIM Control Optim. Calc. Var.* **4**(1999), 419–444. [MR1693904](#); [url](#)
- [11] S. A. MESSAOUDI, Energy decay of solutions of a semilinear wave equation, *Int. J. Appl. Math.* **9**(2000), 1037–1048. [MR1757588](#)
- [12] M. NAKAO, Decay of solutions of some nonlinear evolution equations, *J. Math. Anal. Appl.* **60**(1977), 542–549. [MR0499564](#)
- [13] S. NICAISE, C. PIGNOTTI, Stability and instability results of the wave equation with a delay term in the boundary or internal feedbacks, *SIAM J. Control Optim.* **45**(2006), 1561–1585. [MR2272156](#); [url](#)
- [14] S. NICAISE, J. VALEIN, Stabilization of second order evolution equations with unbounded feedback with delay, *ESAIM Control Optim. Calc. Var.* **16**(2010), 420–456. [MR2654201](#); [url](#)
- [15] S. NICAISE & C. PIGNOTTI, Stabilization of the wave equation with boundary or internal distributed delay, *Differ. Int. Equ.* **21**(2008), 935–958. [MR2483342](#)
- [16] S. NICAISE, J. VALEIN, E. FRIDMAN, Stability of the heat and of the wave equations with boundary time-varying delays, *Discrete Contin. Dyn. Syst. Ser. S* **2**(2009), 559–581. [MR2525768](#); [url](#)
- [17] S. NICAISE, C. PIGNOTTI, J. VALEIN, Exponential stability of the wave equation with boundary time-varying delay, *Discrete Contin. Dyn. Syst. Ser. S* **4**(2011), 693–722. [MR2746429](#); [url](#)
- [18] J. Y. PARK, T. G. HA, Energy decay for nondissipative distributed systems with boundary damping and source term, *Nonlinear Anal.* **70**(2009), 2416–2434. [MR2498341](#); [url](#)
- [19] I. H. SUH, Z. BIEN, Use of time delay action in the controller design, *IEEE Trans. Autom. Control* **25**(1980) 600–603.
- [20] C. Q. XU, S.P. YUNG, L. K. LI, Stabilization of the wave system with input delay in the boundary control, *ESAIM Control Optim. Calc. Var.* **12**(2006), 770–785. [url](#)
- [21] E. ZUAZUA, Stability and decay for a class of nonlinear hyperbolic problems, *Asymptot. Anal.* **1**(1988), 161–185. [MR35B40](#)