



## Center problem for a class of degenerate quartic systems

Bo Sang 

School of Mathematical Sciences, Liaocheng University,  
No. 1 Hunan Road, Liaocheng, 252059, P.R. China

Received 7 March 2014, appeared 11 January 2015

Communicated by László Hatvani

**Abstract.** This paper, using pseudo-division algorithm, introduces a method for computing resonant focus numbers of a class of complex polynomial differential systems, establishes the necessary and sufficient conditions for existence of a center for a class of complex quartic systems with a degenerate resonant singular point.

**Keywords:** complex quartic systems, degenerate resonant singular point, integrability, resonant focus number.

**2010 Mathematics Subject Classification:** 34C05, 34C07.

### 1 Introduction

In the qualitative theory of real planar differential systems, focal values and saddle values are two important detection quantities. In [2], the authors introduce a new efficient computational method, which combines the computation of focal values and saddle values into a unified calculation of singular point quantities for a class of complex planar differential systems. Using pseudo-divisions, Wang [21] gives an improved formal power series method for computing focal values of a class of polynomial differential systems. Using a perturbation technique based on multiple time scales, Yu [25] presents an efficient method for computing focal values of some classes of differential systems.


Żołądek [27] generalizes the notion of center to the case of a  $p : -q$  resonant singular point of the following complex polynomial vector fields

$$\begin{cases} \frac{dx}{dt} = px + X_m(x, y), \\ \frac{dy}{dt} = -qy + Y_m(x, y), \end{cases} \quad (1.1)$$

in  $\mathbb{C}^2$ , where  $p, q \in \mathbb{N}$ ,  $p \leq q$ ,  $(p, q) = 1$ , and

$$X_m(x, y) = \sum_{k=2}^m \sum_{j=0}^k a_{k,j} x^{k-j} y^j, \quad Y_m(x, y) = \sum_{k=2}^m \sum_{j=0}^k b_{k,j} x^{k-j} y^j.$$

---

 Email: [sangbo\\_76@163.com](mailto:sangbo_76@163.com)

**Definition 1.1.** System (1.1) is said to have a  $p : -q$  resonant center at the origin if it admits a local first integral of the form

$$F(x, y) = x^q y^p + \sum_{k=p+q+1}^{\infty} \sum_{j=0}^k B_{k,j} x^{k-j} y^j. \quad (1.2)$$

Although, for system (1.1) with  $p : -q = 1 : -1, 1 : -2, 1 : -3, 2 : -3, 1 : -q$ , the resonant center problems have received intensive attentions, see [3–5, 8, 10–12, 18, 19], very few results are known for systems having high order nonlinearities.

For system (1.1), we can derive a formal power series of the form (1.2) with  $B_{s(p+q),sp} = 0$ ,  $s = 2, 3, \dots$ , such that

$$\left. \frac{dF}{dt} \right|_{(1.1)} = \frac{\partial F}{\partial x}(px + X_m) + \frac{\partial F}{\partial y}(-qy + Y_m) = \sum_{n=1}^{\infty} W_n (x^q y^p)^{n+1}, \quad (1.3)$$

where  $W_n$  are called the  $n$ -th order  $p : -q$  resonant focus numbers. For some computational methods of such quantities, see [14, 19]. For large  $n$ , the computation of  $W_n$  is very complicated, which is the main reason of slow progress in the center problems.

The only way to get the necessary conditions for a center is to compute the  $p : -q$  resonant focus numbers. Before presenting a new algorithm, we start with a precise definition of pseudo-remainder of polynomials. For more details, see [6, 22–24].

Let  $K[x_1, x_2, \dots, x_n]$  denote the ring of polynomials in indeterminates  $x_1, x_2, \dots, x_n$  with coefficients in a field  $K$  of characteristic 0. Consider a fixed ordering on the set of indeterminates:  $x_1 < x_2 < \dots < x_n$ . A polynomial  $f \in K[x_1, x_2, \dots, x_n]$  is said to be of class  $i$  if  $i$  is the maximum index such that  $f$  has a positive degree in  $x_i$ . The class of elements of  $K$  is zero. If  $f$  is of class  $i$  the coefficient of the  $x_i$  of the maximum degree is said to be the initial of polynomial  $f$  and is denoted by  $\text{In}(f)$ .

If  $f$  and  $g$  are two polynomials of class respectively  $i$  and  $j$ , with  $i < j$ , or such that  $i = j$  and the degree in  $x_i$  of  $f$  is less than the degree of  $g$ , then it is possible, using the Euclidean algorithm over  $K(x_1, x_2, \dots, x_{i-1})[x_i]$  to find polynomials  $q$  and  $r$  with  $\deg_{x_i}(r) < \deg_{x_i}(f)$  such that

$$\text{In}(f)^\alpha g = qf + r,$$

with  $\alpha$  bounded by  $\deg_{x_i}(f) - \deg_{x_i}(g) + 1$ . The polynomial  $r$  is called the pseudo-remainder of  $g$  with respect to  $f$ , and it is denoted by  $\text{prem}(g, f)$ . This operation is called pseudo-division.

**Definition 1.2** ([23]). A sequence of polynomials  $AS = [f_1, f_2, \dots, f_r]$  is called a triangular set if  $r = 1$  and  $f_1$  is not identically zero, or  $r > 1$  and  $0 < \text{class}(f_1) < \text{class}(f_2) < \dots < \text{class}(f_r) \leq n$ .

**Definition 1.3** ([22]). Consider a triangular set  $AS = [f_1, f_2, \dots, f_r]$ , and a polynomial  $g \in K[x_1, x_2, \dots, x_r]$ . Let us pseudo-divide  $g$  by  $f_r, f_{r-1}, \dots, f_1$  successively as polynomials in  $x_{c_r}, \dots, x_{c_1}$ ,  $c_i = \text{class}(f_i)$ , and denote the final remainder by  $R$ . Then we shall get an expression of the form:

$$I_1^{s_1} \cdots I_r^{s_r} g = \sum_{i=1}^r Q_i f_i + R,$$

where  $I_i$  is the initial of  $f_i$ ,  $s_i$  assumes the smallest possible power achievable.  $R$  is called the pseudo-remainder of  $g$  with respect to  $AS$ , denoted as  $R = \text{Prem}(g, AS)$ .

Now we are in a position to develop the algorithm for computing  $W_n$  in (1.3). Grouping the like terms in the second expression of (1.3), we get

$$\begin{aligned} \frac{\partial F}{\partial x}(px + X_m) + \frac{\partial F}{\partial y}(-qy + Y_m) &= \sum_{l=p+q+1}^{(p+q)(n+1)-1} \sum_{j=0}^l f_{l,j} x^{l-j} y^j \\ &+ \sum_{j=0, j \neq p(n+1)}^{(p+q)(n+1)} f_{(p+q)(n+1),j} x^{(p+q)(n+1)-j} y^j \\ &+ V_n(x^q y^p)^{n+1} + \dots, \end{aligned}$$

where  $V_n, f_{l,j}, f_{(p+q)(n+1),j}$  are polynomials in  $a_{k,j}, b_{k,j}, B_{k,j}$ .

When computing the  $n$ -th order resonant focus number  $W_n$ , the coefficients  $f_{l,j}, f_{(p+q)(n+1),j}$  have to be zero. Thus in order to eliminate indeterminates  $B_{k,j}$  from  $V_n$ , we use successive pseudo-divisions: first choosing a suitable variable order of  $B_{k,j}$ ; secondly, rearranging some polynomials  $f_{l,j}, f_{(p+q)(n+1),j}$  to get a triangular set  $TS_n$ ; finally, performing successive pseudo-division of  $V_n + v$  by  $TS_n$  to get the pseudo-remainder  $R_n$ , then the  $n$ -th order  $p : -q$  resonant focus number can be written as  $W_n = \frac{R_n}{\text{coeff}(R_n, v)} - v$ , where  $\text{coeff}(R_n, v)$  is the coefficient of  $v$  in the polynomial  $R_n$ , and  $v$  is a new variable.

To illustrate the main idea of the algorithm, we compute the second order  $1 : -2$  resonant focus number  $W_2$  of the family

$$\begin{cases} \frac{dx}{dt} = x(1 + a_1x + a_2x^2 + a_3yx + a_4y^2), \\ \frac{dy}{dt} = y(-2 + b_1y + b_2x^2 + b_3yx + b_4y^2). \end{cases} \quad (1.4)$$

Let

$$F(x, y) = x^2 y + \sum_{k=4}^9 \sum_{j=0}^k B_{k,j} x^{k-j} y^j + \dots$$

and using the same notations as described in the algorithm, we have

$$\begin{aligned} \left. \frac{dF}{dt} \right|_{(1.4)} &= \sum_{l=4}^8 \sum_{j=0}^l f_{l,j} x^{l-j} y^j \\ &+ \sum_{j=0, j \neq 3}^9 f_{9,j} x^{9-j} y^j \\ &+ V_2(x^2 y)^3 + \dots, \end{aligned}$$

where

$$V_2 = 6 B_{7,1} a_4 + B_{7,1} b_4 + 5 B_{7,2} a_3 + 2 B_{7,2} b_3 + 4 B_{7,3} a_2 + 3 B_{7,3} b_2 + 2 B_{8,2} b_1 + 5 B_{8,3} a_1.$$

Under the variable ordering

$$\begin{aligned} B_{4,0} &\prec B_{4,1} \prec B_{4,2} \prec B_{4,3} \prec B_{5,0} \prec B_{5,1} \prec B_{5,2} \prec B_{5,3} \prec B_{6,0} \prec B_{6,1} \\ &\prec B_{6,3} \prec B_{7,1} \prec B_{7,2} \prec B_{7,3} \prec B_{8,2} \prec B_{8,3}, \end{aligned}$$

the following sequence of polynomials

$$TS_2 = [f_{4,0}, f_{4,1}, f_{4,2}, f_{4,3}, f_{5,0}, f_{5,1}, f_{5,2}, f_{5,3}, f_{6,0}, f_{6,1}, f_{6,3}, f_{7,1}, f_{7,2}, f_{7,3}, f_{8,2}, f_{8,3}]$$

is a triangular set, where

$$\begin{aligned}
f_{4,0} &= 4 B_{4,0}, \\
f_{4,1} &= 2 a_1 + B_{4,1}, \\
f_{4,2} &= -2 B_{4,2} + b_1, \\
f_{4,3} &= -5 B_{4,3}, \\
f_{5,0} &= 4 B_{4,0} a_1 + 5 B_{5,0}, \\
f_{5,1} &= 3 B_{4,1} a_1 + 2 B_{5,1} + 2 a_2 + b_2, \\
f_{5,2} &= B_{4,1} b_1 + 2 B_{4,2} a_1 - B_{5,2} + 2 a_3 + b_3, \\
f_{5,3} &= 2 B_{4,2} b_1 + B_{4,3} a_1 - 4 B_{5,3} + 2 a_4 + b_4, \\
f_{6,0} &= 4 B_{4,0} a_2 + 5 B_{5,0} a_1 + 6 B_{6,0}, \\
f_{6,1} &= 4 B_{4,0} a_3 + 3 B_{4,1} a_2 + B_{4,1} b_2 + 4 B_{5,1} a_1 + 3 B_{6,1}, \\
f_{6,3} &= 3 B_{4,1} a_4 + B_{4,1} b_4 + 2 B_{4,2} a_3 + 2 B_{4,2} b_3 + B_{4,3} a_2 + 3 B_{4,3} b_2 + 2 B_{5,2} b_1 + 2 B_{5,3} a_1 - 3 B_{6,3}, \\
f_{7,1} &= 5 B_{5,0} a_3 + 4 B_{5,1} a_2 + B_{5,1} b_2 + 5 B_{6,1} a_1 + 4 B_{7,1}, \\
f_{7,2} &= 5 B_{5,0} a_4 + 4 B_{5,1} a_3 + B_{5,1} b_3 + 3 B_{5,2} a_2 + 2 B_{5,2} b_2 + B_{6,1} b_1 + B_{7,2}, \\
f_{7,3} &= 4 B_{5,1} a_4 + B_{5,1} b_4 + 3 B_{5,2} a_3 + 2 B_{5,2} b_3 + 2 B_{5,3} a_2 + 3 B_{5,3} b_2 + 3 B_{6,3} a_1 - 2 B_{7,3}, \\
f_{8,2} &= 6 B_{6,0} a_4 + 5 B_{6,1} a_3 + B_{6,1} b_3 + B_{7,1} b_1 + 5 B_{7,2} a_1 + 2 B_{8,2}, \\
f_{8,3} &= 5 B_{6,1} a_4 + B_{6,1} b_4 + 3 B_{6,3} a_2 + 3 B_{6,3} b_2 + 2 B_{7,2} b_1 + 4 B_{7,3} a_1 - B_{8,3}.
\end{aligned}$$

By computing pseudo-remainder of  $V_2 + v$  by  $TS_2$ , one gets

$$\begin{aligned}
R_2 = & -9676800 a_1^3 b_1 b_3 - 4838400 a_1^2 b_1^2 b_2 + 921600 a_1^2 a_2 a_4 - 42854400 a_1^2 a_3 b_3 \\
& + 921600 a_1^2 a_4 b_2 - 2073600 a_1^2 b_2 b_4 - 19353600 a_1^2 b_3^2 - 921600 a_1 a_2 a_3 b_1 \\
& - 4147200 a_1 a_2 b_1 b_3 - 20275200 a_1 a_3 b_1 b_2 - 12441600 a_1 b_1 b_2 b_3 - 2073600 a_2 b_1^2 b_2 \\
& - 1382400 b_1^2 b_2^2 - 2764800 a_2 a_3^2 + 1382400 a_2 a_4 b_2 - 691200 a_2 b_2 b_4 + 1382400 a_3^2 b_2 \\
& + 1382400 a_3 b_2 b_3 - 691200 b_2^2 b_4 - 1382400 v.
\end{aligned}$$

Hence the second order 1 :  $-2$  resonant focus number can be written as

$$\begin{aligned}
W_2 &= \frac{R_2}{\text{coeff}(R_2, v)} - v \\
&= 7 a_1^3 b_1 b_3 + \frac{7}{2} a_1^2 b_1^2 b_2 - \frac{2}{3} a_1^2 a_2 a_4 + 31 a_1^2 a_3 b_3 - \frac{2}{3} a_1^2 a_4 b_2 + \frac{3}{2} a_1^2 b_2 b_4 \\
&\quad + 14 a_1^2 b_3^2 + \frac{2}{3} a_1 a_2 a_3 b_1 + 3 a_1 a_2 b_1 b_3 + \frac{44 a_1 a_3 b_1 b_2}{3} + 9 a_1 b_1 b_2 b_3 + \frac{3}{2} a_2 b_1^2 b_2 \\
&\quad + b_1^2 b_2^2 + 2 a_2 a_3^2 - a_2 a_4 b_2 + \frac{1}{2} a_2 b_2 b_4 - a_3^2 b_2 - a_3 b_2 b_3 + \frac{1}{2} b_2^2 b_4.
\end{aligned}$$

A general purposed Maple package *Myvalue* based on our algorithm is developed in Maple V.18 on Intel Core 2 Quad CPU Q8400, 4G RAM, and such Maple package is available for non-commercial purpose via email to: [sangbo\\_76@163.com](mailto:sangbo_76@163.com). Another Maple package *Liuc* based on the method [14] is also developed by us using the same computing platform. For technical comparison of these two packages, let us consider a class of cubic differential systems

$$\begin{cases} \frac{dx}{dt} = x + X_3(x, y), \\ \frac{dy}{dt} = -y + Y_3(x, y), \end{cases}$$

in  $\mathbb{C}^2$ , and

$$X_3(x, y) = \sum_{k=2}^3 \sum_{j=0}^k a_{k,j} x^{k-j} y^j, \quad Y_3(x, y) = \sum_{k=2}^3 \sum_{j=0}^k b_{k,j} x^{k-j} y^j.$$

Computing the first eight  $1 : -1$  resonant focus numbers  $W_j$ ,  $1 \leq j \leq 8$  by *Myvalue* and *Liuc* respectively, we find that the outputs (in expanded form) are the same for these two methods, and get the following experimental results on efficiency, see Table 1.1.

Method	$W_1$	$W_2$	$W_3$	$W_4$	$W_5$	$W_6$	$W_7$	$W_8$
<i>Myvalue</i>	0.015	0.015	0.046	0.203	1.281	5.640	45.968	160.968
<i>Liuc</i>	0.0	0.0	0.0	0.062	0.578	3.546	40.078	592.750

Table 1.1: Computing times (in CPU seconds) for the first eight resonant focus numbers

For computing  $W_n$  with  $n$  large, it is worth noting that the expansion of long polynomials in the last stage of package *Liuc* is pretty time-consuming, whereas the package *Myvalue* does not need any expansions before generating its outputs.

Consider the system of differential equations

$$\begin{cases} \frac{dx}{dt} = P_n(x, y), \\ \frac{dy}{dt} = Q_n(x, y), \end{cases} \quad (1.5)$$

where  $(x, y) \in \mathbb{C}^2$ ,  $P_n, Q_n$  are polynomials of degree  $n$  with  $(P_n, Q_n) = 1$ .

**Definition 1.4.** The polynomial  $f(x, y) \in \mathbb{C}[x, y]$  is called an algebraic partial integral of the system (1.5) if there exists a polynomial  $h \in \mathbb{C}[x, y]$  such that

$$\left. \frac{df}{dt} \right|_{(1.5)} = h(x, y) f(x, y).$$

The polynomial  $h$  is called a cofactor. If  $h \equiv 0$  then  $f(x, y) = \text{const}$  is a first integral of system (1.5).

**Lemma 1.5.** Suppose that system (1.5) admits  $m$  independent algebraic partial integrals  $f_1, f_2, \dots, f_m$  satisfying

$$\left. \frac{df_k}{dt} \right|_{(1.5)} = h_k(x, y) f_k(x, y), \quad k = 1, 2, \dots, m.$$

If there are scalars  $\alpha_1, \alpha_2, \dots, \alpha_m$ , not all zero, such that

$$\alpha_1 h_1 + \alpha_2 h_2 + \dots + \alpha_m h_m = - \left( \frac{\partial P_n}{\partial x} + \frac{\partial Q_n}{\partial y} \right),$$

then the function  $f = f_1^{\alpha_1} f_2^{\alpha_2} \dots f_m^{\alpha_m}$  is an integrating factor of system (1.5).

Mattei and Moussu [16] proved the next result for all isolated singularities.

**Lemma 1.6.** Assume that system (1.1) with an isolated singularity at the origin has a formal first integral  $F(x, y) \in \mathbb{R}[[x, y]]$  around it. Then, there exists an analytic first integral around the singularity.

Another mechanism to prove the integrability of system (1.5) is time-reversibility. From [20], we have the following result.

**Lemma 1.7.** *System (1.5) is time-reversible with respect to a transformation*

$$R: x \mapsto \gamma y, \quad y \mapsto \gamma^{-1}x,$$

where  $\gamma$  is a nonzero scalar, if and only if,

$$\gamma Q_n(\gamma y, \gamma^{-1}x) = -P_n(x, y), \quad \gamma Q_n(x, y) = -P_n(\gamma y, \gamma^{-1}x).$$

## 2 Main result

In the qualitative theory of planar differential systems, there are few works about degenerate singular point. Most of the work focuses on the center problem of the system

$$\begin{cases} \frac{dx}{dt} = y + P(x, y), \\ \frac{dy}{dt} = Q(x, y), \end{cases}$$

where  $P, Q$  are polynomials in  $x$  and  $y$  with degree no less than two, see [1, 7, 9, 17].

Let us consider the real analytic system

$$\begin{cases} \frac{du}{dt_1} = -v(u^2 + v^2)^n + \sum_{k=2n+2}^{\infty} U_k(u, v) = U(u, v), \\ \frac{dv}{dt_1} = u(u^2 + v^2)^n + \sum_{k=2n+2}^{\infty} V_k(u, v) = V(u, v), \end{cases} \quad (2.1)$$

where  $U(u, v), V(u, v)$  are analytic in a sufficiently small neighborhood of the origin,  $U_k(u, v), V_k(u, v)$  are homogeneous polynomials of degree  $k$ , and  $n \geq 0$ . Because the singularity  $(u, v) = (0, 0)$  of system (2.1) has no characteristic directions, it is a center or a focus.

Under the transformation  $u = r \cos(\theta), v = r \sin(\theta)$ , system (2.1) becomes

$$\begin{cases} \frac{dr}{dt_1} = r^{2n+1} \sum_{k=0}^{\infty} \phi_{2n+2+k}(\theta) r^k, \\ \frac{d\theta}{dt_1} = r^{2n} \sum_{k=0}^{\infty} \psi_{2n+2+k}(\theta) r^k. \end{cases} \quad (2.2)$$

It can be written as

$$\frac{dr}{d\theta} = r \sum_{k=0}^{\infty} R_k(\theta) r^k, \quad (2.3)$$

where the function on the right side of (2.3) is convergent in the range  $\theta \in [-4\pi, 4\pi], r < r_0$ , and

$$R_k(\theta + \pi) = (-1)^k R_k(\theta), \quad k = 0, 1, 2, \dots \quad (2.4)$$

For sufficient small  $h$ , let

$$\Delta(h) = r(2\pi, h) - h, \quad r = r(\theta, h) = \sum_{m=1}^{\infty} v_m(\theta) h^m \quad (2.5)$$

be the Poincaré successor function and the solution of (2.3) satisfying the initial value condition  $r|_{\theta=0} = h$ .

By using the homeomorphic transformation

$$z = u + iv, \quad w = u - iv, \quad T = it_1, \quad (2.6)$$

system (2.1) is transformed into

$$\begin{cases} \frac{dz}{dT} = z^{n+1}w^n + \sum_{k=2n+2}^{\infty} Z_k(z, w), \\ \frac{dw}{dT} = -w^{n+1}z^n - \sum_{k=2n+2}^{\infty} W_k(z, w), \end{cases} \quad (2.7)$$

where  $Z_k(z, w) = \sum_{\alpha+\beta=k} a_{\alpha,\beta} z^\alpha w^\beta$ ,  $W_k(z, w) = \sum_{\alpha+\beta=k} b_{\alpha,\beta} w^\alpha z^\beta$  are homogeneous polynomials of degree  $k$ , and  $a_{\alpha,\beta} = \overline{b_{\alpha,\beta}}$ .

Let

$$x = z(zw)^{-\frac{(n+1)}{(2n+3)}}, \quad y = w(zw)^{-\frac{(n+1)}{(2n+3)}}, \quad dt = (zw)^n dT, \quad (2.8)$$

system (2.7) is transformed into

$$\begin{cases} \frac{dx}{dt} = x + x \sum_{k=1}^{\infty} \Phi_{k(2n+3)}(x, y), \\ \frac{dy}{dt} = -y - y \sum_{k=1}^{\infty} \Psi_{k(2n+3)}(x, y), \end{cases} \quad (2.9)$$

where  $\Phi_{k(2n+3)}, \Psi_{k(2n+3)}$  are homogeneous polynomials of degree  $k(2n+3)$ . Because  $z = \bar{w}$ , (2.8) is a homeomorphic transformation in some open neighborhood of the origin  $(z, w) = (0, 0)$ .

**Lemma 2.1** ([14]). For system (2.9), we can derive successively the terms of the following formal series

$$F(x, y) = xy \left[ 1 + \sum_{m=1}^{\infty} f_{m(2n+3)}(x, y) \right], \quad (2.10)$$

such that

$$\left. \frac{dF}{dt} \right|_{(2.9)} = \sum_{m=1}^{\infty} \mu_m (xy)^{m(2n+3)+1}, \quad (2.11)$$

where  $f_{m(2n+3)}$  are homogeneous polynomials of degree  $m(2n+3)$ .

**Lemma 2.2** ([14]). System (2.9) has a complex center at the origin if and only if there exists a non-zero real number  $s$  and a first integral of the form

$$\tilde{F}(x, y) = (xy)^s \left[ 1 + \sum_{m=1}^{\infty} \tilde{f}_{m(2n+3)}(x, y) \right], \quad (2.12)$$

where  $\tilde{f}_{m(2n+3)}$  are homogeneous polynomials of degree  $m(2n+3)$ . The power series in (2.12) has a non-zero convergence radius.

**Definition 2.3** ([13–15]). For any positive integer  $m$ , the number  $\mu_m$  is called the  $m$ -th singular point value of system (2.7) at the origin. And  $v_{2m+1}(2\pi) \sim i\pi\mu_m$  is called the  $m$ -th focal value of system (2.1) at the origin.

**Definition 2.4** ([13–15]). If for all  $m$ ,  $\mu_m = 0$ , then the origin of system (2.7) is called a complex center. If for all  $m$ ,  $v_{2m+1}(2\pi) = 0$ , then the origin of real system (2.1) is a center.

**Lemma 2.5.** *The origin of system (2.7) is a complex center if and only if the origin of system (2.9) is a complex center.*

*Proof. Necessity.* Suppose that system (2.7) has a complex center at the origin, then for all  $m$ ,  $\mu_m = 0$ . Hence system (2.9) has a formal first integral  $F(x, y)$  of the form (2.10), so by Lemma 1.6, it has a complex center at the origin.

*Sufficiency.* Suppose that system (2.9) has a complex center at the origin, then by Lemma 2.2 it has an analytic first integral  $\tilde{F}(x, y)$  of the form (2.12). Thus it also has an analytic first integral of the form  $F(x, y) = [\tilde{F}(x, y)]^{\frac{1}{5}}$ , which implies  $\mu_m = 0$  for all  $m$  by Lemma 2.1, and so that the origin of system (2.7) is a complex center.  $\square$

Because system (2.9) is integrable at the origin if and only if the origin of it is a complex center, we have the following theorem.

**Theorem 2.6.** *The origin of system (2.7) is a complex center if and only if system (2.9) is integrable at the origin.*

As a consequence of Theorem 2.6, we have the following corollary.

**Corollary 2.7.** *The origin of the real system (2.1) is a center if and only if system (2.9) is integrable at the origin.*

The authors of [26] obtain the center conditions of the following system:

$$\begin{cases} \frac{dz}{dt} = z^2 w (1 + a z + b w), \\ \frac{dw}{dt} = -z w^2 (2 + c z + d w). \end{cases}$$

In this paper, we consider the center problem of a class of complex quartic systems

$$\begin{cases} \frac{dz}{dT} = z^2 w + a_1 z^4 + a_2 z^2 w^2 + a_3 w^4, \\ \frac{dw}{dT} = -z w^2 + b_1 z^4 + b_2 z^2 w^2 + b_3 w^4, \end{cases} \quad (2.13)$$

where

$$a_1 = -\bar{b}_3, \quad a_2 = -\bar{b}_2, \quad a_3 = -\bar{b}_1. \quad (2.14)$$

Using the non-linear change (2.8) for  $n = 1$ , system (2.13) becomes

$$\begin{cases} \frac{dx}{dt} = x - \frac{2}{5} b_1 x^6 + \frac{3}{5} a_1 y x^5 - \frac{2}{5} b_2 y^2 x^4 + \frac{3}{5} a_2 y^3 x^3 - \frac{2}{5} b_3 y^4 x^2 + \frac{3}{5} a_3 y^5 x, \\ \frac{dy}{dt} = -y + \frac{3}{5} b_1 y x^5 - \frac{2}{5} a_1 y^2 x^4 + \frac{3}{5} b_2 y^3 x^3 - \frac{2}{5} a_2 y^4 x^2 + \frac{3}{5} b_3 y^5 x - \frac{2}{5} a_3 y^6. \end{cases} \quad (2.15)$$

Applying our method to compute the first thirty  $1 : -1$  resonant focus numbers, we get  $W_1, W_2, \dots, W_{30}$ , where the quantity  $W_k$  is reduced w.r.t. the Gröbner basis of  $\{W_j : j < k\}$ .

$$W_k = 0, \quad k \in \{1, 2, \dots, 25\} \setminus \{10, 15, 20, 25\},$$

$$W_{10} = -\frac{16}{45} a_3 a_1^2 a_2 - \frac{4}{15} a_1 a_2^3 - \frac{4}{25} a_3 a_1 b_2^2 + \frac{4}{25} a_2^2 b_1 b_3 + \frac{16}{45} b_1 b_2 b_3^2 + \frac{4}{15} b_2^3 b_3,$$



$$\begin{aligned}
W_{15} = & \frac{2857}{6750} a_1^3 a_3^2 b_2 + \frac{427}{2700} a_1^2 a_2^3 b_3 + \frac{97}{4500} a_1^2 a_3 b_2^2 b_3 - \frac{169}{200} a_1 a_2^4 b_2 + \frac{3527}{3000} a_1 a_2^3 a_3 b_1 \\
& - \frac{97}{4500} a_1 a_2^2 b_1 b_3^2 - \frac{253}{9000} a_1 a_2 a_3 b_2^3 + \frac{2711}{45000} a_1 a_3^2 b_1 b_2^2 - \frac{427}{2700} a_1 b_2^3 b_3^2 + \frac{27}{100} a_2^5 b_1 \\
& + \frac{253}{9000} a_2^3 b_1 b_2 b_3 - \frac{2711}{45000} a_2^2 a_3 b_1^2 b_3 - \frac{2857}{6750} a_2 b_1^2 b_3^3 + \frac{169}{200} a_2 b_2^4 b_3 - \frac{3527}{3000} a_3 b_1 b_2^3 b_3 \\
& - \frac{27}{100} a_3 b_2^5, \\
W_{20} = & \frac{1235077}{103994800} a_1^3 a_3 b_2^2 b_3^2 - \frac{955578854125547}{4202028745200000} a_1^2 a_2^3 a_3 b_2^2 - \frac{5542584347}{17819109000} a_1 a_2^3 a_3^2 b_1^2 \\
& - \frac{564506438409643}{336162299616000} a_1 a_2^2 a_3 b_2^4 + \frac{24252192229}{386080695000} a_1 a_3^3 b_1^2 b_2^2 - \frac{4890698789}{8236388160} a_1 a_3 b_2^5 b_3 \\
& - \frac{24252192229}{386080695000} a_2^2 a_3^2 b_1^3 b_3 + \frac{5542584347}{17819109000} a_3^2 b_1^2 b_2^3 b_3 + \frac{481204102950071}{882426036492000} a_1^2 a_2^4 b_2 b_3 \\
& - \frac{1235077}{103994800} a_1^2 a_2^2 b_1 b_3^3 + \frac{2683884447677231}{3268244579600000} a_1 a_2^5 b_1 b_3 \\
& + \frac{955578854125547}{2521217247120000} a_1 a_2^3 b_2^3 b_3 - \frac{481204102950071}{882426036492000} a_1 a_2 b_2^4 b_3^2 \\
& + \frac{564506438409643}{336162299616000} a_2^4 b_1 b_2^2 b_3 - \frac{616607103869689}{756365174136000} a_2 a_3 b_1 b_2^4 b_3 \\
& + \frac{955578854125547}{1890912935340000} a_1^2 a_2 a_3 b_2^3 b_3 - \frac{1396275623989}{2316484170000} a_1^2 a_3^2 b_1 b_2^2 b_3 \\
& + \frac{616607103869689}{756365174136000} a_1 a_2^4 a_3 b_1 b_2 + \frac{1396275623989}{2316484170000} a_1 a_2^2 a_3 b_1^2 b_3^2 \\
& + \frac{9548870928976103}{8824260364920000} a_1 a_2 a_3^2 b_1 b_2^3 - \frac{9548870928976103}{8824260364920000} a_2^3 a_3 b_1^2 b_2 b_3 \\
& - \frac{955578854125547}{2521217247120000} a_1^2 a_2^6 - \frac{184074380363917}{156875739820800} a_1 a_2^5 b_2^2 + \frac{136361}{14856400} a_1^3 a_2^3 b_3^2 \\
& - \frac{136361}{14856400} a_1^2 b_2^3 b_3^3 + \frac{1125970354387}{3088645560000} a_2^4 b_1^2 b_3^2 + \frac{184074380363917}{156875739820800} a_2^2 b_2^5 b_3 \\
& - \frac{955578854125547}{1890912935340000} a_1^3 a_2^4 a_3 - \frac{1125970354387}{3088645560000} a_1^2 a_3^2 b_2^4 - \frac{2707696721}{2969851500} a_2^5 a_3 b_1^2 \\
& - \frac{637748287543}{8404057490400} a_2 a_3 b_2^6 + \frac{2707696721}{2969851500} a_3^2 b_1 b_2^5 + \frac{637748287543}{8404057490400} a_2^6 b_1 b_2,
\end{aligned}$$

and  $W_{25}$ ,  $W_{30}$  are very complicated so we do not present these polynomials here, but the interested reader can easily compute them using any computer algebra system.

If condition (2.14) holds, by applying suitable non-degenerate similarity transformation and time scaling, system (2.13) becomes one of the two forms:

$$\begin{cases} \frac{dz}{dT} = z^2 w + a_1 z^4 + z^2 w^2 + a_3 w^4, \\ \frac{dw}{dT} = -z w^2 + b_1 z^4 - z^2 w^2 + b_3 w^4. \end{cases} \quad (2.16)$$

$$\begin{cases} \frac{dz}{dT} = z^2 w + a_1 z^4 + a_3 w^4, \\ \frac{dw}{dT} = -z w^2 + b_1 z^4 + b_3 w^4, \end{cases} \quad (2.17)$$

where

$$a_1 = -\bar{b}_3, \quad a_3 = -\bar{b}_1. \quad (2.18)$$

**Theorem 2.8.** *If condition (2.18) holds, system (2.16) has a complex center at the origin if and only if  $a_1 = -b_3$ ,  $a_3 = -b_1$ .*

**Theorem 2.9.** *If condition (2.18) holds, system (2.17) has a complex center at the origin if and only if one of the following conditions holds:*

(i)  $b_1^3 b_3^5 - a_1^5 a_3^3 = 0$ ,  $a_1 b_3 \neq 0$ ;

(ii)  $a_1 = b_1 = 0$ ;

(iii)  $a_1 = b_3 = 0$ ,  $b_1 \neq 0$ .

### 3 Proof of the Theorems 2.8 and 2.9

Using the transformation (2.8) for  $n = 1$ , system (2.16) becomes

$$\begin{cases} \frac{dx}{dt} = x - \frac{2}{5} b_1 x^6 + \frac{3}{5} a_1 y x^5 + \frac{2}{5} y^2 x^4 + \frac{3}{5} y^3 x^3 - \frac{2}{5} b_3 y^4 x^2 + \frac{3}{5} a_3 y^5 x, \\ \frac{dy}{dt} = -y + \frac{3}{5} b_1 y x^5 - \frac{2}{5} a_1 y^2 x^4 - \frac{3}{5} y^3 x^3 - \frac{2}{5} y^4 x^2 + \frac{3}{5} b_3 y^5 x - \frac{2}{5} a_3 y^6. \end{cases} \quad (3.1)$$

**Lemma 3.1.** *System (3.1) is integrable at the origin if and only if  $a_1 = -b_3$ ,  $a_3 = -b_1$ .*

*Proof. Necessity.* Let  $W_1, W_2, \dots, W_{30}$  be the first thirty  $1 : -1$  resonant focus numbers of system (2.15). By substituting  $a_2 = 1$ ,  $b_2 = -1$  into these numbers respectively, one gets the first thirty  $1 : -1$  resonant focus numbers  $W'_1, W'_2, \dots, W'_{30}$  of system (3.1).

Computing a Gröbner basis of the ideal  $\langle W'_1, W'_2, \dots, W'_{30} \rangle$  with respect to the graded reverse lexicographical order with  $b_1 \succ a_3 \succ b_3 \succ a_1$ , we obtain a list of polynomials  $G = \{b_3 + a_1, b_1 + a_3\}$ . The vanishing of  $G$  leads to  $a_1 = -b_3, a_3 = -b_1$ .

*Sufficiency.* In the case  $a_1 = -b_3, a_3 = -b_1$ , system (3.1) takes the form

$$\begin{cases} \frac{dx}{dt} = x - \frac{2}{5} b_1 x^6 - \frac{3}{5} b_3 x^5 y + \frac{2}{5} x^4 y^2 + \frac{3}{5} x^3 y^3 - \frac{2}{5} b_3 x^2 y^4 - \frac{3}{5} b_1 x y^5, \\ \frac{dy}{dt} = -y + \frac{3}{5} b_1 x^5 y + \frac{2}{5} b_3 x^4 y^2 - \frac{3}{5} x^3 y^3 - \frac{2}{5} x^2 y^4 + \frac{3}{5} b_3 x y^5 + \frac{2}{5} b_1 y^6. \end{cases} \quad (3.2)$$

From Lemma 1.7, we know that system (3.2) is time-reversible w.r.t. the transformation  $x \mapsto y$ ,  $y \mapsto x$ . So by the symmetry principle, the origin of such system is a resonant center, and hence system (3.2) is integrable at the origin.  $\square$

**Lemma 3.2.** *If condition (2.18) holds, system (3.1) is integrable at the origin if and only if  $a_1 = -b_3$ ,  $a_3 = -b_1$ .*

*Proof.* In view of the consistence of condition (2.18) and condition  $a_1 = -b_3, a_3 = -b_1$ , the result follows by Lemma 3.1.  $\square$

Using the transformation (2.8) for  $n = 1$ , system (2.17) becomes

$$\begin{cases} \frac{dx}{dt} = x - \frac{2}{5} b_1 x^6 + \frac{3}{5} a_1 y x^5 - \frac{2}{5} b_3 y^4 x^2 + \frac{3}{5} a_3 y^5 x, \\ \frac{dy}{dt} = -y + \frac{3}{5} b_1 y x^5 - \frac{2}{5} a_1 y^2 x^4 + \frac{3}{5} b_3 y^5 x - \frac{2}{5} a_3 y^6. \end{cases} \quad (3.3)$$

**Lemma 3.3.** *System (3.3) is integrable at the origin if and only if one of the following conditions holds:*

- (1)  $b_1^3 b_3^5 - a_1^5 a_3^3 = 0, a_1 b_3 \neq 0$ ;
- (2)  $a_1 = a_3 = 0, b_1 b_3 \neq 0$ ;
- (3)  $a_1 = b_1 = 0$ ;
- (4)  $a_1 = b_3 = 0, b_1 \neq 0$ ;
- (5)  $a_3 = b_3 = 0, a_1 \neq 0$ ;
- (6)  $b_1 = b_3 = 0, a_1 a_3 \neq 0$ .

*Proof. Necessity.* Let  $W_1, W_2, \dots, W_{30}$  be the first thirty  $1 : -1$  resonant focus numbers of system (2.15). By substituting  $a_2 = b_2 = 0$  into these numbers respectively, one gets the first thirty  $1 : -1$  resonant focus numbers  $W'_1, W'_2, \dots, W'_{30}$  of system (3.3).

Computing a Gröbner basis of the ideal  $\langle W'_1, W'_2, \dots, W'_{30} \rangle$  with respect to the graded reverse lexicographical order with  $b_1 \succ a_3 \succ b_3 \succ a_1$  and we get a list of polynomials

$$G = \left\{ 16504950 a_1^6 a_3^3 b_3 - 6113731 a_1^5 a_3^4 b_1 - 16504950 a_1 b_1^3 b_3^6 + 6113731 a_3 b_1^4 b_3^5, \right. \\ \left. - a_1^7 a_3^3 b_3^2 + a_1^2 b_1^3 b_3^7 \right\}.$$

The vanishing of  $G$  gives rise to six cases in the Lemma.

*Sufficiency.* When condition (1) holds, system (3.3) is of the form

$$\begin{cases} \frac{dx}{dt} = x - \frac{2}{5} b_1 x^6 + \frac{3}{5} a_1 y x^5 - \frac{2}{5} b_3 y^4 x^2 + \frac{3}{5} \frac{b_3^{\frac{5}{3}} b_1}{a_1^{\frac{5}{3}}} y^5 x, \\ \frac{dy}{dt} = -y + \frac{3}{5} b_1 y x^5 - \frac{2}{5} a_1 y^2 x^4 + \frac{3}{5} b_3 y^5 x - \frac{2}{5} \frac{b_3^{\frac{5}{3}} b_1}{a_1^{\frac{5}{3}}} y^6. \end{cases} \quad (3.4)$$

From Lemma 1.7, we know that system (3.4) is time-reversible w.r.t. the transformation

$$x \mapsto \gamma_0 y, \quad y \mapsto \gamma_0^{-1} x, \quad \text{where } \gamma_0 = \left( \frac{-b_3}{a_1} \right)^{\frac{1}{3}},$$

so by the symmetry principle, the origin of it is a resonant center, and hence system (3.4) is integrable.

If condition (2) holds, system (3.3) is reduced to

$$\begin{cases} \frac{dx}{dt} = x - \frac{2}{5} b_1 x^6 - \frac{2}{5} b_3 y^4 x^2, \\ \frac{dy}{dt} = -y + \frac{3}{5} b_1 y x^5 + \frac{3}{5} b_3 y^5 x. \end{cases} \quad (3.5)$$

We will show that for system (3.5) there exists a formal first integral in the form  $F(x, y) = \sum_{n=1}^{\infty} v_n(y) x^n$ , where functions  $v_n(y)$  should satisfy the first-order linear differential equation

$$\frac{dv_n}{dy} = \frac{n}{y} v_n - \frac{2}{5} (n-1) b_3 y^3 v_{n-1} - \frac{2}{5y} (n-5) b_1 v_{n-5} + \frac{3}{5} b_3 y^4 v'_{n-1} + \frac{3}{5} b_1 v'_{n-5}. \quad (3.6)$$

Solving this equation, we obtain

$$\begin{aligned}
v_1(y) &= y, & v_2(y) &= \frac{1}{15} b_3 y^5, & v_3(y) &= \frac{11}{450} b_3^2 y^9, \\
v_4(y) &= \frac{77}{6750} b_3^3 y^{13}, & v_5(y) &= \frac{2387}{405000} b_3^4 y^{17}, \\
v_6(y) &= \frac{1}{30375000} \left( 97867 b_3^5 y^{20} - 1215000 b_1 \right) y, \\
v_7(y) &= \frac{41}{911250000} b_3 \left( 40579 b_3^5 y^{20} - 2430000 b_1 \right) y^5, \\
v_8(y) &= \frac{1}{13668750000} b_3^2 \left( 14498297 b_3^5 y^{20} + 1104435000 b_1 \right) y^9, \\
v_9(y) &= \frac{11}{164025000000} b_3^3 \left( 93579917 b_3^5 y^{20} + 9263160000 b_1 \right) y^{13}, \\
v_{10}(y) &= \frac{11}{2733750000000} b_3^4 \left( 93579917 b_3^5 y^{20} + 10979010000 b_1 \right) y^{17}.
\end{aligned}$$

taking the integration constants for  $n = 1$  and for  $n > 1$  equal to 1 and 0, respectively. We will show by induction that

$$\begin{aligned}
v_{5k+1}(y) &= y P_{k,1}(y^{20}), \\
v_{5k+2}(y) &= y^5 P_{k,2}(y^{20}), \\
v_{5k+3}(y) &= y^9 P_{k,3}(y^{20}), \\
v_{5k+4}(y) &= y^{13} P_{k,4}(y^{20}), \\
v_{5k+5}(y) &= y^{17} P_{k,5}(y^{20}),
\end{aligned}$$

for  $k \geq 1$ , where  $P_{k,j}$  are polynomials of degree  $k$ . Hence we assume that for  $k = s$ , the assertion is true. We then solve the linear differential equation (3.6) for  $n = 5s + 6$  and obtain

$$v_{5s+6}(y) = -\frac{1}{5} y^{(5s+6)} \int y^{-(5s+7)} g_{20s+21}(y) dy, \quad (3.7)$$

taking the integration constant to be 0, where

$$g_{20s+21}(y) = 2(5s+1)b_1 v_{5s+1} + 10(s+1)b_3 y^4 v_{5s+5} - 3y(b_3 y^4 v'_{5s+5} + b_1 v'_{5s+1}).$$

According to the hypothesis for  $k = s$ , we can see that the integrand of (3.7) involves no terms of  $y^{-1}$  and  $g_{20s+21}$  is a polynomial of degree  $20s + 21$ . Consequently,  $v_{5s+6}(y) = v_{5(s+1)+1}(y)$  must be of the form

$$v_{5s+6}(y) = y P_{s+1,1}(y^{20}),$$

where  $P_{s+1,1}$  is a polynomial of degree  $s + 1$ . In a similar way, we can also prove that  $v_{5s+7}(y)$ ,  $v_{5s+8}(y)$ ,  $v_{5s+9}(y)$ ,  $v_{5s+10}(y)$  are of the forms

$$\begin{aligned}
v_{5s+7}(y) &= y^5 P_{s+1,2}(y^{20}), \\
v_{5s+8}(y) &= y^9 P_{s+1,3}(y^{20}), \\
v_{5s+9}(y) &= y^{13} P_{s+1,4}(y^{20}), \\
v_{5s+10}(y) &= y^{17} P_{s+1,5}(y^{20}).
\end{aligned}$$

Hence, we have proved that system (3.5) admits a formal first integral of the form  $F(x, y) = \sum_{n=1}^{\infty} v_n(y) x^n$ . Consequently it has an analytic first integral in some neighborhood of the origin.

If condition (3) holds, system (3.3) is reduced to

$$\begin{cases} \frac{dx}{dt} = x - \frac{2}{5} b_3 y^4 x^2 + \frac{3}{5} a_3 y^5 x, \\ \frac{dy}{dt} = -y + \frac{3}{5} b_3 y^5 x - \frac{2}{5} a_3 y^6. \end{cases} \quad (3.8)$$

We will show that for system (3.8) there exists a formal first integral in the form  $F(x, y) = \sum_{n=1}^{\infty} v_n(x) y^n$ , where functions  $v_n(x)$  should satisfy the first-order linear differential equation

$$\frac{dv_n}{dx} = \frac{n}{x} v_n - \frac{3}{5} (n-4) b_3 v_{n-4} + \frac{2(n-5) a_3}{5x} v_{n-5} + \frac{2}{5} b_3 x v'_{n-4} - \frac{3}{5} a_3 v'_{n-5}. \quad (3.9)$$

Solving this equation, we obtain

$$\begin{aligned} v_k(x) &= x^k, \quad k = 1, 2, 3, 4. \\ v_5(x) &= \frac{1}{15} x^2 (15x^3 + b_3). \end{aligned}$$

taking the integration constants equal to 1. We will show by induction that the functions  $v_n(x)$  are polynomials of degree  $n$ . Hence, we assume that for  $k = 1, 2, \dots, n-1$  there exist  $k$ -th degree polynomials  $v_k(x)$  satisfying (3.9). We then solve the linear differential equation (3.9) for  $k = n$  and obtain

$$v_n(x) = \left( 1 - \frac{1}{5} \int x^{-(n+1)} g_{n-3}(x) dx \right) x^n, \quad (3.10)$$

where

$$g_{n-3}(x) = -2b_3 v'_{n-4} x^2 + (3nb_3 v_{n-4} + 3a_3 v'_{n-5} - 12b_3 v_{n-4})x - 2(n-5)a_3 v_{n-5},$$

taking the integration constant equals to 1. Using the induction hypothesis that

$$\deg(v_{n-5}(x)) = n-5, \quad \deg(v_{n-4}(x)) = n-4,$$

we find that the degree of  $g_{n-3}(x)$  is at most  $n-3$ . Now, we must study whether the integral can give any logarithmic terms. Therefore, we must prove that terms involving  $x^{-1}$  do not appear in the integrand of (3.10). Since the exponents that can appear in the integrand are of the form

$$-(s+4), \quad s = 0, 1, 2, \dots, n-3,$$

there can be no logarithmic terms in (3.10) and  $v_n(x)$  is an  $n$ -th degree polynomial in  $x$ . Hence, we have proved that system (3.8) admits a formal first integral of the form  $F(x, y) = \sum_{n=1}^{\infty} v_n(x) y^n$ . Consequently it has an analytic first integral around the origin.

If condition (4) holds, system (3.3) is of the form

$$\begin{cases} \frac{dx}{dt} = x - \frac{2}{5} b_1 x^6 + \frac{3}{5} a_3 y^5 x, \\ \frac{dy}{dt} = -y + \frac{3}{5} b_1 y x^5 - \frac{2}{5} a_3 y^6. \end{cases} \quad (3.11)$$

From Lemma 1.7, we know that system (3.11) is time-reversible w.r.t. the transformation

$$x \mapsto \gamma_0 y, \quad y \mapsto \gamma_0^{-1} x, \quad \text{where } \gamma_0 = \sqrt[5]{\frac{-a_3}{b_1}},$$

so by the symmetry principle, the origin of system is a resonant center, and hence system (3.11) is integrable.

If condition (5) holds, system (3.3) is of the form

$$\begin{cases} \frac{dx}{dt} = x - \frac{2}{5}b_1x^6 + \frac{3}{5}a_1yx^5, \\ \frac{dy}{dt} = -y + \frac{3}{5}b_1yx^5 - \frac{2}{5}a_1y^2x^4. \end{cases} \quad (3.12)$$

We will show that for system (3.12) there exists a formal first integral in the form  $F(x, y) = \sum_{n=1}^{\infty} v_n(y)x^n$ , where functions  $v_n(y)$  should satisfy the first-order linear differential equation

$$\frac{dv_n}{dy} = \frac{n}{y}v_n + \frac{3}{5}(n-4)a_1v_{n-4} - \frac{2(n-5)b_1}{5y}v_{n-5} - \frac{2}{5}a_1yv'_{n-4} + \frac{3}{5}b_1v'_{n-5}. \quad (3.13)$$

Solving this equation, we obtain

$$\begin{aligned} v_k(y) &= y^k, \quad k = 1, 2, 3, 4. \\ v_5(y) &= \frac{1}{15}y^2(15y^3 - a_1), \end{aligned}$$

taking the integration constants equal to 1. We will show by induction that the functions  $v_n(y)$  are polynomials of degree  $n$ . Hence, we assume that for  $k = 1, 2, \dots, n-1$  there exist  $k$ -th degree polynomials  $v_k(y)$  satisfying (3.13). We then solve the linear differential equation (3.13) for  $k = n$  and obtain

$$v_n(y) = \left(1 - \frac{1}{5} \int y^{-(n+1)} g_{n-3}(y) dy\right) y^n, \quad (3.14)$$

where

$$g_{n-3}(y) = 2a_1v'_{n-4}y^2 + (-3na_1v_{n-4} + 12a_1v_{n-4} - 3b_1v'_{n-5})y + 2(n-5)b_1v_{n-5},$$

taking the integration constant equals to 1. Using the induction hypothesis that

$$\deg(v_{n-5}(y)) = n-5, \quad \deg(v_{n-4}(y)) = n-4,$$

we find that the degree of  $g_{n-3}(y)$  is  $n-3$ . Now, we must study whether the integral can give any logarithmic terms. Therefore, we must prove that terms involving  $y^{-1}$  do not appear in the integrand of (3.14). Since the exponents that can appear in the integrand are of the form

$$-(s+4), \quad s = 0, 1, 2, \dots, n-3,$$

there can be no logarithmic terms in (3.14) and  $v_n(y)$  is an  $n$ -th degree polynomial in  $y$ . Hence, we have proved that system (3.12) admits a formal first integral of the form  $F(x, y) = \sum_{n=1}^{\infty} v_n(y)x^n$ . Consequently it has an analytic first integral around the origin.

If condition (6) holds, system (3.3) is reduced to

$$\begin{cases} \frac{dx}{dt} = x + \frac{3}{5}a_1yx^5 + \frac{3}{5}a_3y^5x, \\ \frac{dy}{dt} = -y - \frac{2}{5}a_1y^2x^4 - \frac{2}{5}a_3y^6. \end{cases} \quad (3.15)$$

By the transformation  $x \mapsto y, y \mapsto x, t \mapsto -t$ , system (3.15) can be transformed into the form of (3.5), and therefore system (3.15) is integrable at the origin.  $\square$

**Lemma 3.4.** *If condition (2.18) holds, system (3.3) is integrable at the origin if and only if one of the following conditions holds:*

(i)  $b_1^3 b_3^5 - a_1^5 a_3^3 = 0$ ,  $a_1 b_3 \neq 0$ ;

(ii)  $a_1 = b_1 = 0$ ;

(iii)  $a_1 = b_3 = 0$ ,  $b_1 \neq 0$ .

*Proof.* Elementary computations shows that condition (2.18) is consistent with conditions (1), (3), (4) in Lemma 3.3, and is inconsistent with the other conditions (2), (5), (6). Hence by Lemma 3.3, the result follows.  $\square$

**The proof for Theorems 2.8 and 2.9.** According to Theorem 2.6 in Section 2, we can arrive at Theorems 2.8 and 2.9 by using Lemmas 3.2 and 3.4, respectively.

## Acknowledgements

This work is supported by a grant from the National Natural Science Foundation of China (No. 11401285) and a grant from the Foundation for Research in Experimental Techniques of Liaocheng University (No. LDSY2014110). The author is grateful to Prof. Colin Christopher, Prof. Xiang Zhang, Prof. Qi Zhang and Prof. Qinlong Wang for their valuable remarks and comments. The author is also grateful to the referees for their valuable remarks which helped to improve the manuscript.

## References

- [1] M. J. ÁLVAREZ, A. GASULL, Monodromy and stability for nilpotent critical points, *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* **15**(2005), 1253–1265. [MR2152073](#); [url](#)
- [2] H. B. CHEN, Y. R. LIU, Linear recursion formulas of quantities of singular point and applications, *Appl. Math. Comput.* **148**(2004), 163–171. [MR2014632](#); [url](#)
- [3] X. W. CHEN, J. GINÉ, V. G. ROMANOVSKI, D. S. SHAFER, The  $1 : -q$  resonant center problem for certain cubic Lotka–Volterra systems, *Appl. Math. Comput.* **218**(2012), 11620–11633. [MR2944006](#); [url](#)
- [4] B. FERČEC, X. W. CHEN, V. G. ROMANOVSKI, Integrability conditions for complex systems with homogeneous quintic nonlinearities, *J. Appl. Anal. Comput.* **1**(2011), 9–12. [MR2881844](#)
- [5] B. FERČEC, J. GINÉ, Y. R. LIU, V. G. ROMANOVSKI, Integrability conditions for Lotka–Volterra planar complex quartic systems having homogeneous nonlinearities, *Acta. Appl. Math.* **124**(2013), 107–122. [MR3029242](#); [url](#)
- [6] A. FERRO, G. GALLO, Automated theorem proving in elementary geometry, *Matematiche (Catania)* **43**(1988), 195–224. [MR1075559](#)
- [7] A. GASULL, J. TORREGROSA, Center problem for several differential equations via Cherkas' method, *J. Math. Anal. Appl.* **228**(1998), 322–343. [MR1663628](#); [url](#)

- [8] J. GINÉ, V. G. ROMANOVSKI, Integrability conditions for Lotka–Volterra planar complex quintic systems, *Nonlinear Anal. Real World Appl.* **11**(2010), 2100–2105. [MR2646619](#); [url](#)
- [9] J. GINÉ, On the degenerate center problem, *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* **21**(2011), 1383–1392. [MR2819828](#); [url](#)
- [10] J. GINÉ, C. CHRISTOPHER, M. PRESERN, V. G. ROMANOVSKI, N. L. SHCHEGLOVA, The resonant center problem for a  $2 : -3$  resonant cubic Lotka–Volterra system, in: *Computer Algebra in Scientific Computing*, Lecture Notes in Computer Science, Vol. 7442 (14th International Workshop, CASC 2012, Maribor, Slovenia, September 3–6, 2012. Proceedings), Springer-Verlag, Berlin Heidelberg, 2012, 129–142. [url](#)
- [11] Z. P. HU, V. G. ROMANOVSKI, D. S. SHAFER,  $1 : -3$  resonant centers on  $\mathbb{C}^2$  with homogeneous cubic nonlinearities, *Comput. Math. Appl.* **56**(2008), 1927–1940. [MR2466695](#); [url](#)
- [12] C. J. LIU, G. T. CHEN, G. R. CHEN, Integrability of Lotka–Volterra type systems of degree 4, *J. Math. Anal. Appl.* **388**(2012), 1107–1116. [MR2869810](#); [url](#)
- [13] Y. R. LIU, Theory of center-focus for a class of higher-degree critical point and infinite points, *Sci. China Ser. A* **44**(2001), 365–377. [MR1828761](#); [url](#)
- [14] Y. R. LIU, J. B. LI, W. T. HUANG, *Singular point values, center problem and bifurcations of limit cycles of two dimensional differential autonomous systems*, Nonlinear Dynamics Series, Vol. 6, Science Press, Beijing, 2008.
- [15] Y. R. LIU, J. B. LI, W. T. HUANG, *Planar dynamical systems: selected classical problems*, Mathematics Monograph Series, Vol. 28, Science Press, Beijing, 2014.
- [16] J. F. MATTEI, R. MOUSSU, Holonomie et intégrales premières (in French) [Holonomy and first integrals], *Ann. Sci. École Norm. Sup. (4)* **13**(1980), 469–523. [MR608290](#)
- [17] V. MAÑOSA, On the center problem for degenerate singular points of planar vector fields, *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* **12**(2002), 687–707. [MR1901192](#); [url](#)
- [18] V. G. ROMANOVSKI, N. L. SHCHEGLOVA, The integrability conditions for two cubic vector fields, *Differ. Equa.* **36**(2000), 108–112. [url](#)
- [19] V. G. ROMANOVSKI, D. S. SHAFER, On the center problem for  $p : -q$  resonant polynomial vector fields, *Bull. Belg. Math. Soc. Simon Stevin* **15**(2008), 871–887. [MR2484138](#)
- [20] V. G. ROMANOVSKI, Time-reversibility in 2-dimensional systems, *Open Syst. Inf. Dyn.* **15**(2008), 359–370. [MR2508125](#); [url](#)
- [21] D. M. WANG, Mechanical manipulation for a class of differential systems, *J. Symbolic Comput.* **12**(1991), 233–254. [MR1125940](#); [url](#)
- [22] D. K. WANG, L. H. ZHI, Software development in MMRC, *Proc. 1st Asian Tech. Confer. in Math., Singapore, 1995*, Assoc. Math. Educators, Singapore, 551–560.
- [23] D. M. WANG, *Elimination methods*, Texts and Monographs in Symbolic Computation, Springer-Verlag, Vienna, 2001. [MR1826878](#); [url](#)
- [24] W. T. WU, *Mathematics mechanization*, Mathematics and its Applications, Vol. 489, Kluwer Academic Publishers Group, Dordrecht, Science Press, Beijing, 2000. [MR1834540](#)



- [25] P. YU, Computation of normal forms via a perturbation technique, *J. Sound Vibration* **211**(1998), 19–38. [MR1615733](#); [url](#)
- [26] Q. ZHANG, W. H. GUI, Y. R. LIU, The generalized center problem of degenerate resonant singular point, *Bull. Sci. Math.* **133**(2009), 198–204. [MR2494467](#); [url](#)
- [27] H. ŻOŁĄDEK, The problem of center for resonant singular points of polynomial vector fields, *J. Differential Equations* **137**(1997), 94–118. [MR1451537](#); [url](#)