



The Dirichlet problem for discontinuous perturbations of the mean curvature operator in Minkowski space

Cristian Bereanu^{1,2}, Petru Jebelean³ and Călin Șerban^{✉3}

¹Faculty of Mathematics and Computer Science, University of Bucharest,
14 Academiei Street, Bucharest, RO-70109, Romania

²Institute of Mathematics “Simion Stoilow” of the Romanian Academy,
21 Calea Griviței, Sector 1, Bucharest, RO-010702, Romania

³Department of Mathematics, West University of Timișoara,
4 Blvd. V. Pârvan, Timișoara, RO-300223, Romania

Received 15 April 2015, appeared 6 July 2015

Communicated by Gabriele Bonanno

Abstract. Using the critical point theory for convex, lower semicontinuous perturbations of locally Lipschitz functionals, we prove the solvability of the discontinuous Dirichlet problem involving the operator $u \mapsto \operatorname{div} \left(\frac{\nabla u}{\sqrt{1-|\nabla u|^2}} \right)$.

Keywords: nonsmooth critical point theory, discontinuous Dirichlet problem, mean curvature operator, Palais–Smale condition.

2010 Mathematics Subject Classification: 34A60, 49J40, 49J52.

1 Introduction

Let Ω be an open bounded set in \mathbb{R}^N ($N \geq 2$) with boundary $\partial\Omega$ of class C^2 and $f: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ be a measurable function satisfying the growth condition

$$|f(x, s)| \leq C(1 + |s|^{q-1}), \quad \text{a.e. } x \in \Omega \text{ and all } s \in \mathbb{R}, \quad (1.1)$$

with some $q \in (1, \infty)$ and C a positive constant. For a.e. $x \in \Omega$ and all $s \in \mathbb{R}$, we denote

$$\underline{f}(x, s) := \lim_{\delta \searrow 0} \operatorname{ess\,inf} \{f(x, t) : |t - s| < \delta\}$$

and

$$\overline{f}(x, s) := \lim_{\delta \searrow 0} \operatorname{ess\,sup} \{f(x, t) : |t - s| < \delta\}.$$

In this paper we consider the discontinuous Dirichlet problem with mean curvature operator in Minkowski space:

$$\mathcal{M}(u) := \operatorname{div} \left(\frac{\nabla u}{\sqrt{1-|\nabla u|^2}} \right) \in [\underline{f}(x, u), \overline{f}(x, u)] \quad \text{in } \Omega, \quad u|_{\partial\Omega} = 0. \quad (1.2)$$

[✉]Corresponding author. Email: cserban2005@yahoo.com

We assume that

$$\underline{f} \text{ and } \bar{f} \text{ are } N\text{-measurable} \quad (1.3)$$

(recall, a function $h: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is called *N-measurable* if $h(\cdot, v(\cdot)): \Omega \rightarrow \mathbb{R}$ is measurable whenever $v: \Omega \rightarrow \mathbb{R}$ is measurable [3]).

By a *solution* of (1.2) we mean a function $u \in W^{2,p}(\Omega)$ for some $p > N$, such that $\|\nabla u\|_\infty < 1$, which satisfies

$$\mathcal{M}(u)(x) \in \left[\underline{f}(x, u(x)), \bar{f}(x, u(x)) \right], \quad \text{a.e. } x \in \Omega$$

and vanishes on $\partial\Omega$. At our best knowledge, this type of solutions, but for differential inclusions was firstly considered by A. F. Filippov [7]. Also, for partial differential inclusions we refer the reader to the pioneering works of I. Massabo and C. A. Stuart [12], J. Rauch [14], C. A. Stuart and J. F. Toland [16].

This work is motivated by the recent advances in the study of boundary value problems involving the operator \mathcal{M} (see [2, 6] and the references therein) and by the seminal paper of K.-C. Chang [4] where the classical critical point theory is extended to locally Lipschitz functionals in order to study the problem

$$\Delta u \in \left[\underline{f}(x, u), \bar{f}(x, u) \right] \quad \text{in } \Omega, \quad u|_{\partial\Omega} = 0.$$

It is worth to point out that the operators \mathcal{M} and Δ have essentially different structures and the theory developed in [4] appears as not being applicable to problem (1.2). Thus, we shall use a more general critical point theory, namely the one concerning convex, lower semicontinuous perturbations of locally Lipschitz functionals, which was developed by D. Motreanu and P. D. Panagiotopoulos [13] (also, see [10, 11]). It should be noticed that, using this theory, various existence results concerning Filippov type solutions for Dirichlet, periodic and Neumann problems involving the “*p*-relativistic” operator

$$u \mapsto \left(\frac{|u'|^{p-2} u'}{(1 - |u'|^p)^{1-1/p}} \right)'$$

were obtained in the recent paper [9].

A first existence result for the Dirichlet problem involving the operator \mathcal{M} was obtained by F. Flaherty in [8], where it is shown that problem

$$\mathcal{M}(u) = 0 \quad \text{in } \Omega, \quad u|_{\partial\Omega} = \varphi,$$

has at least one solution, provided that $\partial\Omega$ has non-negative mean curvature and $\varphi \in C^2(\bar{\Omega})$ with $\|\nabla\varphi\|_\infty < 1$. The result was generalized in [1] by R. Bartnik and L. Simon, proving that problem

$$\mathcal{M}(u) = g(x, u) \quad \text{in } \Omega, \quad u|_{\partial\Omega} = 0 \quad (1.4)$$

is solvable, provided that the Carathéodory function $g: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is bounded. More general, if g satisfies the L^∞ -growth condition:

for each $\rho > 0$ there is some $\alpha_\rho \in L^\infty(\Omega)$ such that

$$|g(x, s)| \leq \alpha_\rho(x) \quad \text{for a.e. } x \in \Omega, \quad \forall s \in \mathbb{R} \text{ with } |s| \leq \rho,$$

it is shown in [2, Theorem 2.1] that (1.4) is still solvable. The approach in [2] relies on Szulkin’s critical point theory [17]. The aim of the present paper is to obtain a similar result for the

discontinuous problem (1.2). Precisely, we show in the main result (Theorem 4.1) that under assumptions (1.1) and (1.3) problem (1.2) always has at least one solution.

The rest of the paper is organized as follows. In Section 2 we recall some notions from nonsmooth analysis which will be needed in the sequel. The variational formulation of problem (1.2) is a key step in our approach and it is given in Section 3. Section 4 is devoted to the proof of the main result.

2 Preliminaries

Let $(X, \|\cdot\|)$ be a real Banach space and X^* its topological dual. A functional $\mathcal{G}: X \rightarrow \mathbb{R}$ is called *locally Lipschitz* if for each $u \in X$, there is a neighborhood \mathcal{N}_u of u and a constant $k > 0$ depending on \mathcal{N}_u such that

$$|\mathcal{G}(w) - \mathcal{G}(z)| \leq k\|w - z\|, \quad \forall w, z \in \mathcal{N}_u.$$

For such a function \mathcal{G} , the *generalized directional derivative* at $u \in X$ in the direction of $v \in X$ is defined by

$$\mathcal{G}^0(u; v) = \limsup_{w \rightarrow u, t \searrow 0} \frac{\mathcal{G}(w + tv) - \mathcal{G}(w)}{t}$$

and the *generalized gradient* (in the sense of Clarke [5]) of \mathcal{G} at $u \in X$ is defined as being the subset of X^*

$$\partial\mathcal{G}(u) = \{\eta \in X^* : \mathcal{G}^0(u; v) \geq \langle \eta, v \rangle, \quad \forall v \in X\},$$

where $\langle \cdot, \cdot \rangle$ stands for the duality pairing between X^* and X . For more details concerning the properties of the generalized directional derivative and of the generalized gradient we refer to [5].

If $\mathcal{I}: X \rightarrow (-\infty, +\infty]$ is a functional having the structure

$$\mathcal{I} = \Phi + \mathcal{G}, \tag{2.1}$$

with $\mathcal{G}: X \rightarrow \mathbb{R}$ locally Lipschitz and $\Phi: X \rightarrow (-\infty, +\infty]$ proper, convex and lower semicontinuous, then an element $u \in X$ is said to be a *critical point* of \mathcal{I} provided that

$$\mathcal{G}^0(u; v - u) + \Phi(v) - \Phi(u) \geq 0, \quad \forall v \in X.$$

The number $c = \mathcal{I}(u)$ is called a *critical value* of \mathcal{I} corresponding to the critical point u . According to Kourogenis *et al.* [10], $u \in X$ is a critical point of \mathcal{I} iff

$$0 \in \partial\mathcal{G}(u) + \bar{\partial}\Phi(u),$$

where $\bar{\partial}\Phi(u)$ stands for the subdifferential of Φ at $u \in X$ in the sense of convex analysis [15], i.e.,

$$\bar{\partial}\Phi(u) = \{\eta \in X^* : \Phi(v) - \Phi(u) \geq \langle \eta, v - u \rangle, \quad \forall v \in X\}.$$

Also, \mathcal{I} in (2.1) is said to *satisfy the Palais–Smale condition* (in short, *PS condition*) if every sequence $(u_n) \subset X$ for which $(\mathcal{I}(u_n))$ is bounded and

$$\mathcal{G}^0(u_n; v - u_n) + \Phi(v) - \Phi(u_n) \geq -\varepsilon_n\|v - u_n\|, \quad \forall v \in X,$$

for a sequence $(\varepsilon_n) \subset \mathbb{R}_+$ with $\varepsilon_n \rightarrow 0$, possesses a convergent subsequence.

Theorem 2.1. ([11, Theorem 1]) *If \mathcal{I} is bounded from below and satisfies the (PS) condition then $c = \inf_X \mathcal{I}$ is a critical value of \mathcal{I} .*

3 The variational setting

In the sequel we shall give the variational formulation of problem (1.2). With this aim, we introduce the set

$$K_0 = \left\{ v \in W^{1,\infty}(\Omega) : \|\nabla v\|_\infty \leq 1, v = 0 \text{ on } \partial\Omega \right\}.$$

Notice that since $W^{1,\infty}(\Omega)$ is continuously (in fact, compactly) embedded into $C(\overline{\Omega})$, the evaluation at $\partial\Omega$ is understood in the usual sense. According to [2], K_0 is compact in $C(\overline{\Omega})$ and one has

$$\|v\|_\infty \leq c(\Omega) \quad \text{for all } v \in K_0, \quad (3.1)$$

with $c(\Omega)$ a positive constant. Also, the functional $\Psi: C(\overline{\Omega}) \rightarrow (-\infty, +\infty]$ given by

$$\Psi(v) = \begin{cases} \int_\Omega \left[1 - \sqrt{1 - |\nabla v|^2} \right], & \text{for } v \in K_0, \\ +\infty, & \text{for } v \in C(\overline{\Omega}) \setminus K_0 \end{cases} \quad (3.2)$$

is proper, convex and lower semicontinuous [2, Lemma 2.4].

Having in view the growth condition (1.1), we define $\widehat{\mathcal{F}}: L^q(\Omega) \rightarrow \mathbb{R}$ by

$$\widehat{\mathcal{F}}(v) = \int_\Omega F(x, v), \quad \forall v \in L^q(\Omega),$$

where

$$F(x, s) = \int_0^s f(x, \xi) d\xi \quad (x \in \Omega, s \in \mathbb{R})$$

and, on account of the embedding $C(\overline{\Omega}) \subset L^q(\Omega)$, we introduce the functional

$$\mathcal{F} = \widehat{\mathcal{F}}|_{C(\overline{\Omega})}. \quad (3.3)$$

From [4, Theorem 2.1], one has that $\widehat{\mathcal{F}}$ is locally Lipschitz in $L^q(\Omega)$ and

$$\partial\widehat{\mathcal{F}}(v) \subset \left[\underline{f}(\cdot, v(\cdot)), \overline{f}(\cdot, v(\cdot)) \right], \quad (3.4)$$

for all $v \in L^q(\Omega)$. Then, by the continuity of the embedding $C(\overline{\Omega}) \subset L^q(\Omega)$ it is clear that \mathcal{F} is locally Lipschitz on $C(\overline{\Omega})$. Also, since $C(\overline{\Omega})$ is dense in $L^q(\Omega)$, the following holds (see [5, p. 47]):

$$\partial\widehat{\mathcal{F}}(v) = \partial\mathcal{F}(v), \quad \forall v \in C(\overline{\Omega}). \quad (3.5)$$

Lemma 3.1. *Let $v \in K_0$. If $\ell \in \partial\mathcal{F}(v)$, then there is some $\zeta_\ell \in L^\infty(\Omega)$ such that $\zeta_\ell(x) \in \left[\underline{f}(x, v(x)), \overline{f}(x, v(x)) \right]$ for a.e. $x \in \Omega$ and*

$$\langle \ell, w \rangle = \int_\Omega \zeta_\ell w \quad (3.6)$$

for all $w \in C(\overline{\Omega})$.

Proof. From (3.5) and (3.4) we infer that there is a function $\zeta_\ell \in L^{q'}(\Omega)$ with $1/q + 1/q' = 1$, such that $\zeta_\ell(x) \in \left[\underline{f}(x, v(x)), \overline{f}(x, v(x)) \right]$ for a.e. $x \in \Omega$ and (3.6) holds true for all $w \in L^q(\Omega)$. To see that $\zeta_\ell \in L^\infty(\Omega)$, from (1.1) and (3.1), one gets

$$-C_1 \leq \underline{f}(x, v(x)) \leq \overline{f}(x, v(x)) \leq C_1, \quad \text{for a.e. } x \in \Omega,$$

with $C_1 = C(1 + c(\Omega)^{q-1})$. This shows that $|\zeta_\ell(x)| \leq C_1$ for a.e. $x \in \Omega$ and the proof is complete. \square

The functional framework of Section 2 fits the following choices: $X = C(\overline{\Omega})$, $\Phi = \Psi$ in (3.2), $\mathcal{G} = \mathcal{F}$ in (3.3) and

$$\mathcal{I} := \Psi + \mathcal{F}.$$

Notice that, the compactness of $K_0 \subset C(\overline{\Omega})$ implies that \mathcal{I} satisfies the (PS) condition.

4 Main result

We have the following theorem.

Theorem 4.1. *Assume that (1.1) and (1.3) hold true. If u is a critical point of \mathcal{I} , then u is a solution of problem (1.2). Moreover, \mathcal{I} is bounded from below and attains its infimum at some $u_0 \in K_0$, which solves problem (1.2).*

Proof. Let u be a critical point of \mathcal{I} . Then $u \in K_0$ and there exist $h_u \in \bar{\partial}\Psi(u)$ and $\ell_u \in \partial\mathcal{F}(u)$ such that

$$\langle h_u, w \rangle + \langle \ell_u, w \rangle = 0, \quad \forall w \in C(\overline{\Omega}).$$

This and the fact that $h_u \in \bar{\partial}\Psi(u)$ yield

$$\Psi(w) - \Psi(u) + \langle \ell_u, w - u \rangle \geq 0, \quad \forall w \in C(\overline{\Omega}). \quad (4.1)$$

Using Lemma 3.1 we deduce that there is some $\zeta_u = \zeta(\ell_u) \in L^\infty(\Omega)$ such that

$$\zeta_u(x) \in \left[\underline{f}(x, u(x)), \bar{f}(x, u(x)) \right], \quad \text{a.e. } x \in \Omega \quad (4.2)$$

and

$$\langle \ell_u, w \rangle = \int_{\Omega} \zeta_u w, \quad \forall w \in C(\overline{\Omega}). \quad (4.3)$$

By virtue of (4.3), inequality (4.1) becomes

$$\Psi(w) - \Psi(u) + \int_{\Omega} \zeta_u (w - u) \geq 0, \quad \forall w \in C(\overline{\Omega}). \quad (4.4)$$

On account of Lemma 2.2 in [6], for each function $e \in L^\infty(\Omega)$, the Dirichlet problem

$$\mathcal{M}(v) = e(x) \quad \text{in } \Omega, \quad v|_{\partial\Omega} = 0$$

has a unique solution $v_e \in W^{2,p}(\Omega)$ for all $1 \leq p < \infty$. Then, from Lemma 2.3 in [2], one has that v_e is the unique solution in K_0 of the variational inequality

$$\int_{\Omega} \left[\sqrt{1 - |\nabla v|^2} - \sqrt{1 - |\nabla w|^2} + e(w - v) \right] \geq 0, \quad \forall w \in K_0$$

and hence,

$$\Psi(w) - \Psi(v_e) + \int_{\Omega} e(w - v_e) \geq 0, \quad \forall w \in C(\overline{\Omega}).$$

From this and (4.4), we infer that $u = v_e$, with $e = \zeta_u$. But, on account of (4.2), this means that u solves problem (1.2).

Next, for arbitrary $u \in K_0$, by (1.1) and (3.1), the primitive F satisfies

$$|F(x, u(x))| \leq C(c(\Omega) + c(\Omega)^q/q) =: C_2, \quad \text{for a.e. } x \in \Omega.$$

Hence,

$$|\mathcal{F}(u)| \leq \int_{\Omega} |F(x, u)| \leq C_2 \text{vol}(\Omega), \quad \forall u \in K_0.$$

We deduce that the functional \mathcal{I} is bounded from below on $C(\overline{\Omega})$. Then, using that \mathcal{I} verifies the (PS) condition and Theorem 2.1, we have that

$$c = \inf_{C(\overline{\Omega})} \mathcal{I} = \inf_{K_0} \mathcal{I}$$

is a critical value of \mathcal{I} and the proof is complete. \square

Acknowledgements

The authors are grateful to the anonymous referee for providing a useful reference for further developments of the subject. The work of Călin Şerban was supported by the strategic grant POSDRU/159/1.5/S/137750, “Project Doctoral and Postdoctoral programs support for increased competitiveness in Exact Sciences research” co-financed by the European Social Fund within the Sectoral Operational Programme Human Resources Development 2007-2013.

References

- [1] R. BARTNIK, L. SIMON, Spacelike hypersurfaces with prescribed boundary values and mean curvature, *Comm. Math. Phys.* **87**(1982), 131–152. [MR0680653](#); [url](#)
- [2] C. BEREANU, P. JEBELEAN, J. MAWHIN, The Dirichlet problem with mean curvature operator in Minkowski space – a variational approach, *Adv. Nonlinear Stud.* **14**(2014), 315–326. [MR3194356](#)
- [3] K.-C. CHANG, The obstacle problem and partial differential equations with discontinuous nonlinearities, *Comm. Pure Appl. Math.* **33**(1980), 117–146. [MR0562547](#); [url](#)
- [4] K.-C. CHANG, Variational methods for nondifferentiable functionals and their applications to partial differential equations, *J. Math. Anal. Appl.* **80**(1981), 102–129. [MR0614246](#); [url](#)
- [5] F.H. CLARKE, *Optimization and nonsmooth analysis*, John Wiley and Sons, New York, 1983. [MR0709590](#); [url](#)
- [6] C. CORSATO, F. OBERSNEL, P. OMARI, S. RIVETTI, Positive solutions of the Dirichlet problem for the prescribed mean curvature equation in Minkowski space, *J. Math. Anal. Appl.* **405**(2013), 227–239. [MR3053503](#); [url](#)
- [7] A. F. FILIPPOV, Right hand side discontinuous differential equations, *Trans. Am. Math. Soc.* **42**(1964), 199–227.
- [8] F. FLAHERTY, The boundary value problem for maximal hypersurfaces, *Proc. Nat. Acad. Sci. USA* **76**(1979), 4765–4767. [MR0545611](#); [url](#)
- [9] P. JEBELEAN, C. ŞERBAN, Boundary value problems for discontinuous perturbations of singular ϕ -Laplacian operator, *J. Math. Anal. Appl.* **431**(2015), 662–681. [MR3357606](#); [url](#)

- [10] N. C. KOUROGENIS, J. PAPADRIANOS, N. S. PAPAGEORGIU, Extensions of nonsmooth critical point theory and applications, *Atti Sem. Mat. Fis. Univ. Modena* **50**(2002), 381–414. [MR1958288](#)
- [11] C. LEFTER, Critical point theorems for lower semicontinuous functionals, *An. Ştiinţ. Univ. Al. I. Cuza Iaşi Mat. (N.S.)* **47**(2001), 189–198. [MR1920202](#)
- [12] I. MASSABO, C. A. STUART, Elliptic eigenvalue problems with discontinuous nonlinearities, *J. Math. Anal. Appl.* **66**(1978), 261–281. [MR0515891](#); [url](#)
- [13] D. MOTREANU, P. D. PANAGIOTOPOULOS, *Minimax theorems and qualitative properties of the solutions of hemivariational inequalities*, Nonconvex Optimization and Its Applications, Vol. 29, Kluwer Academic Publishers, Dordrecht, 1999. [MR1675895](#); [url](#)
- [14] J. RAUCH, Discontinuous semilinear differential equations and multiple valued maps, *Proc. Amer. Math. Soc.* **64**(1977), 277–282. [MR0442453](#); [url](#)
- [15] R. T. ROCKAFFELLAR, *Convex analysis*, Princeton University Press, Princeton, New York, 1970. [MR0274683](#)
- [16] C. A. STUART, J. F. TOLAND, A variational method for boundary value problems with discontinuous nonlinearities, *J. London Math. Soc.* **21**(1980), 319–328. [MR0575391](#); [url](#)
- [17] A. SZULKIN, Minimax principles for lower semicontinuous functions and applications to nonlinear boundary value problems, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **3**(1986), 77–109. [MR0837231](#)