Electronic Journal of Qualitative Theory of Differential Equations 2009, No. 51, 1-1; http://www.math.u-szeged.hu/ejqtde/

Addendum to integrable and continuous solutions of a nonlinear quadratic integral equation

El-Sayed A.M.A and Hashem H.H.G Faculty of Science, Alexandria University, Alexandria, Egypt e.mails : amasayed@hotmail.com, amasayed5@yahoo.com

August 17, 2009

This addendum concerns the paper of the above title found in EJQTDE No. 25 (2008). There are some misprints in that paper:

(i) Page 3, line 5 should be

 $k:[0,1] \times [0,1] \to R_+$ satisfies Carathèodory condition (i.e. measurable in t for all $s \in [0,1]$ and continuous in s for all $t \in [0,1]$) such that $\int_0^1 k(t,s)m_2(s)ds$ is bounded $\forall t \in [0,1]$.

(ii) Page 6, line 6 should be

 $k: [0,1] \times [0,1] \to R_+$ satisfies Carathèodory condition (i.e. measurable in s for all $t \in [0,1]$ and continuous in t for all $s \in [0,1]$) such that $k(t,s)m_2(s) \in L_1 \ \forall t \in [0,1]$.

EJQTDE, 2009 No. 51, p. 1