

## OSCILLATION RESULTS ON MEROMORPHIC SOLUTIONS OF SECOND ORDER DIFFERENTIAL EQUATIONS IN THE COMPLEX PLANE

TING-BIN CAO AND LEI-MIN LI

**ABSTRACT.** The main purpose of this paper is to consider the oscillation theory on meromorphic solutions of second order linear differential equations of the form  $f'' + A(z)f = 0$  where  $A$  is meromorphic in the complex plane. We improve and extend some oscillation results due to Bank and Laine, Kinnunen, Liang and Liu, and others.

### 1. INTRODUCTION AND MAIN RESULTS

Let us define inductively, for  $r \in [0, +\infty)$ ,  $\exp_1 r = e^r$  and  $\exp_{n+1} r = \exp(\exp_n r)$ ,  $n \in \mathbb{N}$ . For all  $r$  sufficiently large, we define  $\log_1 r = \log^+ r = \max\{\log x, 0\}$  and  $\log_{n+1} r = \log(\log_n r)$ ,  $n \in \mathbb{N}$ . We also denote  $\exp_0 r = r = \log_0 r$ ,  $\log_{-1} r = \exp_1 r$  and  $\exp_{-1} r = \log_1 r$ . We assume that the reader is familiar with the fundamental results and the standard notations of the Nevanlinna's value distribution theory of meromorphic functions (e.g. see [8, 21]), such as  $T(r, f)$ ,  $m(r, f)$ , and  $N(r, f)$ . Throughout the paper, a meromorphic function  $f$  means meromorphic in the complex plane  $\mathbb{C}$ . To express the rate of fast growth of meromorphic functions, we recall the following definitions (e.g. see [4, 6, 12, 14, 15, 19]).

**Definition 1.1.** *The iterated  $n$ -order  $\sigma_n(f)$  of a meromorphic function  $f$  is defined by*

$$\sigma_n(f) = \limsup_{r \rightarrow \infty} \frac{\log_n T(r, f)}{\log r} \quad (n \in \mathbb{N}).$$

**Remark 1.1.** *If  $f$  is an entire function, then*

$$\sigma_n(f) = \limsup_{r \rightarrow \infty} \frac{\log_{n+1} M(r, f)}{\log r}.$$

**Definition 1.2.** *The growth index (or the finiteness degree) of the iterated order of a meromorphic function  $f$  is defined by*

$$i(f) = \begin{cases} 0 & \text{if } f \text{ is rational,} \\ \min\{n \in \mathbb{N} : \sigma_n(f) < \infty\} & \text{if } f \text{ is transcendental and } \sigma_n(f) < \infty \\ & \text{for some } n \in \mathbb{N}, \\ \infty & \text{if } f \text{ with } \sigma_n(f) = \infty \text{ for all } n \in \mathbb{N}. \end{cases}$$

---

2000 *Mathematics Subject Classification.* Primary 34M10; Secondary 30D35, 34M05.

*Key words and phrases.* meromorphic function, differential equation, complex oscillation, iterated order.

This work was supported by the NSF of Jiangxi (No. 2008GQS0075) and the YFED of Jiangxi (No. GJJ10050) of China.

**Definition 1.3.** The iterated convergence exponent of the sequence of  $a$ -points of a meromorphic function  $f$  is defined by

$$\lambda_n(f - a) = \limsup_{r \rightarrow \infty} \frac{\log_n N(r, \frac{1}{f-a})}{\log r} \quad (n \in \mathbb{N}).$$

**Definition 1.4.** The growth index (or the finiteness degree) of the iterated convergence exponent of the sequence of  $a$ -points of a meromorphic function  $f$  with iterated order is defined by

$$i_\lambda(f - a) = \begin{cases} 0 & \text{if } n(r, \frac{1}{f-a}) = O(\log r), \\ \min\{n \in \mathbb{N} : \lambda_n(f) < \infty\} & \text{if } \lambda_n(f - a) < \infty \text{ for some } n \in \mathbb{N}, \\ \infty & \text{if } \lambda_n(f - a) = \infty \text{ for all } n \in \mathbb{N}. \end{cases}$$

**Remark 1.2.** Similarly, we can use the notation  $\bar{\lambda}_n(f - a)$  to denote the iterated convergence exponent of the sequence of distinct  $a$ -points, and use the notation  $i_{\bar{\lambda}}(f - a)$  to denote the growth index of  $\bar{\lambda}_n(f - a)$ .

It is well-known that Nevanlinna theory has appeared to be a powerful tool in the field of complex differential equations (see [1-7, 11-18, 20], for example). The active research of the complex oscillation theory of linear differential equations in the complex plane  $\mathbb{C}$  was started to investigate the second order differential equation

$$(1) \quad f'' + A(z)f = 0$$

by Bank and Laine [1, 2]. They investigated this question in the case where  $A$  is an entire function, mainly from the point of determining the distribution of zeros of solutions. In this case all solutions of Eq.(1) are entire. When  $A$  is meromorphic, there are some immediate difficulties. For example, it is possible that no solution of Eq.(1) except the zero solution is single-valued on the plane. This obstacle was handled since Bank and Laine [2] gave necessary and sufficient conditions for all solutions of Eq.(1) to be meromorphic, and hence single-valued, in a simply-connected region. To consider poles as well as zeros, they obtained the following theorems.

**Theorem 1.1.** ([2], Theorem 5) Let  $A$  be a transcendental meromorphic function of order  $\sigma_1(A)$ , where  $0 < \sigma_1(A) \leq \infty$ , and assume that  $\bar{\lambda}_1(A) < \sigma_1(A)$ . Then, if  $f \not\equiv 0$  is a meromorphic solution of Eq.(1), we have

$$\sigma_1(A) \leq \max\{\bar{\lambda}_1(f), \bar{\lambda}_1(\frac{1}{f})\}.$$

**Theorem 1.2.** ([2], Theorem 6) Let  $A$  be a transcendental meromorphic function, and assume that Eq.(1) possesses two linearly independent meromorphic solutions  $f_1$  and  $f_2$  satisfying  $\bar{\lambda}_1(f_1) < \infty$ ,  $\bar{\lambda}_1(f_2) < \infty$ . Then, any solution  $f \not\equiv 0$  of Eq.(1) which is not a constant multiple of either  $f_1$  or  $f_2$  satisfies,

$$\max\{\bar{\lambda}_1(f), \bar{\lambda}_1(1/f)\} = \infty,$$

unless all solutions of Eq.(1) are of finite order (namely, finite iterated 1-order). In the special case where  $\bar{\lambda}_1(\frac{1}{A}) < \infty$  (e.g.  $A$  is of finite order), we can conclude that  $\bar{\lambda}_1(f) = \infty$  unless all solutions of Eq.(1) are finite order.

In [14], Kinnunen obtained some results on the Eq. (1) with entire solutions by using the idea of iterated  $n$ -order.

**Theorem 1.3.** ([14], Theorem 3.2) *Let  $A$  be an entire function with  $i(A) = n$ , assuming  $0 < n < \infty$ . Let  $f_1$  and  $f_2$  be two linearly independent solutions of Eq.(1), and denote  $E := f_1 f_2$ . Then  $i_\lambda(E) \leq n + 1$  and*

$$\lambda_{n+1}(E) = \sigma_{n+1}(E) = \max\{\lambda_{n+1}(f_1), \lambda_{n+1}(f_2)\} \leq \sigma_n(A).$$

*If  $i_\lambda(E) < n$ , then  $i_\lambda(f) = n + 1$  holds for all solutions of type of  $f = c_1 f_1 + c_2 f_2$ , where  $c_1 \neq 0$  and  $c_2 \neq 0$ .*

**Theorem 1.4.** ([14], Theorem 3.3) *Let  $A$  be an entire function with  $0 < i(A) = n < \infty$ , let  $f$  be any non-trivial solution of Eq.(1), and assume  $\bar{\lambda}_n(A) < \sigma_n(A) \neq 0$ . Then  $\lambda_{n+1}(f) \leq \sigma_n(A) \leq \lambda_n(f)$ .*

It is conjectured that a situation where  $\max\{\lambda(f_1), \lambda(f_2)\} < \infty$  for an equation  $f'' + A(z)f = 0$  implies that  $\max\{\lambda(g_1), \lambda(g_2)\} = \infty$  is true for the equation  $g'' + B(z)g = 0$  where  $B \neq A$  is sufficiently close to  $A$  in some sense (see [15], p.109). Kinnunen obtained the following result, of this type corresponding to Theorem 3.1 in [3].

**Theorem 1.5.** ([14], Theorem 3.6) *Let  $A$  be an entire function with  $i(A) = n$  and the iterated order  $\sigma_n(A) = \sigma$ , where  $1 < n < \infty$ . Let  $f_1$  and  $f_2$  be two linearly independent solutions of Eq.(1) such that  $\max\{\lambda_n(f_1), \lambda_n(f_2)\} < \sigma$ . Let  $\Pi \neq 0$  be any entire function for which either  $i(\Pi) < n$  or  $i(\Pi) = n$  and  $\sigma_n(\Pi) < \sigma$ . Then any two linearly independent solutions  $g_1$  and  $g_2$  of the differential equation  $g'' + (A(z) + \Pi(z))g = 0$  satisfy  $\max\{\lambda_n(g_1), \lambda_n(g_2)\} \geq \sigma$ .*

Thus it is interesting to consider the complex oscillation on the meromorphic solutions of the Eq. (1) for the case where  $A$  is meromorphic function in the terms of the idea of iterated order. In 2007, Liang and Liu[17] considered the complex oscillation on the Eq. (1) when  $A$  is a meromorphic function with finite many poles. By using the Wiman-Valiron theory (for an entire function [9, 11], for a meromorphic function [7, 20]), they obtained some results which extend Theorems 1.3 and 1.4. There arises naturally a question:

**Question 1.1.** *What can be said if  $A$  has infinitely many poles?*

Although the Wiman-Valiron theory is a powerful tool to investigate entire solutions, it is only useful for the meromorphic function  $A$  with  $\lambda_1(\frac{1}{A}) < \sigma_1(A)$  if considering the Eq. (1). In this paper we shall make use of a recent result due to Chiang and Hayman (see Lemma 2.3 in the next section) instead of the Wiman-Valiron theory, and thus answer the above question. In fact, we obtain the following results which improve and extend some oscillation results due to Bank & Laine [2], Liang & Liu [17] and others. Furthermore, considering the deficiencies of poles of the coefficient  $A$  and solutions  $f$  of Eq. (1), we obtain some special results. For  $a \in \bar{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ , the deficiency of  $a$  with respect to a meromorphic function  $g$  in  $\mathbb{C}$  is defined by

$$\delta(a, g) = 1 - \limsup_{r \rightarrow \infty} \frac{N(r, \frac{1}{g-a})}{T(r, g)},$$

provided that  $g$  has unbounded characteristic. The first result is the following

**Theorem 1.6.** *Let  $A$  be a meromorphic function with  $0 < i(A) = n < \infty$ , and assume that  $\bar{\lambda}_n(A) < \sigma_n(A) \neq 0$ . Then, if  $f$  is a nonzero meromorphic solution of the Eq. (1) we have*

$$(2) \quad \sigma_n(A) \leq \max\{\bar{\lambda}_n(f), \bar{\lambda}_n(\frac{1}{f})\}.$$

*In the special case where either  $\delta(\infty, f) > 0$  or the poles of  $f$  are of uniformly bounded multiplicities, we can conclude that*

$$(3) \quad \max\{\lambda_{n+1}(f), \lambda_{n+1}(\frac{1}{f})\} \leq \sigma_n(A) \leq \max\{\bar{\lambda}_n(f), \bar{\lambda}_n(\frac{1}{f})\}.$$

Theorem 1.6 improves and extends Theorem 3.3 in [14] and Theorem 5 in [2]. It is obvious from Theorem 1.6 that the following corollary is true, which improves and extends Corollary 3.4 in [14].

**Corollary 1.1.** *Let  $A$  be a meromorphic function with  $1 < i(A) = n < \infty$ . If  $i_{\bar{\lambda}}(A) < n$ , then any nonzero meromorphic solution  $f$  of the Eq. (1) satisfies  $\max\{i_{\bar{\lambda}}(f), i_{\bar{\lambda}}(\frac{1}{f})\} \geq n$ .*

The next result improves and extends Theorem 6 and Corollary 7 in [2].

**Theorem 1.7.** *Let  $A$  be a meromorphic function with  $0 < i(A) = n < \infty$ . Assume that the Eq. (1) possesses two linearly independent meromorphic solutions  $f_1$  and  $f_2$ . Denote  $E := f_1 f_2$ . If  $\bar{\lambda}_n(E) < \infty$ , then any nonzero solution  $f$  of (1) which is not a constant multiple of either  $f_1$  or  $f_2$  satisfies,  $\bar{\lambda}_n(f) = \infty$ , unless all solutions of (1) are of finite iterated  $n$ -order. In the special case where  $\delta(\infty, A) > 0$ , or  $i_{\lambda}(\frac{1}{A}) < n$ , or  $\lambda_n(\frac{1}{A}) < \sigma_n(A)$ , (e.g.  $A$  is an entire function), we can conclude that  $\bar{\lambda}_n(f) = \infty$ .*

We remark that Theorem 1.7 and the following theorem are the improvement and extension of Theorem 3.2 in [14].

**Theorem 1.8.** *Let  $A$  be a meromorphic function satisfying  $0 < i(A) = n < \infty$ . Assume that the Eq. (1) possesses two linearly independent meromorphic solutions  $f_1$  and  $f_2$ . Denote  $E := f_1 f_2$ . If  $\delta(\infty, A) > 0$ , or  $i_{\lambda}(\frac{1}{A}) < n$ , or  $\lambda_n(\frac{1}{A}) < \sigma_n(A)$ , and if either  $\delta(\infty, f) > 0$  or the poles of  $f$  are of uniformly bounded multiplicities, then we have  $i_{\lambda}(E) \leq n + 1$  and have*

$$\begin{aligned} \bar{\lambda}_{n+1}(E) &= \lambda_{n+1}(E) = \sigma_{n+1}(E) = \max\{\bar{\lambda}_{n+1}(f_1), \bar{\lambda}_{n+1}(f_2)\} \\ &\leq \sigma_{n+1}(f_1) = \sigma_{n+1}(f_2) = \sigma_n(A). \end{aligned}$$

From the proof of Theorem 1.8 one can get the following result.

**Corollary 1.2.** *Let  $A$  be a meromorphic function satisfying  $0 < i(A) = n < \infty$ . Assume that the Eq. (1) possesses two linearly independent meromorphic solutions  $f_1$  and  $f_2$ . Denote  $E := f_1 f_2$ . If  $\delta(\infty, A) > 0$ , then we have  $i_{\lambda}(E) \leq n + 1$  and have*

$$\begin{aligned} \bar{\lambda}_{n+1}(E) &= \lambda_{n+1}(E) = \sigma_{n+1}(E) = \max\{\bar{\lambda}_{n+1}(f_1), \bar{\lambda}_{n+1}(f_2)\} \\ &\leq \sigma_{n+1}(f_1) = \sigma_{n+1}(f_2). \end{aligned}$$

The following corollary is immediately obtained from Theorem 1.8 which is an improvement of Theorem 1.3.

**Corollary 1.3.** *Let  $A$  be an entire function with  $0 < i(A) = n < \infty$ . Let  $f_1$  and  $f_2$  be two linear independent solutions of Eq.(1), and denote  $E := f_1 f_2$ . Then  $i_\lambda(E) \leq n + 1$  and*

$$\begin{aligned} \bar{\lambda}_{n+1}(E) &= \lambda_{n+1}(E) = \sigma_{n+1}(E) = \max\{\bar{\lambda}_{n+1}(f_1), \bar{\lambda}_{n+1}(f_2)\} \\ &\leq \sigma_{n+1}(f_1) = \sigma_{n+1}(f_2) = \sigma_n(A). \end{aligned}$$

Finally, we show the following result which extends and improves Theorem 1.5.

**Theorem 1.9.** *Let  $A$  be a meromorphic function with  $1 < i(A) = n < \infty$ . Assume that  $f_1$  and  $f_2$  are two linearly independent meromorphic solutions of the Eq. (1) such that*

$$(4) \quad \max\{\lambda_n(f_1), \lambda_n(f_2)\} < \sigma_n(A).$$

*Let  $\Pi \neq 0$  be any meromorphic function for which either  $i(\Pi) < n$  or  $\sigma_n(\Pi) < \sigma_n(A)$ . Let  $g_1$  and  $g_2$  be two linearly independent solutions of the differential equation  $g'' + (A(z) + \Pi(z))g = 0$ . Denote  $E := f_1 f_2$  and  $F := g_1 g_2$ . If either*

$$\max\{i_\lambda(1/E), i_\lambda(1/F)\} < n \quad \text{or} \quad \max\{\lambda_n(1/E), \lambda_n(1/F)\} < \sigma_n(A),$$

*then  $\max\{\lambda_n(g_1), \lambda_n(g_2)\} \geq \sigma_n(A)$ .*

The remainder of this paper is organized as follows. Section 2 is for some lemmas and the other sections are for the proofs of our main results. The idea and formulations of our main results come from [1, 2, 14]. The proof of Theorem 1.6 is from the proof of Theorem 5 in [2], the proof of Theorem 1.7 is essentially from the proof of Theorem 6 in [2], and the proof of Theorem 1.9 is a parallel to a corresponding reasoning in the proof of Theorem 3.6 in [14].

## 2. SOME LEMMAS

To prove our results, we need the following lemmas.

**Lemma 2.1.** ([10], Theorem 4) *Let  $f$  be a transcendental meromorphic function not of the form  $e^{\alpha z + \beta}$ . Then*

$$T\left(\frac{f}{f'}\right) \leq 3\bar{N}(r, f) + 7\bar{N}\left(\frac{1}{f}\right) + 4\bar{N}\left(\frac{1}{f''}\right) + S\left(r, \frac{f}{f'}\right),$$

*where  $S(r, f) := o(T(r, f))$  as  $r \rightarrow \infty$  outside of a possible exceptional set of finite linear measure.*

**Lemma 2.2.** ([14], Remark 1.3) *If  $f$  is a transcendental meromorphic function, then  $\sigma_n(f) = \sigma_n(f')$ .*

**Lemma 2.3.** ([13], Theorem 6.2) *Let  $f$  be a meromorphic solution of*

$$(5) \quad f^{(k)} + A_{k-1}(z)f^{(k-1)} + \dots + A_1(z)f' + A_0(z)f = 0$$

where  $A_0, \dots, A_{k-1}$  are meromorphic functions in the plane  $\mathbb{C}$ . Assume that not all coefficients  $A_j$  are constants. Given a real constant  $\gamma > 1$ , and denoting  $T(r) := \sum_{j=0}^{k-1} T(r, A_j)$ , we have

$$\log m(r, f) < T(r) \{(\log r) \log T(r)\}^\gamma, \quad \text{if } p = 0;$$

and

$$\log m(r, f) < r^{2p+\gamma-1} T(r) \{\log T(r)\}^\gamma, \quad \text{if } p > 0;$$

outside of an exceptional set  $E_p$  with  $\int_{E_p} t^{p-1} dt < +\infty$ .

We note that in the above lemma,  $p = 1$  corresponds to Euclidean measure and  $p = 0$  to logarithmic measure. Using logarithmic measure not Euclidean measure, we correct here the proof of Theorem 3.2 in [6].

**Lemma 2.4.** ([6], Theorem 3.2) *Let  $A_0, A_1, \dots, A_{k-1}$  be meromorphic functions such that  $0 < \max\{i(A_j) : j = 0, 1, \dots, k-1\} = n < \infty$ . If  $f$  is a meromorphic solution of (5) whose poles are of uniformly bounded multiplicities or  $\delta(\infty, f) > 0$ , then  $\sigma_{n+1}(f) \leq \max\{\sigma_n(A_j) : j = 0, 1, \dots, k-1\}$ .*

*Proof.* It obvious that if  $\sigma_n(f) < \infty$ , then  $\sigma_{n+1}(f) = 0 \leq \sigma := \max\{\sigma_n(A_j) : j = 0, 1, \dots, k-1\}$ . Now we assume  $\sigma_n(f) = \infty$ . By (5) we get that the poles of  $f(z)$  can only occur at the poles of  $A_0, A_1, \dots, A_{k-1}$ . Note that the multiplicities of poles of  $f$  are uniformly bounded, and thus we have

$$N(r, f) \leq M_1 \overline{N}(r, f) \leq M_1 \sum_{j=0}^{k-1} \overline{N}(r, A_j) \leq M \max\{N(r, A_j) : j = 0, 1, \dots, k-1\},$$

where  $M_1$  and  $M$  are some suitable positive constants. This gives

$$(6) \quad T(r, f) = m(r, f) + O(\max\{N(r, A_j) : j = 0, 1, \dots, k-1\}).$$

If  $\delta(\infty, f) := \delta_1 > 0$ , then for sufficiently large  $r$ ,

$$(7) \quad m(r, f) \geq \frac{\delta_1}{2} T(r, f).$$

Applying now (6) or (7) with Lemma 2.3, we obtain

$$\log T(r, f) \leq \log m(r, f) + O(\log T(r)) \leq O\{T(r) \{(\log r) \log T(r)\}^\gamma\}$$

or

$$\log T(r, f) \leq \log\left(\frac{2}{\delta_1} m(r, f)\right) \leq O\{T(r) \{(\log r) \log T(r)\}^\gamma\}$$

outside of an exceptional set  $E_0$  with finite logarithmic measure. Using a standard method to deal with the finite logarithmic measure set, one immediately gets from above inequalities that  $\sigma_{n+1}(f) \leq \max\{\sigma_n(A_j) : j = 0, 1, \dots, k-1\}$ .  $\square$

**Lemma 2.5.** ([6], Lemma 3.6) *Let  $\Phi(r)$  be a continuous and positive increasing function, defined for  $r$  on  $(0, +\infty)$ , with  $\sigma_n(\Phi) = \limsup_{r \rightarrow \infty} \frac{\log_n \Phi(r)}{\log r}$ . Then for any subset  $E$  of  $[0, +\infty)$  that has finite linear measure, there exists a sequence  $\{r_m\}, r_m \notin E$  such that*

$$\sigma_n(\Phi) = \lim_{r_m \rightarrow \infty} \frac{\log_n \Phi(r_m)}{\log r_m}.$$

Replacing the notation  $n(r, f)$  by  $\bar{n}(r, f)$  and following the reasoning of the proof of Lemma 1.7 in [14], one can easily obtain the following lemma.

**Lemma 2.6.** *Let  $g_1$  and  $g_2$  be two entire functions. Then*

$$i_{\bar{\lambda}}(g_1 g_2) = \max\{i_{\bar{\lambda}}(g_1), i_{\bar{\lambda}}(g_2)\}.$$

If  $i_{\bar{\lambda}}(g_1 g_2) := n > 0$ , then

$$\bar{\lambda}_n(g_1 g_2) = \max\{\bar{\lambda}_n(g_1), \bar{\lambda}_n(g_2)\}.$$

We recall here the essential part of the factorization theorem for meromorphic functions of finite iterated order.

**Lemma 2.7.** ([12], Satz 12.4) *A meromorphic function  $f$  for which  $i(f) = n$  can be represented by the form*

$$f(z) = \frac{U(z)e^{g(z)}}{V(z)},$$

where  $U, V$  and  $g$  are entire functions such that

$$\lambda_n(f) = \lambda_n(U) = \sigma_n(U), \quad \lambda_n\left(\frac{1}{f}\right) = \lambda_n(V) = \sigma_n(V)$$

and

$$\sigma_n(f) = \max\{\sigma_n(U), \sigma_n(V), \sigma_n(e^g)\}.$$

The following result plays a key role in the present paper, which is an improvement and extension of Theorem 3.1 in [14] and Theorem 1 in [17].

**Lemma 2.8.** *Let  $A$  be a meromorphic function with  $i(A) = n$  ( $0 < n < \infty$ ), and let  $f$  be a nonzero meromorphic solution of the Eq. (1). Then*

(i) *if either  $\delta(\infty, f) > 0$  or the poles of  $f$  are of uniformly bounded multiplicities, then  $i(f) \leq n + 1$  and  $\sigma_{n+1}(f) \leq \sigma_n(A)$ .*

(ii) *if  $\delta(\infty, A) > 0$ , or  $i_{\lambda}\left(\frac{1}{A}\right) < n$ , or  $\lambda_n\left(\frac{1}{A}\right) < \sigma_n(A)$ , then  $i(f) \geq n + 1$  and  $\sigma_{n+1}(f) \geq \sigma_n(A)$ .*

*Proof.* Assume that  $f$  is a nonzero meromorphic solution of the Eq. (1). It is obvious that (i) is just a special case of Lemma 2.4.

We now assume that  $A$  satisfies  $\delta(\infty, A) > 0$ , or  $i_{\lambda}\left(\frac{1}{A}\right) < n$ , or  $\lambda_n\left(\frac{1}{A}\right) < \sigma_n(A)$ . Then we shall prove  $i(f) \geq n + 1$  and  $\sigma_{n+1}(f) \geq \sigma_n(A)$ . By (1), we get

$$(8) \quad -A(z) = \frac{f''}{f}.$$

By the lemma of the logarithmic derivative and (8), we get that

$$m(r, A) \leq m\left(r, \frac{f''}{f}\right) = O\{\log(rT(r, f))\}$$

holds for all sufficiently large  $r \notin E$ , where  $E \subset (0, \infty)$  has finite linear measure. Hence

$$(9) \quad T(r, A) = m(r, A) + N(r, A) \leq N(r, A) + O\{\log(rT(r, f))\}$$

holds for all sufficiently large  $|z| = r \notin E$ .

If  $\sigma_n(A) = 0$ , and hence  $i_\lambda(\frac{1}{A}) < n$ , then by Lemma 2.5 there exists a sequence  $\{r_m\}$  such that for all  $r_m \notin E_6$ ,

$$(10) \quad T(r_m, A) \geq \exp_{n-2}\{r_m^M\}$$

holds for any sufficiently large constant  $M > 0$ . If  $\sigma_n(A_0) > 0$ , then again by Lemma 2.5 there exists a sequence  $\{r_m\}$  such that for all  $r_m \notin E_6$ ,

$$(11) \quad T(r_m, A) \geq \exp_{n-1}\{r_m^{\sigma-\varepsilon}\}$$

holds for any given  $\varepsilon$  ( $0 < \varepsilon < \sigma$ ).

Now we consider three cases below.

**Case 1.** Assume that  $\delta(\infty, A) := \delta_2 > 0$ . Then for sufficiently large  $r$ ,

$$(12) \quad \frac{\delta_2}{2}T(r, A) \leq m(r, A) = O\{\log(rT(r, f))\}.$$

If  $\sigma_n(A) = 0$ , then from (10) and (12) we get that  $i(f) \geq n + 1$  and  $\sigma_{n+1}(f) \geq \sigma_n(A) = 0$ . If  $\sigma_n(A) > 0$ , then from (11) and (12) we get that  $i(f) \geq n + 1$  and  $\sigma_{n+1}(f) \geq \sigma_n(A)$ .

**Case 2.** Assume that  $i_\lambda(\frac{1}{A}) < n$ . Then

$$(13) \quad N(r, A) \leq \exp_{n-2}(r^{\alpha_1})$$

holds for a positive constant  $\alpha_1 < M$ . By (13), (9) and either (10) or (11), we get that  $i(f) \geq n + 1$  and  $\sigma_{n+1}(f) \geq \sigma_n(A)$ .

**Case 3.** Assume that  $\lambda_n(\frac{1}{A}) < \sigma_n(A) = \sigma$ . Then there holds

$$(14) \quad N(r, A) \leq \exp_{n-1}(r^{\lambda_n(\frac{1}{A})+\varepsilon}),$$

where ( $0 < 2\varepsilon < \sigma - \lambda_n(\frac{1}{A})$ ). Thus by (9), (14) and (10), we get that  $i(f) \geq n + 1$  and  $\sigma_{n+1}(f) \geq \sigma_n(A)$ . □

Following Hayman [10], we shall use the abbreviation "n. e." (nearly everywhere) to mean "everywhere in  $(0, \infty)$  except in a set of finite measure" in the proofs of our main theorems, see the following sections.

### 3. PROOF OF THEOREM 1.6

Since  $f$  is a solution of (1) where  $\sigma_n(A) > 0$ , it is obvious that  $f$  can not be rational, nor be of the form  $e^{az+b}$  for constants  $a$  and  $b$ . Hence, by Lemma 2.1 we have

$$(15) \quad T\left(r, \frac{f}{f'}\right) = O\left(\overline{N}(r, f) + \overline{N}\left(r, \frac{1}{f}\right) + \overline{N}\left(r, \frac{1}{f''}\right)\right) \quad n. e. \quad \text{as } r \rightarrow \infty.$$

In addition, by (1) we have

$$(16) \quad \overline{N}\left(r, \frac{1}{f''}\right) \leq \overline{N}\left(r, \frac{1}{f}\right) + \overline{N}\left(r, \frac{1}{A}\right).$$

By assumption,  $\overline{\lambda}_n(A) < \sigma_n(A)$ . Hence, if we assume that (2) fails to hold, then we deduce by (15) and (16) that  $\sigma_n\left(\frac{f}{f'}\right) < \sigma_n(A)$ . By the first main theorem, we then see that if  $\varphi = \frac{f'}{f}$ , then  $\sigma_n(\varphi) < \sigma_n(A)$ . However, from (1) it easily follows that



$-A = \varphi' + \varphi^2$ , and so we obtain  $\sigma_n(A) \leq \sigma_n(\varphi) < \sigma_n(A)$ , a contradiction. Hence, (2) is true.

In the special case where either  $\delta(\infty, f) > 0$  or the poles of  $f$  are of uniformly bounded multiplicities, by Lemma 2.8 we have

$$\max\{\lambda_{n+1}(\frac{1}{f}), \lambda_{n+1}(f)\} \leq \sigma_{n+1}(f) \leq \sigma_n(A).$$

Hence, we obtain (3).

#### 4. PROOF OF COROLLARY 1.1

Let  $f$  be a nonzero meromorphic solution of Eq.(1). Assume that

$$\max\{\bar{\lambda}_{n-1}(f), \bar{\lambda}_{n-1}(\frac{1}{f})\} < \infty,$$

where  $n = i(A) > 1$ . Then we obtain

$$\bar{N}(r, \frac{1}{f}) = O(\exp_{n-2}(\frac{1}{1-r})^{\alpha_1}) \quad \text{and} \quad \bar{N}(r, f) = O(\exp_{n-2}(\frac{1}{1-r})^{\alpha_2})$$

for some finite constants  $\alpha_1$  and  $\alpha_2$ . Similarly, we get from the assumption  $i_{\bar{\lambda}}(A) < n$  that

$$\bar{N}(r, \frac{1}{A}) = O(\exp_{n-2}(\frac{1}{1-r})^{\alpha_3})$$

for some finite constant  $\alpha_3$ . The relations (15) and (16) now yield

$$T(r, \frac{f}{f'}) = O(\exp_{n-2}(\frac{1}{1-r})^{\alpha})$$

for some finite constant  $\alpha \geq \max\{\alpha_1, \alpha_2, \alpha_3\}$ . Hence we have

$$\sigma_{n-1}(\frac{f'}{f}) = \sigma_{n-1}(\frac{f}{f'}) < \infty.$$

By the relation  $-A = \varphi' + \varphi^2$  where  $\varphi = \frac{f'}{f}$ , we now obtain  $\sigma_{n-1}(A) < \infty$ , a contradiction. Therefore, we obtain the conclusion  $\max\{i_{\bar{\lambda}}(f), i_{\bar{\lambda}}(\frac{1}{f})\} \geq n$ .

#### 5. PROOF OF THEOREM 1.7

Assume that the Eq. (1) possesses two linearly independent meromorphic solutions  $f_1$  and  $f_2$  such that  $\bar{\lambda}_n(E) < \infty$ , where  $E_1 := E = f_1 f_2$ . Let  $f = a f_1 + b f_2$  where  $a$  and  $b$  are nonzero constants, and set  $E_2 := f f_1$ . It is easy to see that any pole of  $f$  is a pole of  $A$ . Since  $\bar{\lambda}_n(\frac{1}{A}) \leq \sigma_n(A) < \infty$ , we thus have  $\bar{\lambda}_n(\frac{1}{f}) \leq \bar{\lambda}_n(\frac{1}{A}) < \infty$ . Assume that  $\bar{\lambda}_n(f) = \infty$  fails to hold, so that  $\bar{\lambda}_n(f) < \infty$  and  $\bar{\lambda}_n(\frac{1}{f}) < \infty$ . From these relations we easily see that  $\bar{\lambda}_n(E_1) < \infty$  and  $\bar{\lambda}_n(E_2) < \infty$ . By Lemma D(e) in [2], there is a constant  $c > 0$  such that n. e. as  $r \rightarrow \infty$ ,

$$(17) \quad T(r, E_j) = O\left(\bar{N}\left(r, \frac{1}{E_j}\right) + T(r, A) + \log r\right) = O(\exp_{n-1}(r^c) + T(r, A))$$

for  $j = 1, 2$ . Since  $E_2 = a f_1^2 + b E_1$ , we thus obtain that n. e. as  $r \rightarrow \infty$ ,

$$(18) \quad T(r, f_1) = O(\exp_{n-1}(r^c) + T(r, A)).$$

Since  $A = -\frac{f''}{f}$ , we deduce by the lemma of logarithmic derivative that n. e. as  $r \rightarrow \infty$ ,

$$(19) \quad m(r, A) = O(\log T(r, f_1) + \log r).$$

From (1), we see that any pole of  $A$  is at most double and is either a zero or pole of  $f$ , we thus have

$$N(r, A) \leq 2 \left( \overline{N} \left( r, \frac{1}{f} \right) + \overline{N}(r, f) \right).$$

Hence by assumption,  $N(r, A) = O(\exp_{n-1}(r^d))$  as  $r \rightarrow \infty$  for some  $d > 0$ . Together with (18) and (19), we obtain  $N(r, A) = O(\exp_{n-1}(r^d))$  n. e. as  $r \rightarrow \infty$ , from which it follows by standard reasoning that  $f_1$  is of finite iterated  $n$ -order. By the identity of Abel, we have

$$(20) \quad \left( \frac{f_2}{f_1} \right)' = \frac{\beta}{f_1^2},$$

where  $\beta$  is equal to the Wronskian of  $f_1$  and  $f_2$ . Hence, by Lemma 2.2 and (20), we obtain

$$\sigma_n(f_2) = \sigma_n \left( f_1 \frac{f_2}{f_1} \right) \leq \max \left\{ \sigma_n \left( \frac{f_2}{f_1} \right), \sigma_n(f_1) \right\} = \sigma_n(f_1).$$

Reversing the roles of  $f_1$  and  $f_2$ , we can conclude that  $\sigma_n(f_1) = \sigma_n(f_2)$ . Hence, all solutions of (1) are of finite iterated  $n$ -order if  $\overline{\lambda}_n(f) < \infty$ .

In special case where  $\delta(\infty, A) > 0$ , or  $i_\lambda(\frac{1}{A}) < n$ , or  $\lambda_n(\frac{1}{A}) < \sigma_n(A)$ , by Lemma 2.8 that all meromorphic solutions  $f \neq 0$  of (1) satisfy  $i(f) \geq n + 1$  and  $\sigma_{n+1}(f) \geq \sigma_n(A)$ . Therefore, we can conclude that  $\overline{\lambda}_n(f) = \infty$  holds for any solution  $f \neq 0$  of (1) which is not a constant multiple of either  $f_1$  or  $f_2$ .

## 6. PROOF OF THEOREM 1.8

It is obvious that (see page 664 in [2])  $\sigma_{n+1}(f_1) = \sigma_{n+1}(f_2)$ . Assume that  $\delta(\infty, A) > 0$ , or  $i_\lambda(\frac{1}{A}) < n$ , or  $\lambda_n(\frac{1}{A}) < \sigma_n(A)$ , and that either  $\delta(\infty, f) > 0$  or the poles of  $f$  are of uniformly bounded multiplicities. Then by Lemma 2.8 we obtain

$$\sigma_{n+1}(E) \leq \max \{ \sigma_{n+1}(f_1), \sigma_{n+1}(f_2) \} \leq \sigma_{n+1}(f_1) = \sigma_{n+1}(f_2) = \sigma_n(A) < \infty.$$

By Lemma D(e) in [2], there is a constant  $c > 0$  such that n. e. as  $r \rightarrow \infty$ ,

$$(21) \quad T(r, E) = O \left( \overline{N} \left( r, \frac{1}{E} \right) + T(r, A) + \log r \right).$$

By the lemma of logarithmic derivative and Lemma 2.8, we have

$$m(r, A) = m \left( r, \frac{f_1''}{f_1} \right) = O(\log r T(r, f_1)) = O(\exp_{n-1}(r^{a_1}))$$

for some  $a_1 < \infty$  outside of a possible exceptional set  $G \subset [0, \infty)$  with finite linear measure. If  $\delta(\infty, A) := \delta_3 > 0$ . Then for sufficiently large  $r$ ,

$$(22) \quad \frac{\delta_3}{2} T(r, A) \leq m(r, A) = O(\exp_{n-1}(r^{a_1})), \quad r \notin G.$$

If either  $i_\lambda(\frac{1}{A}) < n$  or  $\lambda_n(\frac{1}{A}) < \sigma_n(A) < \infty$ , we have

$$N(r, A) = O(\exp_{n-1}(r^{a_2}))$$

for some  $a_2 < \infty$ . Thus

$$(23) \quad T(r, A) = m(r, A) + N(r, A) = O(\exp_{n-1}(r^a)), \quad r \notin G,$$

where  $a = \max\{a_1, a_2\}$ . Hence, together with (21) and either (22) or (23) we obtain

$$(24) \quad T(r, E) = O\left(\overline{N}\left(r, \frac{1}{E}\right) + \exp_{n-1}(r^a)\right), \quad r \notin G.$$

Suppose that  $\overline{\lambda}_{n+1}(E) < \sigma_{n+1}(E)$ , then we have  $\overline{N}\left(r, \frac{1}{E}\right) = O(\exp_n(r^b))$  for some  $b < \sigma_{n+1}(E)$ . Together with (24),  $T(r, E) = O(\exp_n(r^b))$ ,  $r \notin G$ , and then by standard reasoning, we obtain  $\sigma_{n+1}(E) \leq b < \sigma_{n+1}(E)$ . This is a contradiction. Hence, we have  $\overline{\lambda}_{n+1}(E) \geq \sigma_{n+1}(E)$ . Noting that  $\overline{\lambda}_{n+1}(E) \leq \lambda_{n+1}(E) \leq \sigma_{n+1}(E)$ , we obtain  $\overline{\lambda}_{n+1}(E) = \lambda_{n+1}(E) = \sigma_{n+1}(E)$ .

By Lemma D(a) in [2],  $f_1$  and  $f_2$  have no common zeros. Let  $f_j = \frac{g_j}{d_j}$ , where  $g_j$  and  $d_j$  have no common zeros,  $j = 1, 2$ . This implies that  $g_1$  and  $g_2$  have no common zeros, that  $\overline{\lambda}_n(f_j) = \overline{\lambda}_n(g_j)$  for  $j = 1, 2$ , and that  $\overline{\lambda}_n(E) = \overline{\lambda}_n(g_1 g_2)$ . Hence, by Lemma 2.6, we have  $\overline{\lambda}_{n+1}(E) = \max\{\overline{\lambda}_{n+1}(f_1), \overline{\lambda}_{n+1}(f_2)\}$ .

Therefore, we obtain the conclusion

$$\begin{aligned} \overline{\lambda}_{n+1}(E) &= \lambda_{n+1}(E) = \sigma_{n+1}(E) = \max\{\overline{\lambda}_{n+1}(f_1), \overline{\lambda}_{n+1}(f_2)\} \\ &\leq \sigma_{n+1}(f_1) = \sigma_{n+1}(f_2) = \sigma_n(A) < \infty. \end{aligned}$$

## 7. PROOF OF THEOREM 1.9

We denote  $E := f_1 f_2$  and  $F := g_1 g_2$ . By a similar argument, by Lemma 1.7 in [14] we obtain  $\lambda_n(F) = \max\{\lambda_n(g_1), \lambda_n(g_2)\}$ . We assume that  $\lambda_n(F) < \sigma_n(A)$ .

By the assumption (4), we have

$$\overline{N}\left(r, \frac{1}{E}\right) = O(\exp_{n-1}(r^\beta))$$

for some  $\beta < \sigma_n(A)$  and the iterated order of the function  $A$  implies that

$$T(r, A) = O\left(\exp_{n-1}(r^{\sigma_n(A)+\varepsilon})\right).$$

Again by Lemma D(e) in [2], we have also the Eq. (21), and thus we obtain

$$T(r, E) = O(\exp_{n-1}(r^\beta)).$$

So, we obtain  $\sigma_n(E) \leq \sigma_n(A)$ . On the other hand, by Lemma B(iv) in [2] we have

$$(25) \quad 4A = \left(\frac{E'}{E}\right)^2 - 2\frac{E''}{E} - \frac{1}{E^2},$$

which implies that  $\sigma_n(A) \leq \sigma_n(E)$ . Noting that either  $i(\Pi) < n$  or  $\sigma_n(\Pi) < \sigma_n(A)$ . The same reasoning is valid for the function  $F$ , and hence, we have  $\sigma_n(E) = \sigma_n(F) = \sigma_n(A)$ .

By the assumption (4) and Lemma 2.7, we can write

$$(26) \quad E = \frac{Qe^P}{U}, F = \frac{Re^S}{V},$$

where  $\sigma_n(Q) = \lambda_n(E) < \sigma_n(A)$ ,  $\sigma_n(R) = \lambda_n(F) < \sigma_n(A)$ . Together with the assumption that  $\max\{i_\lambda(1/E), i_\lambda(1/F)\} < n$  or  $\max\{\lambda_n(1/E), \lambda_n(1/F)\} < \sigma_n(A)$ , we have

$$\sigma_n(e^P) = \sigma_n(e^S) = \sigma_n(A).$$

Substituting (26) into (25) and following the similar reasoning step by step as in the proof of Theorem 3.1 in [3], one may derive the fact that

$$ce^{2(P-S)} = -\frac{U^2R^2}{V^2Q^2},$$

where  $c \neq 0$ . Hence,

$$(27) \quad \frac{E^2}{F^2} = \frac{V^2Q^2}{U^2R^2}e^{2(P-S)} = -\frac{1}{c}.$$

From the equations (25), (27) and the similar equation for  $F$ ,

$$(28) \quad 4(A + \Pi) = \left(\frac{F'}{F}\right)^2 - 2\frac{F''}{F} - \frac{1}{F^2},$$

we obtain

$$4\left(A + \Pi + \frac{1}{c}A\right) = \left(\frac{F'}{F}\right)^2 - 2\frac{F''}{F} + \frac{1}{c}\left(\frac{E'}{E}\right)^2 - \frac{2E''}{cE}.$$

Since  $\infty > i(A) = n > 1$ , then by the lemma of logarithmic derivative, we obtain

$$\begin{aligned} T\left(r, A\left(1 + \frac{1}{c}\right) + \Pi\right) &= m\left(r, A\left(1 + \frac{1}{c}\right) + \Pi\right) + N\left(r, A\left(1 + \frac{1}{c}\right) + \Pi\right) \\ &= O\left(\exp_{n-2}(r^{\sigma_n(A)+\varepsilon})\right) \end{aligned}$$

n.e. as  $r \rightarrow \infty$ . This implies that  $i\left(A\left(1 + \frac{1}{c}\right) + \Pi\right) < n$  or  $\sigma_n\left(A\left(1 + \frac{1}{c}\right) + \Pi\right) < \sigma_n(A)$ . Hence,  $c$  must be  $-1$ . Thus  $E^2 = F^2$ , and so we have  $\frac{E'}{E} = \frac{F'}{F}$  and  $\frac{E''}{E} = \frac{F''}{F}$ . we can see from the equations (25) and (28) that  $\Pi = 0$ . This is a contradiction.

**Acknowledgement.** The authors would like to thank the referees for making valuable suggestions and comments to improve the present paper.

#### REFERENCES

- [1] S. Bank and I. Laine, On the oscillation theory of  $f'' + Af = 0$  where  $A$  is entire, Tran. Amer. Math. Soc. 273(1982), no. 1, 351-363.
- [2] S. Bank and I. Laine, On the zeros of meromorphic solutions of second-order linear differential equations, Comment. Math. Helv. 58(1983), 656-677.
- [3] S. Bank, I. Laine and J. Langley, Oscillation results for solutions of linear differential equations in the complex domain, Results Math. 16(1989), 3-15.
- [4] L. G. Bernal, On growth  $k$ -order of solutions of a complex homogeneous linear differential equation, Proc. Amer. Math. Soc. 101(1987), no. 2, 317-322.

- [5] B. Belaïdi, Growth and oscillation of solutions to linear differential equations with entire coefficients having the same order, *Electronic J. Diff. Equations* 2009(2009), no. 70, pp 1-10.
- [6] T. B. Cao, J. F. Xu and Z. X. Chen, On the meromorphic solutions of linear differential equations on the complex plane, *J. Math. Anal. Appl.* 364(2010), 130-142.
- [7] Z. X. Chen, On the rate of growth of meromorphic solutions of higher order linear differential equations, *Acta Math. Sinica (Chin. Ser.)* 42(1999), no. 3, 552-558 (in Chinese).
- [8] W. K. Hayman, *Meromorphic Functions*, Clarendon Press, Oxford, 1964.
- [9] W. K. Hayman, The local growth of power series: A survey of Wiman-Valiron method, *Canad. Math. Bull.* 17(1974), no. 3, 317-358.
- [10] W. K. Hayman, Picard values of meromorphic functions and their derivatives, *Ann. of Math.* 70(1959), no. 2, 9-42.
- [11] Y. Z. He and X. Z. Xiao, *Algebroid Functions and Ordinary Differential Equations*, Beijing, Science Press, 1988(in Chinese).
- [12] G. Jank and L. Volkmann, *Meromorphe Funktionen und Differentialgleichungen*, Birkäuser, 1985.
- [13] Y. M. Chiang and W. K. Hayman, Estimates on the growth of meromorphic solutions of linear differential equations, *Comment. Math. Helv.* 79(2004), 451-470.
- [14] L. Kinnunen, Linear differential equations with solutions of finite iterated order, *Southeast Asian Bull. Math.* 22(1998), 385-405.
- [15] I. Laine, *Nevanlinna Theory and Complex Differential Equations*, Walter de Gruyter, Berlin, 1993.
- [16] I. Laine and J. Rieppo, Differential polynomials generated by linear differential equations, *Complex Variables* 49(2004), 897-911.
- [17] J. J. Liang and M. S. Liu, The iterated order of meromorphic solutions of second order homogeneous linear differential equations, *Natural Science Journal of Hainan University*, 25(2007), no. 2, 109-116 (in Chinese).
- [18] K. Liu and L. Z. Yang, On the complex oscillation of higher order linear differential equations, *Bull. Korean Math. Soc.* 46(2009), no. 4, 607-615.
- [19] D. Sato, On the rate of growth of entire functions of fast growth, *Bull. Amer. Math. Soc.* 69(1963), 411-414.
- [20] J. Wang and W. R. Lü, The fixed points and hyper-order of solutions of second order linear differential equations with meromorphic coefficients, *Acta Math. Sinica (Chin. Ser.)* 27(2004), no. 1, 72-80(in Chinese).
- [21] L. Yang, *Value Distribution Theory*, Springer-Verlag, Berlin, 1993, and Science Press, Beijing, 1982.

(Received April 14, 2010)

(Ting-Bin Cao) DEPARTMENT OF MATHEMATICS, NANCHANG UNIVERSITY, NANCHANG, JIANGXI 330031, CHINA

*E-mail address:* [tbcao@ncu.edu.cn](mailto:tbcao@ncu.edu.cn), [ctb97@163.com](mailto:ctb97@163.com)

(Lei-Min Li) DEPARTMENT OF MATHEMATICS, NANCHANG UNIVERSITY, NANCHANG, JIANGXI 330031, CHINA

*E-mail address:* [leiminli@hotmail.com](mailto:leiminli@hotmail.com)