



# Iterative technique for a third-order differential equation with three-point nonlinear boundary value conditions

Xiuli Lin and Zengqin Zhao 

School of Mathematical Sciences, Qufu Normal University, Qufu 273165, People's Republic of China

Received 3 October 2015, appeared 18 March 2016

Communicated by Nickolai Kosmatov

**Abstract.** In this paper, we study the existence of extremal solutions for a nonlinear third-order differential equation with three-point nonlinear boundary value conditions. By means of the method of upper and lower solutions and different monotone iterative techniques, the sufficient conditions which guarantee the existence of extremal solutions are given. An example illustrates the main results.

**Keywords:** third-order differential equation, nonlinear boundary value conditions, the method of upper and lower solutions, iterative technique.

**2010 Mathematics Subject Classification:** 34B15.

## 1 Introduction

Nonlinear boundary value conditions in differential equations can describe many phenomena in applied mathematics, engineering, physical or biological processes. In this paper, we consider the following third-order differential equation with three-point nonlinear boundary value conditions

$$\begin{cases} -u'''(t) = f(t, u(t)), & t \in [0, 1], \\ u(0) = u''(0) = 0, \\ p(u(1), u(\xi)) = 0, \end{cases} \quad (1.1)$$

where  $\xi \in (0, 1)$ ,  $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$  and  $p : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  are continuous.

In recent years, third-order differential equations with nonlinear boundary value conditions have been discussed in many papers (see [1–15] and the references therein). For example, [4] considered a class of two-point nonlinear boundary value conditions by using a priori estimate, Nagumo condition, upper and lower solutions and Leray–Schauder degree. Papers [7–9, 13] considered some nonlinear nonlocal boundary conditions. However, according to our knowledge, for third-order differential equation, the three-point nonlinear boundary

---

 Emails: lin-xiuli78@163.com (Xiuli Lin), zqzhaoy@163.com (Zengqin Zhao)

value conditions in problem (1.1) are never discussed in literature. Hence the aim of this paper is to discuss this issue.

The main contributions are as follows: (a) we present problems with linear boundary value conditions, and on this basis we obtain the existence of the extremal solutions for problem (1.1) by applying the method of upper and lower solutions and monotone iterative technique; (b) the iterative technique is not unique and an example illustrates the result.

## 2 Notations and preliminaries

In this section, we present some definitions and lemmas that will be used throughout the paper.

**Definition 2.1.** Assume  $\xi \in (0, 1)$ ,  $f : [0, 1] \times R \rightarrow R$  is continuous. A function  $u(t)$  is called an upper solution for problem (1.1) if it satisfies

$$\begin{cases} u'''(t) + f(t, u(t)) \leq 0, & t \in [0, 1], \\ u(0) = u''(0) = 0, \\ p(u(1), u(\xi)) \geq 0. \end{cases}$$

Similarly, a function  $u(t)$  is called a lower solution for problem (1.1) if it satisfies

$$\begin{cases} u'''(t) + f(t, u(t)) \geq 0, & t \in [0, 1], \\ u(0) = u''(0) = 0, \\ p(u(1), u(\xi)) \leq 0. \end{cases}$$

**Lemma 2.2.** Assume that  $\alpha\xi \neq 1$ ,  $b \in R$  and  $h : [0, 1] \rightarrow R$  is continuous. Then the boundary value problem

$$\begin{cases} -u'''(t) = h(t), & t \in [0, 1], \\ u(0) = u''(0) = 0, \\ u(1) = \alpha u(\xi) + b, \end{cases}$$

has a unique solution

$$u(t) = \frac{b}{1 - \alpha\xi}t + \frac{\alpha t}{1 - \alpha\xi} \int_0^1 G(\xi, s)h(s)ds + \int_0^1 G(t, s)h(s)ds,$$

where

$$G(t, s) = \frac{1}{2} \begin{cases} (1-t)(t-s)^2, & 0 \leq s \leq t \leq 1, \\ t(1-s)^2, & 0 \leq t \leq s \leq 1. \end{cases} \quad (2.1)$$

*Proof.* Integrating the equation

$$-u'''(t) = h(t), \quad t \in [0, 1],$$

over  $[0, t]$  for three times, we have

$$u(t) = -\frac{1}{2} \int_0^t (t-s)^2 h(s)ds + \frac{1}{2} u''(0)t^2 + u'(0)t + u(0).$$

Due to the boundary conditions  $u(0) = u''(0) = 0$ ,  $u(1) = \alpha u(\xi) + b$ , it follows that

$$-\frac{1}{2} \int_0^1 (1-s)^2 h(s) ds + u'(0) = \alpha u(\xi) + b.$$

And then

$$\begin{aligned} u(t) &= t(\alpha u(\xi) + b) + \frac{1}{2} t \int_0^1 (1-s)^2 h(s) ds - \frac{1}{2} \int_0^t (t-s)^2 h(s) ds \\ &= t(\alpha u(\xi) + b) + \frac{1}{2} \int_0^t t(1-s)^2 h(s) ds - \frac{1}{2} \int_0^t (t-s)^2 h(s) ds + \frac{1}{2} \int_t^1 t(1-s)^2 h(s) ds \\ &= t(\alpha u(\xi) + b) + \frac{1}{2} \int_0^t (1-t)(t-s^2) h(s) ds + \frac{1}{2} \int_t^1 t(1-s)^2 h(s) ds \\ &= t(\alpha u(\xi) + b) + \int_0^1 G(t,s) h(s) ds. \end{aligned} \quad (2.2)$$

Putting  $t = \xi$ , we have

$$u(\xi) = \frac{1}{1-\alpha\xi} \left( \xi b + \int_0^1 G(\xi,s) h(s) ds \right).$$

Substituting it into (2.2), we get

$$u(t) = \frac{b}{1-\alpha\xi} t + \frac{\alpha t}{1-\alpha\xi} \int_0^1 G(\xi,s) h(s) ds + \int_0^1 G(t,s) h(s) ds.$$

□

**Remark 2.3.** It is easy to see that  $G(t,s) > 0$  for all  $(t,s) \in (0,1) \times (0,1)$  and

$$u'(t) = \frac{b}{1-\alpha\xi} + \frac{\alpha}{1-\alpha\xi} \int_0^1 G(\xi,s) h(s) ds + \int_0^1 G'(t,s) h(s) ds, \quad (2.3)$$

where

$$G'(t,s) = \frac{1}{2} \begin{cases} 1-2t+s^2, & 0 \leq s \leq t \leq 1, \\ (1-s)^2, & 0 \leq t \leq s \leq 1. \end{cases} \quad (2.4)$$

**Remark 2.4.** For  $(t,s) \in [0,1] \times [0,1]$ ,

$$0 \leq t(1-t)\varphi(s) \leq G(t,s) \leq \varphi(s), \quad (2.5)$$

where  $\varphi(s) = \frac{1}{8}(1+s)^2(1-s)^2$ .

In fact, for  $t \in [s,1]$ ,  $G(\cdot,s) = \frac{1}{2}(1-t)(t-s^2)$  attains its maximum at  $t = \frac{1}{2}(1+s^2) \in [s,1]$ , so that

$$G(\cdot,s) \leq \varphi(s) = \frac{1}{8}(1+s)^2(1-s)^2.$$

For  $t \in [0,s]$ , clearly  $G(\cdot,s) \leq \frac{1}{2}s(1-s)^2 \leq \varphi(s) = \frac{1}{8}(1+s)^2(1-s)^2$ . That is,  $G(t,s) \leq \varphi(s)$  for  $(t,s) \in [0,1] \times [0,1]$ .

Also, for  $s \in [0,t]$ ,

$$\frac{G(t,s)}{\varphi(s)} = \frac{\frac{1}{2}(1-t)(t-s^2)}{\frac{1}{8}(1+s)^2(1-s)^2} \geq \frac{4(1-t)(t-st)}{(1+s)^2(1-s)^2} = \frac{4t(1-t)}{(1+s)^2(1-s)} \geq t(1-t).$$

If for  $s \in [t, 1]$ ,

$$\frac{G(t, s)}{\varphi(s)} = \frac{\frac{1}{2}t(1-s)^2}{\frac{1}{8}(1+s)^2(1-s)^2} = \frac{4t}{(1+s)^2} \geq t.$$

Since  $\min\{t, t(1-t)\} = t(1-t)$ , we have

$$0 \leq t(1-t)\varphi(s) \leq G(t, s) \leq \varphi(s).$$

**Lemma 2.5.** Assume that  $0 < \alpha\bar{\xi} < 1$ . If  $u(t) \in C[0, 1]$ , satisfying  $u'''(t) \in C[0, 1]$  and

$$\begin{cases} u'''(t) \leq 0, & t \in [0, 1], \\ u(0) = u''(0) = 0, \\ u(1) \geq \alpha u(\bar{\xi}), \end{cases}$$

then  $u(t) \geq 0$ ,  $t \in [0, 1]$ .

*Proof.* Let  $-u'''(t) = h(t) \geq 0$ ,  $t \in [0, 1]$  and  $u(1) = \alpha u(\bar{\xi}) + b$ ,  $b \geq 0$ . Then by Lemma 2.2, we get

$$\begin{aligned} u(t) &= \frac{b}{1-\alpha\bar{\xi}}t + \frac{\alpha t}{1-\alpha\bar{\xi}} \int_0^1 G(\bar{\xi}, s)h(s)ds + \int_0^1 G(t, s)h(s)ds \\ &\geq 0. \end{aligned}$$

□

**Lemma 2.6.** Assume that  $\kappa(t), \mu(t) \in C[0, 1]$  and

$$\int_0^1 (1+s)^2(1-s)^2 |\kappa(s)| ds < \frac{8(1-\alpha\bar{\xi})}{\alpha+1-\alpha\bar{\xi}}. \quad (2.6)$$

Then the following linear boundary value problem

$$\begin{cases} -u'''(t) = \kappa(t)u(t) + \mu(t), & t \in [0, 1], \\ u(0) = u''(0) = 0, \\ u(1) = \alpha u(\bar{\xi}) + b, \end{cases} \quad (2.7)$$

has a unique solution  $u(t) \in C[0, 1]$ , where  $0 < \alpha\bar{\xi} < 1, b \geq 0$ .

*Proof.* By Lemma 2.2, problem (2.7) is equivalent to the following integral equation

$$\begin{aligned} u(t) &= \frac{b}{1-\alpha\bar{\xi}}t + \frac{\alpha t}{1-\alpha\bar{\xi}} \int_0^1 G(\bar{\xi}, s) (\kappa(s)u(s) + \mu(s)) ds \\ &\quad + \int_0^1 G(t, s) (\kappa(s)u(s) + \mu(s)) ds =: Tu(t). \end{aligned}$$

Obviously,  $T : C[0, 1] \rightarrow C[0, 1]$ . Note that by (2.5), we have for any  $u, v \in C[0, 1]$ ,

$$\begin{aligned} |Tu(t) - Tv(t)| &\leq \frac{\alpha t}{1-\alpha\bar{\xi}} \int_0^1 G(\bar{\xi}, s) |\kappa(s)| |u(s) - v(s)| ds + \int_0^1 G(t, s) |\kappa(s)| |u(s) - v(s)| ds \\ &\leq \frac{1}{8} \left( 1 + \frac{\alpha t}{1-\alpha\bar{\xi}} \right) \|u - v\| \int_0^1 (1+s)^2(1-s)^2 |\kappa(s)| ds \\ &\leq \frac{1}{8} \left( \frac{\alpha+1-\alpha\bar{\xi}}{1-\alpha\bar{\xi}} \right) \|u - v\| \int_0^1 (1+s)^2(1-s)^2 |\kappa(s)| ds = L \|u - v\|, \end{aligned}$$

where  $L := \frac{1}{8} \left( \frac{\alpha+1-\alpha\zeta}{1-\alpha\zeta} \right) \int_0^1 (1+s)^2(1-s)^2 |\kappa(s)| ds < 1$ , which is easy to see from (2.6). Therefore, the operator  $T$  is a contraction map in the space  $C[0,1]$  and  $T$  has a unique fixed point in  $C[0,1]$ .  $\square$

**Lemma 2.7.** *Assume that  $0 < \alpha\zeta < 1$ ,  $\kappa(t) \in C[0,1]$ ,  $\kappa(t) > 0$  and satisfies (2.6). If  $u(t) \in C[0,1]$ , satisfying  $u'''(t) \in C[0,1]$  and*

$$\begin{cases} u'''(t) + \kappa(t)u(t) \leq 0, & t \in [0,1], \\ u(0) = u''(0) = 0, \\ u(1) \geq \alpha u(\zeta) \end{cases}$$

then  $u(t) \geq 0$ ,  $t \in [0,1]$ .

*Proof.* Let  $-u'''(t) = \kappa(t)u(t) + \mu(t)$ ,  $t \in [0,1]$ ,  $\kappa(t) > 0$ ,  $\mu(t) \geq 0$ ,  $u(1) = \alpha u(\zeta) + b$ ,  $b \geq 0$ .

Suppose that the inequality  $u(t) \geq 0$ ,  $t \in [0,1]$  is not true. It means that there exists at least a  $t^* \in [0,1]$  such that  $u(t^*) < 0$ . Without loss of generality, we assume  $u(t^*) = \min\{u(t) : t \in [0,1]\} = \rho$ ,  $\rho < 0$ . Then by Lemma 2.2 and (2.5), we have

$$\begin{aligned} u(t) &= \frac{b}{1-\alpha\zeta}t + \frac{\alpha t}{1-\alpha\zeta} \int_0^1 G(\zeta, s)(\kappa(s)u(s) + \mu(s))ds \\ &\quad + \int_0^1 G(t, s)(\kappa(s)u(s) + \mu(s))ds \\ &\geq \frac{\alpha t}{1-\alpha\zeta} \int_0^1 G(\zeta, s)\kappa(s)u(s)ds + \int_0^1 G(t, s)\kappa(s)u(s)ds \\ &\geq u(t^*) \left( \frac{\alpha t}{1-\alpha\zeta} \int_0^1 \varphi(s)\kappa(s)ds + \int_0^1 \varphi(s)\kappa(s)ds \right). \end{aligned}$$

Let  $t = t^*$ , and note that  $\rho < 0$ ,  $0 < \alpha\zeta < 1$ ,  $0 < \zeta < 1$ , it follows that

$$\begin{aligned} \rho &\geq \rho \left( \frac{\alpha t^*}{1-\alpha\zeta} \int_0^1 \varphi(s)\kappa(s)ds + \int_0^1 \varphi(s)\kappa(s)ds \right) \\ &\geq \rho \left( \frac{\alpha}{1-\alpha\zeta} \int_0^1 \varphi(s)\kappa(s)ds + \int_0^1 \varphi(s)\kappa(s)ds \right). \end{aligned}$$

And then

$$1 \leq \frac{\alpha + 1 - \alpha\zeta}{8(1-\alpha\zeta)} \int_0^1 (1+s)^2(1-s)^2 \kappa(s) ds.$$

That is

$$\int_0^1 (1+s)^2(1-s)^2 \kappa(s) ds \geq \frac{8(1-\alpha\zeta)}{\alpha + 1 - \alpha\zeta},$$

which is in contradiction to (2.6). Hence  $u(t) \geq 0$  for all  $t \in [0,1]$ .  $\square$

### 3 Main result

In this section, we shall apply the method of upper and lower solutions and monotone iterative technique to consider the existence of extremal solutions for problem (1.1).

**Theorem 3.1.** *Assume that the following conditions hold.*

(A<sub>1</sub>)  $f(t, u)$  is increasing with respect to  $u$ .

(A<sub>2</sub>)  $v_0(t), w_0(t) \in C[0, 1]$  are lower and upper solutions of problem (1.1), respectively, and  $v_0(t) \leq w_0(t)$ ,  $t \in [0, 1]$ .

(A<sub>3</sub>) There exist constants  $\varsigma, \tau$  such that  $0 < \tau\bar{\xi} < \varsigma$  and for  $v_0(1) \leq x \leq y \leq w_0(1)$ ,  $v_0(\bar{\xi}) \leq \bar{x} \leq \bar{y} \leq w_0(\bar{\xi})$ ,

$$p(y, \bar{y}) + \tau(\bar{y} - \bar{x}) \leq p(x, \bar{x}) + \varsigma(y - x).$$

Then problem (1.1) has extremal solutions in the sector  $[v_0, w_0]$ , where

$$[v_0, w_0] = \{u \in C[0, 1] : v_0(t) \leq u(t) \leq w_0(t), t \in [0, 1]\}.$$

*Proof.* For  $n = 0, 1, \dots$ , define

$$\begin{aligned} v_{n+1}(t) &= \frac{b_n}{1 - \alpha\bar{\xi}}t + \frac{\alpha t}{1 - \alpha\bar{\xi}} \int_0^1 G(\bar{\xi}, s)f(s, v_n(s))ds + \int_0^1 G(t, s)f(s, v_n(s))ds, \\ w_{n+1}(t) &= \frac{c_n}{1 - \alpha\bar{\xi}}t + \frac{\alpha t}{1 - \alpha\bar{\xi}} \int_0^1 G(\bar{\xi}, s)f(s, w_n(s))ds + \int_0^1 G(t, s)f(s, w_n(s))ds, \end{aligned}$$

where

$$\alpha = \frac{\tau}{\varsigma}, \quad b_n = v_n(1) - \alpha v_n(\bar{\xi}) - \frac{1}{\varsigma}g(v_n(1), v_n(\bar{\xi})), \quad c_n = w_n(1) - \alpha w_n(\bar{\xi}) - \frac{1}{\varsigma}g(w_n(1), w_n(\bar{\xi})).$$

Then due to Lemma 2.2, it is easy to show that  $v_{n+1}(t), w_{n+1}(t)$  are solutions of the following boundary value problems, respectively:

$$\begin{cases} -v_{n+1}'''(t) = f(t, v_n(t)), & t \in [0, 1], \\ v_{n+1}(0) = v_{n+1}''(0) = 0, \\ 0 = p(v_n(1), v_n(\bar{\xi})) + \varsigma(v_{n+1}(1) - v_n(1)) - \tau(v_{n+1}(\bar{\xi}) - v_n(\bar{\xi})), \end{cases} \quad (3.1)$$

and

$$\begin{cases} -w_{n+1}'''(t) = f(t, w_n(t)), & t \in [0, 1], \\ w_{n+1}(0) = w_{n+1}''(0) = 0, \\ 0 = p(w_n(1), w_n(\bar{\xi})) + \varsigma(w_{n+1}(1) - w_n(1)) - \tau(w_{n+1}(\bar{\xi}) - w_n(\bar{\xi})). \end{cases} \quad (3.2)$$

Moreover, from (2.3) we have

$$v'_{n+1}(t) = \frac{b_n}{1 - \alpha\bar{\xi}} + \frac{\alpha}{1 - \alpha\bar{\xi}} \int_0^1 G(\bar{\xi}, s)f(s, v_n(s))ds + \int_0^1 G'(t, s)f(s, v_n(s))ds, \quad (3.3)$$

$$w'_{n+1}(t) = \frac{c_n}{1 - \alpha\bar{\xi}} + \frac{\alpha}{1 - \alpha\bar{\xi}} \int_0^1 G(\bar{\xi}, s)f(s, w_n(s))ds + \int_0^1 G'(t, s)f(s, w_n(s))ds, \quad (3.4)$$

where  $G'(t, s)$  is given as in (2.4).

**Claim 1.** The sequences  $v_n(t), w_n(t) (n \geq 1)$  are lower and upper solutions of problem (1.1), respectively and the following relation holds

$$v_0(t) \leq v_1(t) \leq \dots \leq v_n(t) \leq \dots \leq w_n(t) \leq \dots \leq w_1(t) \leq w_0(t), \quad t \in [0, 1]. \quad (3.5)$$

First, we prove that

$$v_0(t) \leq v_1(t) \leq w_1(t) \leq w_0(t), \quad t \in [0, 1].$$

Let  $x(t) = w_0(t) - w_1(t)$ . From (3.2) and  $(A_1)$ , we have

$$\begin{cases} x'''(t) \leq 0, & t \in [0, 1], \\ x(0) = x''(0) = 0, \\ x(1) \geq \frac{\tau}{\zeta}x(\xi), & 0 < \tau\xi < \zeta. \end{cases}$$

In view of Lemma 2.5, we have  $x(t) \geq 0, t \in [0, 1]$ , that is  $w_0(t) \geq w_1(t)$ . Similarly, it can be obtained that  $v_0(t) \leq v_1(t), t \in [0, 1]$ .

Now, let  $x(t) = w_1(t) - v_1(t)$ . From  $(A_1)$  and  $(A_2)$ , it follows that

$$x'''(t) = f(t, v_0(t)) - f(t, w_0(t)) \leq 0.$$

Also,  $x(0) = x''(0) = 0$  and

$$\begin{aligned} 0 &= p(w_0(1), w_0(\xi)) - p(v_0(1), v_0(\xi)) + \zeta(w_1(1) - w_0(1) - v_1(1) + v_0(1)) \\ &\quad - \tau(w_1(\xi) - w_0(\xi) - v_1(\xi) + v_0(\xi)) \\ &\leq \zeta(w_0(1) - v_0(1)) - \tau(w_0(\xi) - v_0(\xi)) + \zeta(w_1(1) - w_0(1) - v_1(1) + v_0(1)) \\ &\quad - \tau(w_1(\xi) - w_0(\xi) - v_1(\xi) + v_0(\xi)) \\ &= \zeta x(1) - \tau x(\xi). \end{aligned}$$

That is,

$$x(0) = x''(0) = 0, \quad x(1) \geq \frac{\tau}{\zeta}x(\xi).$$

By Lemma 2.5, we have  $w_1(t) \geq v_1(t), t \in [0, 1]$ . And then, by induction, (3.5) holds.

In what follows, we show that  $v_1(t), w_1(t)$  are lower and upper solutions of problem (1.1), respectively. From (3.1), (3.2) and  $(A_1), (A_2)$ , it follows that

$$-v_1'''(t) = f(t, v_0(t)) \leq f(t, v_1(t)).$$

Also,  $v_1(0) = v_1''(0) = 0$  and

$$\begin{aligned} 0 &= -p(v_0(1), v_0(\xi)) + p(v_1(1), v_1(\xi)) - p(v_1(1), v_1(\xi)) - \zeta[v_1(1) - v_0(1)] + \tau[v_1(\xi) - v_0(\xi)] \\ &\leq \zeta[v_1(1) - v_0(1)] - \tau[v_1(\xi) - v_0(\xi)] - p(v_1(1), v_1(\xi)) - \zeta[v_1(1) - v_0(1)] + \tau[v_1(\xi) - v_0(\xi)] \\ &= -p(v_1(1), v_1(\xi)), \end{aligned}$$

which prove that  $v_1(t)$  is a lower solution of problem (1.1). Similarly, it can be obtained that  $w_1(t)$  is an upper solution of problem (1.1).

Analogously to the above arguments, using the induction method, we can show that the sequences  $v_n(t), w_n(t)$  ( $n \geq 1$ ) are lower and upper solutions of problem (1.1), respectively and the following relation holds

$$v_0(t) \leq v_1(t) \leq \dots \leq v_n(t) \leq \dots \leq w_n(t) \leq \dots \leq w_1(t) \leq w_0(t), \quad t \in [0, 1].$$

**Claim 2.** The sequences  $\{v_n(t)\}, \{w_n(t)\}$  uniformly converge to their limit functions  $v(t), w(t)$ , respectively.

We need to show that the sequences are bounded and equicontinuous on  $[0, 1]$ . Indeed,

$$C_1 \leq v_0(t) \leq \cdots \leq v_n(t) \leq \cdots \leq \cdots \leq w_n(t) \cdots \leq w_0(t) \leq C_2 \quad (3.6)$$

for  $t \in [0, 1]$  and  $n = 1, 2, \dots$ . That is to say that the sequences  $\{v_n(t)\}, \{w_n(t)\}$  are uniformly bounded with respect to  $t$ . Note that  $\{v'_n(t)\}, \{w'_n(t)\}$  are bounded on  $[0, 1]$  by  $C_3 > 0$  because (3.3), (3.4), (3.6) and  $|f(t, v_n)|, |p(v_n, v_n)|$  is bounded. Hence  $\{v_n(t)\}, \{w_n(t)\}$  are equicontinuous because for  $\forall \varepsilon > 0, t_1, t_2 \in [0, 1]$  such that  $|t_1 - t_2| < \frac{\varepsilon}{M_3}$ , we have

$$|v_n(t_1) - v_n(t_2)| = |v'_n(\gamma)| |t_1 - t_2| < \varepsilon, \quad |w_n(t_1) - w_n(t_2)| < \varepsilon, \quad \gamma \in [0, 1].$$

Therefore, by the Arzelà–Ascoli theorem, the sequences  $\{v_n(t)\}, \{w_n(t)\}$  have subsequences  $\{v_{n_k}(t)\}, \{w_{n_k}(t)\}$  which uniformly converge to their continuous limit functions  $v(t), w(t)$ , respectively.

**Claim 3.** The limit functions  $v(t), w(t)$  are the minimal solution and maximal solution of problem (1.1), respectively.

Let  $u(t) \in [v_0(t), w_0(t)]$  be any solution of problem (1.1). We assume that the following relation holds for some  $k \in N$ :

$$v_k(t) \leq u(t) \leq w_k(t), \quad t \in [0, 1].$$

Let  $y(t) = u(t) - v_{k+1}(t), z(t) = w_{k+1}(t) - u(t)$ . Then

$$\begin{cases} y'''(t) \leq 0, & t \in [0, 1], \\ y(0) = y''(0) = 0, \\ y(1) \geq \frac{\tau}{\zeta} y(\xi), & 0 < \tau\xi < \zeta, \end{cases}$$

and

$$\begin{cases} z'''(t) \leq 0, & t \in [0, 1], \\ z(0) = z''(0) = 0, \\ z(1) \geq \frac{\tau}{\zeta} z(\xi), & 0 < \tau\xi < \zeta. \end{cases}$$

This and Lemma 2.5 show  $v_{k+1}(t) \leq u(t) \leq w_{k+1}(t)$ . By induction,  $v_n(t) \leq u(t) \leq w_n(t)$ ,  $t \in [0, 1]$  for all  $n \in N$ . Taking the limit as  $n \rightarrow \infty$ , we get  $v(t) \leq u(t) \leq w(t)$ ,  $t \in [0, 1]$ .  $\square$

**Theorem 3.2.** Assume that all assumptions of Theorem 3.1 hold. In addition, we assume that there exists  $q(t) > 0$  ( $\in C[0, 1]$ ) such that

$$f(t, x) - f(t, y) \leq q(t)(x - y), \quad v_0(t) \leq x(t) \leq y(t) \leq w_0(t), \quad t \in [0, 1]$$

and

$$\int_0^1 (1+s)^2(1-s)^2 q(s) ds < \frac{8(\zeta - \tau\xi)}{\zeta + \tau - \tau\xi}.$$

Then the sequences  $\{v_n(t)\}, \{w_n(t)\}$  ( $n \geq 1$ ) satisfying

$$\begin{cases} -v''_{n+1}(t) = f(t, v_n(t)) + q(t)[v_{n+1}(t) - v_n(t)], & t \in (0, 1), \\ v_{n+1}(0) = v''_{n+1}(0) = 0, \\ 0 = p(v_n(1), v_n(\xi)) + \zeta(v_{n+1}(1) - v_n(1)) - \tau(v_{n+1}(\xi) - v_n(\xi)), \end{cases}$$



and

$$\begin{cases} -w_{n+1}'''(t) = f(t, w_n(t)) + q(t)[w_{n+1}(t) - w_n(t)], & t \in (0, 1), \\ w_{n+1}(0) = w_{n+1}''(0) = 0, \\ 0 = p(w_n(1), w_n(\xi)) + \zeta(w_{n+1}(1) - w_n(1)) - \tau(w_{n+1}(\xi) - w_n(\xi)), \end{cases}$$

also uniformly converge to their continuous limit functions  $v(t), w(t)$ , respectively. That is,  $v(t), w(t)$  are also extremal solutions for problem (1.1).

*Proof.* Using Lemmas 2.6 and 2.7, we can complete the proof by the same way as in Theorem 3.1.  $\square$

**Example 3.3.** Consider the following third-order boundary value problem

$$\begin{cases} -u'''(t) = u(t) \sin t - \frac{113}{8}t^{\frac{1}{2}}, & t \in [0, 1], \\ u(0) = u''(0) = 0, \\ u(1) - u(\frac{1}{2}) - \frac{1}{8}u(1)u(\frac{1}{2}) = 0, \end{cases} \quad (3.7)$$

where

$$f(x, y) = y \sin x - \frac{113}{8}x^{\frac{1}{2}}, \quad p(x, y) = x - y - \frac{1}{8}xy, \quad \xi = \frac{1}{2}.$$

It is not difficult to show that  $v_0 = 0, w_0(t) = t^{\frac{7}{2}}$  are lower and upper solutions of problem (3.7), respectively. Moreover, for  $t \in [0, 1]$ ,  $f(t, u)$  is increasing with respect to  $u$ , and for  $v_0(1) \leq x \leq y \leq w_0(1), v_0(\eta) \leq \bar{x} \leq \bar{y} \leq w_0(\eta)$ ,

$$p(y, \bar{y}) - p(x, \bar{x}) = (y - x) - (\bar{y} - \bar{x}) - \frac{1}{8}(y\bar{y} - x\bar{x}) \leq (y - x) - (\bar{y} - \bar{x}).$$

Choose  $\zeta = \tau = 1$  in Theorem 3.1 or  $\zeta = \tau = 1, q(t) = \sin t$  in Theorem 3.2, problem (3.7) has extremal solutions in  $[v_0(t), w_0(t)]$ .

## Acknowledgements

The authors would like to thank the referee and Professor Kosmatov for their careful reading and kind suggestions. This work was supported by Program for Scientific Research Innovation Team in Colleges and Universities of Shandong Province, the Doctoral Program Foundation of Education Ministry of China (20133705110003), the Natural Science Foundation of Shandong Province of China (ZR2014AM007), the National Natural Science Foundation of China (11571197).

## References

- [1] M. AKTAŞ, D. ÇAKMAK, A. TIRYAKI, On the Lyapunov-type inequalities of a three-point boundary value problem for third order linear differential equations, *Appl. Math. Lett.* **45**(2015), 1–6. [MR3316952](#); [url](#)
- [2] D. R. ANDERSON, Green's function for a third-order generalized right focal problem, *J. Math. Anal. Appl.* **288**(2003), 1–14. [MR2019740](#); [url](#)
- [3] A. BOUCHERIF, N. AL-MALKI, Nonlinear three-point third-order boundary value problems, *Appl. Math. Comput.* **190**(2007), 1168–1177. [MR2339710](#); [url](#)

- [4] Z. J. DU, W. G. GE, X. J. LIN, Existence of solutions for a class of third-order nonlinear boundary value problems, *J. Math. Anal. Appl.* **294**(2004), 104–112. [MR2059792](#); [url](#)
- [5] Z. J. DU, W. G. GE, M. R. ZHOU, Singular perturbations for third-order nonlinear multi-point boundary value problem, *J. Differential Equations* **218**(2005), 69–90. [MR2174967](#); [url](#)
- [6] X. F. FENG, H. Y. FENG, D. L. BAI, Eigenvalue for a singular third-order three-point boundary value problem, *Appl. Math. Comput.* **219**(2013), 9783–9790. [MR3049599](#); [url](#)
- [7] C. S. GOODRICH, Positive solutions to boundary value problems with nonlinear boundary conditions, *Nonlinear Anal.* **75**(2012), 417–432. [MR2846811](#); [url](#)
- [8] J. R. GRAEF, J. R. L. WEBB, Third order boundary value problems with nonlocal boundary conditions, *Nonlinear Anal.* **71**(2009), 1542–1551. [MR2524369](#); [url](#)
- [9] J. R. GRAEF, B. YANG, Positive solutions of a third order nonlocal boundary value problem, *Discrete Contin. Dyn. Syst. Ser. S.* **1**(2008), 89–97. [MR2375585](#)
- [10] L. J. GUO, J. P. SUN, Y. H. ZHAO, Multiple positive solutions for nonlinear third-order three-point boundary value problems, *Electron. J. Differential Equations* **2007**, No. 112, 1–7. [MR2349940](#)
- [11] L. J. GUO, J. P. SUN, Y. H. ZHAO, Existence of positive solutions for nonlinear third-order three-point boundary value problems, *Nonlinear Anal.* **68**(2008), 3151–3158. [MR2404825](#); [url](#)
- [12] J. HENDERSON, N. KOSMATOV, Three-point third-order problems with a sign-changing nonlinear term, *Electron. J. Differential Equations* **2014**, No. 175, 1–10. [MR3262046](#)
- [13] E. ROVDEROVA, Third-order boundary-value problem with nonlinear boundary conditions, *Nonlinear Anal.* **25**(1995), 473–485. [MR1338798](#); [url](#)
- [14] Y. P. SUN, Positive solutions for third-order three-point nonhomogeneous boundary value problems, *Appl. Math. Lett.* **22**(2009), 45–51. [MR2483159](#); [url](#)
- [15] Z. L. WEI, Some necessary and sufficient conditions for existence of positive solutions for third order singular sublinear multi-point boundary value problems, *Acta Math. Sci. Ser. B Engl. Ed.* **34**(2014), No. 6, 1795–1810. [MR3260754](#); [url](#)