

BMJ Open Associations between highly active antiretroviral therapy and the presence of HPV, premalignant and malignant cervical lesions in sub-Saharan Africa, a systematic review: current evidence and directions for future research

Sonia Menon,^{1,2} Rodolfo Rossi,³ Natasha Zdraveska,⁴ Mbabazi Kariisa,² Sushama D Acharya,² Davy Vanden Broeck,^{1,5} Steven Callens⁶

To cite: Menon S, Rossi R, Zdraveska N, *et al.* Associations between highly active antiretroviral therapy and the presence of HPV, premalignant and malignant cervical lesions in sub-Saharan Africa, a systematic review: current evidence and directions for future research. *BMJ Open* 2017;7:e015123. doi:10.1136/bmjopen-2016-015123

► Prepublication history and additional material are available. To view these files please visit the journal online (<http://dx.doi.org/bmjopen-2016-015123>)

SM and RR contributed equally.

Received 10 November 2016

Revised 10 April 2017

Accepted 10 April 2017



CrossMark

For numbered affiliations see end of article.

Correspondence to

Rodolfo Rossi;
rossirodolfo@yahoo.co.uk

ABSTRACT

Objectives In sub-Saharan Africa, substantial international funding along with evidence-based clinical practice have resulted in an unparalleled scale-up of access to antiretroviral treatment at a higher CD4 count. The role and timing of highly active antiretroviral therapy (HAART) in mediating cervical disease remains unclear. The aim of this article is to systematically review all evidence pertaining to Africa and identify research gaps regarding the epidemiological association between HAART use and the presence of premalignant/malignant cervical lesions.

Method Five databases were searched until January 2017 to retrieve relevant literature from sub-Saharan Africa. Publications were included if they addressed prevalence, incidence or clearance of human papillomavirus (HPV) infection in women undergoing HAART as well as cytological or histological neoplastic abnormalities.

Results 22 studies were included, of which seven were prospective studies. Women receiving HAART are less likely to develop squamous intraepithelial lesions (SILs). There is evidence that duration of HAART along with the CD4 count may reduce the prevalence of high-risk HPV (HR-HPV), suggesting that without HAART, severe immunosuppression increases the risk of becoming or remaining infected with HR-HPV. Furthermore, according to existent literature, the CD4 count, rather than HAART coverage or its duration, plays a central role in the prevalence of cervical intraepithelial neoplasia (CIN) 2 and CIN 3.

Conclusion Our findings suggest a positive impact of HAART duration, in conjunction and interaction with CD4 count, on reducing the prevalence of HR-HPV. The greatest treatment effect might be seen among women starting at the lowest CD4 count, which may have a more instrumental role in cervical oncogenesis than either HAART use or the treatment duration on the prevalence of CIN 2 and CIN 3. There is still insufficient evidence to show a clear association between HAART coverage and the incidence of invasive cervical cancer. Enhanced surveillance on the impact of HAART treatment is crucial.

Strengths and limitations of this study

- Comprehensive review within five databases on a major public health concern for women's health.
- Contributes to and summarises the evidence of the role of CD4 count and highly active antiretroviral therapy (HAART) use in the progression to cervical disease for women who are HIV-positive in Africa.
- High variability in the sample size of the retrieved studies, therefore difficult to give the right weight to the current evidence.
- Heterogeneity in the sensitivity or specificity of screening methods may also render results less comparable; in some studies, HIV status of participants may not have been accurately measured, leaving the possibility of misclassification.
- Lack of prospective studies, with a clear strategy to deal with the loss to follow-up, retrieved by the literature search using the same definition of regression and progression of cervical disease; therefore the criterion of temporality for causality is not well fulfilled.

INTRODUCTION

Cervical cancer is the second most common female malignancy and the leading cause of death from cancer in women worldwide, of which 85% of cases occur in low and middle income countries.¹ The human papillomavirus (HPV), one of the most common sexually transmitted infections, is now well established as the aetiological agent. 'High-risk' (HR) HPV DNA has been found to be present in 99.7% of cervical cancers worldwide.² Over 200 HPV genotypes have been identified and are divided into HR and low-risk (LR) carcinogens, depending on their potential to cause cancer.

Invasive cervical cancer (ICC) and its precursor, cervical intraepithelial neoplasia (CIN), are associated with persistent infection with HR HPV genotypes. Products of HPV oncogenes E6 and E7 alter normal genetic and cellular functions and induce malignant transformation. HPV infections are usually transient, and even those that persist for a few months are usually cleared naturally, however women infected with HIV are at higher risk of contracting HPV and developing squamous intraepithelial lesions (SILs), the precursor of cervical cancer.^{3 4} In this population, this type of lesion tends to be more aggressive, persistent and more likely to recur following cervical disease treatment.^{5 6}

In sub-Saharan Africa, the region with the highest prevalence of HIV one and HIV two in the world, the current standard recommendations for first-line adult antiretroviral therapy include two nucleoside reverse transcriptase inhibitors (NRTIs) and one non-nucleoside reverse transcriptase inhibitor (NNRTI).⁷ Protease inhibitors (PIs) are primarily reserved for second-line treatment due to issues of cost, dosing frequency, drug–drug interactions, potential for long-term side effects and higher pill burden. Persons not responding to first-line regimens are usually switched to a cocktail of two NRTIs plus a boosted PI.⁸

While the use of HAART decreased the risk of Kaposi sarcoma and non-Hodgkin's lymphoma, both AIDS-induced cancers,⁹ there was no significant change in the incidence of cervical cancer.^{10–12} Moreover, studies on the impact of HAART on the natural history of cervical SILs have yielded inconsistent results.^{13 14} Moreover, most studies have been conducted in industrialised settings and not in sub-Saharan Africa, the area hardest hit by the global HIV/AIDS pandemic, home to nearly 25 million people living with HIV/AIDS.¹⁵

While the use of HAART might be expected to reduce the incidence of cervical cancer and progression of cervical dysplasia as a result of improved immunological function, as women are accessing HAART, their life expectancy will increase, due to less competing risk of dying from an opportunistic infection, which in turn may put them more at risk for ICC.¹⁶ These two reasons will augment their risks of a premalignant condition to transform into an ICC. Inversely, prolonged use of HAART might potentially prevent the acquisition of HR-HPV and/or evolution to ICC.

In the HAART policy landscape, there have been recent shifts towards earlier initiation of treatment. Until 2009, the WHO recommendations for sub-Saharan Africa stipulated initiation of antiretroviral treatment at a CD4 count threshold of 200 cells/mm³ in order to further reduce the risk of HIV transmission to others. However, this guideline was revised in 2009 and initiation of antiretroviral therapy recommended at CD4 counts of 350 cells/mm³, only to be revised in 2013 and the treatment threshold set to 500 cells/mm³ or less.¹⁷ The policy was again recently revised and treatment initiation recommended for all adults with HIV infection adults, regardless of WHO

clinical stage or CD4 cell count, following the results of the START trial.¹⁸

As a research gap, it remains to be determined whether HAART could improve control of and reduce the burden of HPV infection and the associated precancerous states if initiated even earlier in the course of HIV infection.¹⁹ The role and timing of HAART in mediating this relationship has been difficult to study and remains unclear.^{20 21} The objectives of this paper are to review the existing evidences on the association between the use of HAART and presence of HPV infection and premalignant/malignant cervical lesions.

METHODOLOGY

We conducted this systematic review based on a pre-defined search protocol that conformed to the criteria set out by the PRISMA statement (see Research Checklist).²² CINAHL, SCOPUS, Global Health, PubMed and Embase databases were searched ([figure 1](#)).

Search terms

The domains of the search terms were antiretroviral therapy, human papillomavirus, abnormal cytology, cervical dysplasia, cervical cancer and sub-Saharan Africa (see online supplementary file 1).

Inclusion criteria

Article selection criteria included any clinic-based randomised controlled trials (RCTs), meta-analyses/systematic reviews, observational or population-based linkage studies documenting both HAART status and HPV/CIN/ICC rates in sub-Saharan Africa. The five databases mentioned above were searched in January 2017 for articles published from January 1997 onwards to limit the publications to the combined HAART era. The following types of articles were excluded from our search: case reports, articles that did not include combination antiretroviral therapy, risk factors for CIN 2+ or did not stratify data by HAART users. We limited our search to English language literature.

Study selection

We considered the following four components (Population, Intervention, Comparison, Outcome—PICO) to assess and categorise studies to be included in this review.

Population: the population eligible for this study includes women with HIV infection or women with AIDS with concomitant HPV infection and/or with various degrees of abnormal cytology diagnosed through a recognised method such as PCR HPV genotyping, cytology, visual inspection with acetic acid (VIA), visual inspection with Lugol's iodine (VILI) and histologically confirmed biopsies, and living in sub-Saharan countries.

Intervention: the intervention examined in this study is the administration of HAART, defined as at least three antiretroviral drugs belonging to two drug classes (NRTI, NNRTI or PI) and (if possible) the duration of treatment.

Comparison: comparison is made between women not receiving HAART and women receiving HAART.




Figure 1 Flowchart with steps to retrieve the studies to include in the review.

Outcome(s): The wide spectrum of cervical diseases, ranging from the HPV infection (and its potential clearing) through cervical oncogenesis and precursor lesions at the lower end of the neoplastic spectrum, to invasive cervical dysplasia.

All potentially relevant publications were evaluated by two reviewers (SM and RR) and were included in this review if they addressed any of the following outcomes: (1) prevalence, incidence of onset or clearance of HPV infection in women on HAART; (2) prevalence of cervical cytological or histological neoplastic abnormalities or ICC in HAART users. Possible cases of disagreement about the inclusion or exclusion of articles were resolved on an individual basis via thorough discussion among the authors of the justification for exclusion/inclusion.

With the exception of steps concerning the quantitative synthesis (meta-analysis), the PRISMA 2009 checklist and flow diagram²³ were used to screen and include studies. We also enriched the search by reviewing reference lists from retrieved publications to identify additional manuscripts not captured by the searches. In reporting the results, we used the original terminology as stated in the original articles.

Duplication removal: the original searches from all the databases were combined in an EndNote library (EndNote version 8, San Francisco, California, USA) and all duplicates removed.

Extraction of relevant information from the selected studies

For all publications, SM and RR recorded the following variables: study location, year of study, study design, study population, main exposure of interest, main outcome of interest, confounding factors adjusted for and main findings.

Data assessment

Because of the lack of randomised trials retrieved (due to ethical considerations), we considered prospective/

longitudinal studies of the highest quality. We reported an overall score (from 3 to 1) based on our assessment of the overall quality of evidence for the following parameters: (1) minimisation of selection bias: (score '3' = strong) the participants were likely, (score '2' = moderate) somewhat likely, (score '1' = weak) not likely to represent the target population. (2) Study design: (score '3' = strong) prospective study and large sample size (higher than 200)²³; a prospective study with a sample size lower than 200 and a cross-sectional study with a sample size >200 (score '2' = moderate); and a cross-sectional study with a sample size under 200 (score '1' = weak). (3) Diagnostic accuracy: (score '3') strong histopathological diagnosis; moderate (score '2') VIA/VILI using colposcopy with biopsy solely in cases when lesions appeared suspicious for CIN; and weak (score '1') cytological diagnosis. (4) Confounders: (score '3') strong, controlled for all relevant confounders; (score '2') moderate, controlled for some confounders; (score '1') weak, control for confounders not specified. (5) Withdrawals, loss to follow-up and dropouts: (score '3') strong, description of a strategy to minimise the loss to follow-up, reported withdrawals and dropouts (80–100% of participants completed the study)²⁴; (score '2') moderate, reported withdrawals and dropouts; (60% of participants completed the study); (score '1') weak, withdrawal and dropout rates not specified.

We assessed the methodological quality of evidence by calculating the mean of each study and attributing the mean one of the qualitative attributes, strong, moderate, and weak.

Results

On 15 January 2017, we included 22 studies in this review (figure 1). No disagreement appeared between the two main coauthors about the studies to be included. A summary of the study is presented in table 1.

**Table 1** Summary of the quality of evidence assessment for the included studies

First author	Minimisation of selection bias	Study design	Diagnostic accuracy	Control of confounders	Loss to follow-up	Overall methodological score and quality
Mogtomo M	strong	weak	weak	weak	N/A	1.5 (weak to moderate)
Ononogbu U	strong	moderate	weak	weak	N/A	1.8 (weak to moderate)
Ezechi O. C	strong	moderate	weak	strong	N/A	2.3 (moderate to strong)
Ezechi O. C	strong	moderate	strong	strong	N/A	2.8 (moderate to strong)
McKenzie KP	strong	moderate	weak	weak	N/A	1.8 (weak to moderate)
Huscko M	strong	moderate	moderate	strong	N/A	2.5 (moderate to strong)
De Vuyst H	strong	moderate	strong	strong	N/A	2.8 (moderate to strong)
Gedefaw A	strong	moderate	weak	strong	N/A	2.3 (moderate to strong)
Mutyaba I *	N/A	N/A	N/A	N/A	N/A	N/A
Firnhaber C	strong	strong	weak	strong	weak	2.2 (moderate to strong)
Memiah P	strong	moderate	moderate	weak	N/A	2.0 (moderate)
Rositch A	strong	Strong	strong	strong	weak	2.6 (moderate to strong)
Omar T	strong	strong	weak	strong	strong	2.6 (moderate to strong)
Adler DH	strong	strong	weak	strong	weak	2.2 (moderate to strong)
Zeier MD	strong	strong	strong	strong	weak	2.6 (moderate to strong)
Bekolo CP	strong	moderate	moderate	strong	N/A	2.5 (moderate to strong)
Chichi Capo CV	strong	weak	strong	weak	N/A	2.0 (moderate)
van Bogaert LJ	strong	moderate	strong	weak	N/A	2.0 (moderate)
Mwakigonja AM	strong	moderate	weak	weak	N/A	1.8 (weak to moderate)
Jacquet A	strong	moderate	moderate	strong	N/A	2.5 (moderate to strong)
Katumba AC	strong	moderate	weak	weak	N/A	1.8 (weak to moderate)
Kelly HA	strong	strong	strong	strong	strong	3.0 (strong)

*Ecological study, not amenable to quality assessment since this study is population based and not individual based.

Study population

A total of 19 345 women with HIV infection were included in the studies retrieved and included in this review (one study was an ecological study which did not report the individual sample size). The sample sizes for these studies ranged from 70 to 3241 women with HIV infection.

Study design

The published literature on HAART and HR HPV presence and cervical dysplasia is characterised by a lack of standardisation in study designs. Twelve studies were cross sectional, two case-control, seven were prospective cohorts and one was a time-series ecological study.

Geographical location

Two studies were conducted in Nigeria; four in Kenya; seven in South Africa; one in Ethiopia; two in Uganda; two in Cameroon; one in Benin; one in Tanzania; one in Ivory Coast; and one in Burkina Faso (see table 2).

Quality assessment of studies

The overall quality of evidence ranged from weak/moderate (score of 1.5) to strong (score of 3) (table 1). Five studies had an average score below 2 and the quality of their evidence was considered as 'weak to moderate'; 15 studies had a score above 2 but less than 3 and the quality of their evidence was considered as 'moderate' or

'moderate to strong'. Just one study had a full score of 3 and was considered as 'strong'. One study was ecological in design and could not be assessed in the same way as the other studies that analysed individual data. The quality of evidence was mostly downgraded due to the cross-sectional nature of the studies as well as the lack of studies where samples have been histopathologically confirmed.

We discuss these differences by stage of disease: (1) HAART and HR HPV infection; (2) HAART and SIL/CIN; (3) HAART and ICC, and outline trends and differences.

Summary of studies on the epidemiological association between HAART and HR HPV infection/prevalence

Four studies found a significant and inverted association between HAART use and HR HPV infection/prevalence. De Vuyst *et al* (Kenya, 2012)²⁵ reported that the prevalence of HR HPV linearly decreased with use and duration of HAART ($p=0.011$). This association becomes more evident when this analysis is stratified by CD4 count categories. While the prevalence of HR HPV among patients not receiving HAART linearly increased with each decrease in CD4 count category (CD4 <200; CD4 200–349; CD4 350–499; and CD4 ≥500; PR for linear trend 5.03, $p=0.025$), this trend was only marginally significant



among women receiving HAART for <2 years (prevalence ratio (PR) for linear trend 2.98, $p=0.084$) and insignificant for women on HAART therapy for ≥ 2 years (PR for linear trend 1.19, $p=0.275$). Zejer *et al* (South Africa 2015)²⁶ reported a significant reduction in risk of HPV in women on HAART (OR 0.23, 95% CI 0.15 to 0.37) in the 204 patients eligible for HAART, and for every additional month of HAART since initiation, the risk of detection of any HPV type decreased by 9%.

Kelly *et al* (Burkina Faso and South Africa, 2017)²⁷ observed that, compared with long-duration antiretroviral therapy users (>2 years), HR HPV prevalence was higher among those on short duration HAART in both countries. However, when adjusted for CD4 cell count, this association was observed in Burkina Faso only (65.1 vs 52.1% for <2 years compared with >2 years; aPR 1.24, 95% CI 1.04 to 1.47). Ezechi *et al* (Nigeria, 2014) reported lower HR HPV prevalence in women who were HIV positive on antiretroviral drugs (OR=0.4; 95% CI: 0.3 to 0.5).

Whereas four studies found a significant epidemiological association between HAART and HR HPV genotypes, one did not. Rositch *et al*. (Uganda, 2013)²⁸ did not find any impact on monthly HPV DNA detection (PR 1.0; 95% CI 0.96 to 1.08), regardless of immune reconstitution or HIV viral suppression. Almost all test subjects (92%) had detectable HPV in the 6 months preceding HAART initiation and the cumulative prevalence remained high following initiation of therapy (90%) after a follow-up of 6 months (table 2).

The review revealed heterogeneity between studies in the use of the screening/diagnostic methods for cervical pathology (table 3). Nine studies used cytological changes with reporting based on the revised Bethesda classification, two studies used histopathological confirmation, and five studies used VIA/VILL screening methods of which three ascertained disease using colposcopy with biopsy solely in cases when lesions appeared suspicious for CIN.

Seven cross-sectional studies reported on the frequency of SIL and risk factors for SIL. Memiah *et al*. (Kenya, 2012)²⁹ observed that patients not receiving HAART treatment were 2.2 times more likely to have precancerous lesions than patients receiving HAART (adjusted OR=2.21, 95% CI 1.28 to 3.83). Similarly, Mogtomo *et al* (Cameroon, 2009)³⁰ reported that, of 22 high-grade SIL (HSIL)-positive women included in the study, 63.6% (14/22) were not on HAART, whereas 36.4% (8/22) were receiving HAART. The study revealed that total cervical abnormalities (HSIL and low-grade SIL (LSIL)) had a prevalence of 48.6% ($n=17$) among HAART users compared with 62.9% among women not receiving HAART ($p=0.034$).

In line with the association between the absence of HAART treatment and development of SIL, O. Ezechi *et al* (Nigeria, 2013) showed an increased risk of both SIL (adjusted OR=2.0; 95% CI 1.2 to 3.5) and HSIL (adjusted OR=2.6; 95% CI 1.0 to 6.7)³¹ among 1140 participants not on HAART, adjusted for age, marital

status, age at first intercourse and lifetime sexual partners. Nevertheless, Ononogbu *et al* (Nigeria, 2013) found only marginal or insignificant associations with different HAART regimens.³² Their study reported that 2501 women on tenofovir-containing HAART had a borderline significant RR of 1.4 (95% CI 1.0 to 2.0), and a RR of 1.5 (95% CI 1.0 to 2.2) on zidovudine-containing HAART, whereas women on stavudine-containing HAART showed a RR of 0.9 (95% CI 0.4 to 2.11) of having cervical disease compared with women not receiving HAART.²⁹

Hushko *et al* (Kenya, 2014)³³ observed that combined oral contraceptives remained significantly associated with detection of CIN 2+ in women on HAART (adjusted OR=1.84; 95% CI 1.20 to 2.82), compared with in women not HAART (adjusted OR=1.72; 95% CI 1.08 to 2.73), whereas the use of a progesterone implant was associated with increased detection of CIN 2+ (adjusted OR=9.43; 95% CI 2.85 to 31.20) only among women not on HAART. McKenzie *et al* (Kenya, 2011)³⁴, who explored risk factors for SIL development in women on HAART, reported no significant difference in the use of nevirapine-based regimens compared with efavirenz-based regimens between women with normal cytology and those with SIL (74% vs 72%; $p=0.80$). Similarly, there was no difference in the prevalence of nucleoside reverse transcriptase inhibitors backbones containing zidovudine compared with those containing stavudine between women with SIL and those without (12% vs 15%; $p=0.41$).

Bekolo *et al* (Cameroon, 2016)³⁵ also reported a decreased risk of cervical lesions among HAART receivers. After controlling for age and other covariates, women in the HAART group had a 67% reduction in the odds of cervical lesions compared with the community group (adjusted OR=0.33; 95% CI 0.15 to 0.73; $p=0.006$).

Mwakigonja *et al* (Tanzania, 2012)³⁶ identified immunodeficiency as the main determinant of the presence of CIN+ in HIV-infected women: CD4 count >350 cells/mm³ (OR=0.3; 95% CI 0.2 to 0.6) or ≥ 200 –350 cells/mm³ (OR=0.6; 95% CI 0.4 to 1.0) (Ref: <200 cells/mm³ CD4).

One study, Katumba *et al* (South Africa, 2010) reported that women who did not use HAART had more abnormal results compared with those who used HAART ($p<0.028$), although no confounders were adjusted for.

Two longitudinal studies reporting on progression/regression of cervical disease in women with HIV infection on HAART were identified. There was a lack of harmony in how regression/ progression was considered. A prospective observational study conducted in South Africa by Firnhaber *et al* (2010)³⁷ suggested that HAART was associated with a robust reduction in the incidence and progression of cervical disease (adjusted IRR=0.55; 95% CL 0.37 to 0.80). The study sensitivity analyses confirmed that the result was not dependent on the length of HAART exposure and this significant effect of HAART was only seen in women with CD4 counts of <350 cells/mm³.³⁴ ASC-US (atypical squamous cells of undetermined significance) results were classified as

**Table 2** Summary of studies exploring the epidemiological association between HAART and HR HPV

First author	Year of publication and setting	Study design and sample size	Main exposure(s) and outcome(s) of interest	Main results concerning HAART and remarks	Diagnosis	Confounding factors adjusted for
Ezechi OC	Nigeria 2014	Cross-sectional study among 231 women who were HIV positive and 305 HIV -negative	HIV infection and HAART on HR HPV prevalence and distribution	A multivariate logistic regression analysis showed a lower HR HPV prevalence in women who were HIV positive on antiretroviral drugs (OR=0.4; 95% CI 0.3 to 0.5)	HPV PCR	Age, type of community, marital status and lifetime sexual partners
De Vuyst H*	Kenya 2012	Cross-sectional study of 498 women who were HIV positive	HR HPV genotypes and CIN 2+ in HIV infected women	HAART users (≥ 2 year) had lower hr HPV prevalence (PR vs non-users=0.77, 95% CI 0.61 to 0.96) and multiple infections (PR=0.68, 95% CI 0.53 to 0.88)	HPV PCR	Age
Rositch A	Uganda 2013	Longitudinal study of HIV and HSV-2 coinfecting individuals (n=440)	Detection of HPV before and after initiation of HAART	No effect noticed of HAART on monthly HPV DNA detection (PR=1.0; 95% CI 0.96 to 1.08), regardless of immune reconstitution or HIV viral suppression	HPV PCR	CD4 counts and HIV viral load were analysed using both time-invariant pre-HAART and time-varying measurements based on the corresponding pre- and post-HAART period values
Zeier MD	South Africa 2015	HIV-infected women on HAART (n=204) prospective cohort study	Effect of the initiation of HAART for HPV genotype detection on cervical samples in HIV-infected South African women	HAART significantly reduced the risk for detection of HPV by 77% (OR=0.23, 95% CI 0.15 to 0.37)	HPV PCR	All models were adjusted for excision treatment of cervical, sexual activity and the CD4 cell count as time-dependent variables. Age was included as a non-time-dependent variable due to the relatively short follow-up time

Continued

Table 2 Continued

First author	Year of publication and setting	Study design and sample size	Main exposure(s) and outcome(s) of interest	Main results concerning HAART and remarks	Diagnosis	Confounding factors adjusted for
Kelly HA*	Burkina Faso and South Africa (2017)	Prospective cohort of 1238 women with HIV infection in Burkina Faso and South Africa	Effect of HAART on HR HPV in women with HIV-infection	HR HPV prevalence was higher among those on short duration HAART in both countries. However, when adjusted for CD4 cell count, this association was observed in Burkina Faso only (65.1 vs 52.1% for <2 years compared with >2 years (adjusted PR=1.24, 95% CI 1.04 to 1.47)	HPV PCR	The site and sociodemographic and behavioural factors that were independently associated in univariate analyses ($p<0.10$) were adjusted for

*Studies with more than one outcome of interest.

CIN, cervical intraepithelial neoplasia; HAART, highly active antiretroviral therapy; HPV, human papillomavirus; HR, high risk; HSV, herpes simplex virus; PR, prevalence ratio.

LSIL results, while ASC-H results were classified as HSIL results.

A smaller magnitude of effect was found by Omar *et al* (South Africa, 2011)³⁸ (HR=0.72; 95% CL 0.52 to 0.99) using time-varying CD4 counts as a covariate in the multivariate model of progression. Progression was defined either as a subsequent smear with a cytological diagnosis of ASCH, HSIL or worse in women who had a previous normal or LSIL smear with an interval between smears of >5.5 months.

One study was identified on HAART and regression of cervical disease. A large prospective cohort study from Soweto, South Africa undertaken by Adler *et al* (2012)³⁹ reported that HAART positively affected the natural history of cervical disease in women with HIV infection and a multivariate marginal models analysis identified a significantly increased likelihood (OR=2.61; 95% CI 1.75 to 3.89; $p<0.0001$) of regression of cervical lesions among women on HAART. Regression was defined as a subsequent improvement in cytological results from normal, ASC-US, LSIL, ASC-H, HSIL and cancer.

Other studies reported specifically on the duration of HAART and SIL. Kelly *et al* (South Africa, 2017)²⁷ reported that CIN 2+ prevalence was higher among short-duration antiretroviral therapy users <2 years (adjusted OR=1.99; 95% CI 1.12 to 3.54) and naive participants (adjusted OR=1.87; 95% CI 1.11 to 3.17) in South Africa. De Vuyst *et al* (Kenya, 2012)²⁵ reported that the prevalence of CIN 2 and CIN 3 did not vary across HAART non-users, HAART users for <2 years or ≥ 2 years (p for linear trend=0.416). However, when stratified for CD4 count categories, the prevalence of CIN 2/3 among HAART non-users linearly increased per each decrease of CD4 count category, CD4 <200; CD4 200–349; CD4 350–499; and CD4 ≥ 500 (p for linear trend=0.013) but this is not seen among women receiving HAART for <2 years and ≥ 2 years (p for linear trend=0.9 and 0.5, respectively). Still, CD4 nadir before initiating HAART was unknown and HAART may have been started at a CD4 count too low to prevent or reverse CIN 2/3. Similarly, Huchko *et al* (Kenya, 2014)³⁰ detected a non-significant association between the duration of HAART and CIN 2+ (adjusted OR=0.98; 95% CI 0.95 to 1.00), although it was not adjusted for CD4 count. McKenzie (Kenya, 2011)³¹ did not find any significant correlation between the duration and type of HAART and SIL and the median number of months on HAART did not differ between women with normal cytology and those with SIL (14 vs 11 months; $p=0.17$).

Nevertheless, Mogtomo *et al* (Cameroon, 2009)³⁰ observed that while LSIL incidence in women with HIV infection under HAART therapy decreased from 8.6% (three cases) to 5.7% (two cases) the first 10 months of HAART use, it increased to 11.4% (four cases) after 10 months. In contrast, HSIL cases decreased from 14.3% (n=5) to 5.7% to 2.8% (n=1) after 10 months.

Capo Chichi *et al* (2016)⁴⁰ in Benin reported that cervical dysplasia was observed in 4/86 (5%) of women living with HIV on HAART, in whom after 2 years no new

cases of cervical dysplasia were reported, nor were the HPV genotypes present at baseline there after 2 years.

A cross-sectional study from Ethiopia (Gedefaw *et al* 2013) showed that the presence of precancerous lesions was inversely associated with HAART: patients under HAART at the moment of the study (not specifying the duration of the treatment) were 48% less likely to have precancerous cervical cancer lesions than those who were not on HAART (adjusted OR=0.52, 95% CI 0.35 to 0.92; $p=0.015$).⁴¹

One prospective study assessed CIN 2+ incidence. Kelly *et al* (South Africa, 2017)²⁷ reported that CIN 2+ incidence was reduced among women on HAART (adjusted OR=0.39; 95% CI 0.15 to 1.01).

Four studies contrasted in their reported results concerning the protective effect of higher CD4+ count nadir, prior to HAART initiation, on the onset of cervical disease. Huchko *et al* (Kenya, 2014)³⁰ reported that CD4+ nadir over 500 cells/mm³ was associated with a reduced detection of CIN 2+ (adjusted OR=0.61, 95% CI 0.38 to 0.97) in the overall group, but current CD4+ was only associated with reduced detection of CIN 2+ among HAART non-users (adjusted OR=0.42; 95% CI 0.22 to 0.80).⁴² This finding is compatible with the immune driven trend between CD4+ and a positive cervical screening among women with HIV infection observed by Ononogbu *et al* (Nigeria, 2013),⁴³ which reported a decreasing risk with increasing CD4 count: RR=0.5 (95% CI 0.3 to 0.7) for CD4 count 300 to <450; RR=0.5 (95% CI 0.3 to 0.7) for 450 to <650; and RR=0.3 (95% CI 0.2 to 0.6) for ≥ 650 .

A prospective observational study of 601 HIV-seropositive women conducted by Firnhaber *et al* (South Africa, 2012) suggested that the effect of baseline HAART exposure was strongly modified by baseline CD4 count when initiating HAART in women with CD4 counts <350 cells/mm³. A RR=0.47 (95% CI 0.30 to 0.73) was found in women initiating HAART with CD4 counts <350 cells/mm³ compared with RR=0.84 (95% CI 0.42 to 1.72) in women with baseline CD4 counts ≥ 350 cells/mm³.

Only one study reported baseline CD4 count not to be significant. Jacquet *et al* (Ivory Coast, 2014)⁴⁴ reported that the baseline CD4 count at the time of first clinical follow-up, which could be interpreted as a proxy for nadir CD4 count, was not found to be significantly associated with CIN+ in the multivariate model (table 3).

Epidemiological association between HAART and ICC

Only two studies were found that assessed the effect of HAART on cervical cancer (table 3). An age-adjusted population-level evaluation of the effect of antiretroviral therapy on cancer incidence in Kyadondo County, Uganda (1999–2008), revealed that the increase in population-level HAART coverage over time did not affect the incidence of invasive cervical cancer (IRR=1.02; 95% CI 0.98 to 1.05).⁴⁵ However, Bogaert *et al* (South Africa, 2013) reported that patients on HAART had not only less CIN 1 ($p=0.018$) and CIN 2 ($p=0.18$) but also less ICC ($p=0.019$) than their untreated counterparts, despite similar mean CD4 count.

DISCUSSION

The findings of our systematic review suggest a positive impact of HAART duration, in conjunction and interaction with CD4 count, on reducing the prevalence of HR HPV. Our review also illustrates how untreated people living with HIV with low CD4 count may be at increased risk of being or remaining infected with HR HPV compared with those treated with HAART. Moreover, our findings support the hypothesis that the greatest treatment effect might be seen among women starting at the lowest CD4 count, since higher CD4 count might be associated with lower incidence of CIN 2+, thereby reducing the possible treatment effect of HAART.

The literature identified in this systematic review suggests that CD4 count may have a more instrumental role in cervical oncogenesis or the integration of the latent reservoir throughout the body than either HAART use or the treatment duration on the prevalence of CIN 2 and CIN 3.

Surprisingly, we were unable to retrieve any studies that assess the impact of HAART on cervical disease and HPV infection across different types of HIV infections. It therefore remains unknown whether the effect of HAART differs between patients with HIV-1, HIV-2 or dual infection. HIV-2 infection and dual infection are highly prevalent in sub-Saharan Africa, predominantly in Western Africa in countries like Guinea Bissau, Ivory Coast, Mali and Burkina Faso.⁴⁶

The sample size of the studies included in this review greatly varies, rendering it difficult to assess and compare strength of associations. Due to various difficulties, such as financial and ethical considerations linked to the conduct of RCTs, evidence is mostly based on retrospective or cross-sectional studies, which do not fulfil the causal criterion of temporality. Heterogeneity in the sensitivity or specificity of screening methods may also render results less comparable. Cervical cytology, which has a clinical sensitivity between 55% and 65% for detection of histopathologically confirmed 'true disease status'⁴⁷ along with low interobserver or intraobserver true correlation can result in even one grade of misclassification introducing a bias in studies assessing regression or incidence of SIL in women on HAART. Moreover, a recent study in Kenya has reported that the sensitivity of VIA is affected in women on HAART and by CD4 count,⁴⁸ which in studies ascertaining disease using colposcopy with biopsy solely in cases when lesions appeared suspicious for CIN 2+, may have led to under-ascertainment of true disease: sensitivity 89.9% (95% CI 82.2% to 95.0%) in women under 35 and 78.6% (95% CI 63.2% to 89.7%) in women above 35.⁴⁷

Our findings derived from the sub-Saharan context contrast with the conclusion of a systematic review of studies from industrialised settings suggesting a largely inconsistent impact of HAART on reducing the incidence and progression and facilitating the regression of HPV infection and cervical abnormalities.⁴⁹ This discrepancy may be attributed to the greater access in industrialised

Table 3 Summary of studies exploring the association between HAART and SIL/CIN and ICC

First author	Year of publication & setting	Study design and sample size	Main exposure(s) and outcome(s) of interest	Main results concerning HAART and Remarks	Diagnosis	Confounding factors adjusted for
Mogtomo M	Cameroon 2009	A cross-sectional design of 70 women with HIV infection	Cervical abnormalities in women on HAART and without HAART	Among the 22 HSIL-positive women, 63.6% (14/22) were not on HAART, while 36.4% (8/22) were on HAART. Women with HIV infection on HAART with positive HSIL showed a median CD4+ count of 253.7±31.7 higher than those without therapy (164.7±26.1)	Pap smears were interpreted and classified according to the Bethesda system	N/A (univariate analysis)
Ononogbu U	Nigeria 2013	Cross-sectional study of 2501 women who were HIV positive	Explore risk factors leading to VIA/VILI positivity	HAART and positive cervical screening: TDF-containing HAART OR 1.4 (95% CI 1.0–2.0); DDV-containing HAART OR 1.5 (95% CI 1.0–2.2); d4T-containing HAART OR 0.9 (95% CI 0.4–2.1)	Results of VIA or VILI were classified according to the IARC manual and recorded after each test	N/A (univariate analysis)
Ezechi O C	Nigeria 2014	Cross-sectional study among 1140 women (531 HIV positive; 609 HIV negative)	Explore the association among HIV, HAART use and SIL	Not using HAART was found to be associated with an increased risk of SIL (aOR 2.1; 95% CI 1.4 to 3.5) and HSIL (aOR:2.6; 95% CI 1.1 to 6.4)	Pap smears were interpreted and classified according to the Bethesda system. 31.4% of slides were reviewed for cytology quality control	Age, viral load, marital status, CD4
McKenzie K P	Kenya 2011	Cross-sectional study of 267 women with HIV infection	Risk factors for SIL	SIL is prevalent among women on HAART and is associated with immunosuppression. Duration (14 vs 11 months; p=0.17) and type (NVP based (74%) vs EFV based (72%); p=0.8) of antiretroviral regimen were not significantly associated with SIL	Pap smears were interpreted and classified according to the Bethesda system	N/A (no OR calculated)
Huscko M	Kenya 2014	Cross-sectional design. 3241 women with HIV infection were screened for cervical cancer	Explore risk factors for CIN 2+	Combined oral contraceptives were associated with detection of CIN 2+ among women on HAART (aOR 1.84, 95% CI 1.20 to 2.82), and not on HAART (aOR 1.72, 95% CI 1.08 to 2.73), use of a progesterone implant was associated with increased detection of CIN 2+ (aOR 9.43, 95% CI 2.85 to 31.20) only among women not on HAART	Those with a positive VIA underwent immediate colposcopy with biopsy for confirmation of any lesions suspicious for CIN 2+	Age adjusted
De Vuyst H	Kenya 2012	Cross-sectional study of 498 HIV-positive women	Explore risk factors for CIN 2+ in women with HIV infection	The prevalence of CIN 2 and CIN 3 did not vary across HAART non-users, HAART users for <2 years or ≥2 years (p for linear trend=0.416). However, when stratified for CD4 count categories, the prevalence of CIN 2 or 3 among HAART non-users linearly increased per each decrease in CD4 count category, CD4<200; CD4 200–349; CD4 350–499; and CD4≥500 (p for linear trend=0.013) but this is not seen among women receiving HAART for <2 years and ≥2 years (p for linear trend=0.9 and 0.5, respectively)	Histologically confirmed cytology	Age adjusted

Continued

Table 3 Continued

First author	Year of publication & setting	Study design and sample size	Main exposure(s) and outcome(s) of interest	Main results concerning HAART and Remarks	Diagnosis	Confounding factors adjusted for
Gedefaw A	Ethiopia 2013	Cross-sectional study (hospital-based) among 448 women with HIV infection	Risk factors for precancerous cervical cancer lesion among women with HIV infection	Being currently on HAART had a protective effect for precancerous lesions (aOR=0.52, 95% CI 0.35 to 0.92)	Results of VIA were classified as negative, positive, or suspicious for ICC	Adjusted for age, education, occupation, parity, lifetime history of pelvic infection, STD, ulcerative genital lesion, age at first marriage, age at first sexual intercourse, and lifetime number of sexual partners
Firnhaber C	South Africa 2012	Prospective cohort study of 601 HIV-seropositive women	Incidence and progression of cervical lesions among women who are HIV positive	HAART use was associated with reduction of incidence and progression of cervical lesions, adjusted IRR=0.55 (95% CI 0.37 to 0.80)	Pap smears were interpreted and classified according to the Bethesda system	Adjusted for age, CD4 count, age at first intercourse, lifetime number of sexual partners, history of sexual transmitted diseases, hormonal contraception, condom use, employment, smoking and education
Memiah P	Kenya 2012	Cross-sectional design, 715 women with HIV infection	Prevalence and risk factors associated with precancerous cervical cancer lesions among women with HIV infection	Non-HAART users were 2.21 times more likely to have precancerous lesions than HAART patients (95% CI 1.28 to 3.83)	VILI was used as the screening technique. A positive VILI test necessitated a cervical biopsy	Not adjusted
Omar T	South Africa 2012	Longitudinal study 2325 women with HIV infection	Progression and regression of cervical dysplasia in women with HIV infection in Soweto	HAART reduced the risk of SIL progression (aHR 0.72; 95% CI 0.52 to 0.99)	Conventional Pap smears are performed	CD4, baseline weight, smoking exposure, age, baseline smear results
Adler DH	South Africa 2012	HIV-infected women (n=1123) from Soweto; prospective cohort study	Risks of progression, regression, and incidence of HPV-related cervical lesions between those on HAART and those not on HAART	Increased likelihood (aOR 2.61; 95% CI 1.75 to 3.89; p<0.0001) of regression of cervical lesions among women on HAART	Pap smear were interpreted and classified according to the Bethesda system and verified by a second reader	BMI, number of sexual partners, number of STIs

Continued

Table 3 Continued

First author	Year of publication & setting	Study design and sample size	Main exposure(s) and outcome(s) of interest	Main results concerning HAART and Remarks	Diagnosis	Confounding factors adjusted for
Bekolo CP	Cameroon 2016	Cross-sectional study of 302 women of whom 131 (43.4%) had HIV infection and were receiving HAART	Risks for cervical disease in HAART-experienced women compared with those in women in the general population of Cameroon	After controlling for age and other covariates, women in the HAART group had a 67% reduction in the odds of cervical lesions compared with the community group (aOR=0.33, 95% CI 0.15 to 0.73, p=0.006). The authors could not, however, account for differences in methods and quality assurance in the ascertainment of cervical lesions as well as other risk factors for cervical cancer that must have changed over time	Each woman was screened using three different methods in series: visual inspection VIA, VILI and conventional Pap smear cytology	Age, place of residence, occupation, religion, education, marital status, smoking, and family history of cancer
Capo Chichi CV	Benin 2016	86 female cases and 86 female controls	Association of HR HPV to cervical dysplasia among women under stringent HAART treatment and in controls without HIV	Cervical dysplasia was observed in 4/86 (5%) of women living with HIV on HAART. Among the HAART group, no new cases of cervical dysplasia were reported after 2 years	VILI and video colposcopy	No adjustment for potential confounders
van Bogaert LJ	South Africa 2013	Case control study of 1023 women with HIV infection and 1023 uninfected women	Influence of the CD4 cell count and of HAART on the relative distribution of cervical pathology	Patients on HAART had less CIN 1 (p=0.018), 2 (p=0.18)	Histologically confirmed cytology	No adjustment for potential confounders
Mwakigonja AM	Tanzania 2012	Prospective unmatched, case-control study of 120 HIV-seropositive cases on HAART and 50 seronegative controls	Frequency of cervical cancer and precancerous lesions in the general population compared with the HIV-infected populations in Tanzania	The cytological findings in this study suggest that the frequency of SIL and carcinoma appeared to be higher among women with HIV infection on HAART compared with seronegative controls and as expected increased with age. SIL and carcinoma constituted 28.3% (34/120), 38.3% (46/120) and 5.8% (7/120) among cases, and 28% (14/50), 34% (17/50) and 2% (1/50) for controls respectively (p=0.6)	Pap smears were interpreted and classified according to the revised Bethesda system	Not adjusted
Jacquet A	Ivory Coast 2014	Cross-sectional study of 2998 women with HIV infection, of which 76% were on HAART	Preventable determinants of SIL in women who were HIV positive	Non-significant (p=0.26) decrease of CIN with increased exposure to HAART	Positively screened women by VIA and VILI were scheduled for colposcopy. Directed biopsies were performed in case of positive findings with colposcopy	Stepwise descending procedure for adjustment of confounders in the multivariate analysis. Variables adjusted for not mentioned
Katumba AC	South Africa 2016	A cross-sectional descriptive study of 390 women who were HIV positive	Correlation between HAART and abnormal Pap smear results of women with HIV infection	Participants who did not use HAART had more abnormal results compared with those who used HAART (p<0.028, 95% CI 0.28 to 0.93)	Pap smears were interpreted and classified according to the Bethesda system	No adjustment for potential confounders

Continued



Table 3 Continued

First author	Year of publication & setting	Study design and sample size	Main exposure(s) and outcome(s) of interest	Main results concerning HAART and Remarks	Diagnosis	Confounding factors adjusted for
De Vuyst H*	Kenya 2012	Cross-sectional study of 498 women who were HIV positive	CIN 2+ in women with HIV infection	The prevalence of CIN 2 and CIN 3e did not vary across HAART non-users, HAART users for < 2 years or ≥2 years (p for linear trend=0.416). However, the prevalence of CIN 2/3 among HAART non-users linearly increased per each decrease in CD4 count category, CD4<200; CD4 200–349; CD4 350–499; and CD4≥500 (p for linear trend=0.013) but this is not seen among women receiving HAART for <2 years and ≥2 years (p for linear trend=0.9 and 0.5, respectively)	Histologically confirmed cytology	Age adjusted
Kelly HA*	Burkina Faso and South Africa 2017	Prospective cohort of 1238 women with HIV infection in Burkina Faso and South Africa	Effect of HAART on CIN 2+ in women with HIV infection	CIN 2+ prevalence was higher among short-duration HAART users (aOR= 1.99, 95% CI 1.12 to 3.54) and HAART-naïve participants (aOR 1.87, 95% CI 1.11 to 3.17) in South Africa	Biopsies were analysed by histology	Stratification by site, HAART use and duration (or >2 years), HIV-1 viral suppression (≤1000 copies/ml) and CD4 cell counts
Mutyaba I	Uganda 2015	Ecological population based study	Correlation between % of HAART coverage and incidence of AIDS-related malignancies	HAART coverage was not associated with incidence of invasive cervical cancer	Unknown	Age adjusted
Van Bogaert LJ *	South Africa 2013	Case control study of 1023 women with HIV infection and 1023 uninfected women	Relative distribution of ICC among women with HIV infection (treated or not) and uninfected women	There was a significantly higher proportion of CIN 1 (p=0.012) and 2 (p=0.01) but a lower proportion of ICC (p=0.015) among women with HIV infection. Patients on HAART had less CIN 1 (p=0.018), 2 (p=0.18) and ICC (p=0.019) than their untreated counterparts	Histologically confirmed cytology	No adjustment for potential confounders

*Studies with more than one outcome of interest.

AOR, adjusted OR; CIN, cervical intraepithelial neoplasia; EFV, efavirenz; HAART, highly active antiretroviral therapy; HPV, human papillomavirus; HR, high risk; HSIL, high-grade SIL; IARC, International Agency for Research on Cancer; ICC invasive cervical cancer; NVP, nevirapine; PR, prevalence ratio; SIL, squamous intraepithelial lesion; STD, sexually transmitted disease; STI, sexually transmitted infection; TDF, tenofovir; VIA, visual inspection with acetic acid; VILI, visual inspection with Lugol's iodine.



settings to frequent cytological screening, thereby making the potential benefits of HAART less apparent.

One of the few retrieved prospective studies exploring the impact of HAART on SIL incidence and progression only reported a significant association in women with CD4 counts <350 cells/mm³. Apart from the inclusion criteria providing heterogeneity in baseline profiles of women, different potential confounders have been adjusted for in studies addressing the same outcome. Not all studies control for hormonal/barrier contraception use, HIV viral load, CD4 count baseline or the number of sexual partners, therefore residual confounding cannot be excluded. In addition, the duration of follow-up may be too short and not always detailed in articles.

Clinical and public health implications

The lack of statistically significant associations observed in this review between duration of HAART intake and reduction of CIN 2+ in women initiating HAART at an earlier CD4 count initiation suggests the importance of continued follow-up with cervical cytological and/or histological screening. Even after HAART initiation and immune reconstitution, lesions remain prevalent and progression of disease should be studied further to allow best timing of a screening schedule.

Although current HAART guidelines in industrialised countries promote biannual cervical cancer screening in the first year following HIV diagnosis and annually thereafter, in the absence of evidence-based knowledge in sub-Saharan African women, it may be advisable to address the following research gaps before suggesting the relevance of the guideline in this population. In the mean time, a cautionary approach may be employed and biannual cervical screening continued, if feasible.

Research gaps

The dearth of studies on duration and initiation of HAART at an earlier CD4 count and the consensus on findings preclude policy makers from developing local guidelines. Moreover, due to the changes in recommendations for HAART start, women in sub-Saharan Africa at the time of the studies were more likely to have low CD4 counts, whereas in the future they will be initiating HAART in a far less immunosuppressed state.

Future studies should preferably be prospective in design. This will allow a better follow-up of patients at different points in time and fulfil the criterion of temporality for causality. Moreover, including nested case-control studies within cohort follow-up will allow robust intermediary evidence for such research questions. Studies should also elucidate the association between cervical HIV DNA and HPV incidence/prevalence, or cervical dysplasia.

The exploration of an association between HAART and cervical cancer warrants large cancer epidemiological studies. Previously, the lack of effect of HAART on the prevalence on cervical cancer can be ascribed to the high mortality associated with AIDS. Despite improved longevity, and in settings with a life expectancy that exceeds the average age of cervical cancer onset, one study to date

from our review suggested that, thanks to extensive HAART programmes, decrease in cervical cancer incidence can be expected.⁵⁰

Finally, women with HIV-HPV coinfections may be at risk for immune-modulating helminthic, tuberculosis and malarial infections⁵¹, which may compromise the positive impact that HAART may have on clearing cervical lesions. Their synergistic interactions in the post-HAART era should be elucidated.

CONCLUSION

Access to HAART in sub-Saharan Africa has dramatically increased over the past decade, improving life expectancy for women living with HIV. This in turn requires urgent clarification of the impact of HAART on the development of cervical precancerous states and cancer.

Whilst the preponderance of studies suggests that women on HAART are less likely to have SIL, for ICC, evidence remains scarce and inconclusive. However, our review imputes a more significant role of CD4 count or the CD4 count at which HAART is initiated, the timing of HAART in relation to immunosuppression on HR HPV genotype and CIN 2/3 prevalence, than to HAART use, irrespective of its duration. The scarce evidence shows that the positive impact of HAART is only felt in women with CD4 counts of <350 cells/mm³.³

Given the current limited evidence available to assist policy makers in developing local guidelines and prevention and treatment strategies for cervical cancer among women with HIV infection on HAART, a tailored screening protocol cannot be established. Longer term surveillance data on women who are HIV positive at different levels of immunosuppression is crucial to determine the effect of HAART on CIN 2+.

Author affiliations

¹International Centre for Reproductive Health (ICRH), Ghent University, Ghent, Belgium

²CDC Foundation, Atlanta, Georgia, USA

³Laboratory for Cell Biology & Histology, University of Antwerp, Antwerp, Belgium

⁴Department of Clinical Pharmacy, Saints Cyril and Methodius, Republic of Macedonia

⁵AMBIOR, Laboratory for Cell Biology & Histology, University of Antwerp, National Reference Centre for HPV, Laboratory of Molecular Pathology, Antwerp, Belgium

⁶Department of Internal Medicine & Infectious diseases, University Hospital, Ghent, Belgium

Contributors MS and RR contributed equally to the study: conception of the study, coordination of the study, screening of studies, interpretation of the findings of included studies, writing and editing the paper, validation of the final version. ZN: interpretation of the findings, writing and editing the paper, validation of the final version. KM: editing the paper, validation of the final version. DS: editing the paper, validation of the final version. BD: editing and validation of the final version. CS: conception, interpretation and validation of the final version.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially,

and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

1. WHO. Fact sheet: human papilloma and cervical cancer. 2015 <http://www.who.int/mediacentre/factsheets/fs380/en/> (accessed 17 May 2016).
2. Walboomers JM, Jacobs MV, Manos MM, *et al*. Human papillomavirus is a necessary cause of invasive cervical Cancer worldwide. *J Pathol* 1999;189:12–19.
3. Rabkin CS, Biggar RJ, Baptiste MS, *et al*. Cancer incidence trends in women at high risk of human immunodeficiency virus (HIV) infection. *Int J Cancer* 1993;55:208–12.
4. Sun XW, Kuhn L, Ellerbrock TV, *et al*. Human papillomavirus infection in women infected with the human immunodeficiency virus. *N Engl J Med* 1997;337:1343–9.
5. Ellerbrock TV, Chiasson MA, Bush TJ, *et al*. Incidence of cervical squamous intraepithelial lesions in HIV-infected women. *JAMA* 2000;283:1031–7.
6. Rowhani-Rahbar A, Hawes SE, Sow PS, *et al*. Dembele B critchlow CW, N'doye I, Kiviat NB. the impact of HIV status and type on the clearance of human papillomavirus infection among Senegalese women. *J Infect Dis* 2007;196:887–94.
7. World Health Organization. Antiretroviral therapy for HIV infection in adults and adolescents: recommendations for a public health approach—2010 revision. http://whqlibdoc.who.int/publications/2010/9789241599764_eng.pdf (accessed 15 May 2016).
8. Adult combination antiretroviral therapy in sub-Saharan Africa: lessons from Botswana and future challenges, C William Wester, Hermann Bussmann, John Koethe, Claire Moffat, Sten Vermund, Max Essex, Richard G Marlink. *HIV Ther* 2009.
9. Pipkin S, Scheer S, Okeigwe I, *et al*. The effect of HAART and calendar period on Kaposi's sarcoma and non-Hodgkin lymphoma: results of a match between the AIDS and cancer registry. *AIDS* 2011;25:463–71.
10. Ledergerber B, Telenti A, Egger M. Risk of HIV related Kaposi's sarcoma and non-Hodgkin's lymphoma with potent antiretroviral therapy: prospective cohort study. Swiss HIV Cohort Study. *BMJ* 1999;319:23–4.
11. Jacobson LP, Yamashita TE, Detels R, *et al*. Impact of potent antiretroviral therapy on the incidence of kaposi's sarcoma and non-Hodgkin's lymphomas among HIV-1-infected individuals. Multicenter AIDS Cohort Study. *J Acquir Immune Defic Syndr* 1999;21:S34–S41.
12. International Collaboration on HIV and Cancer. Highly active antiretroviral therapy and incidence of Cancer in human immunodeficiency virus-infected adults. *J Natl Cancer Inst* 2000;92:1823–30.
13. Palefsky JM. Cervical human papillomavirus infection and cervical intraepithelial neoplasia in women positive for human immunodeficiency virus in the era of highly active antiretroviral therapy. *Curr Opin Oncol* 2003;15:382–8.
14. Heard I, Palefsky JM, Kazatchkine MD. The impact of HIV antiviral therapy on human papillomavirus (HPV) infections and HPV-related diseases. *Antivir Ther* 2004;9:13–22.
15. Overview HIV/STI. <http://www.afro.who.int/en/clusters-a-programmes/dpc/acquired-immune-deficiency-syndrome/overview.html> (accessed 15 May 2016).
16. Adler DH, Kakinami L, Modisenyane T, *et al*. Increased regression and decreased incidence of human papillomavirus-related cervical lesions among HIV-infected women on HAART. *AIDS* 2012;26:1645–52.
17. WHO. WHO issues new HIV recommendations calling for earlier treatment. 2013. (accessed 1 Apr 2016).
18. WHO. Guidelines on when to start antiretroviral therapy and on pre exposure prophylaxis for HIV. 2015. http://apps.who.int/iris/bitstream/10665/186275/1/9789241509565_eng.pdf (accessed 17 May 2016).
19. Palefsky JM. Cervical human papillomavirus infection and cervical intraepithelial neoplasia in women positive for human immunodeficiency virus in the era of highly active antiretroviral therapy. *Curr Opin Oncol* 2003;15:382–8.
20. Heard I, Potard V, Costagliola D. Limited impact of immunosuppression and HAART on the incidence of cervical squamous intraepithelial lesions in HIV-positive women. *Antivir Ther* 2006;11:1091–6.
21. Adler DH, Kakinami L, Modisenyane T, *et al*. Increased regression and decreased incidence of human papillomavirus-related cervical lesions among HIV-infected women on HAART. *AIDS* 2012;26:1645–52.
22. PRISMA statement. <http://www.prisma-statement.org/statement.htm> (accessed 2 Jan 2016).
23. BMJ CLinical evidence what is Grade. <http://clinicalevidence.bmj.com/x/set/static/ebm/learn/665072.html> (accessed 21 Jan 2017).
24. Altman DG. Statistics in medical journals: some recent trends. *Stat Med* 2000;19:3275–89.
25. De Vuyst H, Mugo NR, Chung MH, *et al*. Prevalence and determinants of human papillomavirus infection and cervical lesions in HIV-positive women in Kenya. *Br J Cancer* 2012;107:1624–30.
26. Zeier MD, Botha MH, Engelbrecht S, *et al*. Combination antiretroviral therapy reduces the detection risk of cervical human papilloma virus infection in women living with HIV. *AIDS* 2015;29:59–66.
27. Kelly HA, Sawadogo B, Chikandiwa A, *et al*. Epidemiology of high-risk human papillomavirus and cervical lesions in African women living with HIV/AIDS: effect of anti-retroviral therapy. *AIDS* 2017;31:273–85.
28. Rositch AF, Gravitt PE, Tobian AA, *et al*. Frequent detection of HPV before and after initiation of antiretroviral therapy among HIV/HSV-2 co-infected women in Uganda. *PLoS One* 2013;8:e55383.
29. Memiah P, Muthia W, Kiiru G, *et al*. Prevalence and risk factors associated with precancerous cervical Cancer lesions among HIV-Infected women in Resource-Limited Settings. *AIDS Res Treat* 2012;2012:1–7.
30. Mogtomo ML, Malieugoue LC, Djiepgang C, *et al*. Incidence of cervical disease associated to HPV in human immunodeficiency infected women under highly active antiretroviral therapy. *Infect Agent Cancer* 2009;4:9.
31. Ezechi OC, Ostergren PO, Nwaokorie FO, *et al*. The burden, distribution and risk factors for cervical oncogenic human papilloma virus infection in HIV positive Nigerian women. *Viro J* 2014;11:5.
32. Ononogbu U, Almutaba M, Modibbo F, *et al*. Cervical Cancer risk factors among HIV-infected Nigerian women. *BMC Public Health* 2013;13:582.
33. Huchko MJ, Leslie H, Sneden J, *et al*. Risk factors for cervical precancer detection among previously unscreened HIV-infected women in western Kenya. *Int J Cancer* 2014;134:740–5.
34. McKenzie KP, Rogers RK, Njoroge JW, *et al*. Cervical squamous intraepithelial lesions among HIV-positive women on antiretroviral therapy in Kenya. *Curr HIV Res* 2011;9:180–5.
35. Bekolo CE, O'Bryan G, Tchago FE, *et al*. Integrating cervical Cancer screening with HIV care in Cameroon: comparative risk analysis of cervical disease in HIV-infected women receiving antiretroviral therapy to women in the General Population. *PLoS One* 2016;11:e0149152.
36. Mwakigonja AR, Torres LM, Mwakiyoma HA, *et al*. Cervical cytological changes in HIV-infected patients attending care and treatment clinic at Muhimbili National Hospital, Dar es Salaam, Tanzania. *Infect Agent Cancer* 2012;7:3.
37. Firnhaber C, Westreich D, Schulze D, *et al*. Highly active antiretroviral therapy and cervical dysplasia in HIV-positive women in South Africa. *J Int AIDS Soc* 2012;15:17382.
38. Omar T, Schwartz S, Hanrahan C, *et al*. Progression and regression of premalignant cervical lesions in HIV-infected women from Soweto: a prospective cohort. *AIDS* 2011;25:87–94.
39. Adler DH, Kakinami L, Modisenyane T, *et al*. Increased regression and decreased incidence of human papillomavirus-related cervical lesions among HIV-infected women on HAART. *AIDS* 2012;26:1645–52.
40. Capo-Chichi CD, Aguida B, Chabi NW, *et al*. Diversity of high risk human papilloma viruses in women treated with antiretroviral and in healthy controls and discordance with cervical dysplasia in the South of Benin. *Infect Agent Cancer* 2016;11.
41. Gedefaw A, Astatkie A, Tessema GA. The prevalence of precancerous cervical Cancer lesion among HIV-infected women in southern Ethiopia: a cross-sectional study. *PLoS One* 2013;8:e84519.
42. Huchko MJ, Leslie H, Sneden J, *et al*. Risk factors for cervical precancer detection among previously unscreened HIV-infected women in western Kenya. *Int J Cancer* 2014;134:740–5.
43. Ononogbu U, Almutaba M, Modibbo F, *et al*. Cervical Cancer risk factors among HIV-infected nigerian women. *BMC Public Health* 2013;13:582.



- 44 Jaquet A, Horo A, Ekouevi DK, *et al.* Risk factors for cervical intraepithelial neoplasia in HIV-infected women on antiretroviral treatment in Côte D'ivoire, West Africa. *PLoS One* 2014;9:e90625.
- 45 Mutyaba I, Phipps W, Krantz EM, *et al.* A Population-Level evaluation of the effect of antiretroviral therapy on Cancer incidence in Kyadondo County, Uganda, 1999-2008. *J Acquir Immune Defic Syndr* 2015;69:481–6.
- 46 Balestre E, Ekouevi DK, Tchounga B, *et al.* Immunologic response in treatment-naïve HIV-2-infected patients: the leDEA West Africa cohort. *J Int AIDS Soc* 2016;19:20044.
- 47 Nanda K, McCrory DC, Myers ER, *et al.* Accuracy of the Papanicolaou test in screening for and follow-up of cervical cytologic abnormalities: a systematic review. *Ann Intern Med* 2000;132:810–9.
- 48 Huchko MJ, Sneden J, Sawaya G, *et al.* Accuracy of visual inspection with acetic acid to detect cervical Cancer precursors among HIV-infected women in Kenya. *Int J Cancer* 2015;136:392–8.
- 49 Bratcher LF, Sahasrabudde VV. The impact of antiretroviral therapy on HPV and cervical intraepithelial neoplasia: current evidence and directions for future research. *Infect Agent Cancer* 2010;5:8.
- 50 Hawes SE, Critchlow CW, Faye Niang MA, *et al.* Increased risk of high-grade cervical squamous intraepithelial lesions and invasive cervical cancer among African women with human immunodeficiency virus type 1 and 2 infections. *J Infect Dis* 2003;188:555–63.
- 51 Anastos K, Hoover DR, Burk RD, *et al.* Risk factors for cervical precancer and cancer in HIV-infected, HPV-positive Rwandan women. *PLoS One* 2010;5:e13525.

BMJ Open

Associations between highly active antiretroviral therapy and the presence of HPV, premalignant and malignant cervical lesions in sub-Saharan Africa, a systematic review: current evidence and directions for future research

Sonia Menon, Rodolfo Rossi, Natasha Zdraveska, Mbabazi Kariisa, Sushama D Acharya, Davy Vanden Broeck and Steven Callens

BMJ Open 2017 7:

doi: 10.1136/bmjopen-2016-015123

Updated information and services can be found at:

<http://bmjopen.bmj.com/content/7/8/e015123>

These include:

References

This article cites 42 articles, 1 of which you can access for free at: <http://bmjopen.bmj.com/content/7/8/e015123#ref-list-1>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Epidemiology](#) (2247)
[Infectious diseases](#) (595)
[Oncology](#) (459)
[Sexual health](#) (158)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>