

Aniline-Hydroxylation Activity of a Flavin-linked α -Type Polypeptide Packing an Iron Porphyrin

著者	Tomizaki Kin-ya, Nishino Hidekazu, Arai Toru, Kato Tamaki, Nishino Norikazu
journal or publication title	Chemistry Letters
volume	32
number	1
page range	6-7
year	2003-01-05
URL	http://hdl.handle.net/10228/00006510

doi: info:doi/10.1246/cl.2003.6

Aniline-Hydroxylation Activity of a Flavin-linked $\beta\alpha\beta\alpha$ -Type Polypeptide Packing an Iron Porphyrin

Kin-ya Tomizaki, Hidekazu Nishino, Toru Arai,[†] Tamaki Kato,^{††} and Norikazu Nishino*^{††}

Faculty of Engineering, Kyushu Institute of Technology, Kitakyushu 804-8550

[†]Institute for the Fundamental Research of Organic Chemistry, Kyushu University, Fukuoka 812-8581

^{††}Graduate School of Life Science and Systems Engineering, Kyushu Institute of Technology, Kitakyushu 808-0196

(Received September 12, 2002; CL-020783)

Polypeptide containing iron porphyrin and flavin showed aniline-hydroxylation activity in the presence of 1-benzyl-1,4-dihyronicotinamide and O₂. The increased activity at pH 5.5 compared with that at pH 7.0 suggests that the dissociation of His from the iron porphyrin favored the hydroxylation.

One of the key steps in the oxidation with cytochrome P-450 is the electron transfer to this enzyme from cytochrome P-450 reductase.¹ Flavin derivatives reduce nearby hemes during the oxidation reaction catalyzed by enzymes² or artificial 4α -helix bundle polypeptides.³ Unlike 4α -helix bundle, which seems to surround porphyrin too closely, we recently synthesized a $\beta\alpha\beta\alpha$ -type polypeptide loosely packing porphyrin and reported its peroxide-dependent oxidation catalysis.⁴ In order to utilize O₂ as oxidant, we have attached flavin to this $\beta\alpha\beta\alpha$ -type polypeptide containing iron porphyrin, which could successfully hydroxylate aniline under the aerobic conditions (Figure 1).




Figure 1. (a) Polypeptides 1–3 with the iron porphyrin and/or flavin. A, Ala; a, D-Ala; E, Glu; G, Gly; H, His; K, Lys; L, Leu; P, Pro, and V, Val. (b) Supposed $\beta\alpha\beta\alpha$ -structure of 1.

Polypeptides 1 consist of β -strand segments and α -helix segments linked by spacers (Figure 1a). The β -strand segment (≈ 1.2 nm) are of alternative hydrophobic (X₁, Val, and X₂)/hydrophilic (Glu/Lys ion pair) sequences⁵ and the α -helix segments (≈ 1.5 nm) are of hydrophobic (four Leu)/hydrophilic (two pairs of Glu/Lys) residues on the opposite side of the helix,⁵ both designed to cover the porphyrin (≈ 1.0 nm). When the hydrophobic faces of the helices come close to the porphyrin to coordinate with iron by His, the hydrophobic Val in the β -strands would also come close to the porphyrin and the whole molecule would be arranged to give a structure with hydrophobic interior (porphyrin, Val, and Leu) and hydrophilic (Glu/Lys) exterior (Figure 1b). The flavin moiety exists at the molecular surface (see below), therefore, the reductant in the solution is expected to reduce flavin, and then the intramolecular electron transfer from the reduced flavin to iron porphyrin takes place efficiently in the

globular structure.

Polypeptide 1 and the control compounds (2, 3) were synthesized via the condensations of the peptide fragments as reported.⁴ Flavin (7 α -bromoacetyl-10-methylisalloxazine) was attached to Cys side chain of the deprotected 49-mer polypeptide⁶ before iron insertion. After the final purification with HPLC using a Superdex Peptide HR 10/30 column, polypeptides 1–3 were characterized by MALDI-TOF-MS. Unfortunately, ¹H NMR signals of the polypeptides were too broad to assign even before iron insertion.

UV/vis spectrum of 1 at pH 7.0 was characteristic for the low-spin Fe(III) complex with a narrow Soret band at 420 nm and broad Q bands at 550 and 575 nm (sh) (Figure 2a),⁷ which indicate the bis-His coordination as fifth and sixth ligands. The UV/vis spectra of 1 was concentration-independent from 0.6 μ M ($M = \text{mol dm}^{-3}$) to 15 μ M, indicate its monomeric structure. At pH 2.7, the Soret band shifted to 416 nm with the Q bands appeared at 570 and 615 nm, which indicate the formation of high-spin Fe(III) species.⁷ The His side chains were protonated at pH 2.7 and did not coordinate to iron porphyrin. However, these spectral changes showed no clear isosbestic points and showed a bimodal profile in the Abs₄₂₀ vs pH plot (Figure 2b), which suggests some intermediate species at around pH 5.5. The CD spectra of 1 at pH 7.0 showed negative bands at 222 and 208 nm, characteristic of the polypeptide containing α -helices. The CD spectrum is indeed ambiguous in the polypeptides containing β -strand due to the overlapping of the small Cotton effect of β -strand. However, when the $[\theta]_{222}$ value of 1 ($-12500 \text{ deg cm}^2 \text{ dmol}^{-1}$) was employed to analyze the α -helicity,⁴ 1 was evaluated to be 34% α -helix, which coincided with the 37% (18/49) value assuming the α -helical structure for the nine residues of each α -segment. We have no direct evidences for the conformation of β -strands at present, therefore, we wish to denominate 1 as the $\beta\alpha\beta\alpha$ -‘type’ polypeptide. However, CD spectra showed that the β -strand are not helical, consequently, it is at least sure that the porphyrin was loosely surrounded by the polypeptide at pH 7.0. The peptide 2 without flavin showed a CD spectrum ($[\theta]_{222} = -14000$) similar




Figure 2. UV/vis spectra of 1 (3.5 μ M) (a) at pH 7.0, 4.0, and 2.7. (b) Absorbance at 420 nm of 1 vs pH.

to **1**, indicating that the flavin attachment induce only slight changes to the conformation of the polypeptide. This fact supports the hypothesis that the flavin moiety is at the surface of the molecule (Figure 1b) due to the hydrophilic nature of the isoalloxazine ring. The bis-His coordination to Fe(III) in **1** and **2** stabilize their folded structures. However, polypeptide **3** without a porphyrin ($[\theta]_{222} = -9500$) and the polypeptide **1** at pH 2.7 with protonated His ($[\theta]_{222} = -10200$) show smaller helicities.

Next, we examined aniline hydroxylation in the presence of O₂ (air) catalyzed by **1**–**3** with 30 equivalents of the reductant (1-benzyl-1,4-dihydropyridinamide, BNAH) at pH 7.0 (Eq 1).⁸ The amount of the product *p*-aminophenol formed after the 15 min reaction, the only one product detected by HPLC, was determined by the literature method.⁸ The formation of *p*-aminophenol with **1** was six times greater than that with **2** at pH 7.0 (Table 1), which showed efficient electron relay via the attached flavin in **1**. To verify the electron relay through flavin we monitored the UV/vis spectra of the solution of **1** and **2** after addition of BNAH under Ar atmosphere. For polypeptide **1**, the absorptions at 432, 536, and 568 nm increased after the addition of BNAH, which correspond to the peaks of Fe(II) porphyrin.⁷ However, the Fe(II) species formation was significantly slow for **2** in the presence of BNAH. This clearly illustrates the importance of the flavin moiety in the polypeptide for the efficient electron transfer. Aniline hydroxylation hardly occurred with **3**, indicating that the hydroxylation was actually catalyzed by the iron porphyrin. Although hydroxylation of aniline occurs with Fe(III) porphyrin/H₂O₂,⁹ the reactivity of **1** was not changed by the addition of an H₂O₂ scavenger, catalase (350 unit cm⁻³), or an O₂⁻ scavengers, superoxide dismutase (100 unit cm⁻³). These results showed that **1** utilize O₂ and electrons derived from BNAH via flavin and not using H₂O₂ and O₂⁻ generated from the reducing system. Although excess amounts of BNAH was added, ≈20 equivalents of BNAH was consumed after 15 min in the reactions including **1** and **3**. However, BNAH was less consumed in the reaction with **2** (≈5 equivalents after 15 min). Although BNAH reduces flavin at the surface of **1**, aerobic reoxidation of the reduced flavin may occur to consume excess BNAH (Eq 2). This may be the reason of the low conversion of aniline and the low yield of *p*-aminophenol.




Table 1. Aniline hydroxylation catalyzed by **1**–**3** at 37 °C^a

Peptide	pH	[<i>p</i> -Aminophenol] after 15 min/μM
1	7.0	0.36
	5.5	0.63
2	7.0	0.06
	5.5	0.10
3	7.0	no reaction

^a[Peptide] = 7.0 μM, [aniline] = 10 mM, [BNAH] = 210 μM.

Saturation kinetics for aniline hydroxylation was observed for **1** at the ≈40 mM aniline concentration (Figure 3a). The Michaelis-Menten parameters of **1** are, $k_{\text{cat}} = 1.2 \times 10^{-2} \text{ min}^{-1}$ and $K_M = 2.6 \times 10^{-2} \text{ M}$. Such small k_{cat} and large K_M values of **1** are attributed to bis-His coordination in **1**, which prevents the




Figure 3. *p*-Aminophenol formation ([**1** or **2**] = 7.0 μM, [BNAH] = 210 μM, 37 °C). (a) **1** or **2** vs [aniline] at pH 7.0, (b) **1** or **2** vs pH, [aniline] = 10 mM).

access of aniline. In natural hemes, the distal side is exchangeable or vacant and the substrate is easily accessible showing the large k_{cat} and small K_M , for instance in myoglobin, $k_{\text{cat}} = 0.14$ and $K_M = 8.4 \times 10^{-3}$.^{2b} Thus, the aniline hydroxylation is carried out in acidic buffer solutions in which the protonated His dissociated from the iron porphyrin (Figure 3b). The maximal activity of **1** was at pH 5.5, where *p*-aminophenol formation was 175% compared to that at pH 7.0 (Table 1). It is interesting to note that the maximal absorption of the Soret band was obtained at pH 5.5 (Figure 2b). Although the pK_a of His is around 6.0, His might be in the mid of the deprotonation/protonation and coordination/dissociation at pH 5.5, which implied the formation of some catalytic active species (for instance, the 5-coordinated iron porphyrin). At high pH, two His coordinated to the iron porphyrin, therefore, the substrate was not easily accessible. At low pH such as pH 4, His was protonated and dissociated from the iron porphyrin to collapse the molecular structure and then the porphyrin and flavin might be separated. Thus, **1** showed a bell-shaped activity depending on the pH and its maximum activity was observed at pH 5.5. The addition of a denaturant, guanidine HCl, caused a decrease in the hydroxylation activity of **1** to ≈10%. This is probably because the conformational change by the denaturant increased the distance between the porphyrin and flavin. Further spectroscopic studies of the catalytic active species are underway.

This work was partly supported for N. N. by a Grant-in-Aid for Scientific Research (10480153) from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

References

- M. Sono, M. P. Roach, E. D. Coulter, and J. H. Dawson, *Chem. Rev.*, **96**, 2841 (1996); I. Tabushi, *Coord. Chem. Rev.*, **86**, 1 (1988).
- a) I. F. Sevrioukova, H. Li, H. Zhang, J. A. Peterson, and T. L. Poulos, *Proc. Natl. Acad. Sci. U.S.A.*, **96**, 1863 (1999). b) I. Hamachi, A. Fujita, and T. Kunitake, *Chem. Lett.*, **1995**, 657.
- R. E. Sharp, C. C. Moser, F. Rabanal, and P. L. Dutton, *Proc. Natl. Acad. Sci. U.S.A.*, **95**, 10465 (1998); H. Mihara, K. Tomizaki, T. Fujimoto, S. Sakamoto, H. Aoyagi, and N. Nishino, *Chem. Lett.*, **1996**, 187.
- K.-y. Tomizaki, H. Nishino, T. Kato, A. Miike, and N. Nishino, *Chem. Lett.*, **2000**, 648.
- J. P. Schneider and J. W. Kelly, *Chem. Rev.*, **95**, 2169 (1995).
- H. L. Levine and E. T. Kaiser, *J. Am. Chem. Soc.*, **100**, 7670 (1978).
- M. W. Makinen and A. K. Churg, in "Iron Porphyrins, Part 1," ed. by A. B. P. Lever and H. B. Gray, Addison-Wesley, Reading, M. A. (1983), pp 141–235.
- O. Takikawa, R. Yoshida, and O. Hayaishi, *J. Biol. Chem.*, **258**, 6808 (1983).
- T. Sasaki and E. T. Kaiser, *J. Am. Chem. Soc.*, **111**, 380 (1989); T. Mori, T. Santa, and M. Hirobe, *Tetrahedron Lett.*, **26**, 5555 (1985).