

## RESEARCH ARTICLE

## Open Access



# Vitamin D nutritional status and vitamin D regulated antimicrobial peptides in serum and pleural fluid of patients with infectious and noninfectious pleural effusions

Carlos A. Amado<sup>1\*</sup>, María T. García-Unzueta<sup>2</sup>, M. Carmen Fariñas<sup>3</sup>, Francisca Santos<sup>2</sup>, María Ortiz<sup>2</sup>, Pedro Muñoz-Cacho<sup>4</sup> and José A. Amado<sup>5</sup>

## Abstract

**Background:** Vitamin D and vitamin D dependent antimicrobial peptides such as Cathelicidin (LL-37) and  $\beta$ -defensin 2 have an important role in innate and adaptative immunity, but their role in pleural effusions has not been studied before.

**Methods:** Serum and pleural fluid samples from 152 patients with pleural effusion were collected, corresponding to 45 transudates and 107 exudates, 51 infectious effusions (14 complicated and 37 non-complicated), 44 congestive heart failure effusions and 38 malignant effusions. The levels of 25 OH-vitamin D, 1,25-(OH)<sub>2</sub>-vitamin D, Vitamin D Binding Protein (VDBP), LL-37 and  $\beta$ -defensin 2, both in serum and pleural fluid were evaluated in this prospective study. Differences between groups were analysed using unpaired t tests or Mann–Whitney tests. Correlations between data sets were examined using Pearson correlation coefficient or Spearman rank correlation coefficient. Diagnostic accuracy was estimated using ROC curve analysis.

**Results:** Low serum 25 OH vitamin D levels were found in all groups. Infectious effusions (IE) had higher serum and pleural fluid LL-37 levels compared to congestive heart failure or malignant effusions. Among IE, complicated had higher serum and pleural fluid LL-37 levels, and lower serum  $\beta$ -defensin-2 levels. Positive correlations were found between serum 25 OH-vitamin D levels and serum or pleural 1,25-(OH)<sub>2</sub>-vitamin D levels, and between 1,25-(OH)<sub>2</sub>-vitamin D and LL-37 serum. Diagnostic accuracy of the different molecules was moderate at best.

**Conclusions:** Hypovitaminosis D is highly prevalent in pleural effusions. LL-37 is produced intrapleurally in IE. This production is higher in complicated IE. No evidence of pleural production of  $\beta$ -defensin 2 was found in any of the groups. Diagnostic accuracy of the different molecules is at the best moderate for discriminating different types of effusions.

**Keywords:** Pleural effusion, LL-37, Cathelicidin,  $\beta$ -defensin-2, 25 OH vitamin D

\* Correspondence: [camado@humv.es](mailto:camado@humv.es)

<sup>1</sup>Division of Pneumology, Universidad de Cantabria, IDIVAL, Santander, Spain  
Full list of author information is available at the end of the article

## Background

Vitamin D modulates maturation, proliferation and function of different cells of both innate and adaptive immune system [1]. Vitamin D nutritional status is evaluated measuring serum 25 OH vitamin D, the stable metabolite of vitamin D. This molecule is the precursor of calcitriol or 1,25 (OH)<sub>2</sub> vitamin D, the active metabolite of vitamin D [2]. Hypovitaminosis D is a highly prevalent situation in hospitalized patients [3]. Multiple studies have shown that vitamin D and its antimicrobial dependent molecules play a defensive role in different infectious diseases such as tuberculosis, bacterial or viral infections [4–6]. Liu et al. [7] demonstrated that some *M. tuberculosis* antigens were recognized by Toll-Like 2/1 receptors of monocytes and innate immunity cells (skin, respiratory, digestive and urinary epithelium, etc.). This was followed by an intracrine activation of 1 $\alpha$ -hydroxylase, synthesis of 1,25 (OH)<sub>2</sub> vitamin D, and activation of the vitamin D receptor, inducing the synthesis and release of LL-37 (also known as human cathelicidin) and  $\beta$ -defensin-2. These molecules have important antimicrobial effects against virus, bacteria, and fungus in the lung [5, 6]. On the other hand, Vitamin D-binding protein (VDBP) is a multifunctional serum protein that binds vitamin D and its metabolites and transports them to target tissues [8]. VDBP is also a precursor of a macrophage activating factor [8].

There are no data in the literature regarding the prevalence of hypovitaminosis D in pleural effusions. An old study by Barnes et al. [9] showed that pleural levels of calcitriol were significantly higher in tuberculous pleural effusions as compared with cardiac effusions, although the number of patients studied was very small. On the other hand, in tuberculous patients pleural levels of 1,25 (OH)<sub>2</sub> vitamin D were significantly higher than serum levels. These data suggested intrapleural synthesis of 1,25 (OH)<sub>2</sub> vitamin D in tuberculosis, followed by release to blood. Otherwise, these aspects have never been studied in non-tuberculous pleural effusions.

One study measured VDBP in pleural fluid and serum [10]. Pleural fluid VDBP and VDBP pleural fluid/serum ratio were significantly higher in bacterial effusions compared to tuberculous or malignant. However, no differences were found in the serum VDBP/total protein ratio.

The possible defensive role of LL-37 in infectious pleural effusions has never been studied before. On the other hand, it is known that  $\beta$ -defensin-2 can be produced in vitro by pleural mesothelial cells when stimulated by *Staphylococcus* peptidoglycan [11]. This interesting finding suggests that these cells may have an important role in innate immunity of the pleural cavity, but there are no data in vivo in humans.

In this study we aimed to measure serum and pleural fluid levels of 25 OH vitamin D, 1,25 (OH)<sub>2</sub> vitamin D, VDBP, LL-37 and  $\beta$ -defensin-2 in patients with different etiological types of pleural effusions. We hypothesized that infectious effusions should have higher levels of serum or pleural calcitriol, LL-37 and  $\beta$ -defensin-2, reflecting a higher inflammatory response.

## Methods

### Study subjects


This is a prospective study approved by the Ethics Committee of Cantabria (CEIC Cantabria 25/2012). All patients gave informed consent to obtain samples of blood and pleural fluid at diagnosis time. We collected pleural fluid and serum at time of diagnosis of patients who were admitted in our hospital with the clinical diagnosis of pleural effusion along a period of 18 months. Patients with diseases or drugs known to alter vitamin D metabolism or immunological responses and patients with pleural effusion of unknown etiology were excluded. In this way, one hundred and fifty two patients managed by the same medical team were recruited as explained in Fig. 1.

### Classification

Patients were diagnosed of: 1) Congestive heart failure effusion (CHFE),  $n = 44$ , based on clinical grounds, according to the AHA guidelines, 2) Malignant effusion (ME),  $n = 38$ , when malignant cells were found in pleural effusion or pleural biopsy, 3) Infectious effusion (IE),  $n = 51$ , when clinical findings were compatible with pneumonia and there was response to antimicrobial treatment and/or positive culture results. There was also a miscellaneous group of 19 patients with pleural effusion due to other less frequent causes (pulmonary thromboembolism  $n = 7$ , pancreatitis  $n = 6$ , postoperative  $n = 5$ , and nephrotic syndrome  $n = 1$ , that was not separately analyzed. Pleural fluids were also classified as exudates  $n = 107$  or transudates  $n = 45$ , according to the cause of the effusion in consonance with Light criteria [12]. Patients with infectious effusions who received chest tubes for the treatment were classified as having complicated effusions (CIE). The decision to place a chest tube was left to the discretion of the attending physician, although it is likely that this decision was at least partially influenced by the biochemical characteristics of the fluid. Finally, we studied 37 noncomplicated (NCIE) and 14 CIE.

### Biochemical analyses

Serum and pleural effusion samples were collected in the appropriate way for the different assays. Routine determinations were done using the standard methods of our Hospital. Serum and pleural fluid aliquots for this specific study were stored at  $-80^{\circ}\text{C}$  until assayed.



25 OH vitamin D levels were measured by the automated chemiluminescence competitive assay IDS iSYS (Immunodiagnostic Systems Ltd, Boldon, UK), as previously reported [13]. Our laboratory is DEQAS (Vitamin D External Quality Assessment Scheme) certified. We considered hypovitaminosis D when serum levels of 25 OH vitamin D were lower than 20 ng/ml, according to the 2011 Food and Nutrition Board dietary references intakes for vitamin D and calcium. 1,25(OH)<sub>2</sub> vitamin D levels were measured by radioimmunoassay (DiaSorin Inc, Stillwater MN USA). VDBP, LL-37 and  $\beta$ -defensin-2 levels were measured by ELISA using the following kits: Quantikine Human Vitamin D Binding Protein (R&D Systems Europe, Abingdon, UK), HK321 Human LL-37 (Hycult Biotech, Uden, Holland) and human  $\beta$ -defensin-2 (Alpha Diagnostic International, San Antonio, TX USA).

#### Statistical analysis

Data are presented as mean  $\pm$  SD values for normally distributed data or median (interquartile range, IQR) for nonparametric data. Differences between proportions were calculated using chi-square test. Differences between groups were analyzed using unpaired t tests or Mann-Whitney tests. Correlations between data sets were examined using Pearson (r) correlation coefficient or Spearman rank (rs) correlation coefficient. Differences were considered significant for all statistical tests at *p* values of less than 0.05. All reported *p* values are two-sided. Analysis was performed using statistical computer Software (IBM SPSS Statistics) version 20.00 for Windows. ROC curves analysis was performed using software MED-CALC version 11.6.1.0 (MedCalc Software, Mariakerke, Belgium). The database and legend of the database supporting the conclusions of this article are included within its additional files.

#### Results

Tables 1 and 2 show the principal results.

#### Blood and pleural fluid cytologic and biochemical parameters

As expected, IE had higher levels of neutrophils and Lactate dehydrogenase (LDH), and lower pH and glucose levels in the pleural fluid compared to CHF. Exudates had higher levels of pleural fluid leukocytes, neutrophils and LDH and lower glucose and pH levels than transudates. CIE had a higher level of LDH and a lower pH than the NCIE.

#### Serum and pleural fluid vitamin D and vitamin D related molecules

There were no seasonal differences in the dates of obtaining the samples of the different groups (*p* = 0.102).

34 patients (77 %) in the CHF group, 33 (64 %) in the IE group and 20 (52 %) in the malignant group had vitamin D deficiency (*p* < 0.001). There were 35 (77 %) patients with transudative effusion presenting serum 25 OH levels below 20 ng/ml and 75 (65 %) in the exudative group (*p* = NS). There were 10 patients with 25 OH levels below 20 ng/ml in the CIE group (71 %) and 23 (62 %) in the NCIE group (*p* = NS).

ME had higher levels of serum 25OH vitamin D compared to CHF and IE (*p* < 0.001 and *p* = 0.01). Similar differences were found in 1,25 (OH)<sub>2</sub> vitamin D levels (*p* = 0.041 and *p* = 0.019). Exudative effusions had higher levels of 25 OH vitamin D compared to transudative effusions (*p* = 0,019). Pleural fluid levels of 1,25 (OH)<sub>2</sub> vitamin D were lower in CHF compared with infectious or malignant effusions (*p* = 0.033 and *p* < 0.001). No significant difference was found in serum or pleural fluid levels of 25 OH vitamin D, 1,25 (OH)<sub>2</sub> vitamin D or VDBP between CIE and NCIE.

**Table 1** Biochemical Parameters in Pleural effusion

Parameter		IE	ME	CHFE	P Value	Exudate	Transudate	P Value	CIE	NCIE	P Value
WBC (cells/ $\mu$ L)	Serum	10715 $\pm$ 5088	9435 $\pm$ 3451	9453 $\pm$ 5403	0.32	10156 $\pm$ 4575	9278 $\pm$ 5456	0.368	10969 $\pm$ 4457	10609 $\pm$ 5396	0.834
	Pleural	555 (160, 960)	400 (240, 680)	240 (160, 640)	0.149	410 (240, 800)	240 (160, 640)	0.046	560 (160, 800)	550 (160, 960)	0.761
N, (%)	Serum	74 (68, 80)	71 (61, 76.7)	75 (66, 81)	0.102	74 (68, 80)	74 (62, 81)	0.55	71.5 (68, 80)	74 (65, 81)	0.718
	Pleural	50 (17.5, 72.5)	15 (10, 20)	15 (5, 25)	0.001	20 (10, 55)	15 (5, 25)	0.026	58 $\pm$ 24	40 $\pm$ 29	0.111
pH	Pleural	7.23 $\pm$ 0.31	7.37 $\pm$ 0.07	7.4 $\pm$ 0.11	0.03	7.28 $\pm$ 0.26	7.4 $\pm$ 0.11	0.016	7.03 $\pm$ 0.32	7.30 $\pm$ 0.29	0.028
Glucose (mg/dL)	Pleural	99.47 $\pm$ 51.95	109 $\pm$ 42	133.67 $\pm$ 45.4	0.01	105.05 $\pm$ 50.98	133.67 $\pm$ 45.43	0.07	78.00 $\pm$ 58	107.68 $\pm$ 47.61	0.080
LDH (IU/L)	Pleural	270 (140, 735)	329 (144, 436)	123 (88, 270)	0.001	283 (134, 528)	119 (84, 261)	0.001	682 (164, 1771)	243 (121, 414)	0.035

IE Infectious effusions, CHFE Congestive heart failure effusions, ME Malignant effusions, CIE Complicated infectious effusions, NCIE Non complicated infectious effusions, WBC White blood cells, N Neutrophils, LDH Lactate Dehydrogenase. Data presented as mean  $\pm$  SD for parametric results and mean (interquartile range)

**Table 2** Vitamin D related molecules in Pleural effusion

Parameter		IE	ME	CHFE	P Value	Exudate	Transud.	P Value	CIE	NCIE	P Value
25OH Vitamin D (ng/mL)	Serum	12 (5, 84)	18.4 (6.9, 41)	10.4 (3, 39.6)	0,001	12.8 (5, 84)	10.6 (3, 39,6)	0,019	11.85 (9.2, 16.2)	13.5 (7.6, 22.4)	0.731
	Pleura	16 (1, 63)	17 (5.5, 47)	12.2 (1, 30.5)	0,016	15 (1, 63)	12 (1, 25)	0,048	16 (9.65, 20.65)	15 (7.75, 25.32)	0.551
1,25 (OH) <sub>2</sub> Vitamin D (pg/mL)	Serum	25.65 (7, 113)	46 (9, 140)	22.5 (5, 116)	0,043	25.65 (5, 140)	22 (8, 66)	0,708	22 (14.5, 35.5)	32 (8.5, 41.5)	0.751
	Pleura	22 (4, 132)	31.5 (0.4, 1.28)	12 (2, 52)	0,001	18 (4, 132)	12 (2, 50)	0,030	22 (12, 34)	21.5 (7.2, 32.5)	0.627
VDBP (mg/L)	Serum	203 (29, 494)	275.10 (13.13, 730)	197 (27, 1122)	0,1	236 (13, 730)	190 (27, 1122)	0,307	139 (29, 285)	166 (42, 338)	0.71
	Pleura	134 (4, 834)	191 (5, 464)	95 (10, 446)	0,001	155 (4, 834)	95 (10, 446)	0,48	85 (10, 183)	217 (30, 258)	0.112
LL-37 (ng/mL)	Serum	0.84 (0.01, 153)	0.47 (0.01, 12.62)	0.25 (0.01, 10.34)	0.017	0.67 (0.01, 153)	0.24 (0.01, 10.3)	0.017	0.51 (0.28, 1.29)	1.43 (0.95, 1.85)	0.029
	Pleura	1.42 (0.05, 153)	0.52 (0.01, 13.51)	0.3 (0.01, 3.3)	0.001	0.91 (0.01, 153)	0.3 (0.01, 3.3)	0.001	1.05 (0.47, 4.28)	3.11 (1.62, 5.62)	0.014
β-defensin-2 (pg/mL)	Serum	232 (14, 2100)	224 (10, 3000)	284 (9, 1834)	0.966	213 (9, 3000)	258 (9, 1834)	0.651	431 (136, 1015)	112 (71, 233)	0.023
	Pleura	265.2 (6, 1293)	276 (25, 3000)	241 (5, 2206)	0.706	240 (6, 3000)	214 (5, 2206)	0.484	383 (72, 804)	152 (16, 352)	0.091

IE Infectious effusions, CHFE Congestive heart failure effusions, ME Malignant effusions, CIE Complicated infectious effusions, NCIE Non complicated infectious effusions, WBC White blood cells, N Neutrophils, LDH Lactate Dehydrogenase, VDBP Vitamin D Binding Protein, LL-37 Leucin leucin 37 (Cathelicidin). Data presented as mean ± SD for parametric results and mean (interquartile range)

IE had higher levels of serum cathelicidin compared to CHFPE or ME ( $p = 0.005$  and  $p = 0.017$ ). Pleural levels of LL 37 were also higher in the IE group ( $p = 0.001$  and  $p = 0.001$ ). Exudative effusions had higher serum levels of LL-37 compared to transudative effusions ( $p = 0.017$ ). CIE had significantly higher levels of LL-37, both in serum ( $p = 0.029$ ) and pleural fluid ( $p = 0.014$ ) compared with the NCIE group. No differences in serum or pleural  $\beta$ -defensin-2 levels were found among the different etiological groups or between exudate and transudate groups. In contrast,  $\beta$ -defensin-2 serum levels were lower in CIE ( $p = 0.023$ ); otherwise, a non-significant trend for lower levels was found in CIE pleural fluid compared with the NCIE group ( $p = 0.091$ ).

#### Relationships among vitamin D related molecules

When all patients were considered, significant positive correlations were found between serum and pleural levels of 25 OH Vitamin D ( $rs = 0.596$ ,  $p < 0.001$ ), 1,25 (OH)<sub>2</sub> vitamin D ( $rs = 0.516$ ,  $p = 0.001$ ), VDBP ( $rs = 0.870$ ,  $p < 0.001$ ), LL-37 ( $rs = 0.707$ ,  $p < 0.001$ ) and  $\beta$ -defensin-2 ( $rs = 0.870$ ,  $p < 0.001$ ).

Interestingly, serum levels of 25 OH vitamin D also showed significant positive correlations with pleural ( $rs = 0.54$ ,  $p < 0.001$ ) 1,25 (OH)<sub>2</sub> vitamin D levels. Furthermore, serum 1,25 (OH)<sub>2</sub> vitamin D levels were significantly correlated with serum LL-37 levels ( $rs = 0.388$ ,  $p = 0.05$ ), but not with serum  $\beta$ -defensin-2 levels.

On the other hand, vitamin D metabolites and related molecules did not correlate with glucose, pH, or LDH, either in serum or in pleural effusion.

#### Diagnostic accuracy of vitamin D related molecules

Evaluation of ROC plots in serum and pleural fluid samples taken from patients with effusions for the discrimination between infectious origin and other origin were similar for pleural fluid LL-37 [AUC = 0.732 (95 % CI, 0.657-0.798),  $p < 0.001$ ] and pleural fluid 1,25 (OH)<sub>2</sub> vitamin D [AUC = 0.754 (95 % CI, 0.654-0.838),  $p < 0.001$ ].

Pleural fluid LL-37 [AUC = 0.657 (95 % CI, 0.575-0.733),  $p = 0.0016$ ] was superior to serum LL-37 [AUC = 0.586 (95 % CI, 0.503-0.667),  $p = 0.1181$ ] for the differentiation between exudative and transudative effusions.

Pleural fluid LL-37 [AUC = 0.731 (95 % CI, 0.585-0.847),  $p = 0.0012$ ] and serum  $\beta$ -defensin2 [AUC = 0.731 (95 % CI, 0.562-0.861),  $p = 0.0085$ ] were superior to serum LL-37 [AUC = 0.708 (95 % CI, 0.557-0.831),  $p = 0.0015$ ] for the differentiation between CIE and NCIE.

#### Discussion

In 1998 Thomas et al. [3] showed in an influential paper that the prevalence of hypovitaminosis D was high in non-selected inpatient subjects from the medical wards of the Massachusetts General Hospital in Boston. A

multivariate study demonstrated different factors related to hypovitaminosis, such as advanced age, insufficient ingestion of vitamin D or sun exposure, presence of specific diseases (nephrotic syndrome, malabsorption, hepatic cirrhosis, morbid obesity etc.) and use of drugs that increase vitamin D catabolism. In that study no patients with pleural effusion were included, and since then the prevalence of hypovitaminosis D in pleural effusions has not been specifically studied. Our study shows, as expected, that all kinds of pleural effusion are associated frequently with hypovitaminosis D defined as 25 OH vitamin D serum levels lower than 20 ng/ml.

Significant lower levels of serum 1,25 (OH)<sub>2</sub> vitamin D were found in CHFPE. This might probably be due to lower levels of 25 OH vitamin D in this group. This is supported by the positive correlation between 25 OH vitamin D and 1,25 (OH)<sub>2</sub> vitamin D levels. This correlation is typical of hypovitaminosis D, and it means that the substrate (25 OH vitamin D) is really a limiting factor for the production of 1,25 (OH)<sub>2</sub> vitamin D. If 25 OH vitamin D levels were normal (plenty of substrate) this correlation should not be found. No significant differences were found in either serum or pleural fluid levels of 25 OH vitamin D, 1,25 (OH)<sub>2</sub> vitamin D or VDBP in CIE vs. NCIE. We had hypothesized that pleural 1,25 (OH)<sub>2</sub> vitamin D could be increased in complicated patients, but our data does not support this hypothesis. This can be due, at least in part, to the fact that low levels of the precursor molecule 25 OH vitamin D limit the production of 1,25 (OH)<sub>2</sub> vitamin D. Furthermore, the paracrine/intracrine production of calcitriol in response to the infection, may not be enough to induce measurable changes in pleural fluid or serum.

On the other hand, we found higher levels of LL-37 both in serum and pleural fluid in IE, specially in CIE, suggesting that, despite hypovitaminosis D, innate immune cells are able to produce higher quantities of this natural antimicrobial molecule when they are severely challenged. The positive correlation between serum 1,25 (OH)<sub>2</sub> vitamin D and LL-37 suggests that higher levels of the active metabolite in fact potentiate LL-37 synthesis, even in the presence of hypovitaminosis D. On the contrary serum  $\beta$ -defensin-2 levels were decreased in CIE and did not correlate with 1,25 (OH)<sub>2</sub> levels. The different behavior of these antibiotic molecules suggests that, although both are modulated by vitamin D, other factors are also importantly involved in their response. For example, it has been demonstrated in vitro that  $\beta$ -defensin-2 responses to vitamin D are lower than those of LL-37 [7]. In vivo, quantity and time course of response of these molecules to different challenges are also different [14, 15]. Furthermore it has been reported that  $\beta$ -defensin-2 decreases in sepsis [16] and lower plasma level of this substance is an independent predictor

of adverse outcomes in patients with community-acquired pneumonia [17]. Further time course studies are needed to analyze whether this is due to consumption of  $\beta$ -defensin-2, or because certain patients have lower response of this molecule leading to susceptibility to sepsis.

Significant positive correlations were found between serum and pleural levels of all the molecules studied, suggesting there is good diffusion of these molecules through the pleural membranes. Otherwise, no correlations were found with other established pleural biomarkers, such as pH, glucose or LDH.

Diagnostic accuracy of the different molecules evaluated by ROC analysis is similar or lower to standardized methods used in clinical practice. These results suggest that although some vitamin D related molecules could be markers with relatively good sensitivity and specificity for pleural effusion evaluation, the cost and time needed do not support its use as a routine laboratory test.

Our study has limitations. The number of patients studied is relatively small in some of the groups, especially in the CIE group, and we have only one determination, at the time of diagnosis. Subsequent samples should give a clearer picture of the responses and interactions among all molecules. Likewise, the definition of CIE and NCIE is never clear-cut. In the present study CIE were defined as those that received chest tubes. This decision was made with the knowledge of the pleural fluid pH, glucose and LDH, and this could have influenced which patients were given chest tubes.

## Conclusion

In conclusion hypovitaminosis D is highly prevalent in all kinds of pleural effusions. LL-37 is higher both in pleural fluid and blood of the more inflammatory effusions, but there is a weaker response of  $\beta$ -defensin-2 in blood. The pathophysiological significance of these findings merits to be studied in further detail. Also, a clinical trial aimed at improving the vitamin D nutritional status should test if vitamin D boosts the innate antibiotic and clinical response in these patients.

## Abbreviations

VDBP Vitamin D-binding protein; LL-37 human cathelicidin; CHF Congestive heart failure effusion; ME Malignant effusion; IE Infectious effusion; CIE complicated effusions; NCIE noncomplicated effusions; DEQAS Vitamin D External Quality Assessment Scheme; LDH Lactate dehydrogenase

## Acknowledgments

We gratefully acknowledge the laboratory work done by Blanca Paule and Ana Berja.

## Funding

This study was funded by IDIVAL and University of Cantabria. All Authors meet conditions listed in the international committee of medical journal editors.

## Availability of data and materials

The database supporting the conclusions of this article is included within the article and its additional files. A legend is provided as part of the file upload.

## Authors' contributions

CAA participated in the design of the study, drafted the manuscript, and participated in the statistical analysis. MGU participated in the design of the study and carried out the immunoassays. MCF participated in the design of the study and coordination and helped to draft the manuscript. FS Participated in the design of the study and performed the immunoassays. MO Participated in the design of the study and performed the immunoassays. PMC participated in the design of the study and performed the statistical analysis. JAA conceived of the study, and participated in its design and coordination and helped to draft the manuscript. All authors read and approved the final manuscript.

## Author's information

CAA Division of Pneumology HUMV. PhD. MGU Division of Clinical Biochemistry. PhD. MCF Head of Infectious Diseases Division HUMV, Professor of the University of Cantabria, member of IDIVAL, Santander. FS Division of Clinical Biochemistry. MO Division of Clinical Biochemistry. PMC Statistics at Primary Care management. Servicio Cántabro de Salud, Santander. PhD JAA Head of the Division of Endocrinology, HUMV, Professor at University of Cantabria, member of IDIVAL, Santander.

## Competing interests

The authors declare that they have no competing interests.

## Consent for publication

Not applicable.

## Ethics approval and consent to participate

This is a prospective study approved by the Ethics Committee of Cantabria (CEIC Cantabria 25/2012). All patients gave informed consent to obtain samples of blood and pleural fluid at diagnosis time. This statement is included in Methods section of the manuscript.

## Author details

<sup>1</sup>Division of Pneumology, Universidad de Cantabria, IDIVAL, Santander, Spain. <sup>2</sup>Clinical Biochemistry, Universidad de Cantabria, IDIVAL, Santander, Spain. <sup>3</sup>Infectious Diseases Hospital Universitario Marqués de Valdecilla (HUMV), Universidad de Cantabria, IDIVAL, Santander, Spain. <sup>4</sup>Gerencia Atención Primaria, Servicio Cántabro de Salud, Santander, Spain. <sup>5</sup>Division of Endocrinology, Hospital Universitario Marqués de Valdecilla, Universidad de Cantabria, IDIVAL, Santander, Spain.

Received: 2 December 2015 Accepted: 28 June 2016

Published online: 08 July 2016

## References

- Adams JS, Hewison M. Update in vitamin D. *J Clin Endocrinol Metab.* 2010; 95:471–8.
- Holick MF. Vitamin D, deficiency. *N Engl J Med.* 2007;357:266–81.
- Thomas MK, Lloyd-Jones DM, Thadhani RI, Shaw A, Deraska DJ, Kitchi BT, Vamvakas EC, Dick IM, Prince RL, Finkelstein JS. Hypovitaminosis D in medical inpatients. *N Engl J Med.* 1998;338:777–83.
- Khoo AL, Chai L, Koenen H, Joosten I, Netea M, van der Ven A. Translating the role of vitamin D3 in infectious diseases. *Crit Rev Microbiol.* 2012;38:122–35.
- Vandamme D, Landuyt B, Luyten W, Schoofs L. A comprehensive summary of LL-37, the factotum human cathelicidin peptide. *Cell Immunol.* 2012;280:22–35.
- Teclé T, Tripathi S, Hartshorn KL. Review: defensins and cathelicidins in lung immunity. *Innate Immun.* 2010;16:151–9.
- Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzik SR, Ochoa MT, Schaubert J, Wu K, Meinken C, Kamen DL, Wagner M, Bals R, Steinmeyer A, Zügel U, Gallo RL, Eisenberg D, Hewison M, Hollis BW, Adams JS, Bloom BR, Modlin RL. Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science.* 2006;311:1770–3.
- Chishimba L, Thickett DR, Stockley RA, Wood AM. The vitamin D axis in the lung: a key role for vitamin D-binding protein. *Thorax.* 2010;65:456–62.

9. Barnes PF, Modlin RL, Bikle DD, Adams JS. Transpleural gradient of 1,25-dihydroxyvitamin D in tuberculous pleuritis. *J Clin Invest.* 1989;83:1527–32.
10. Hamashima Y, Kanazawa T, Hirata A, Yamai Y, Fujihara H, Sekine K, Nagao K. Measurement of vitamin D-binding protein in pleural fluids and sera by means of a turbidimetric immunoassay measuring system. *Clin Chim Acta.* 2002;321:23–8.
11. Hussain T, Nasreen N, Lai Y, Bellew BF, Antony VB, Mohammed KA. Innate immune responses in murine pleural mesothelial cells: toll-like receptor-2 dependent induction of  $\beta$ -defensin-2 by staphylococcal peptidoglycan. *Am J Physiol Lung Cell Mol Physiol.* 2008;295:461–70.
12. Light RW, Macgregor I, Luchsinger PC, Ball Jr WC. Pleural effusions: the diagnostic separation of transudates and exudates. *Ann Intern Med.* 1972;77:507–13.
13. Alvarez RL, Lopez HM, Garcia Unzueta MT, Amado JA, Cacho PM, Martinez Taboada VM. Age and low levels of circulating vitamin D are associated with impaired innate immune function. *J Leukoc Biol.* 2012;91:829–38.
14. Lippross S, Klueter T, Steubesand N, Oestern S, Mentlein R, Hildebrandt F, Podschum R, Pufe T, Seekamp A, Varoga D. Multiple trauma induces serum production of host defence peptides. *Injury.* 2011;43:137–42.
15. Barbeiro DF, Barbeiro HV, Zampieri FG, Machado MCC, Filho FT, Cunha DMG, Goulart AC, Velasco IT, Neto LM, de Souza HP, da Silva FP. Cathelicidin LL-37 bloodstream surveillance is down regulated during septic shock. *Microb Infect.* 2013;15:342–6.
16. Olbrich P, Pavón A, Rosso ML, Molinos A, De Felipe B, Sanchez B, Praena-Fernandez JM, Jimenez F, Obando I, Neth O. Association of human beta-defensin 2 levels and sepsis in preterm neonates. *Pediatr Crit Care Med.* 2013;14:796–800.
17. Liu S, He LR, Wang W, Wang GH, He ZY. Prognostic value of plasma human  $\beta$ -defensin-2 level on short-term clinical outcomes in patients with community-acquired pneumonia: a preliminary study. *Respir Care.* 2013;58:655–61.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

