

Supplement of Biogeosciences, 15, 1029–1045, 2018
<https://doi.org/10.5194/bg-15-1029-2018-supplement>
© Author(s) 2018. This work is distributed under
the Creative Commons Attribution 3.0 License.



Supplement of

Simultaneous shifts in elemental stoichiometry and fatty acids of *Emiliana huxleyi* in response to environmental changes

Rong Bi et al.

Correspondence to: Meixun Zhao (maxzhao@ouc.edu.cn)

The copyright of individual parts of the supplement might differ from the CC BY 3.0 License.

1 **Table S1.** Measured dissolved inorganic carbon (DIC) and total alkalinity (TA), and
 2 calculated $p\text{CO}_2$ (mean \pm SE) at the end of the experiments in the cultures of
 3 *Emiliana huxleyi*. N:P: N:P supply ratios. Outliers in the data of $p\text{CO}_2$ were excluded
 4 in the table.

Treatment			DIC ($\mu\text{mol kg}^{-1}$)	TA ($\mu\text{mol kg}^{-1}$)	$p\text{CO}_2$ (μatm)
12 °C	Low $p\text{CO}_2$	N:P = 10:1	1302 \pm 54	1269 \pm 57	1509 \pm 35
		N:P = 24:1	1328 \pm 18	1292 \pm 27	1564 \pm 149
		N:P = 63:1	1374 \pm 25	1349 \pm 24	1412 \pm 21
	High $p\text{CO}_2$	N:P = 10:1	1956 \pm 46	1962 \pm 50	1357 \pm 14
		N:P = 24:1	2042 \pm 17	2053 \pm 17	1357 \pm 76
		N:P = 63:1	1829 \pm 22	1801 \pm 49	1041 \pm 191
18 °C	Low $p\text{CO}_2$	N:P = 10:1	763 \pm 15	793 \pm 4	552 \pm 118
		N:P = 24:1	885 \pm 6	922 \pm 12	567 \pm 84
		N:P = 63:1	1065 \pm 3	1108 \pm 8	633 \pm 44
	High $p\text{CO}_2$	N:P = 10:1	1415 \pm 154	1454 \pm 121	1113 \pm 489
		N:P = 24:1	1278 \pm 13	1196 \pm 18	2944 \pm 330
		N:P = 63:1	1613 \pm 35	1620 \pm 32	1507 \pm 332
24 °C	Low $p\text{CO}_2$	N:P = 10:1	785 \pm 13	808 \pm 10	845 \pm 256
		N:P = 24:1	809 \pm 10	682 \pm 11	-
		N:P = 63:1	1243 \pm 16	1231 \pm 10	1734 \pm 163
	High $p\text{CO}_2$	N:P = 10:1	1266 \pm 22	1240 \pm 20	2079 \pm 406
		N:P = 24:1	1596 \pm 63	1691 \pm 36	1163 \pm 190
		N:P = 63:1	1616 \pm 27	1550 \pm 34	3295 \pm 171

5 **Table S2.** Results of Akaike information criterion corrected (AICc) in GLMMs for the
6 observed maximal growth rate (μ_{\max}), elemental stoichiometry and fatty acid
7 proportions and contents in response to temperature, N:P supply ratios and $p\text{CO}_2$ in
8 *Emiliana huxleyi*. The selected models are shown in bold, the results of which are
9 shown in Table 1. POC: particulate organic carbon; PON: particulate organic nitrogen;
10 POP: particulate organic phosphorus; PIC: particulate inorganic carbon; TFAs: total
11 fatty acids; SFA: saturated fatty acid; MUFA: monounsaturated fatty acid; PUFA:
12 polyunsaturated fatty acid; DHA: docosaheptaenoic acid. Effect builder of main:
13 models containing first order effects of the three factors; effect builder of main, two
14 way: models containing first order effects and second order interactions of the three
15 factors; effect builder of main, two way and three way: models containing first order
16 effects, second and third order interactions of the three factors.

Variable	Effect builder	AICc
μ_{\max}	Main, two way and three way	23.856
	Main, two way	18.473
	Main	5.471
POC (pg cell^{-1})	Main, two way and three way	336.081
	Main, two way	333.586
	Main	339.852
POC ($\mu\text{g ml}^{-1}$)	Main, two way and three way	304.408
	Main, two way	280.234
	Main	235.488
POC production ($\text{pg cell}^{-1} \text{d}^{-1}$)	Main, two way and three way	88.022
	Main, two way	59.365
	Main	5.219
PON (pg cell^{-1})	Main, two way and three way	125.664
	Main, two way	123.586
	Main	125.405
POP (pg cell^{-1})	Main, two way and three way	-139.184
	Main, two way	-140.161

	Main	-136.986
PIC (pg cell ⁻¹)	Main, two way and three way	285.804
	Main, two way	284.025
	Main	299.364
PIC (µg ml ⁻¹)	Main, two way and three way	300.200
	Main, two way	276.029
	Main	231.545
PIC production (pg cell ⁻¹ d ⁻¹)	Main, two way and three way	92.222
	Main, two way	64.188
	Main	9.065
POC:PON (mol mol ⁻¹)	Main, two way and three way	220.755
	Main, two way	218.755
	Main	213.130
POC:POP (mmol mol ⁻¹)	Main, two way and three way	613.955
	Main, two way	611.731
	Main	606.395
PON:POP (mol mol ⁻¹)	Main, two way and three way	362.508
	Main, two way	359.671
	Main	356.018
PIC:POC	Main, two way and three way	56.147
	Main, two way	26.690
	Main	-36.148
SFA proportion (% of TFAs)	Main, two way and three way	304.845
	Main, two way	302.115
	Main	304.984
MUFA proportion (% of TFAs)	Main, two way and three way	300.697
	Main, two way	278.543
	Main	264.319
PUFA proportion (% of TFAs)	Main, two way and three way	359.132
	Main, two way	336.555
	Main	318.057
DHA proportion (% of TFAs)	Main, two way and three way	304.197
	Main, two way	301.625
	Main	310.200
TFA content (µg mg ⁻¹ C ⁻¹)	Main, two way and three way	554.949
	Main, two way	536.499
	Main	512.664
SFA content (µg mg ⁻¹ C ⁻¹)	Main, two way and three way	437.382
	Main, two way	416.262
	Main	393.592
MUFA content (µg mg ⁻¹ C ⁻¹)	Main, two way and three way	421.162
	Main, two way	400.009

	Main	374.298
PUFA content ($\mu\text{g mg}^{-1} \text{C}^{-1}$)	Main, two way and three way	485.817
	Main, two way	465.876
	Main	432.787
DHA content ($\mu\text{g mg}^{-1} \text{C}^{-1}$)	Main, two way and three way	449.256
	Main, two way	428.583
	Main	391.542

17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 **Table S3.** The nature (synergism or antagonism) and magnitude (the difference
53 between observed combined effect and predicted additive effect) of the observed
54 interactive effects of warming, N and P deficiency (-N and -P), and enhanced $p\text{CO}_2$
55 (HCO_2) on cellular contents of particulate organic carbon (POC), particulate organic
56 nitrogen (PON), particulate organic phosphorus (POP) and particulate inorganic
57 carbon (PIC), and proportions of saturated fatty acids (SFAs), and docosahexaenoic
58 acid (DHA) in *Emiliana huxleyi*. TFAs: total fatty acids.

Variable	Treatment	Interaction		<i>n</i>
		Nature	Magnitude \pm SE	
POC (pg cell ⁻¹)	Warming \times -N	Synergism	19.056 \pm 0.392	12
	Warming \times -P	Synergism	39.644 \pm 2.854	12
PON (pg cell ⁻¹)	Warming \times -N	Synergism	0.948 \pm 0.039	12
	Warming \times -P	Synergism	3.586 \pm 0.327	12
POP (pg cell ⁻¹)	Warming \times -N	Synergism	0.154 \pm 0.004	12
	Warming \times -P	Synergism	0.237 \pm 0.019	12
	Warming \times HCO_2	Synergism	0.315 \pm 0.023	18
PIC (pg cell ⁻¹)	Warming \times -N	Antagonism	-2.010 \pm 0.524	12
	Warming \times -P	Synergism	9.511 \pm 2.264	12
	Warming \times HCO_2	Synergism	17.640 \pm 1.495	18
SFAs (% of TFAs)	-N \times HCO_2	Synergism	28.746 \pm 1.070	9
	-P \times HCO_2	Synergism	24.096 \pm 0.840	9
DHA (% of TFAs)	Warming \times -N	Synergism	4.622 \pm 0.873	12
	Warming \times -P	Synergism	4.316 \pm 0.671	12
	Warming \times HCO_2	Synergism	5.013 \pm 0.912	18

59

60 **Table S4.** Fatty acid profiles of *Emiliana huxleyi* under three temperatures (12, 18 and 24 °C), three N:P supply ratios (molar ratios 10:1, 24:1
61 and 63:1) and two $p\text{CO}_2$ levels (560 and 2400 μatm). Data are expressed as fatty acid contents ($\mu\text{g mg C}^{-1}$) and percentages of total fatty acids (%
62 of TFAs) (mean \pm SE). SFAs, saturated fatty acids; MUFAs, monounsaturated fatty acids; PUFAs, polyunsaturated fatty acids; TFAs, total fatty
63 acids.

	12 °C											
	Low $p\text{CO}_2$						High $p\text{CO}_2$					
	N:P = 10:1		N:P = 24:1		N:P = 63:1		N:P = 10:1		N:P = 24:1		N:P = 63:1	
	Content	%	Content	%	Content	%	Content	%	Content	%	Content	%
14:0	31 \pm 1	19 \pm 1	26 \pm 4	20 \pm 0	23 \pm 3	16 \pm 1	22 \pm 0	17 \pm 1	24 \pm 1	20 \pm 2	12 \pm 0	15 \pm 0
16:0	11 \pm 0	7 \pm 0	10 \pm 2	7 \pm 0	9 \pm 2	6 \pm 0	11 \pm 2	8 \pm 0	9 \pm 2	8 \pm 1	4 \pm 0	5 \pm 0
16:1n-7	1 \pm 0	1 \pm 0	1 \pm 0	1 \pm 0	2 \pm 0	1 \pm 0	1 \pm 0	1 \pm 0	1 \pm 0	1 \pm 0	1 \pm 0	1 \pm 0
18:0	3 \pm 0	2 \pm 0	4 \pm 1	3 \pm 0	6 \pm 2	4 \pm 1	6 \pm 2	4 \pm 1	4 \pm 2	3 \pm 1	2 \pm 0	2 \pm 0
18:1n-9	33 \pm 1	20 \pm 0	27 \pm 4	20 \pm 0	25 \pm 3	17 \pm 1	23 \pm 1	18 \pm 1	22 \pm 2	19 \pm 0	11 \pm 0	13 \pm 0
18:1n-7	6 \pm 0	3 \pm 0	5 \pm 1	4 \pm 0	7 \pm 1	5 \pm 0	5 \pm 0	4 \pm 0	5 \pm 0	4 \pm 0	4 \pm 0	4 \pm 0
18:2n-6	9 \pm 0	5 \pm 0	7 \pm 1	6 \pm 0	5 \pm 1	4 \pm 0	5 \pm 0	4 \pm 0	5 \pm 1	5 \pm 0	2 \pm 0	3 \pm 0
18:3n-6	0 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0
18:3n-3	11 \pm 1	7 \pm 0	8 \pm 1	6 \pm 0	10 \pm 2	7 \pm 0	9 \pm 1	7 \pm 0	7 \pm 1	6 \pm 0	6 \pm 0	7 \pm 0
18:4n-3	7 \pm 0	4 \pm 0	5 \pm 1	4 \pm 0	6 \pm 1	4 \pm 0	6 \pm 1	4 \pm 0	5 \pm 1	4 \pm 0	4 \pm 0	5 \pm 0
20:2n-6	1 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	1 \pm 0	1 \pm 0	1 \pm 0	0 \pm 0	0 \pm 0	0 \pm 0	1 \pm 0	1 \pm 0
20:3n-6	0 \pm 0	0 \pm 0	1 \pm 0	0 \pm 0	1 \pm 0	1 \pm 0	0 \pm 0	0 \pm 0	1 \pm 1	1 \pm 0	0 \pm 0	0 \pm 0
22:0	1 \pm 0	1 \pm 0	1 \pm 0	1 \pm 0	2 \pm 0	1 \pm 0	1 \pm 0	1 \pm 0	1 \pm 0	1 \pm 0	0 \pm 0	1 \pm 0

20:5n-3	2 ± 0	1 ± 0	1 ± 0	1 ± 0	2 ± 0	1 ± 0	1 ± 0	1 ± 0	1 ± 0	1 ± 0	1 ± 0	1 ± 0
23:0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	1 ± 1	1 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
24:0	1 ± 0	0 ± 0	0 ± 0	0 ± 0	1 ± 0	0 ± 0	1 ± 0	1 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
22:5n-3, 24:1n-9	1 ± 0	1 ± 0	1 ± 0	1 ± 0	2 ± 1	1 ± 0	1 ± 0	1 ± 0	1 ± 0	1 ± 0	1 ± 0	1 ± 0
22:6n-3	29 ± 2	18 ± 1	23 ± 4	17 ± 1	33 ± 7	22 ± 2	26 ± 3	20 ± 1	20 ± 3	17 ± 1	22 ± 1	27 ± 0
Unidentified	14 ± 1	9 ± 0	10 ± 2	7 ± 0	12 ± 3	8 ± 1	10 ± 1	7 ± 0	9 ± 1	7 ± 0	10 ± 0	12 ± 0
∑SFAs ^a	47 ± 0	29 ± 1	43 ± 7	32 ± 1	41 ± 6	28 ± 1	41 ± 5	32 ± 1	38 ± 5	32 ± 1	20 ± 1	24 ± 1
∑MUFAs ^b	41 ± 1	26 ± 0	35 ± 5	26 ± 0	35 ± 4	25 ± 1	30 ± 1	23 ± 1	30 ± 4	25 ± 1	16 ± 0	20 ± 0
∑PUFAs ^c	59 ± 4	37 ± 1	47 ± 8	35 ± 1	58 ± 11	39 ± 1	49 ± 5	38 ± 1	41 ± 5	35 ± 1	36 ± 1	44 ± 1
∑TFAs ^d	162 ± 5		134 ± 22		146 ± 24		130 ± 12		118 ± 14		82 ± 2	

64 ^a also includes 20:0 present at < 0.5% of TFAs in all treatments. ^b also includes 14:1, 20:1n-9 and 22:1n-9 present at < 0.5% of TFAs in all
65 treatments. ^c also includes 16:3n-4, 20:4n-6, 20:3n-3, 20:4n-3, 22:2n-6 present at < 0.5% of TFAs in all treatments. ^d also includes the
66 unidentified FA component.

67
68
69
70
71
72
73
74
75
76
77
78

79 Table S4. Continued.

	18 °C											
	Low $p\text{CO}_2$						High $p\text{CO}_2$					
	N:P = 10:1		N:P = 24:1		N:P = 63:1		N:P = 10:1		N:P = 24:1		N:P = 63:1	
	Content	%	Content	%	Content	%	Content	%	Content	%	Content	%
14:0	27 ±2	18 ±1	17 ±0	17 ±0	21 ±0	15 ±1	32 ±3	18 ±1	20 ±3	18 ±1	16 ±0	16 ±0
16:0	9 ±0	6 ±0	5 ±0	5 ±0	7 ±0	5 ±0	12 ±1	7 ±0	8 ±2	7 ±0	6 ±0	6 ±0
16:1n-7	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0
18:0	2 ±0	2 ±0	1 ±0	1 ±0	2 ±0	2 ±0	3 ±0	2 ±0	3 ±1	3 ±0	2 ±0	2 ±0
18:1n-9	23 ±1	15 ±0	12 ±0	13 ±0	16 ±0	11 ±0	31 ±2	17 ±1	17 ±3	16 ±1	15 ±0	15 ±0
18:1n-7	6 ±0	4 ±0	3 ±0	3 ±0	6 ±0	4 ±0	6 ±0	4 ±0	4 ±1	4 ±0	4 ±0	4 ±0
18:2n-6	4 ±0	3 ±0	3 ±0	3 ±0	3 ±0	2 ±0	5 ±0	3 ±0	4 ±1	4 ±0	3 ±0	2 ±0
18:3n-6	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0
18:3n-3	10 ±0	7 ±0	8 ±0	8 ±0	10 ±0	7 ±0	12 ±0	7 ±0	7 ±1	6 ±0	8 ±0	8 ±0
18:4n-3	10 ±0	7 ±0	8 ±0	8 ±0	10 ±0	7 ±0	10 ±1	6 ±0	7 ±1	6 ±0	6 ±0	6 ±0
20:2n-6	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	1 ±0	1 ±0
20:3n-6	0 ±0	0 ±0	0 ±0	0 ±0	1 ±0	0 ±0	0 ±0	0 ±0	1 ±0	1 ±0	0 ±0	0 ±0
22:0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0
20:5n-3	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0
23:0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0
24:0	1 ±0	0 ±0	1 ±0	1 ±0	0 ±0	0 ±0	1 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0
22:5n-3, 24:1n-9	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	2 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0
22:6n-3	40 ±2	26 ±1	25 ±1	26 ±0	40 ±3	29 ±1	42 ±1	25 ±1	25 ±5	23 ±2	27 ±1	26 ±1
Unidentified	13 ±1	9 ±0	11 ±0	11 ±0	16 ±1	12 ±0	13 ±0	8 ±0	9 ±2	8 ±0	10 ±0	10 ±0

Σ SFAs ^a	41 ±2	27 ±1	24 ±1	25 ±0	32 ±0	23 ±1	49 ±4	28 ±1	33 ±5	29 ±1	26 ±1	25 ±1
Σ MUFAs ^b	32 ±1	21 ±0	17 ±0	18 ±0	25 ±1	18 ±0	40 ±2	23 ±0	24 ±4	22 ±1	22 ±0	22 ±0
Σ PUFAs ^c	67 ±2	44 ±1	45 ±1	47 ±0	67 ±4	48 ±1	73 ±0	42 ±2	46 ±9	41 ±2	45 ±1	44 ±1
Σ TFAs ^d	153 ±6		97 ±3		140 ±5		176 ±7		112 ±19		103 ±0	

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100 Table S4. Continued.

	24°C											
	Low $p\text{CO}_2$						High $p\text{CO}_2$					
	N:P = 10:1		N:P = 24:1		N:P = 63:1		N:P = 10:1		N:P = 24:1		N:P = 63:1	
	Content	%	Content	%	Content	%	Content	%	Content	%	Content	%
14:0	17 ±0	17 ±0	15 ±0	18 ±1	23 ±3	18 ±1	18 ±0	19 ±1	7 ±2	16 ±1	12 ±1	14 ±0
16:0	7 ±0	7 ±0	6 ±1	7 ±0	10 ±1	8 ±1	8 ±0	8 ±1	4 ±1	10 ±1	7 ±1	8 ±0
16:1n-7	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0
18:0	2 ±0	2 ±0	2 ±1	3 ±1	4 ±1	3 ±1	2 ±0	2 ±1	3 ±0	7 ±1	3 ±1	4 ±1
18:1n-9	11 ±0	11 ±0	7 ±0	8 ±0	11 ±1	8 ±0	12 ±0	13 ±0	5 ±1	11 ±1	8 ±0	10 ±0
18:1n-7	4 ±0	3 ±0	3 ±0	3 ±0	8 ±1	7 ±0	4 ±0	4 ±0	2 ±1	5 ±0	7 ±0	8 ±0
18:2n-6	3 ±0	3 ±0	4 ±0	5 ±0	5 ±0	4 ±0	3 ±0	3 ±0	2 ±1	4 ±0	3 ±0	4 ±0
18:3n-6	0 ±0	0 ±0	1 ±0	1 ±0	1 ±0	0 ±0	0 ±0	0 ±0	1 ±0	2 ±1	0 ±0	0 ±0
18:3n-3	6 ±0	6 ±0	4 ±0	6 ±0	6 ±1	5 ±0	5 ±0	5 ±0	2 ±1	4 ±0	4 ±0	5 ±0
18:4n-3	10 ±1	10 ±1	10 ±1	12 ±0	11 ±1	9 ±0	8 ±1	9 ±0	4 ±1	8 ±1	7 ±1	8 ±0
20:2n-6	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0
20:3n-6	0 ±0	0 ±0	1 ±0	1 ±0	1 ±0	1 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0
22:0	1 ±0	1 ±0	0 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	2 ±1	1 ±0	1 ±0
20:5n-3	1 ±0	1 ±0	0 ±0	0 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	2 ±0	0 ±0	1 ±0
23:0	0 ±0	0 ±0	1 ±0	1 ±0	0 ±0	0 ±0	0 ±0	1 ±0	1 ±0	2 ±1	0 ±0	0 ±0
24:0	0 ±0	0 ±0	1 ±0	1 ±0	1 ±0	1 ±0	0 ±0	0 ±0	1 ±0	2 ±1	1 ±0	1 ±0
22:5n-3, 24:1n-9	1 ±0	1 ±0	0 ±0	1 ±0	1 ±0	1 ±0	1 ±0	1 ±0	0 ±0	0 ±0	1 ±0	1 ±0
22:6n-3	30 ±1	30 ±0	21 ±2	26 ±1	31 ±4	25 ±1	23 ±4	25 ±2	8 ±3	17 ±3	21 ±1	25 ±1
Unidentified	6 ±0	6 ±0	4 ±0	5 ±0	9 ±1	7 ±0	5 ±1	5 ±0	2 ±1	4 ±0	6 ±0	7 ±0

Σ SFAs ^a	27 ±1	27 ±0	25 ±2	30 ±1	39 ±3	31 ±2	30 ±0	32 ±2	16 ±2	39 ±4	24 ±2	29 ±1
Σ MUFAs ^b	16 ±0	16 ±0	11 ±1	13 ±0	21 ±2	17 ±0	17 ±1	19 ±0	8 ±2	18 ±1	17 ±1	21 ±1
Σ PUFAs ^c	51 ±1	51 ±0	42 ±4	51 ±1	56 ±7	45 ±1	41 ±5	44 ±2	17 ±5	39 ±3	36 ±2	43 ±0
Σ TFA ^d	100 ±1		81 ±7		125 ±12		93 ±6		42 ±9		82 ±5	

Table S5. Results of the selected GLMMs testing for the effects of temperature, N:P supply ratios and $p\text{CO}_2$ on population yield and production of particulate organic carbon (POC) and particulate inorganic carbon (PIC), and fatty acid contents in *Emiliana huxleyi*. Significant p values are shown in bold. T: temperature; N:P: N:P supply ratio; TFA: total fatty acid; MUFA: monounsaturated fatty acid; PUFA: polyunsaturated fatty acid; DHA: docosahexaenoic acid.

Variable	Factor	Coefficient \pm SE	t	p
POC population yield ($\mu\text{g ml}^{-1}$)	Intercept	13.456 \pm 1.007	13.360	<0.001
	T	-0.096 \pm 0.047	-2.045	0.046
	$p\text{CO}_2$	<0.001 \pm <0.001	-0.361	0.719
	N:P	-0.035 \pm 0.010	-3.436	0.001
POC production ($\text{pg cell}^{-1} \text{d}^{-1}$)	Intercept	-0.261 \pm 0.101	-2.587	0.013
	T	0.023 \pm 0.005	4.895	<0.001
	$p\text{CO}_2$	<0.001 \pm <0.001	1.631	0.109
	N:P	0.007 \pm 0.001	6.899	<0.001
PIC population yield ($\mu\text{g ml}^{-1}$)	Intercept	6.922 \pm 0.968	7.149	<0.001
	T	0.201 \pm 0.045	4.442	<0.001
	$p\text{CO}_2$	-0.002 \pm <0.001	-8.955	<0.001
	N:P	-0.034 \pm 0.010	-3.404	0.001
PIC production ($\text{pg cell}^{-1} \text{d}^{-1}$)	Intercept	-0.689 \pm 0.105	-6.581	<0.001
	T	0.047 \pm 0.005	9.589	<0.001
	$p\text{CO}_2$	<0.001 \pm <0.001	-5.294	<0.001
	N:P	0.007 \pm 0.001	6.339	<0.001
TFA content ($\mu\text{g mg}^{-1} \text{C}^{-1}$)	Intercept	202.099 \pm 17.745	11.389	<0.001
	T	-3.444 \pm 0.827	-4.164	<0.001
	$p\text{CO}_2$	-0.014 \pm 0.004	-3.038	0.004
	N:P	-0.188 \pm 0.182	-1.033	0.307
SFA content ($\mu\text{g mg}^{-1} \text{C}^{-1}$)	Intercept	58.540 \pm 5.265	11.119	<0.001
	T	-0.978 \pm 0.245	-3.986	<0.001
	$p\text{CO}_2$	-0.003 \pm 0.001	-2.240	0.030
	N:P	-0.118 \pm 0.054	-2.182	0.034
MUFA content ($\mu\text{g mg}^{-1} \text{C}^{-1}$)	Intercept	53.910 \pm 4.324	12.468	<0.001
	T	-1.361 \pm 0.202	-6.755	<0.001
	$p\text{CO}_2$	-0.002 \pm 0.001	-1.882	0.066
	N:P	-0.074 \pm 0.044	-1.675	0.100
PUFA content ($\mu\text{g mg}^{-1} \text{C}^{-1}$)	Intercept	71.361 \pm 7.854	9.086	<0.001

	T	-0.664 ± 0.366	-1.813	0.076
	<i>p</i> CO ₂	-0.007 ± 0.002	-3.626	0.001
	N:P	-0.024 ± 0.081	-0.292	0.772
DHA content (µg mg ⁻¹ C ⁻¹)	Intercept	36.201 ± 5.156	7.021	<0.001
	T	-0.248 ± 0.240	-1.031	0.308
	<i>p</i> CO ₂	-0.004 ± 0.001	-3.034	0.004
	N:P	0.021 ± 0.053	0.392	0.697

Table S6. The changes in cellular elemental contents (as pg cell^{-1}), population yield and production of particulate organic carbon (POC) and particulate inorganic carbon (PIC) (as $\mu\text{g ml}^{-1}$ and $\text{pg cell}^{-1} \text{d}^{-1}$, respectively), elemental molar ratios, and the proportions and contents of major fatty acid groups and docosahexaenoic acid (DHA) (as % of total fatty acids and $\mu\text{g mg C}^{-1}$, respectively) in response to warming, N and P deficiency and enhanced $p\text{CO}_2$ in *Emiliana huxleyi*. Here, only significant changes are shown based on GLMM results in Table 1 and Table S5. Red and blue arrows indicate a mean percent increase and decrease in a given response, respectively. SFAs, saturated fatty acids; MUFAs, monounsaturated fatty acids; PUFAs, polyunsaturated fatty acids.

Response	Effect				Interactions
	Warming	-N	-P	Enhanced $p\text{CO}_2$	
POC cellular content	↓ -8%	↓ -39%	↑ 50%	-	T×N:P supply
PON cellular content	↑ 5%	↓ -53%	↑ 52%	-	T×N:P supply
POP cellular content	↑ 9%	↓ -32%	↓ -8%	↑ 29%	T×N:P supply T×CO ₂
PIC cellular content	↑ 28%	↓ -31%	↑ 65%	↓ -36%	T×N:P supply T×CO ₂
POC population yield	↓ -6%	↓ -11%	↓ -23%	-	
POC production	↑ 68%	↓ -44%	↑ 63%	-	
PIC population yield	↑ 36%	↓ -48%	↑ 1%	↓ -24%	
PIC production	↑ 161%	↓ -37%	↑ 79%	↓ -35%	
POC:PON	↓ -6%	↑ 33%	-	-	
POC:POP	-	↓ -15%	↑ 60%	-	
PON:POP	-	↓ -36%	↑ 62%	-	
PIC:POC	↑ 41%	-	-	↓ -49%	
SFA proportion	-	↓ -7%	↓ -15%	↑ 7%	N:P supply×CO ₂
MUFA proportion	↓ -20%	-	-	↑ 7%	
PUFA proportion	↑ 13%	-	-	↓ -7%	
DHA proportion	↑ 16%	↑ 14%	↑ 22%	↓ -7%	T×N:P supply T×CO ₂
SFAs content	↓ -21%	↑ 36%	↑ 11%	↓ -14%	
MUFAs content	↓ -35%	-	-	-	
PUFAs content	-	-	-	↓ -24%	
DHA content	-	-	-	↓ -24%	


 Changes $\geq 25\%$

 Changes $< 25\%$
 - No significant change

Fig. S1




Fig. S1 Time course of pH (mean \pm SE) under three temperature, three N:P supply ratios and two target $p\text{CO}_2$ levels (low CO_2 : 560 μatm ; high CO_2 : 2400 μatm) in the semi-continuous cultures of *Emiliana huxleyi*.

Fig. S2




Fig. S2 Responses of the proportion of saturated fatty acids (SFAs) (mean \pm SE) to N:P supply ratios and $p\text{CO}_2$ in *Emiliana huxleyi*.

Fig. S3




Fig. S3 Responses of (a, d) cellular contents of particulate inorganic carbon (PIC), (b, e) PIC population yield ($\mu\text{g ml}^{-1}$) and (c, f) PIC production ($\text{pg cell}^{-1} \text{d}^{-1}$) (mean \pm SE) to temperature, N:P supply ratios and $p\text{CO}_2$ in *Emiliana huxleyi*. For cellular PIC content, the selected model contains the first order effects and second order interactions of the three environmental factors, while those for PIC population yield and production contain only the first order effects. The results of AICc are shown in Table S2.

Fig. S4




Fig. S4 Responses of (a, d) cellular contents of particulate organic carbon (POC), (b, e) POC population yield and (c, f) POC production (mean \pm SE) to temperature, N:P supply ratios and $p\text{CO}_2$ in *Emiliana huxleyi*. For cellular POC content, the selected model contains the first order effects and second order interactions of the three environmental factors, while those for POC population yield and production contain only the first order effects. The results of AICc are shown in Table S2.