

TRANSYLVANIAN REVIEW

VoL XXIV, No. 13, 2017



Transylvanian Review

Centrul de Studii Transilvane | str. Mihail Kogalniceanu nr. 12-14, et.5, Cluj-Napoca

Email: transylvanianreview@gmail.com /

Online Submission System: <http://transylvanianreviewjournal.org/>

New Method for Increasing Sustainable Agricultural Yield

¹ F. Leiva Gea F, ¹ A. Cano Ortiz, ² C.M. Musarella, ¹ J.C. Piñar Fuentes, ³ C.J. Pinto Gomes and ^{*1} E. Cano

¹Department of Animal Biology, Plant and Ecology, Botany, University of Jaén, Jaén, Spain, ²Department of AGRARIA, University "Mediterranea" of Reggio Calabria, Località Feo di Vito, 89122 Reggio Calabria, Italy and ³Department of Landscape, Environment and Planning/Institute of Agricultural and Environmental Sciences. Mediterranean (ICAAM), University of Évora (Portugal), Rua Romão Ramalho, Portugal.

Abstract

The main objective of this study is to look for a new method, to obtain the maximum production, and the minimum economic and environmental cost.

Scientific advances in bioclimatology have brought a greater understanding of the functioning of plant individuals, populations and communities, and the study and interpretation of bioclimatic indices reveals the close relationship between the values of these indices and crop yields.

In cases where insufficient meteorological data are available to allow the various formulas to be applied, thermoclimatic and ombroclimatic bioindicators must be used, some of which have been collected by us in previous works.

This paper contains a bioclimatic study of the southern Iberian Peninsula in relation to the main olive varieties grown in these territories. We establish the values of some key indices for cultivation: continentality index (Ic), which determines the annual thermal range and allows us to establish the period of vegetative activity (PAV); the ombrothermic index (Io), which measures the ombrotype in the different sites; and the thermicity and compensated thermicity index, which denote the thermotype in the territory (It/Itc). In the relationship between the value of the bioclimatic indices and olive production, good fits are obtained in the regression analysis with $R^2 > 0.90$, and is very positive influence of the different parameters used.

In conclusion the greatest influence on the production of Ic is between 18.5-19.5, Io 3.5-4.5 and It / Itc 320-350. This allows to obtain areas of maximum production with a lower economic cost.

Keywords: Bioclimatology, olive cultivation, yield, sustainable.

* Corresponding author: ecano@ujaen.es. Department of Animal Biology, Plant and Ecology, Botany, University of Jaén, Jaén, Spain,

Introduction

Advances in bioclimatic research have brought a greater understanding of the functioning of plant individuals, populations and communities. By applying bioclimatic indices, Rivas-Martínez and Loidi (1999) were able to establish a close relationship between bioclimatic data and plant communities, thus enabling the characterisation of each territory. However, these bioclimatic approaches are also applicable to different crops. The correlation between the indices for the Mediterranean macrobioclimate and the distribution of olive plantations clearly highlights the typically Mediterranean character of *Olea europaea* L. In the southern Iberian Peninsula, olive cultivation covers vast areas and has major socio-economic importance. The main varieties are Picual, Cornicabra, Hojiblanca, Lechin, Morisca, Manzanilla, Gordal and Verdial.

Under this climate classification, almost all the olive plantations are located in areas with a Mediterranean pluviseasonal oceanic ($Ic < \text{or} = 24, I_o > 2.0$), Mediterranean pluviseasonal continental ($Ic > 24, I_o > 2.2$) or Mediterranean continental xeric ($Ic > 24, 1.0 < I_o < 2.2$)

bioclimate: there are very few olive plantations under the Mediterranean oceanic xeric bioclimate ($Ic < \text{or} = 24, 1.0 < I_o < 2.2$).

Practically all the olive plantations are located in bioclimatic belts with a thermomediterranean ($I_{tc} 350-450$) or mesomediterranean ($I_{tc} 240-350$) thermotype, and a semiarid ($I_o 1.0-2.0$), dry ($I_o 2.0-3.6$) or subhumid ($I_o 3.6-7.0$) ombroclimate. Only in isolated cases do we find non-extensive cultivations in the suprasediterranean ($I_{tc} 80-240$), and exceptionally in the humid mesomediterranean.

There is scarce representation of the suprasediterranean thermotype due to the relatively low presence of olive plantations in this thermotype. These sites have between 4 to 6 months of frost and can therefore be regarded as unproductive, even causing trees to die due to excessively low temperatures, as occurred in 2005 (Figure 1). This year had a very high number of days with temperatures below $-10\text{ }^{\circ}\text{C}$, and thus any plantations that were not in their bioclimatic optimum were affected by frost. This occurs in upper mesomediterranean and suprasediterranean areas, and even in enclosed valleys where thermal inversion causes the territory to behave as upper meso- or suprasediterranean.



Fig. 1: Olive crop showing frost damage caused by failure to take into account the bioclimatic features of the territory

Although various factors have conditioned the cultivation of the different olive varieties in each zone, they are not always located in their ecological optimum from the bioclimatic point of view. Recent studies (Cano *et al.*, 2001a; Ighbareyeh *et al.*, 2014a,b,c) have highlighted the influence of bioclimatology on olive yields; however this is the first time the bioclimatic characterisation of the different varieties has been undertaken. One of the aims of the present study is to relate the distribution of the

different olive varieties with various bioclimatic indices, and characterise each one according to its bioclimatic requirements. The goal is to contribute new criteria for the agricultural planning of olive cultivation. This also requires biogeographical and soil studies of potential sites for olive cultivation. All this information can be used to propose models for agricultural management, according to Cano *et al.* (2003). The influence of climate change on agriculture (Peters *et al.*, 2014; Kumar *et al.*, 2016) is mitigated by the

application of agricultural management bioclimatology. With this work we correlate the science of Bioclimatology with agricultural crops, and demonstrate the influence of bioclimatic factors on production. A study that completes the Fakhar-u-Zaman Akhtar *et al.* (2016) experiment with the application of nutrients. Which together with the studies on poverty and its relation with the greater or lesser agricultural growth (Abdul Rehman *et al.*, 2016a, b) justify this work.

Material and Methods

Initially we developed bioclimatic index territories where there is olive cultivation in southern Spain.

The present work is based on data from 57 meteorological stations, selected due to their location either in or near areas of olive cultivation, or –in a few cases– far from olive plantations. We used the following formulas to calculate the values of the various indices (Rivas Martínez, 1996), (Rivas Martínez and Loidi, 1999): ombrothermic index, $I_o = P_p/T_p$; bi-monthly summer ombrothermic index, $I_{s2} = P_{July+August}/T_{July+August}$; tri-monthly summer ombrothermic index, $I_{s3} = P_{June+July+August}/T_{June+July+August}$; and continentality index, $I_c = T_{max}-T_{min}$; thermicity index, or –where applicable– compensated thermicity index, $I_t/I_{tc} = (T+M+m)/10$. P_p = positive precipitation and T_p = positive temperature (in this case equivalent to annual precipitation and average annual temperature divided by 12, as all the months have an average temperature above 0°); P = precipitation of the months indicated; T = average temperature of the months indicated; T_{max} = maximum temperature of the averages of the warmest month of the year; T_{min} = minimum temperature of the averages of the coldest month of the year; T = average annual temperature; M = average of the maximum temperature of the coldest month of the year; and m = average of the minimum temperature of the coldest month of the year.

The I_{tc} is applied only to stations with $I_c > 18$, and is obtained by adding a correction factor C to the I_t , thus establishing the PAV = period of vegetative activity for each station. The I_o reveals the ombrotype and determines how much rain falls in a territory and its vegetation response; whereas the I_c shows its continentality, and the I_t/I_{tc} its thermicity. Summer ombrothermic indices are important in agriculture, as the greater the difference between I_{s2} and I_{s3} , the greater the compensation. In order to determine the predominance of particular bioclimatic parameters such as I_o , I_c and I_t/I_{tc} in a territory and establish the dominant thermotypes and ombrotypes, we grouped the meteorological stations according to the number of months with vegetative activity, as all the varieties respond well to this criterion. We defined three major areas throughout the whole territory: those where vegetative activity occurs for a) 12 months, b) 10-11 months, and c) 8-9 months. We calculated the averages of the indices mentioned and obtained the I_{om} ,

I_{cm} and $I_{t/cm}$. We also used previous data on yields collected in Cano *et al.* (1997), as well as the agricultural characteristics of varieties described by authors such as Barranco *et al.* (1998) and Guerrero (1994). We compared each variety with the value of the indices expressed, and contrasted the cultivation with each vegetation series (Rivas Martínez, 1987), (Rivas Martínez *et al.* 2014). To test the influence of bioclimatic parameters (P_p , I_t / I_{tc} , I_c , I_o) and potential evapotranspiration (ETP) in the production of olive, do a statistical multiple regression analysis. For this face mean values (I_{it} / I_{tc} , I_c , I_o) from an area with the productions.

The interpretation of vegetation based on climate domains (sigmetum) is very important for the subsequent treatment of the vegetation. In order to implement agricultural or any other types of actions, it is essential to have a thorough knowledge of the vegetation series, the catenal contacts between the different landscape units, and the vegetation dynamic. Vegetation dynamic is understood as the successions that occur in the plant cover with the appearance of the initial, intermediate, transitional and final stages. The final stage is reached at the point of the stable biological maximum in harmony with itself and with the ecological factors in the environment. Braun-Blanquet (1979) distinguishes between progressive successions directed towards the climax, and regressive successions which move away from the climax. Regressive successions are normally caused by human and animal intervention, and also by environmental factors. Climatophilous series depend on the general characteristics of the environment – particularly the territorial microclimate–, and the location of the olive plantation (Cano and Cano Ortiz, 2016).

Resultados

Table 1 was created with data from 57 meteorological stations, obtaining values for T , P , I_o , I_c , I_t/I_{tc} , T_{max} , T_{min} , Alt (altitude), I_{s2} , I_{s3} and PAV (period of vegetative activity) for each one. The analysis of the climate and bioclimatic parameters revealed that most of the territory in the study has 12 months of PAV, with no interruption due to cold –considered as occurring when the average monthly T falls below 7.5 °C (Montero Burgos and González Rebollar, 1983). This precisely coincides with the territories lying more to the south and south-west of the province, which have mainly a thermomediterranean thermotype with $I_t/I_{tcm} = 346$, and a dry ombrotype with values of $I_{om} = 2.81$ and $I_{cm} = 49.36$. The sites with a PAV of 8-9 months are located predominantly in the northwest of the province of Jaén, with the particularity that both territories have a upper mesomediterranean thermotype with $I_t/I_{tcm} = 264.5$, and an I_{cm} (average value of the I_c) of 19.43. However the value for I_{om} (average value of I_o) is 4.42, due to the fact that the mountain ranges of Segura, Las Villas and Cazorla –among others– act as a screen against low pressure areas, causing the stations to have a high I_o . However (Table 1) shows there are stations with

Table 1: General table of stations and bioclimatic indices. P. Average annual precipitation. Io. Ombrothermic index. Ic. Continentality index (annual thermic index). It/Itc. Thermicity index/Compensated thermicity index. Alt. Altitude. Is2, Is3. Compensated ombrothermic indices. PAV. Period of vegetative activity.

Meteorological station	P	Io	Ic	It/Itc	Alt	Is2	Is3	Is3/Is2	PAV (months)
Vadillo-Castril (J)	1182.3	8.42	17.6	199	970	0.48	1.10	2.29	8
S.Pontones (J)	1148.7	6.98	18.0	214	740	0.53	0.92	1.73	9
La Iruela (J)	850.1	4.78	19.2	343	933	0.27	0.71	2.62	10
B. Moraleda (J)	612.1	3.33	19.7	370	887	0.36	0.66	1.83	10
B. Segura-Perales (J)	612.7	3.52	20.7	253	760	0.43	0.64	1.48	9
Siles (J)	785.7	4.33	21.6	260	826	0.39	0.75	1.92	9
V. del Arzobispo (J)	698.2	3.54	19.4	297	685	0.23	0.54	2.34	12
Villacarrillo (J)	610.8	3.26	20.4	284	794	0.26	0.52	2.00	10
P. del Dañador (J)	612.3	3.75	18.8	260	700	0.40	0.73	1.85	9
B. E.-Centenillo (J)	679.1	3.87	19.4	271	824	0.18	0.45	2.50	9
P. Rumblar (J)	657.4	3.28	18.6	353	300	0.18	0.45	2.50	9
Torredonjimeno (J)	648.9	3.25	19.7	329	591	0.33	0.55	1.66	12
P. Tranco de Beas (J)	849.6	4.65	18.2	298	600	0.41	0.81	1.97	10
Beas de Segura (J)	658.0	3.20	19.7	322	577	0.30	0.58	1.93	12
Cazorla. ICONA (J)	792.4	4.68	19.2	257	885	0.32	0.74	2.31	9
Huelma-Solera	526.9	3.20	20.8	293	1084	0.40	0.74	1.85	9
P. de la Bolera (J)	653.7	3.83	17.0	284	980	0.50	0.90	1.80	10
Jimena (J)	600.3	3.20	20.0	303	590	0.32	0.69	2.15	11
A. Real- Charilla (J)	681.0	4.05	17.3	270	920	0.42	0.68	1.61	10
P. del Jándula (J)	505.2	2.50	19.5	339	360	0.17	0.41	2.43	12
Arjona (J)	609.9	2.97	19.5	406	410	0.25	0.41	1.64	12
Bailen (J)	581.7	2.70	20.1	369	369	0.14	0.32	2.34	12
Andújar (J)	463.9	2.13	19.1	371	212	0.17	0.28	1.66	12
Jaén. Instituto (J)	578.3	2.85	18.8	345	510	0.20	0.40	2.00	12

Transylvanian Review: Vol XXIV, No. 13, 2017									
Gea F <i>et al.</i> Linares (J)	642.2	3.12	18.3	339	419	0.21	0.52	2.48	12
L. Torrubias (J)	490.4	2.34	20.3	355	290	0.14	0.31	2.26	12
Mancha Real (J)	551.6	3.02	18.1	299	753	0.37	0.69	1.87	11
P. Guadalmena (J)	517.9	2.74	20.0	308	602	0.37	0.55	1.48	11
L. Fuente Higuera (J)	471.0	2.36	18.6	342	300	0.15	0.30	2.00	12
Ubeda (J)	579.6	3.03	18.4	313	748	0.22	0.44	2.00	11
La P. de Segura (J)	674.7	3.55	19.9	305	584	0.35	0.64	1.82	10
Cabra de s. Cristo (J)	449.9	2.62	17.7	275	938	0.35	0.65	1.86	10
Ubeda. P. Guadiana (J)	404.6	2.24	19.6	285	420	0.23	0.46	2.00	9
P. Guadalmeñato (CO)	698.4	3.42	18.0	349	200	0.40	0.64	1.60	12
C. La Jarosa (CO)	831.1	4.46	16.5	315	340	0.25	0.31	1.24	12
Pantano P. Nuevo (CO)	760.6	4.03	17.9	321	410	0.16	0.59	3.68	12
Villaralto (CO)	501.4	2.66	18.8	310	583	0.42	0.60	1.42	11
Pozoblanco (CO)	514.4	2.66	19.9	311	649	0.34	0.66	1.94	10
Aldea de Cuenca (CO)	559.0	2.98	18.1	307	571	0.23	0.67	2.91	12
H. del Duque. (CO)	476.9	2.68	18.4	284	540	0.42	0.68	1.61	11
Pedroche (CO)	506.8	2.60	19.2	322	621	0.38	0.63	1.65	12
La Rambla (CO)	527.4	2.81	17.9	308	200	0.30	0.41	1.37	12
Castro del Río (CO)	470.7	2.45	20.1	305	210	0.17	0.36	2.12	11
Montoro (CO)	572.4	2.72	18.1	360	195	0.19	0.37	1.94	12
Pozoblanco. Cerro (CO)	594.7	2.98	19.0	339	500	0.32	0.52	1.62	12
C. Miraflores(CO)	634.4	2.81	18.8	394	120	0.24	0.44	1.83	12
C. Aeropuerto (CO)	630.2	3.00	17.6	365	92	0.14	0.34	2.42	12
Baena. S.E.A. (CO)	519.5	2.60	20.7	326	463	0.25	0.39	1.56	11
P. Pueblonuevo (CO)	524.1	2.49	19.9	356	550	0.16	0.42	2.62	12
P. de Baena (CO)	598.6	2.93	18.2	345	150	0.16	0.37	2.31	12
Posadas (CO)	686.9	3.23	18.4	359	88	0.30	0.50	1.66	12
P. de Bembezar (CO)	667.5	3.21	17.7	363	100	0.24	0.51	2.12	12

Lucena. C. Malpa. (CO)	410.0	2.07	16.8	345	280	0.26	0.44	1.69	12
Carcabuey. Algar (CO)	795.8	4.77	16.3	291	740	0.26	0.60	2.30	12
Rute. S. E. A (CO)	636.4	3.31	17.6	327	639	0.27	0.47	1.74	12
Lucena. CdeA (CO)	679.4	3.37	18.1	344	586	0.21	0.46	2.19	12
Puente Genil. C. (CO)	435.2	2.02	17.9	373	200	0.24	0.39	1.62	12

The interpretation of Is2 is particularly important in this analysis of the bioclimatic parameters affecting olive cultivation, as it has low values compared to Is3. This is due to the June rains, which is compensated plant cover in stations where Is3 is very high compared to Is2, thereby reducing water stress. Evidently the higher the ratio of Is3/Is2, the greater the compensation; however this does not imply that waters stress disappears, as the rains may be abundant during winter and spring and minimal in June, and not offset the deficit in July and August. Thus for example in Table 1 the Santiago-Pontones station has rainfall of 1148, Is2 = 0.27 and Is3 = 0.74, with an Is3/Is2 ratio of 2.62; we can contrast this with the Pantano de Guadalmena station, which has Is2 = 0.37 and Is3 = 0.55, and a ratio of Is3/Is2 = 1.48. For these situations it is important to take into account another parameter of vital importance in the management of olive plantations: the ETP (potential evapotranspiration) and the CR (soil water retention capacity), which –as we highlighted in Cano *et al.* (2001b) – depends fundamentally on texture, structure, organic material and

in order to determine the influence of the parameters (Pp, ETP, It / Itc, Ic, Io) in the production of olive cultivation, face values of these parameters olive production. We use three types of regression to see which best fits the data.

For linear regression, the value R^2 is below 0.90 in the case of ETP and Io, and logarithmic regression with only have value less than 0.90 if the parameter ETP (Table 2). However in all cases R^2 is greater than 0.90 when we apply a polynomial fit with a very high setting in the three types of regression in the case of Ic, demonstrating the influence of the parameters used in the production

For positive precipitation Pp is practically linear adjustment, and R^2 is greater than 0.90 in all three types of regression. For potential evapotranspiration ETP polynomial regression analysis with value $R^2 > 0.90$ fits better (Figure 2). For It / Itc with $R^2 > 0.90$ the impact on production is negative when It / Itc < 300, and very positive when It / Itc ranges from 340-360 (upper thermomediterranean and lower mesomediterranean). It is also very positive influence on the production continentality index Ic, which fits very well in the three types of regression $R^2 = 0.9989$ values very close to 1 (Figure 3). The Io ombrothermic index has a great influence on production, fitting well in the regression analysis polimomial $R^2 = 0.9733$; existing low productions when the value of Io < 3.5, these productions are increased to values between 3.5 and 4.5 Io, and are depleted for values of Io > 4.5 (Figure 4).

Table 2: - Values of the coefficient of linear, logarithmic and polynomial regression

	R^2 (Linear)	R^2 (logarithmic)	R^2 (Polynomial)
Pp	0,9314	0,9518	0,9669
ETP	0,8602	0,8369	0,9039
It/Itc	0,9494	0,9321	0,982
Ic	0,9976	0,9981	0,9989
Io	0,8631	0,909	0,9733
			0,90 < R^2 < 1




Fig. 2: - Multiple regression analysis. Positive precipitation and potential evapotranspiration. Pp and ETP: Ppolynomial regression (Pp) $y = - 2E-0.5x^2 + 0.0293x - 12.137$ ($R^2 = 0.9669$). Linear regression (Pp) $y = 0.0069x - 4.466$ ($R^2 = 0.9314$). Logarithmic regression (Pp) $y = 4.8457\ln(x) - 31.307$ ($R^2 = 0.9518$). Ppolynomial regression (ETP) $y = 2E-0.5x^2 - 0.0286x + 6.8911$ ($R^2 = 0.9038$). Linear regression (ETP) $y = 0.0095x - 8.4747$ ($R^2 = 0.8602$). Logarithmic regression (ETP) $y = 7.6236\ln(x) - 51.736$ ($R^2 = 0.8369$).




Fig. 3: - Multiple regression analysis. Thermicity and continentality index It/Ic and Ic: Ppolynomial regression (It/Ic) $y = 0.0002 x^2 - 0.0709 x + 6.2259$ ($R^2 = 0.982$). Linear regression (It/Ic) $y = 0.0238 x - 7.2948$ ($R^2 = 0.9494$). Logarithmic regression (It/Ic) $y = 6.7967\ln(x) - 38.85$ ($R^2 = 0.9321$). Ppolynomial regression (Ic) $y = - 0.2799x^2 + 12.906x - 144.5$ ($R^2 = 0.9989$). Linear regression (Ic) $y = 2.2334x - 42.806$ ($R^2 = 0.9976$). Logarithmic regression (Ic) $y = 42.581\ln(x) - 125.74$ ($R^2 = 0.9981$).




Fig. 4: Multiple regression analysis. Ombrothermic index I_o . Pplynomial regression $y = -0.4304 x^2 + 4.4035 x - 9.7441$ ($R^2 = 0.9733$). Linear regression $y = 0.7338 x - 2.6448$ ($R^2 = 0.8634$). Logarithmic regression $y = 3.0762 \ln(x) - 3.819$ ($R^2 = 0.909$).

Discussion

The results reveal a great correlation between production and bioclimatic parameters, these bioclimatic parameters also condition the vegetation series present in the territory. Consequently, each type of vegetation corresponds to a type of crop, which Cano and Cano Ortiz (2016) establishes for the cultivation of olive trees.

The interpretation of vegetation based on climate domains (sigmetum) is very important for the subsequent treatment of the vegetation. In order to implement agricultural or any other types of actions, it is essential to have a thorough knowledge of the vegetation series, the catenal contacts between the different landscape units, and the vegetation dynamic. Climatophilous series depend on general features of the environment and act as precise indicators for the optimisation of agricultural cultivation. In this study we propose a set of series that are suitable for olive cultivation, and others that are not. * Series where olive cultivation can occur. ** Series where olive cultivation is optimum (Cano and Cano Ortiz, 2016).

Based on the studies conducted by Cano *et al.* (2007), olive varieties respond to different values in the various bioclimatic indices, and are located in the area of a particular vegetation series. Cano *et al.* (2003) thus propose a specific model for agricultural and forestry management.

The Picual variety has its optimum in the upper thermomediterranean and lower mesomediterranean belt with values of $I_o = 3.6-4.6$, $I_c = 18-20$ and $I_t/I_c = 280/400$. It is somewhat tolerant of frost, and thus grows well in the

lower mesomediterranean, where frosts occur for 1-2 months. However as this variety has a good tolerance for soil humidity and is sensitive to drought, it should be grown in areas with an ombrotype ranging from dry to subhumid. This variety is not particular about soil, but grows poorly on sites with a high proportion of clay (vertisols), as these soils have a low CR and a high ETP, and behave as though they were semiarid unless soil management practices are applied to reduce the ETP. It does not grow well in places with an upper subhumid and humid ombrotype; the I_o should not be above 4.6, as in these environments it is easily attacked by peacock spot – *Spilocaea oleagina*, known locally as "repilo"–, and thus plantations in these territories require a significant economic and environmental cost to maintain their yield. This variety is widespread in the provinces of Jaén, Cordoba and Granada. To achieve its ecological optimum it should be sited in areas occupied by the following series *Pyro bourgaeanae-Quercus rotundifoliae s. thermophilous faciation with Myrtus communis, Paeonio coriaceae-Quercus rotundifoliae s. thermophilous faciation with Pistacia lentiscus, Viburno tini-Queco fagineae s, Myrto communis-Quercus rotundifoliae s, Rhamno-Quercus rotundifoliae s.*

The Cornicabra variety has its optimum in the upper mesomediterranean belt extending into the lower supramediterranean, with values of $I_o = 2.6-3.6$, $I_c = 20-22$ and $I_t/I_c = 210/280$. It is resistant to cold, and therefore grows well with 3-4 months of frosts. It withstands drought well and has a dry ombrotype. Because it is sensitive to peacock spot it should not be located in places with an I_o of over 3.6. This variety is widely grown in Ciudad Real,

Gea F *et al.*

Toledo and parts of Albacete. Its optimum is in the series 1) *Pyro-Quercus rotundifoliae s.*) *Quercus rotundifoliae s.*

The Hojiblanca variety has its optimum in the upper thermomediterranean with a dry ombrotype, and with $I_o = 2.6-3.6$, $I_c = 17-18$ and $I_t/I_{tc} = 350/400$. It does not tolerate cold well, and should therefore be grown on sites with a maximum of 1-2 months of frost. It may extend to the lower mesomediterranean. It grows well in dry ombrotypes, and does not resist $I_o > 3.6$ as it is sensitive to peacock spot, tuberculosis and verticilosis. Although it grows well in most soils, it prefers limy substrates. Its optimum is in the series 1) *Rhamno-Quercus rotundifoliae s.*) *Myrto communis-Quercus rotundifoliae s.*, and also in *Paeonio-Quercus rotundifoliae s. thermophilous faciation with Pistacia lentiscus.*

Lechin, this variety is widespread throughout the provinces of Seville, Cordoba and Cadiz, and has its optimum in the dry-subhumid thermomediterranean with $I_o = 4-6$, $I_c = 15-17$ and $I_t/I_{tc} = 350/450$. It is sensitive to frost, and should therefore be located in frost-free sites or where frosts occur for no more than one month. As it is very resistant to peacock spot, it grows well in subhumid and even humid thermomediterranean locations. It should be located in the thermomediterranean cork oak and Portuguese oak series of *Myrto communis-Quercus suberis s.*, *Oleo-Quercus broteroi s.*, *Rhamno-Quercus rotundifoliae s.*, *Myrto-Quercus rotundifoliae s.* and in mesophytic faciations of holm oak.

The Morisca variety has its optimum in the dry ombrotype and upper thermomediterranean thermotype, and may extend to the lower mesomediterranean. It cannot withstand extreme cold, but can resist a maximum of one month's frost when located in places with values of $I_o = 2.6-3.6$, $I_c = 16-18$, and $I_t/I_{tc} = 280/400$. This plant tolerates drought and limy soils, and therefore grows well in the region of Tierra de Barros (BA), where there is a degree of basicity, and a pH of nearly 7 (neutral). It should be located in the holm oak series of *Pyro-Quercus rotundifoliae s. basophilous faciation and thermophilous faciation.* This variety can also be grown in the area occupied by the series of *Myrto-Quercus rotundifoliae s.*

Manzanilla, this is an eating variety that grows in areas of Seville and Badajoz, extending towards territories in Cáceres, where it is known by the name of "manzanilla cacereña". It is sensitive to frost, and has a thermomediterranean optimum which may extend to the lower mesomediterranean. This plant is sensitive to peacock spot, tuberculosis and verticilosis, and should therefore be located in areas with a dry ombrotype, in this case with values of $I_o = 2.6-3.6$, $I_c = 15-17$ and $I_t/I_{tc} = 280/450$. It can occupy the area of the Morisca variety and is therefore located in the same series.

Gordal, unlike the Manzanilla, this variety is resistant to cold and also requires a certain number of hours of frost to flower; its optimum is therefore mesomediterranean. It is resistant to peacock spot and can therefore be grown in the dry-subhumid ombrotype, where it requires sites with

values of $I_o = 3.6-4.6$, $I_c = 18-20$ and $I_t/I_{tc} = 210/350$. It should be grown in the area occupied by the same series as the Picual and Hojiblanca varieties. This suggests the need to reconsider the cultivation of the Gordal variety in thermomediterranean areas, as this variety is not currently located in its ecological optimum, which –among other causes– may lead to alternate fruiting.

The Verdial variety is located in lower thermo- and mesomediterranean areas in territories in Huelva, Seville, Badajoz, Malaga and Portugal, as reported by Barranco *et al.* (1998) and Guerrero (1994), where –according to the first author– different varieties of Verdial occur. They all share the fact that their fruits do not turn totally black, except in the case of the Verdial from Vélez-Málaga, whose high olive oil yield may make it attractive to cultivate. In general terms it can be said to have its optimum in places with $I_o = 2.6-3.6$, $I_c = 12-15$ and $I_t/I_{tc} = 280-450$, and is present in practically the same series as the Morisca and Manzanilla varieties. As it is sensitive to frost and peacock spot, its cultivation should be planned in thermomediterranean sites with a dry ombrotype.

Arbequina, this cultivar has a mesomediterranean optimum with values of $I_t/I_{tc} = 210-350$, $I_c =$ or > 24 , and $I_o =$ or > 3.6 (dry/subhumid). It is resistant to cold, has relatively underdeveloped trees, and its fruit is small and tightly attached. This is a variety that does not grow well in very limy soils, as this may cause the occurrence of iron chlorosis. It is tolerant of peacock spot and verticilosis, and therefore grows well in zones with $I_o =$ or > 3.6 .

Farga, typical of the provinces of Castellón and Valencia, this plant has its optimum in the upper thermomediterranean to lower mesomediterranean, with $I_t/I_{tc} = 350$, $I_c = 16-17$ and $I_o < 3.6$. It is not frost resistant, although it can withstand cold weather. As it is sensitive to peacock spot it is advisable to grow it in areas with a dry ombrotype.

In view of the considerable increase in the area currently occupied by olive cultivation –and as highlighted by Barranco (1998)–, it is necessary to carry out adaptation tests on all the varieties that are sited far from their habitual cultivation areas. For example in the case of the Empeltre variety –whose cultivation area has traditionally been in the Ebro Valley– in the hypothetical case of its possible introduction in Andalusia, it is worth noting that this variety does not tolerate frost and is somewhat resistant to tuberculosis and verticilosis, which in our opinion would point to the advisability of its cultivation in thermomediterranean areas with an I_t/I_{tc} of around 350, I_c values ranging between 12-15, and $I_o > 3.6$. The Arbequina variety –whose cultivation is proliferating in Andalusia– can withstand low temperatures, does not tolerate very limy soils, and is resistant to peacock spot and verticilosis. Guerrero (1994) reports that it performed well in tests and can thus be grown in Andalusia. In our opinion, these varieties should be planned on soils that are not excessively limy, in the mesomediterranean bioclimatic

Conclusions

The research carried out so far allows points to the need for national and European policies to establish scientific and technological criteria when planning olive cultivation, in order to enable sustainable development and minimise economic and environmental cost. We have therefore carried out a general bioclimatic study of areas of olive cultivation to serve as a model for sustainable development in which each variety occupies its own bioclimatic niche. This calls for the territorial ordination of traditional crops and new known botanical resources. In the case of olive cultivation, we establish the limits of bioclimatic tolerance for each of the main varieties. The bioclimatic parameters that most affect production are Ic, Io and It / Itc. This work is demonstrated the need for bioclimatic bases in agricultural management, to achieve sustainable development to increase production and quality, by placing each crop in its bioclimatic optimum.

References

Abdul Rehman, Luan Jingdong, Rafia Khatoon, Imran Hussain (2016a). Effect of Agricultural Growth on Poverty Reduction, its Importance and Suggestions. *Transylvanian Review* 7 (5):449-454

Abdul Rehman, Luan Jingdong, Muhammad Shahid Iqbal, Imran Hussain (2016b). A Study on Agricultural Development in China and its Comparison with India and Pakistan. *Transylvanian Review* 24 (6):603-610

Barranco, D., Fernández Escobar, R. and Rallo, L. 1998. El cultivo del olivo. Ed. Mundi-Prensa. Junta Andalucía. pp. 1-651.

Braun-Blanquet, J. 1979. *Fitosociología*. Ed. Blume. Madrid

Cano, E., García Fuentes, A., Torres, J.A., Salazar, C., Melendo, M., Pinto Gomes, C.J. and Valle, F. 1997. *Phytosociologie appliquée a la planification agricole*. *Colloques Phytosociologiques* 25 : 1007-1022.

Cano, E., Ruiz, L. and Cano-Ortiz, A. 2001a. Influencia de la Bioclimatología en la producción del olivar. *Aldaba* 11: 151-155.

Cano, E., Ruiz, L. and Cano-Ortiz, A. 2001b. Análisis de las técnicas de cultivo en el olivar. *Aldaba* 11:157-162.

Cano, E., Ruiz, L., Cano-Ortiz, A. and Nieto, J. 2003. Bases para el establecimiento de modelos de gestión agrícola y forestal in Memoriam al prof. Dr. Isidoro Ruiz Martínez. *Serv. Publ. Univ. De Jaén*. Pp.131-142

Cano, E., Cano-Ortiz, A. and Montilla, R.J. 2007. *Bioclimatología y olivar en la provincia de Jaén: Establecimiento de áreas de cultivo para algunas variedades de olivo*. I congreso de cultura del olivo. IEG. pp. 517-528.

Cano, E. and Cano-Ortiz, A. 2016. *Bioclimatología aplicada a agronomía in Nuevas Tendencias en Olivicultura*.

Cano-Ortiz, A. 2007. *Bioindicadores ecológicos y manejo de cubiertas vegetales como herramienta para la implantación de una agricultura sostenible*. Tesis Doctoral. Universidad de Jaén.

Cano-Ortiz, A., Del Río González, S. and Pinto Gomes, C.J. 2013. Impact of soil texture on plant communities of *Raphanus raphanistrum* L. *Plant Sociology* 50 (2): 39-46. doi:10.7338/pls2013502/02

Fakhar-u-Zaman Akhtar, M., Muhammad Munir Anjum, Azhar Hussain (2016). Impact of integrated use of inorganic phosphorus and farm yard manure for improving growth of maize (*Zea mays* L.) under desert climate of Bahawalpur, Pakistan. *Transylvanian Review* 24 (10):2736-2744

Guerrero García, A. 1991. *Nueva Olivicultura*. Ed. Mundi-Prensa. pp. 1-271

Montero Burgos, J.L. and González Rebollar, J.L. 1983. *Diagramas Bioclimáticos*. Ministerio de Agricultura, Pesca y Alimentación. ICONA. pp. 1-379.

Ighbareyeh, J.M.H., Cano-Ortiz, A. and Cano, E. 2014a. Case Study: Analysis of the Physical Factors of Palestinian Bioclimatic. *American Journal of Climate Change* 3:223-231. Published Online June 2014 in *SciRes*. <http://www.scirp.org/journal/ajcc>; <http://dx.doi.org/10.4236/ajcc.2014.32021>

Ighbareyeh, J.M.H., Cano-Ortiz, A. and Cano, E. 2014b. Biological and bioclimatic basis to optimize plant production: Increased economic areas of Palestine. *Agricultural Science Research Journal* 4 (1):10-20. Available online at <http://www.resjournals.com/ARJ>

Ighbareyeh, J.M.H., Cano-Ortiz, A., Suliemeh, A.A.A., Ighbareyeh, M.M.M. and Cano, E. 2014c. *Phytosociology with other characteristic biologically and ecologically of plant in Palestine*. *American Journal of Plant Sciences* 5:3104-3118. Published Online September 2014 in *SciRes*. <http://www.scirp.org/journal/ajps>; <http://dx.doi.org/10.4236/ajps.2014.520327>

Kumar, A., Sharma, P. And Joshe, S. 2016. Assessing the Impacts of Climate Change on Land Productivity in Indian Crop Agriculture: An Evidence from Panel Data Analysis. *J. Agr. Sci. Tech.* 18: 1-13

Peters, K., Breitsameter, L. and Gerowitt, B. 2014. Impacto of climate change on weeds in agriculture: a review. *Agrom. Sustain. Dev.* 34:707-721. doi: 10.1007/s13593-014-0245-2.

Rivas Martínez, S. 1987. *Mapa de series de vegetación de España a escala 1:400.000*. Ministerio de Agricultura, Pesca y Alimentación. ICONA. pp. 1-208.

Rivas Martínez, S. 1996. *Clasificación Bioclimática de la Tierra*. *Folia Botánica Matritensis* 16: 1-32.

Rivas Martínez, S. and Loidi Arregui, J. 1999. *Bioclimatología of the Iberian Peninsula*. *Itinera Geobotanica* 13: 41-47.

Gea F *et al.*

Rivas Martínez *et al.* (2014). Mapa de series, geoseries y geopermaseries de vegetación de España. Parte II. - Itinera Geobotanica 18 (1): 5-424.

Transylvanian Review: Vol XXIV, No. 13, 2017