



Performance of women with fibromyalgia in walking up stairs while carrying a load

Daniel Collado-Mateo¹, José C. Adsuar¹, Pedro R. Olivares^{2,3}, Francisco J. Dominguez-Muñoz¹, Cristina Maestre-Cascales¹ and Narcis Gusi¹

¹ Faculty of Sport Science, University of Extremadura, Cáceres, Spain

² Instituto de Actividad Física y Salud, Universidad Autónoma de Chile, Talca, Chile

³ Instituto Superior de Educación Física, Universidad de la República, Uruguay

ABSTRACT

Background. Fibromyalgia is a chronic disease characterized by widespread pain and other associated symptoms. It has a relevant impact on physical fitness and the ability to perform daily living tasks. The objective of the study was to analyze the step-by-step-performance and the trunk tilt of women with fibromyalgia in the 10-step stair climbing test compared with healthy controls.

Methods. A cross-sectional study was carried out. Twelve women suffering from fibromyalgia and eight healthy controls were recruited from a local association. Participants were asked to climb 10 stairs without carrying a load and 10 stairs carrying a load of 5 kg in each hand. Mediolateral trunk tilt was assessed using the “Functional Assessment of Biomechanics (FAB)” wireless motion capture device, and the time between steps was assessed via weight-bearing insoles.

Results. Trunk tilt in the stair-climbing task carrying a load was significantly higher in women with fibromyalgia when compared to the healthy controls (2.31 (0.63) vs. 1.69 (0.51) respectively). The effect of carrying a load was significantly higher for women with fibromyalgia compared with healthy controls at the intermediate and final part of the task.

Discussion. Trunk tilt during stair climbing while carrying a load was higher in women with FM, which could increase the risk of falling. Additionally, women with FM experienced a higher pace slowdown as a consequence of the load, which supports the need of including specific strength and resistance training to physical therapies for this population.

Subjects Kinesiology, Women’s Health

Keywords Fibromyalgia, Postural balance, Women, Daily living, Stair ascent, Motor control

INTRODUCTION

Fibromyalgia (FM) is a chronic disease of unknown aetiology. It is characterized by widespread pain and several associated symptoms, such as fatigue, non-restorative sleep, mood alterations, and poor health-related quality of life (HRQoL) (*Segura-Jimenez et al., 2014; Wolfe, 2015*). In Spain, approximately 4.2% of the women suffer from this disorder, whereas only 0.2% of the men are affected (*Mas et al., 2008*).

Submitted 27 October 2015

Accepted 13 January 2016

Published 1 February 2016

Corresponding author

Daniel Collado-Mateo,
dcolladom@unex.es

Academic editor

Rodrigo Ramírez-Campillo

Additional Information and
Declarations can be found on
page 8

DOI 10.7717/peerj.1656

 Copyright

2016 Collado-Mateo et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Physical fitness of women with FM has been previously studied. Among the physical impairments of FM patients, low levels of strength, endurance, flexibility, and poor balance are extremely relevant (Aparicio et al., 2015; Muto et al., 2015). In fact, it is suggested that FM may have a larger impact on physical health rather than on psychological health (Segura-Jimenez et al., 2015). In this regard, physical fitness has been associated with a number of FM symptoms, such as pain (Carbonell-Baeza et al., 2011), HRQoL (Latorre-Roman et al., 2014), anxiety and depression (Sener et al., 2013).

There is a growing interest in quantifying the impact which FM has on the performance of ADLs. The work of Huijnen et al. (2015) demonstrated that FM may have a significant impact on the performance and perception of ADLs. Specifically, the capacity to ascend and descend stairs with and without a load can be an important and physically challenging ADL of women with FM and can be potential source of injury due to falls. In this regard, a recent study also reported that women with FM have more than three times more falls than healthy controls during the period of one year (1.45 and 0.40 respectively). These women also reported higher fear of falling and lower levels of objective and perceived balance ability (Collado-Mateo et al., 2015). The circumstances of falls among women with FM were investigated by Rutledge et al. (2013). Most falls occurred at home (67%), during the day (89%) and while walking (44%) or on stairs (39%). Additionally, most of the falls on stairs occurred during stair ascent. Therefore, stair climbing could be a focus for healthcare providers.

Climbing stairs is an extremely important ADL for women with FM. The Fibromyalgia Impact Questionnaire (FIQ) (Bennett, 2005) and its Revised version (FIQ-R) (Bennett et al., 2009b) include an item asking whether patients were able to climb stairs and the difficulty that they experienced while climbing stairs, respectively. Climbing stairs can be measured through a physical test. Previous studies utilized a physical test in which patients with FM had to climb ten stairs as fast as possible (Valkeinen et al., 2004). Most of the studies utilized this method to evaluate the effects of a training program proving that physical exercise could prevent the progressive loss of performance climbing stairs weightless or carrying loads (Tomas-Carus et al., 2008). Two studies utilized it to compare the performance of FM patients and healthy controls (Huijnen et al., 2015; Valkeinen et al., 2008). These two studies concluded that patients with FM needed more time to complete the stair-climbing task. However, to our knowledge, no studies have explored the step-by-step-performance of FM patients in a stair-climbing task.

It is a well-known fact that stair-climbing is a common ADL, but this task is often performed while carrying loads. In fact, the FIQ-R included one item asking about how much FM influenced the difficulty to “lift and carry a bag full of groceries.” Previous studies with healthy people investigated the effects of carrying a load while ascending stairs or walking (Hall et al., 2013; Tseng & Liu, 2011). Among load-carrying methods, hand-held loads seems to cause major effects on the heart-rate, perceived exertion, and postural control (Tseng & Liu, 2011). Therefore, the most common load-carrying method in FM women according with FIQ-R is also the most physically demanding method. However, to our knowledge no studies have investigated the impact of carrying a load on the motor pattern of stair ascent in FM patients.

FM is related to some neurological symptoms that may cause poor balance and coordination, i.e., vision problems, ringing ears, vertigo, dizziness, etc. ([Watson et al., 2009](#)). Maintaining postural control and balance while climbing stairs is crucial to avoid falls. In this regard and to the best of our knowledge, no studies have evaluated the stability (through trunk or pelvic tilt, displacement or velocity of the center of pressure, or other measure) of women with FM in the stair-climbing task.

The aim of the current paper was to analyze the step-by-step-performance and the trunk tilt of women with FM in the 10-step stair-climbing test compared with healthy controls.

MATERIALS & METHODS

Participants

A total of 12 women with FM and eight healthy controls underwent the stair ascent tasks. All participants were recruited from a local association. Some inclusion and exclusion criteria were set: (a) be able to climb 10 stairs while carrying a load of 5 kg in each hand (participants were excluded if they tried but were not able), (b) be aged between 18 and 75 years. Additionally, women with FM were included if they had been diagnosed with FM by a rheumatologist accordingly with the criteria of the American College of Rheumatology ([Wolfe et al., 1990](#)). From a sample of 24 women, three women with FM and one healthy woman were excluded because they were not able to climb 10 stairs while carrying the load. Finally, 20 women provided written informed consent accordingly to the updated Declaration of Helsinki. The Bioethical Committee of the University of Extremadura (Spain) approved this study to be conducted (Registration number: 113/2014).

Procedure

In first place, anthropometric measurements were collected and entered in the software of the motion capture device. Participants were also measured and weighed using Tanita BC-418 MA body composition analyzer. Sensors of the motion capture device were placed across standard body locations ([Rahimi et al., 2011](#)). Once the sensors were placed and the device calibrated, participants climbed the 10 stairs without carrying a load, they rested for 3 min and then they repeated the task, but this time carrying a load of 5 kg in each hand. The riser height of each stair was 17 cm, and its tread depth was 28 cm for safety reasons ([Jackson & Cohen, 1995](#); [Tseng & Liu, 2011](#)). Finally, women with FM completed the revised version of the FM Impact Questionnaire (FIQ-R).

Materials and measures

Kinematic data was assessed using the Functional Assessment of Biomechanics™ (FAB) System (Biosyn Systems Inc., Surrey, BC, Canada). It is wireless motions capture system that consists of 13 sensors, which contain accelerometers, magnetometers, and gyroscopes. Data was collected at 100 Hz. The FAB assesses angles, positions, velocities, and accelerations of body segments. The FAB also collects foot sole pressure using weight-bearing insoles. This novel device was previously used to evaluate whole body mobility in Parkinson's disease ([Rahimi et al., 2014](#); [Rahimi et al., 2011](#)), and velocity and range of motion in cyclists ([Corbett, Peer & Ridgel, 2013](#)).

In the stair-climbing tasks, the mean of the absolute value of the medio-lateral sway of the trunk, that is, trunk tilt was assessed. This parameter is directly assessed by the FAB. Time spent between stairs was also calculated using data from weight bearing sensors. First and last stairs were excluded from the analysis as they cannot be compared with the rest because they start or finish with both feet on the ground.

In order to characterize the sample, the impact of FM was assessed using both the FM Impact Questionnaire (FIQ) and the Revised FM Impact Questionnaire (FIQ-R). FIQ is a 10-item, self-administered instrument (Bennett, 2005). The items consist on physical impairment, feel good, work missed, do job, pain, fatigue, rested, stiffness, anxiety, and depression. A consensus version for Spanish population was developed by Esteve-Vives *et al.* (2007). The FIQ was revised in 2009, originating a new questionnaire called FIQ-R (Bennett *et al.*, 2009b). The validation of the Spanish version of FIQ-R was developed by Salgueiro *et al.* (2013).

Statistical analysis

Characteristics of women with FM and healthy controls were compared using the unpaired Student's *t* test. The Kolmogorov Smirnov and Shapiro tests were used to determine normality. Normal distribution was assumed when the *p*-value was higher than .05. Differences between groups in trunk tilt were evaluated by comparing mean and median trunk tilt in the task of carrying a load and in the task without carrying a load using the independent sample Student's *t*-test or Mann–Whitney *U* test, for parametric and non-parametric variables respectively. The eight analyzed stairs were grouped into: (a) initial stairs (stairs 1, 2 and 3); (b) intermediate stairs (stairs 4 and 5); and (c) final stairs (stairs 6, 7 and 8). A load effect was calculated in the step-by-step analysis. It was defined as: Load effect (%) = (step duration while carrying a load-step duration without carrying a load)/step duration without carrying a load * 100. It represents the percentage rate for the increase as consequence of carrying a load. Load effect coefficient was compared between the two groups using the independent sample Student's *t*-test or Mann–Whitney *U* test, where appropriate. Cohen's *d* was calculated as the difference between means divided by the pooled standard deviation appropriately weighed for groups with non-equal sample sizes (Cohen, 1988). An effect size <0.5 was considered small, between 0.5 and 0.8 was considered medium, and >0.8 was considered large (Cohen, 1988).

RESULTS

Main characteristics of participants are depicted in Table 1. No significant between-group differences were observed in age, height, weight, muscular mass, fat mass, and body mass index (BMI).

As can be seen in Table 2, mean trunk tilt in the stair-climbing task carrying a load was significantly higher in women with FM compared to healthy controls (2.31 (0.63) vs. 1.69 (0.51) from midline respectively). Similar results were observed for the median trunk tilt. There were not differences in trunk tilt in the same task without carrying a load. The effect size was large (>0.8) when carrying a load and small without carrying a load.

Table 1 Main characteristics of fibromyalgia women and healthy women.

	Fibromyalgia women (<i>n</i> = 12)	Healthy women (<i>n</i> = 8)	<i>p</i> -value*
Age (years)	5.25 (9.71)	59.00 (13.72)	.741
Height (cm)	156.67 (3.03)	157.75 (4.13)	.506
Body mass (kg)	71.31 (9.24)	70.05 (7.43)	.752
Muscular mass (%)	57.26 (4.53)	57.65 (4.00)	.845
Fat mass (%)	39.71 (4.81)	39.28 (1.54)	.840
Body mass index (kg/m ²)	29.15 (4.43)	28.19 (3.39)	.610
Years since fibromyalgia symptoms started	17.42 (10.78)	NA	NA
Years since fibromyalgia diagnosis	8.75 (2.63)	NA	NA
FIQ-R function domain	13.39 (6.35)	NA	NA
FIQ-R overall Impact domain	5.83 (4.22)	NA	NA
FIQ-R t symptom domain	30.29 (8.13)	NA	NA
FIQ-R total score	49.51 (14.80)	NA	NA
FIQ total score	50.44 (11.62)	NA	NA
FIQ pain (from 0 to 10)	5.08 (3.03)	NA	NA

Notes.

Values expressed as mean (SD).

NA, Not applicable.

**p*-value of the Student's *t* test.

Table 2 Load effect in women with and without fibromyalgia.

	Shapiro Wilk (<i>p</i> -value)	Fibromyalgia women (<i>n</i> = 12)	Healthy women (<i>n</i> = 8)	<i>p</i> -value*	Effect size (Cohen's <i>d</i>)
Trunk tilt					
Mean tilt carrying a load (°)	.342	2.31 (0.63)	1.69 (0.51)	.034	1.11 ^c
Mean tilt without carrying a load (°)	.653	2.16 (0.60)	2.05 (0.79)	.712	0.17 ^a
Median tilt carrying a load (°)	.025	2.22 (0.78)	1.43 (0.47)	.007	1.23 ^c
Median tilt without carrying a load (°)	.868	1.98 (0.57)	1.80 (1.03)	.624	0.24 ^a
Load effect (%) = (step duration carrying a load-step duration without carrying a load)/step duration without carrying a load* 100					
Initial stairs	.538	14.20 (16.07)	4.44 (7.40)	.127	0.77 ^b
Intermediate stairs	.322	18.32 (19.78)	-0.12 (10.94)	.028	1.15 ^c
Final stairs	.057	18.61 (12.32)	7.12 (6.73)	.028	1.15 ^c
Full task	.169	16.89 (13.73)	4.30 (6.84)	.015	1.15 ^c

Notes.

Values expressed as mean (SD).

**p*-value of the independent sample Student's *t*-test or Mann-Whitney *U* test, where appropriate.

^aSmall effect size (Cohen's *d* < 0.5).

^bMedium effect size (Cohen's *d* between 0.5 and 0.8).

^cLarge effect size (Cohen's *d* > 0.8). Initial stairs: stairs 1, 2 and 3; Intermediate stairs: stairs 4 and 5; Final stairs: stairs 6, 7 and 8. Full task: all eight stairs.

Load effect was significantly different in the two groups. Women with FM showed higher increase in time as a consequence of carrying a load. There were differences in the full task ($p = .015$), the intermediate and the final part of the 10-stairs climbing tasks ($p = .028$). On the other hand, there were no statistically significant differences in the initial stairs.




Figure 1 Mean time required for each step.

The effect size was medium (>0.5 and <0.8) in the initial stairs and large (>0.8) in the rest of the stairs.

Figure 1 depicts the evolution of the time required to complete each step. This time was generally on a rising trend in the two groups. The highest rising trend was observed in the group with FM when they were carrying a load of 5 kg in each hand.

DISCUSSION

The main finding of the current paper was that women with FM altered their motor pattern climbing stairs with loads more than the healthy controls. In more detail, postural sway and the time required to climb stairs were modified differently. Firstly, women with FM significantly increased the trunk tilt when they were carrying a load of 5 kg in each hand compared with the healthy controls. This finding may be extremely relevant as mediolateral stability is closely related to fall risk (*Bergland, Jarnlo & Laake, 2003*). One previous study evaluated different acceleration parameters in the mediolateral axis during stair ascent using accelerometer data (*Wang et al., 2014*). They placed an inertial measurement Opal sensor in the centre of the lower back and observed that acceleration parameters in the mediolateral axis were related with physiological risk of falling. Although most of the studies only assess mediolateral acceleration or deviation in static tasks, it is proved that both measures are relevant in dynamic tasks like walking, or stair ascent and descent. The high risk of fall and the serious consequences of falling in stair-climbing make it necessary to assess the balance stability of populations with altered balance, in order to design specific treatments. In this regard, different treatments may be effective in improving the

aforementioned mediolateral sway, like whole-body vibration (*Sanudo et al., 2013; Sanudo et al., 2012*), physical exercise (*Jeon et al., 2014*), or proprioception training programs (*Martinez-Amat et al., 2013*). Future studies on the identification of the best treatment in improving mediolateral sway in women with FM are required.

Load effect was calculated as percentage of the increased time in each step as a consequence of the load. The observed slowdown when women with FM were carrying loads might be a consequence of the lack of strength or resistance of the lower limb in women with FM, reported by previous studies (*Aparicio et al., 2015; Valkeinen et al., 2008*). In fact, previous studies showed that an aquatic program including leg squats improved 10-stair climbing time while carrying loads and leg strength, whereas FM controls progressively worsened their performance and postural balance (*Tomas-Carus et al., 2008*). Therefore, an implication of the results reported in the current study may be the need of adding resistance and strength training to physical therapies for women with FM.

The step-by-step performance in the stair-climbing tasks is depicted in [Fig. 1](#). The time required to complete each step was on a rising trend for both groups in the two tasks, progressively increasing until the last step. However, this tendency is higher when participants were carrying a load, compared to the weightless task. In addition, this tendency was higher for women with FM. This could indicate that fatigue highly affects as a consequence of the load. This finding is consistent with previous studies that showed the progressive worsening of time climbing 10-stairs, especially while carrying loads (*Tomas-Carus et al., 2008; Tomas-Carus et al., 2007*). Although pain is a major symptom in FM, no significant linear correlation was found between pain and any of the explored variables. In other words, the slowdown seems to be associated in a greater scale with physical deconditioning than with pain. However, there is a need for further studies testing this hypothesis. Similarly, no significant linear correlation was observed between the explored variables and the question about stair climbing in FIQ and FIQ-R. In the FIQ, the question is “Were you able to climb stairs?” Since all the participants in the current study were able to climb stairs (it was an inclusion criteria), it seems reasonable that correlation was not significant. However, the question in the FIQ-R is “how much your fibromyalgia made it difficult to perform each of the following activities during the past 7 days” and significant correlation was expected. This supports that physical fitness tests are more capable to capture differences in the ability of conducting ADL than self-reported questionnaires in women with FM (*von Bulow et al., 2015; Waehrens et al., 2012*). Further studies should investigate the relation between the perceived ability to climb stairs and the actual performance in a physical test.

All the reported differences could be even higher, based on characteristics of the participants of the current study. The mean age was almost two years higher for healthy controls compared to women suffering from FM. The severity of FM according to the score in FIQ was “moderate,” as it was between 39 and 59 (*Bennett et al., 2009a*). Finally, the mean impact of FM on the function domain was low (13.39). Therefore, it could be expected that the differences reported in the current study would be higher with a “severe” score in the FIQ.

Study limitations

The current study encountered several limitations. Firstly, the cross-sectional nature of the study and its relatively small sample size limited the capability to identify the causes of the reported differences between women with and without FM. Therefore, there is a need for larger studies that could explain the causes of the higher slowdown and trunk tilt. Secondly, the current study did not compare directly and statistically the trunk tilt while carrying and without carrying a load because the motor pattern of arms is inherently different due to the action of carrying weight, which add other source of variability. Therefore, a trunk tilt comparison in the two situations would require a specific design and a larger sample size. The third limitation could be due to the selected tasks limited the muscle fatigue as participants were asked to climb only 10 stairs, which is mainly related to non-fully aerobic mechanisms. Future studies focused on the onset of fatigue could also include higher number of stairs. The fourth limitation could be consequence of the dimensions of the stairs, which may vary from stairs in other countries. In this regard, higher steps could result in greater sway given the greater need for hip flexion. Finally, risk of falling is also high during stair descent (*Bosse et al., 2012*), but the assessment of the impact of FM (using FIQ, FIQ-R or physical tests) is usually conducted during stair climbing. Despite these limitations, the current study contributes significantly to the research focused on the performance of women with FM during the stair ascent.

CONCLUSIONS

Two main conclusions can be obtained from the results of the current study. In the first place, trunk tilt during the stair climbing while carrying a load was significantly higher for women with FM, which could increase the risk of falling. Second, women with FM experienced a slowdown as a consequence of the load. This slowdown was especially observed at the intermediate and the final part of the 10-stairs climbing test. Therefore, there is a need to prevent the loss of leg resistance and strength, in order to reduce the impact of FM on the ability to perform activities of daily living in women with FM.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

In the framework of the Spanish National R+D+i Plan, the current study has been co-funded by the Spanish Ministry of Economy and Competitiveness with reference DEP2012-39828. This study has been also co-funded by the Government of Extremadura and European Union Regional Development Funds (FEDER), a way of making Europe (ref. GR10127). DCM is supported by a grant from the Spanish Ministry of Education, Culture and Sport (FPU14/01283). He was previously supported by a Predoctoral Fellowship from the Fundación Tatiana Pérez de Guzmán el Bueno. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Spanish Ministry of Economy and Competitiveness: DEP2012-39828.
Government of Extremadura and European Union Regional Development Funds (FEDER): GR10127.
Spanish Ministry of Education, Culture and Sport: FPU14/01283.
Fundación Tatiana Pérez de Guzmán el Bueno.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Daniel Collado-Mateo conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- José C. Adsuar conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.
- Pedro R. Olivares and Narcis Gusi conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.
- Francisco J. Dominguez-Muñoz and Cristina Maestre-Cascales conceived and designed the experiments, performed the experiments, contributed reagents/materials/analysis tools, reviewed drafts of the paper.

Human Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

Bioethical Committee of the University of Extremadura (Spain). Registration number: 113/2014.

Data Availability

The following information was supplied regarding data availability:

The raw data is supplied as a [Dataset S1](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.1656#supplemental-information>.

REFERENCES

- Aparicio VA, Segura-Jimenez V, Alvarez-Gallardo IC, Soriano-Maldonado A, Castro-Pinero J, Delgado-Fernandez M, Carbonell-Baeza A. 2015. Fitness testing in the fibromyalgia diagnosis: the al-andalus project. *Medicine and Science in Sports and Exercise* 47:451–459 DOI 10.1249/MSS.0000000000000445.

- Bennett R. 2005.** The Fibromyalgia Impact Questionnaire (FIQ): a review of its development, current version, operating characteristics and uses. *Clinical and Experimental Rheumatology* **23**:S154–S162.
- Bennett RM, Bushmakina AG, Cappelleri JC, Zlateva G, Sadosky AB. 2009a.** Minimal clinically important difference in the fibromyalgia impact questionnaire. *Journal of Rheumatology* **36**:1304–1311 DOI [10.3899/jrheum.081090](https://doi.org/10.3899/jrheum.081090).
- Bennett RM, Friend R, Jones KD, Ward R, Han BK, Ross RL. 2009b.** The Revised Fibromyalgia Impact Questionnaire (FIQR): validation and psychometric properties. *Arthritis Research and Therapy* **11**:R120 DOI [10.1186/ar2783](https://doi.org/10.1186/ar2783).
- Bergland A, Jarnlo GB, Laake K. 2003.** Predictors of falls in the elderly by location. *Aging Clinical and Experimental Research* **15**:43–50 DOI [10.1007/BF03324479](https://doi.org/10.1007/BF03324479).
- Bosse I, Oberlander KD, Savelberg HH, Meijer K, Bruggemann GP, Karamanidis K. 2012.** Dynamic stability control in younger and older adults during stair descent. *Human Movement Science* **31**:1560–1570 DOI [10.1016/j.humov.2012.05.003](https://doi.org/10.1016/j.humov.2012.05.003).
- Carbonell-Baeza A, Aparicio VA, Sjostrom M, Ruiz JR, Delgado-Fernandez M. 2011.** Pain and functional capacity in female fibromyalgia patients. *Pain Medicine* **12**:1667–1675 DOI [10.1111/j.1526-4637.2011.01239.x](https://doi.org/10.1111/j.1526-4637.2011.01239.x).
- Cohen J. 1988.** *Statistical power analysis for the behavioral sciences*. 2nd edition. Mahwah: Academic Press.
- Collado-Mateo D, Gallego-Diaz J, Adsuar JC, Domínguez-Muñoz F, Olivares PR, Gusi N. 2015.** Fear of falling in women with fibromyalgia and its relation with number of falls and balance performance. *BioMed Research International* **2015**:Article 589014 DOI [10.1155/2015/589014](https://doi.org/10.1155/2015/589014).
- Corbett DB, Peer KS, Ridgel AL. 2013.** Biomechanical muscle stimulation and active-assisted cycling improves active range of motion in individuals with Parkinson's disease. *NeuroRehabilitation* **33**:313–322 DOI [10.3233/NRE-130961](https://doi.org/10.3233/NRE-130961).
- Esteve-Vives J, Rivera Redondo J, Isabel Salvat Salvat M, De Gracia Blanco M, De Miquel CA. 2007.** Proposal for a consensus version of the Fibromyalgia Impact Questionnaire (FIQ) for the Spanish population. *Reumatología' Clinica* **3**:21–24 DOI [10.1016/S1699-258X\(07\)73594-5](https://doi.org/10.1016/S1699-258X(07)73594-5).
- Hall M, Boyer ER, Gillette JC, Mirka GA. 2013.** Medial knee joint loading during stair ambulation and walking while carrying loads. *Gait Posture* **37**:460–462 DOI [10.1016/j.gaitpost.2012.08.008](https://doi.org/10.1016/j.gaitpost.2012.08.008).
- Huijnen IP, Verbunt JA, Meeus M, Smeets RJ. 2015.** Energy expenditure during functional daily life performances in patients with fibromyalgia. *Pain Practice* **15**:748–756 DOI [10.1111/papr.12245](https://doi.org/10.1111/papr.12245).
- Jackson PL, Cohen HH. 1995.** An in-depth investigation of 40 stairway accidents and the stair safety literature. *Journal of Safety Research* **26**:151–159 DOI [10.1016/0022-4375\(95\)00014-H](https://doi.org/10.1016/0022-4375(95)00014-H).
- Jeon SY, Han SJ, Jeong JH, Fregni F. 2014.** Effect of exercise on balance in persons with mild cognitive impairment. *NeuroRehabilitation* **35**:271–278 DOI [10.3233/NRE-141120](https://doi.org/10.3233/NRE-141120).

- Latorre-Roman P, Santos-Campos M, Heredia-Jimenez J, Delgado-Fernandez M, Soto-Hermoso V. 2014.** Analysis of the performance of women with fibromyalgia in the six-minute walk test and its relation with health and quality of life. *Journal of Sports Medicine and Physical Fitness* **54**:511–517.
- Martinez-Amat A, Hita-Contreras F, Lomas-Vega R, Caballero-Martinez I, Alvarez PJ, Martinez-Lopez E. 2013.** Effects of 12-week proprioception training program on postural stability, gait, and balance in older adults: a controlled clinical trial. *Journal of Strength and Conditioning Research* **27**:2180–2188 DOI [10.1519/JSC.0b013e31827da35f](https://doi.org/10.1519/JSC.0b013e31827da35f).
- Mas AJ, Carmona L, Valverde M, Ribas B, Group ES. 2008.** Prevalence and impact of fibromyalgia on function and quality of life in individuals from the general population: results from a nationwide study in Spain. *Clinical and Experimental Rheumatology* **26**:519–526.
- Muto LH, Sauer JF, Yuan SL, Sousa A, Mango P, Marques AP. 2015.** Postural control and balance self-efficacy in women with fibromyalgia: are there differences? *European Journal of Physical and Rehabilitation Medicine* **51**:149–154.
- Rahimi F, Bee C, Duval C, Boissy P, Edwards R, Jog M, Group E. 2014.** Using ecological whole body kinematics to evaluate effects of medication adjustment in Parkinson disease. *Journal of Parkinsons Disease* **4**:617–627 DOI [10.3233/JPD-140370](https://doi.org/10.3233/JPD-140370).
- Rahimi F, Duval C, Jog M, Bee C, South A, Jog M, Edwards R, Boissy P. 2011.** Capturing whole-body mobility of patients with Parkinson disease using inertial motion sensors: expected challenges and rewards. *Conference Proceedings of the IEEE Engineering in Medicine and Biology Society* **2011**:5833–5838 DOI [10.1109/IEMBS.2011.6091443](https://doi.org/10.1109/IEMBS.2011.6091443).
- Rutledge DN, Martinez A, Traska TK, Rose DJ. 2013.** Fall experiences of persons with fibromyalgia over 6 months. *Journal of Advanced Nursing* **69**:435–448 DOI [10.1111/j.1365-2648.2012.06026.x](https://doi.org/10.1111/j.1365-2648.2012.06026.x).
- Salgueiro M, Garcia-Leiva JM, Ballesteros J, Hidalgo J, Molina R, Calandre EP. 2013.** Validation of a Spanish version of the Revised Fibromyalgia Impact Questionnaire (FIQR). *Health and Quality of Life Outcomes* **11**:132 DOI [10.1186/1477-7525-11-132](https://doi.org/10.1186/1477-7525-11-132).
- Sanudo B, Carrasco L, De Hoyo M, Oliva-Pascual-Vaca A, Rodriguez-Blanco C. 2013.** Changes in body balance and functional performance following whole-body vibration training in patients with fibromyalgia syndrome: a randomized controlled trial. *Journal of Rehabilitation Medicine* **45**:678–684 DOI [10.2340/16501977-1174](https://doi.org/10.2340/16501977-1174).
- Sanudo B, De Hoyo M, Carrasco L, Rodriguez-Blanco C, Oliva-Pascual-Vaca A, McVeigh JG. 2012.** Effect of whole-body vibration exercise on balance in women with fibromyalgia syndrome: a randomized controlled trial. *Journal of Rehabilitation Medicine* **18**:158–164 DOI [10.1089/acm.2010.0881](https://doi.org/10.1089/acm.2010.0881).
- Segura-Jimenez V, Alvarez-Gallardo IC, Carbonell-Baeza A, Aparicio VA, Ortega FB, Casimiro AJ, Delgado-Fernandez M. 2014.** Fibromyalgia has a larger impact on physical health than on psychological health, yet both are markedly affected: the al-Andalus project. *Seminars in Arthritis and Rheumatism* **44**:563–570 DOI [10.1016/j.semarthrit.2014.09.010](https://doi.org/10.1016/j.semarthrit.2014.09.010).

- Segura-Jimenez V, Alvarez-Gallardo IC, Carbonell-Baeza A, Aparicio VA, Ortega FB, Casimiro AJ, Delgado-Fernandez M. 2015.** Fibromyalgia has a larger impact on physical health than on psychological health, yet both are markedly affected: the al-Andalus project. *Seminars in Arthritis and Rheumatism* **44**:563–570 DOI [10.1016/j.semarthrit.2014.09.010](https://doi.org/10.1016/j.semarthrit.2014.09.010).
- Sener U, Ucok K, Ulasli AM, Genc A, Karabacak H, Coban NF, Simsek H, Cevik H. 2013.** Evaluation of health-related physical fitness parameters and association analysis with depression, anxiety, and quality of life in patients with fibromyalgia. *International Journal of Rheumatic Diseases* Epub ahead of print Nov 30 2013 DOI [10.1111/1756-185X.12237](https://doi.org/10.1111/1756-185X.12237).
- Tomas-Carus P, Gusi N, Hakkinen A, Hakkinen K, Leal A, Ortega-Alonso A. 2008.** Eight months of physical training in warm water improves physical and mental health in women with fibromyalgia: a randomized controlled trial. *Journal of Rehabilitation Medicine* **40**:248–252 DOI [10.2340/16501977-0168](https://doi.org/10.2340/16501977-0168).
- Tomas-Carus P, Hakkinen A, Gusi N, Leal A, Hakkinen K, Ortega-Alonso A. 2007.** Aquatic training and detraining on fitness and quality of life in fibromyalgia. *Medicine and Science in Sports and Exercise* **39**:1044–1050 DOI [10.1249/01.mss.0b0138059aec4](https://doi.org/10.1249/01.mss.0b0138059aec4).
- Tseng HY, Liu BS. 2011.** Effects of load carrying methods and stair slopes on physiological response and postures during stairs ascending and descending. *Industrial Health* **49**:30–36 DOI [10.2486/indhealth.MS1100](https://doi.org/10.2486/indhealth.MS1100).
- Valkeinen H, Alen M, Hannonen P, Hakkinen A, Airaksinen O, Hakkinen K. 2004.** Changes in knee extension and flexion force, EMG and functional capacity during strength training in older females with fibromyalgia and healthy controls. *Rheumatology* **43**:225–228 DOI [10.1093/rheumatology/keh027](https://doi.org/10.1093/rheumatology/keh027).
- Valkeinen H, Hakkinen A, Alen M, Hannonen P, Kukkonen-Harjula K, Hakkinen K. 2008.** Physical fitness in postmenopausal women with fibromyalgia. *International Journal of Sports Medicine* **29**:408–413 DOI [10.1055/s-2007-965818](https://doi.org/10.1055/s-2007-965818).
- von Bulow C, Amris K, La Cour K, Danneskiold-Samsoe B, Ejlersen Waehrens E. 2015.** Differences in ability to perform activities of daily living among women with fibromyalgia: a cross-sectional study. *Journal of Rehabilitation Medicine* **47**:941–947 DOI [10.2340/16501977-2021](https://doi.org/10.2340/16501977-2021).
- Waehrens EE, Bliddal H, Danneskiold-Samsoe B, Lund H, Fisher AG. 2012.** Differences between questionnaire- and interview-based measures of activities of daily living (ADL) ability and their association with observed ADL ability in women with rheumatoid arthritis, knee osteoarthritis, and fibromyalgia. *Scandinavian Journal of Rheumatology* **41**:95–102 DOI [10.3109/03009742.2011.632380](https://doi.org/10.3109/03009742.2011.632380).
- Wang K, Lovell NH, Del Rosario MB, Liu Y, Wang J, Narayanan MR, Brodie MA, Delbaere K, Menant J, Lord SR, Redmond SJ. 2014.** Inertial measurements of free-living activities: assessing mobility to predict falls. *Conference Proceedings of the IEEE Engineering in Medicine and Biology Society* **2014**:6892–6895 DOI [10.1109/EMBC.2014.6945212](https://doi.org/10.1109/EMBC.2014.6945212).

Watson NF, Buchwald D, Goldberg J, Noonan C, Ellenbogen RG. 2009. Neurologic signs and symptoms in fibromyalgia. *Arthritis and Rheumatism* **60**:2839–2844 DOI [10.1002/art.24772](https://doi.org/10.1002/art.24772).

Wolfe F. 2015. Editorial: the status of fibromyalgia criteria. *Arthritis and Rheumatism* **67**:330–333 DOI [10.1002/art.38908](https://doi.org/10.1002/art.38908).

Wolfe F, Smythe HA, Yunus MB, Bennett RM, Bombardier C, Goldenberg DL, Tugwell P, Campbell SM, Abeles M, Clark P, Fam AG, Farber SJ, Fiechtner JJ, Franklin CM, Gatter RA, Hamaty D, Lessard J, Lichtbroun AS, Masi AT, McCain GA, Reynolds WJ, Romano TJ, Russell IJ, Sheon RP. 1990. The American College of Rheumatology 1990 Criteria for the Classification of Fibromyalgia. Report of the Multicenter Criteria Committee. *Arthritis and Rheumatism* **33**:160–172 DOI [10.1002/art.1780330203](https://doi.org/10.1002/art.1780330203).