



Walden University  
**ScholarWorks**

---

Walden Dissertations and Doctoral Studies

Walden Dissertations and Doctoral Studies  
Collection

---

2017

# Using an APN-Led Transitional Care Program to Reduce 30-Day Hospital Readmissions

Miaozhen Li  
*Walden University*

Follow this and additional works at: <https://scholarworks.waldenu.edu/dissertations>

 Part of the [Nursing Commons](#), and the [Public Health Education and Promotion Commons](#)

---

This Dissertation is brought to you for free and open access by the Walden Dissertations and Doctoral Studies Collection at ScholarWorks. It has been accepted for inclusion in Walden Dissertations and Doctoral Studies by an authorized administrator of ScholarWorks. For more information, please contact [ScholarWorks@waldenu.edu](mailto:ScholarWorks@waldenu.edu).

# Walden University

College of Health Sciences

This is to certify that the doctoral study by

Miaozhen Li

has been found to be complete and satisfactory in all respects,  
and that any and all revisions required by  
the review committee have been made.

## Review Committee

Dr. Diane Whitehead, Committee Chairperson, Nursing Faculty

Dr. Anna Valdez, Committee Member, Nursing Faculty

Dr. Eric Anderson, University Reviewer, Nursing Faculty

Chief Academic Officer

Eric Riedel, Ph.D.

Walden University

2017

Abstract

Using an APN-Led Transitional Care Program to Reduce 30-Day Hospital Readmissions

by

Miaozhen Li

Project Submitted in Partial Fulfillment  
of the Requirements for the Degree of  
Doctor of Nursing Practice

Walden University

August 2017

## Abstract

Heart failure (HF) is a serious public health problem associated with high mortality rates, hospital readmissions, and health care costs. Transitional care has emerged as a disease management model used to reduce readmissions for hospital-discharged patients with HF. However, the efficacy of an advanced practice nurse (APN)-led transitional care program (TCP) in readmission reduction is under debate. The practice question for this project examined the extent to which an APN-led TCP was effective in reducing 30-day all-cause readmissions for hospital-discharged HF patients. The logic model was the framework guiding this program evaluation. An analysis of quality improvement HF data from September 2015 to August 2016 was reviewed for one hospital in southern California. The APN-led TCP included 47 patients and had 7 patients with 30-day readmissions. The physicians' group included 298 patients and had 53 patients with 30-day readmissions. The results of chi-square analysis revealed a nonsignificant association between 30-day readmissions and post-discharge care providers [ $\chi^2(1, N = 345) = 0.236, p = 0.627$ ], and the HF 30-day readmission rates were the same between two groups. The APN-led TCP served a large proportion of Medi-Cal patients (48.94%) who had less primary care access, while the majority of patients in the physicians' group were Medicare (51%) who had primary care providers. This project highlights the positive social changes that advanced practice nurses affect via their critical leadership and clinical roles in increasing care access for the low-income population. Further studies on payer sources and readmissions are recommended on the efficacy of APN-led TCP in readmission reduction.

Using an APN-Led Transitional Care Program to Reduce 30-Day Hospital Readmissions

by

Miaozhen Li

Project Submitted in Partial Fulfillment

of the Requirements for the Degree of

Doctor of Nursing Practice

Walden University

August 2017

## Dedication

I dedicate this paper to my wonderful daughters, Kathryn and Elizabeth, and my husband, Richard Lin. To my daughters, thank you for giving up some of your mommy time so that I could finish my homework. To Elizabeth, thank you for spending many nights with me when I was working on this paper because you could not go to bed by yourself. To my husband, thank you for your understanding, patient, and support along this DNP journey.

## Acknowledgments

I want to thank my preceptors Cindy and Pat for providing me a wonderful clinical experience and my supervisor Michael Millan for his support and accommodating my schedule requests. I want to extend my gratitude to Dr. Brenda Hudson for her instructions on my writing skills and Dr. Sarah Inkpen for her assistance in my statistical analysis. I especially want to thank my DNP committee: Chair, Dr. Diane Whitehead; committee chair Dr. Anna Valdez; university research review, Dr. Eric Anderson, for their support, guidance, and feedback.

## Table of Contents

List of Tables .....	iii
List of Figures .....	iv
Section 1: Nature of the Project .....	1
Introduction.....	1
Problem Statement.....	4
Purpose Statement.....	7
Nature of the Doctoral Project .....	9
Significance.....	10
Summary.....	11
Section 2: Background and Context .....	12
Introduction.....	12
Concepts, Models, and Theories.....	12
Operational Definitions.....	14
Relevance to Nursing Practice .....	15
HF Regulations and Rules .....	16
HF Guidelines Supporting Disease Management Programs.....	17
Systematic Reviews Supporting TC Programs .....	20
Peer-Reviewed Publications Supporting TC Programs .....	21
Peer-Reviewed Publications Supporting Nurse-or APN-led TC Programs.....	25
Local Background and Context .....	30
Role of the DNP Student.....	31



Summary .....	32
Section 3: Collection and Analysis of Evidence.....	34
Introduction.....	34
Practice-Focused Question.....	35
Source of Evidence .....	37
Archival and Operational Data .....	39
Evidence Generated for the Doctoral Project .....	41
Analysis and Synthesis .....	44
Summary.....	45
Section 4: Findings and Recommendations.....	46
Introduction.....	46
Findings and Implications.....	47
Recommendations.....	57
Section 5: Dissemination Plan .....	61
Dissemination Plan .....	61
Analysis of Self.....	61
Summary.....	63
References.....	64
Appendix: Literature Summary with Level of Evidence.....	75

## List of Tables

Table 1. Hierarchy of Evidence .....	30
Table 2. Sources of Evidence.....	38
Table 3. Explanation of the Basic Logic Model .....	43
Table 4. Cross Tabulations and Chi-Square Results for 30-Day Readmissions With Post- Discharge Care Providers .....	49

## List of Figures

Figure 1. The basic logic model components. ....	13
Figure 2. Flowchart for data collection process.....	48
Figure 3. Bar graph showing the sex distribution for each group.....	50
Figure 4. Bar graph showing HF types for each group.....	51
Figure 5. Bar graph showing insurance types for each group.....	52
Figure 6. Bar graph showing comparisons of 30-day all-cause readmission rates from home in Medicare patients with HF from Health Services Advisory Group's reports .....	53

## Section 1: Nature of the Project

### **Introduction**

Heart failure (HF) is a serious medical condition and a pressing public health problem (Roger, 2013). Data from the 2009-2012 National Health and Nutrition Examination Survey revealed that HF affected 5.7 million U.S. residents age 20 years old and over (Mozaffarian et al., 2016). The prevalence rate of HF has been projected to have a 46% increase by 2030, which means that more than 8 million individuals 18 years old and over would progress to having HF (Mozaffarian et al., 2016). Each year, 915,000 people are newly diagnosed with HF (Mozaffarian et al., 2016). HF resulted in one in nine deaths in 2013, and the 2000-2010 HF data from Olmsted County, Minnesota, indicated that nearly half of all individuals with HF would die in 5 years (Centers for Disease Control and Prevention [CDC], 2015a). HF management is one of the most costly medical conditions (Mozaffarian et al., 2016). The direct and nondirect cost for HF management has been estimated at \$30.7 billion annually, with direct medical cost accounting for 68% of this amount (CDC, 2015a; Mozaffarian et al., 2016). HF management costs continue to grow, and total costs are projected to increase by more than twice the current cost to \$69.7 billion by 2030 (Mozaffarian et al., 2016).

HF has remained as the primary diagnosis of hospital discharge since 2000; however, HF hospitalizations are common in patients with HF, and the HF hospital readmission rates are higher in patients who were previously hospitalized (Mozaffarian et al., 2016). The overall admission rate in HF patients age 55 years and over is 11.6 per 1000 patients annually, and the reoccurring hospitalization rate is 6.6 per 1000 patients

yearly (Mozaffarian et al., 2016). The HF mortality rate is associated with hospital admissions. The 28-day and 1-year mortality rates following hospital discharges in patients with HF are 10.4% and 29.5%, respectively (Mozaffarian et al., 2016). The Centers for Medicare and Medicaid services' (CMS, 2015b) 30-day All-Cause Hospital Readmission measure includes data on Medicare patients who had an unplanned acute care hospitalization for any cause within 30-days of discharge from an acute care hospital. The CMS's data demonstrated that the 30-day all-cause readmission rate for patients with HF was 25%, with the admitting diagnosis of HF contributing to 35% of those readmissions (Roger, 2013). For patients with HF, managing cardiac condition and no cardiac comorbidities after initial hospitalizations is a vital approach to reducing hospital readmissions (Roger, 2013).

Hospital readmissions in HF patients are preventable, and the Transitional Care Program (TCP) had evolved as a new model in HF management, with a goal to reduce 30-day hospital readmissions (Albert et al., 2015). According to the Medicare Payment Advisory Commission (2007) report, 30% of readmission costs were spent on seven medical conditions including HF, with 75% of readmissions being considered avoidable. The transition of care in HF management refers to "individual interventions and programs with multiple activities that are designed to improve shifts or transitions from one setting to the next, most often from hospital to home" (Albert et al., 2015, p. 1). A TCP that is patient-centered and integrated with multidisciplinary collaboration is imperative to provide the care continuum and to reduce hospital readmissions (Albert et al., 2015). Nurses and Advanced Practice Nurses (APNs) often perform various

leadership roles in facilitating care transitions include those of case managers, communicators, liaisons, and coordinators (Albert et al., 2015). The concept of Transitional Care (TC) in HF was validated mostly by small observations or quasi-experimental study designs in a single center or a few centers. The studies excluded patients who were non-English speaking or in the end stage of renal disease (Albert et al., 2015). With these limitations, the findings are not broadly applicable, and Albert et al. (2015) have noted that further studies are needed to determine the efficiency and cost-savings of TC interventions in managing HF patients.

Nurse practitioners (NPs) have played crucial roles in managing HF patients. Echeverry, Lamb, and Miller (2015) examined the impact of NPs home visits in improving quality measures of HF in homebound patients. In the project, NPs visited 40 patients with Class III or IV HF monthly for 3 months (Echeverry et al., 2015). The project findings demonstrated that hospitalizations, emergency room visits, and 30-day readmissions significantly decreased by 64%, 85%, and 95%, respectively (Echeverry et al., 2015). The results indicated the vital roles of NPs in improving the patient outcome of HF management.

In this Doctor Nursing Practice (DNP) doctoral project, I analyzed the existing data to evaluate the effectiveness of using an APN-led TCP in reducing 30-day readmissions for patients with HF. The positive social change implications of the DNP project potentially include decreased HF mortality rates, reduced medical costs, improved physical functions and quality of life, and prolonged life expectancy in HF patients (see Albert et al., 2015; Echeverry et al., 2015).

### **Problem Statement**

HF is an epidemic public health problem associated with high mortality rates and extensive health care spending. The incidence of HF for people older than 65 is 10 per 1000 of the population, and the risk of HF development is 20% after age 40 for both men and women (Mozaffarian et al., 2016). The lifetime risk of HF is twice that number in patients with hypertension (Mozaffarian et al., 2016). The mortality of HF has remained high. The 1-year and 5-year mortality rates are 29.6% and 52.6% respectively, and HF caused 300,122 deaths in 2013 (Mozaffarian et al., 2016). HF hospitalizations are associated with the highest mortality rates and hospital readmissions in patients with a history of HF. HF impacted the health care cost crisis. In 2012, HF accounted for 1,774,000 physician office visits, 553,000 ED visits, and 257,000 outpatient clinic visits (Mozaffarian et al., 2016). Under the Affordable Care Act (ACA), the CMS (2016) implemented the Hospital Readmissions Reduction Program (HRRP) in 2012 to reduce payments to hospitals with an excess amount of readmissions within 30 days of discharge on certain measured conditions, including HF.

Hospitalizations for HF have remained high and unchanged due to the aging population. According to the National Hospital Discharge Survey (NHDS) data from 2000 to 2010, 1 million HF hospitalizations occurred in 2000, and 2010 had the same number of HF hospitalizations (Hall, Levant, & DeFrances, 2012). The majority of hospitalized patients with HF are 65 years old and over; however, there were significantly increased hospitalizations from 23% in 2000, to 29% in 2010 for patients under 65 years of age (Hall et al., 2012).

After-discharge health services have increased, but the effect on reducing hospitalizations for HF have been minimal. Age plays a role in hospital discharge. Patients 65 years and over are commonly released to the long-term care facilities, while patients under 65 years are discharged to home (Hall et al., 2012). In the past 10 years, more than 80% of hospitalized HF patients under age 65 have been discharged to home, while 62% of hospitalized HF patients age 65 and over have been discharged to home (Hall et al., 2012). The discharge proportion for patients aged 65 and over who were discharged to long-term care facilities increased from 17% in 2000, to 21% in 2010 (Hall et al., 2012). The 30-day readmission rate for HF patients with home health agency care was as high as 26%, with 42% of them having hospital readmissions for cardiac issues (Madigan et al., 2012). With a significant proportion of discharged home patients with HF, there has been a great need for improving post-discharge care to reduce their hospital readmissions.

HF hospitalizations are associated with many factors including hospitals, health care providers, and patients. One national study on hospital strategies to reduce 30-day HF readmissions indicated that less than half of participating hospitals (537 hospitals) collaborated with community physicians to manage high-risk patients for readmissions (Bradley et al., 2013). Only 28.9% of hospitals electronically linked their inpatient and outpatient prescriptions records, and only 25.5% of hospitals set up work process to send the discharge summary to patients' primary care providers (PCPs) upon discharge (Bradley et al., 2013). HF patients had higher levels of unmet psychosocial needs than physiological needs after discharge (Davidson, Cockburn, & Newton, 2008).



The post-discharge care barriers and concerns include medication management (cost and medication reconciliation), follow-up appointment (lack of transportation and patient's awareness), PCP communication (poor hand-off and insufficient patient education), and non-medication related signs and symptoms management including non-adherence to diet, activity, exercise, and fluid management (Albert et al., 2015). Albert et al. (2015) have noted that post-discharge disease management programs share some common features, which included telephone follow-up, patient education, self-management, weight monitoring, sodium and fluid restriction or diet advice, exercise recommendations, medication review, and social and psychological support. Among post-discharge care programs, case management with telephone follow-up and home visits and multidisciplinary care programs with care coordination from admission to discharge have improved the HF mortality and hospital readmissions (Takeda, 2012). A patient-centered TCP with a multidisciplinary team approach could potentially prevent hospital readmissions (Albert et al., 2015). HF specialist nurse-led programs have decreased the mortality and readmission rates of HF patients (Takeda, 2012). However, Price (2012) found that less than 50% of HF patients are referred to HF nurses.

The Institute of Medicine (IOM, 2001), has stated the need for a change of care coordination to include "coordination of care across patient-conditions, services, and sites of care over time" (p. 2). The IOM (2010) has emphasized nurses' roles in redesigning the health care delivery system, claiming, "Nurses should be full partners, with physicians and other healthcare professionals, in redesigning health care in the United States" (p. 1). In this DNP project, I analyzed existing archival data to seek the evidence

of the effectiveness of using an APN-led TCP to reduce 30-day readmission, and to validate APNs' roles in leading TC interventions in HF management.

### **Purpose Statement**

The gap in practice that I addressed in this DNP project was to adopt an evidence-based disease management model to improve after discharge care for hospital-discharged patients with HF. The purpose of this DNP project was to validate the efficacy of an APN-led TCP to reduce 30-day readmissions in patients with HF. Hospital-discharged patients are at high risk for hospital readmissions due to the lack of follow-up appointments after discharge. In addition to this lack of follow-up appointments, there is a variety of other reasons that lead to high readmissions among hospital-discharged patients with HF. First, the lack of collaboration and communication from one setting to another setting is common among healthcare providers for patients with HF, especially in regard to medication reconciliation; hence. Such failures of communication are one of the primary reasons for hospitalizations in HF patients (Bradley et al., 2013). Second, the lack of adequate nurses' knowledge in HF management contributes to hospital readmissions in patients with HF. Nurses play pivotal roles in educating patients and improving patients' self-management skills before patients are discharged to home. For example, Guirguis-Blake (2016) pointed out that nurses' knowledge in HF management promote patients' self-management behaviors, which is one of the critical strategies to prevent hospital readmissions. However, Mahramus et al., (2013) stated that overall nurses' knowledge of HF is weak. As a result, patients are not adequately prepared to perform self-management at home before discharge. Third, patients' unaddressed post-

discharge needs precipitate readmissions. Albert et al., (2015) reported that patients' unmet needs (including psychosocial, physical, culture, and individual) contribute to HF hospitalizations. Post-discharge TC is an evolving disease management model in HF management and a guideline-recommended care approach in addressing post-discharge care needs (Yancy et al., 2013). Patient education and early post-discharge follow-up appointment are essential elements of TC programs (Yancy et al., 2013). However, the efficacy of TC programs to reduce readmissions has been disputed (Albert et. 2015; Kociol et al., 2012, 2016; Stamp et al., 2014), and it is not utilized widely. For instance, Nelson and Pulley (2015) stated that less than half of patients could follow up with their PCPs within 2weeks of discharge. The practice-focused question for this DNP project was:

For hospital-discharged HF patients, was an APN-led TCP effective in reducing hospital 30-day all-cause readmissions?

This DNP project had potential to address the lack of post-discharge care by evaluating the efficacy of an APN-led TCP in reducing 30-day readmissions in patients with HF, thereby strengthening the efficiency of APNs' leadership roles in TC programs, and potentially prompting the adoption of an APN-led TCP at my practicum site. The TCP is emerging as a new disease management model used to address the lack of collaboration during a transition of care and prevent hospital readmissions in patients with HF. Researchers have demonstrated that the standards of a TCP include patient education, early telephone or clinic or office follow-ups, assessment, medication reconciliation, home visits, hand off discharge summary, caregivers' engagement, and a

designated person to assume the leadership role in collaborative interventions (Guirguis-Blake, 2016). The evidence-based practice (EBP) guidelines specify the timeframe for patient education and post-discharge follow-ups by phone and clinic or office visits. The American Heart Association's (AHA, 2016) Get With The Guidelines- Heart Failure (GWTG-HF) showed that starting patient education in the hospital with a teach-back method, early telephone follow-up within 24 to 72 hours post-discharge to check patient status at home, and early follow-up appointments within the first week of post-discharge is the standard of post-discharge follow-up care. Furthermore, discharge assessment should be conducted at the beginning of the hospitalization. In particular, the first discharge assessment should begin right after admissions (Guirguis-Blake, 2016). In this DNP project, I used program evaluation results to validate the efficacy of an APN-led TCP as a new disease management approach to prevent readmissions in patients with HF. Meanwhile, the program evaluation results also could assist hospital administrators in deciding whether to adopt the APN-led TCP hospital-wide for managing chronic diseases such as HF and stroke.

### **Nature of the Doctoral Project**

I reviewed evidence from public and private agencies including the CDC, AHA, and the Heart Failure Society of America (HFSA). To search for scholarly and clinical literature, I used online databases including CINAHL and Medline, which I accessed via the Walden University Library. In the searches, I used keywords such as *heart failure*, *transitional care program*, *30-day readmissions*, *mortality*, *quality of life*, *patient education*, *cost reduction*, and *nurse- or Advanced Practice Nurse (APN) or Nurse*

*Practitioner (NP)*. I used the Melnyk and Fineout-Overholt's hierarchy of evidence (2005) to critically evaluate studies (Burns, Rohrich, & Chung, 2011; Grove, Burns, & Gray, 2013), and the health literature review matrix to organize the research and studies.

The purpose of this DNP project was to evaluate the efficacy of an APN-led TCP in reducing 30-day all-cause hospital readmission rates in patients with HF. The practice-focused question was:

For hospital-discharged HF patients, was an APN-led TCP effective in reducing hospital 30-day all-cause readmissions?

### **Significance**

The stakeholders of this DNP project were patients, caregivers, primary care providers, cardiologists, social workers, discharge planners, pharmacists, nurses including APNs, and administrators. All stakeholders worked collaboratively to provide a patient-centered continuum of care and reduce hospital readmissions in HF patients. Albert et al. (2015) and Guirguis-Blake (2016) stated that the TC model is effective in reducing hospital readmissions, mortality rates, and health care spending in patients with HF; however, there has been limited evidence supporting the broad application of the TC model and APNs' leadership roles in a TCP (Albert et al., 2015). This DNP project highlighted the nurse leadership roles in redesigning the American health care delivery system. It also could inspire APNs to use their training to provide an effective and efficient method of care to improve population health (IOM, 2010; 2001). There was limited evidence supporting the wide use of TC model. This DNP project strengthened the evidence regarding the efficacy of APN-led TCP in reducing 30-day readmissions.

This DNP project could result in positive social change including lowering the mortality rates in HF patients, reducing costs of care, and improving quality of life (see Albert et al., 2015; Guirguis-Blake, 2016).

### **Summary**

HF is a complex and costly clinical symptom and a severe population health problem in the United States. HF-related hospitalizations and hospital readmissions have remained high, and people diagnosed with HF often die within 5 years. The poor collaboration and communications among health care providers across all settings contributes to a higher rate of hospitalizations in HF patients.

Under the quality-driven renovated reimbursement system, a TC model is the recommended standard of post-discharge care model to reduce 30-day readmissions and mortality rates in patients with HF (Yancy et al., 2013). Nurses, especially APNs, have been taking leadership roles in directing transitional care interventions and programs (Albert et al., 2015; Mozaffarian et al., 2016). However, the evidence that using an APN-led TCP to reduce readmissions and mortalities in HF patients is limited. I designed this DNP project to assess the effect of using an APN-led TCP to reduce hospital readmissions in patients with HF. The project findings could potentially validate evidence of the effectiveness of an APN-led TCP in reducing 30-day readmissions and highlight nurse leadership roles in directing the health care delivery system (see Albert et al., 2015; IOM, 2010; Mozaffarian et al., 2016).

## Section 2: Background and Context

### **Introduction**

In my practicum hospital, two HF nurse practitioners initiated the APN-led TCP as a quality improvement project to help improve post-discharge follow-up care and reduce 30-day hospital readmissions for patients with HF. In this DNP project, I evaluated the efficacy of an APN-led TCP in reducing HF 30-day all-cause readmissions in patients who were discharged from the hospital with an index admission of HF (see Albert et al., 2015; Guirguis-Blake, 2016). The practice-focused question was:


For hospital-discharged HF patients, was an APN-led TCP effective in reducing hospital 30-day readmissions?

In this section, I discuss the study's concepts, models, and theories; operational definitions; relevance to nurse practice; local background and context; and my role as DNP student.

### **Concepts, Models, and Theories**

I selected the logic model (Hodges & Videto, 2011) to serve as a theoretical framework for this project examining the effectiveness of an APN-led TCP in reducing 30-day hospital readmission in patients with HF. The logic model offers “a picture of how your organization does its work—the theory and assumptions underlying the program” (W.K. Kellogg Foundation, 2004, p. 1). The logic model serves as a roadmap of a program, which systemically and visually demonstrates the relationships and sequences among program resources, activities, and intended results that included outputs, both short- and long-term outcomes, and impacts (W.K. Kellogg Foundation,

2004). The logic model is an instrumental tool for both program planning and evaluation, which helps program evaluators determine whether a project is implemented according to the plan and identify any barriers to the implementation of the project (Hodges & Videto, 2011; W.K. Kellogg Foundation, 2004). The components of the basic logic model were presented in Figure 1.



*Figure 1.* The basic logic model components. Reprinted from “The logic model development guide,” by W.K. Kellogg Foundation, (2004). Reprinted with permission.

The logic model has been widely applied to health care program evaluation. Ladd, Sitaker, Patanian, and Jernigan (2008) used the logic model to guide their evaluation of the Washington State Heart Disease and Stroke Prevention Program (WaHDSPP). Ladd et al. (2008) reported that the logic model was a useful program evaluation tool to help them develop evaluation questions and measurement indicators, monitor program progresses, and track activities and outcomes. Hodges and Videto (2011) reported that partnering with stakeholders was vital in using the logic model for program evaluation. The logic model serves as a program evaluation framework that holds stakeholders accountable for program operations and outcomes, demonstrates program results systematically, and can be used to develop a better program (CDC, 2010).



## Operational Definitions

*Heart failure:* A complex clinical symptom that involves an insufficient heart pumping of blood, and is commonly caused by structural and functional defects of ventricular filling and ejection (Yancy et al., 2013). HF patients typically experienced dyspnea, fatigue, exercise intolerance, and fluid overload (Yancy et al., 2013).

*The 30-day readmission:* According to the CMS (2016), 30-day readmission refers to “an admission to a subsection (d) hospital within 30 days of discharge from the same or another subsection (d) hospital” (para 2). The 30-day All-Cause Hospital Readmission quality measure applies to unplanned acute care hospital readmissions for any cause within 30 days following an acute care hospital discharge for Medicare patients (CMS, 2015b). Under the ACA, the CMS implemented the HRRP to penalize hospitals with poor performances on their 30-day readmissions for HF, pneumonia, and acute myocardial infarction (MI) in 2012 (Naylor, Aiken, Kurtzman, Olds, & Hirschman, 2011).

*Transitional care (TC):* In HF management, transitional care consists of multiple activities and or programs to provide safe and efficient care transfers from one setting to another, and the most frequent transitions took place from hospital to home (Albert et al., 2015).

*Heart failure comprehensive disease management:* The integration of the evidence-based practice, clinical improvement measures, and resources and tools for managing patients with HF to reduce health care costs and improve outcomes (Lindenfeld et al., 2010). Disease management interventions consist of inpatient and outpatient patient

education and self-care instruction, case management, multidisciplinary outpatient care, telephone support, telemedicine, and clinic visits (Feltner et al., 2014). TC is an emerging phrase describing disease management for patients transferred from one setting to another (Feltner et al., 2014). The AHA's statement on the disease management domains encompassed patient population, intervention recipient, intervention content, delivery personnel, communication method, intensity and complexity, environment, and clinical outcomes (Krumholz et al., 2006).

### **Relevance to Nursing Practice**

For the literature review, I used databases available through Walden University Library. These databases included Cochrane Database of Systematic Reviews, MEDLINE, and Cumulative Index of Nursing and Allied Health Literature (CINAHL) simultaneous search, Pub Med, and Walden University's ScholarWorks. The combinations of keywords I used to limit the search included *transitional care* and *heart failure*, *transition of care* and *heart failure*, and *readmission* and *nurse- or nurse practitioner, or advanced practice nurse-led*. The search criteria were also limited by age (adult population), year (from January 2011 to August 2016), and language (English). The majority of articles I used for the literature review were published within 5 years, and only a few articles were published more than 5 years ago. I used the keyword *heart failure* to guide searches of government websites (e.g. CDC, CMS AHRQ), and professional websites (AHA, AAHFN, ACC, HFSA, and TJC) provided rules and regulations on HF management, guidelines, and statistical data.

## **HF Regulations and Rules**

The Medicare Payment Advisory Commission (2007) reported that HF was one of the 10 most common hospital diagnoses for Medicare patients, that HF patients have a high risk for hospital readmissions, and that HF significantly impacts the quality of life of patients with frequent hospitalizations. Hospital readmissions cost the Medicare program \$15 billion annually, but \$12 billion of these costs are potentially preventable (Medicare Payment Advisory Commission, 2007). According to the 2007-2009 Medicare fee-for-service claims data, the 30-day readmission rates for HF, pneumonia, and acute MI were 24.8%, 18.3%, and 19.9%, respectively (Dharmarajan et al., 2013). Among all HF readmissions, 35% of them were readmitted for the same index HF hospitalization (Dharmarajan et al., 2013). High percentages of hospital readmission rates reflected the poor communication among care providers and poor transition of care after discharge (Medicare Payment Advisory Commission, 2007). Therefore, The CMS and Hospital Quality Alliance (HQA) established a public reporting system (including 30-day all-cause readmissions) to help consumers make informed decisions, improve safe and quality care, improve population health, and reduce healthcare costs (CMS, 2015c).

The HF 30-day readmission was the government's quality control measure, and the CMS used it to determine reimbursement rates for hospitals' payments based on their readmission status. The HF 30-day risk-standardized mortality and HF 30-day risk standardized readmission are two required quality reporting measures for hospitals (CMS, 2015c). Under the CMS's Reporting Hospital Quality Data for Annual Payment Update (RHQDAPU) program, hospitals receive higher payments for their compliances

of reporting quality measures (CMS, 2015 b). Under the ACA, the CMS implemented the HRRP to reduce payments to hospitals with excess readmissions for certain conditions in 2012, and the HF 30-day risk standardized readmission was one of the reimbursement measures (CMS, 2016; National Quality Measures Clearinghouse, 2015). HF is a severe and costly health condition. HF 30-day all-cause readmission is a quality measure that drives a value-based reimbursement for Medicare patients. In my practicum hospital, hospital leaders and clinicians have adopted HF 30-day all-cause readmission rates as the quality measures for HF management.

### **HF Guidelines Supporting Disease Management Programs**

Several HF professional organizations have recommended disease management programs and transitional care programs as evidence-based practices in HF management. They have laid out detailed strategies and activities for disease management programs. The sources of HF evidence-based guidelines include the HFSA's 2010 Comprehensive Heart Failure Practice Guideline, 2013 ACCF/AHA's guideline for the management of HF, the joint AHA and TJC advanced certification in HF, and AHA's GWTG-HF.

Comprehensive disease management is a critical approach for patients with HF. According to the HFSA 2010 Comprehensive Heart Failure Practice Guideline, HF comprehensive disease management programs are recommended for patients with a recent hospitalization for an index admission of HF, or for patients at high risk for developing decompensated HF (Lindenfeld et al., 2010). Comprehensive HF disease management programs include patient and family education, counseling, self-care management (including medication titrations), close follow-up post hospital discharge,

care access to health care providers and care coordination, optimization of drug therapy, early detection of signs and symptoms of fluid overload, and addressing social and financial needs (Lindenfeld et al., 2010). Coordination of care between the primary care physicians and HF experts and between different facilities or settings is one of the essential components in HF disease management programs (Lindenfeld et al., 2010).

A patient-centered, multidisciplinary team approach is an essential element of HF disease management programs. The 2013 ACCF/AHA's guideline for the management of HF recommended that for patients at high risk for readmissions, disease management should include a patient-centered multidisciplinary team approach (Yancy et al., 2013). The benefits of a patient-centered multidisciplinary team approach include readmission reduction and the implementation of guideline-directed medical therapy (GDMT; Yancy et al., 2013). The development and implementation of effective systems of care coordination and care transitions is essential for all HF patients to improve the quality of care (Yancy et al., 2013). Effective communications between healthcare providers, medication reconciliation, appropriate hand-off discharge summaries, and transition of care processes between settings are necessary safety practice when releasing HF patients from hospital to home (Yancy et al., 2013). In conclusion, TC is the core of disease management program. An adequate TC involves a patient-centered, multidisciplinary, and systematic approach. The evidence-based TC processes consist of care coordination and communication at all levels of care, medication reconciliations, and appropriate hand-off discharge summaries.

There is a great need for a well-planned transition of care for achieving a safe patient transition from one setting to another. According to the 2013 ACCF /AHA HF management guideline, the HF TC should consist of extensive patient education, medication reconciliation, hand-off updated care summary to follow-up providers, and early post-discharge follow-up arrangement (Yancy et al., 2013). Patient and family educational topics should include follow-up visits at the next care setting, recognizing worsening signs and symptoms of HF, medications, activity and diet (Yancy et al., 2013). A follow-up visit occurring within 7 to 14 days and a telephone follow-up made within three days of post-discharge were considered the optimal care for hospital- discharged patients with HF (Yancy et al., 2013). The guideline also stated that the standard of inpatient and outpatient HF management incorporated initiating the GDMT appropriately, addressing risk factors of HF and its comorbidities, identifying barriers for optimal care transitions and after discharge care support, assessing volume status, titrating HF drug therapy, assessing of renal function and electrolytes, reinforcing patient and family education, and considering palliative care or hospice care in specified patients (Yancy et al., 2013). However, the evidence of the guideline-recommended TC was limited, and the research was needed to support the efficacy of TC in HF management (Yancy et al., 2013). In summary, a well-planned TC is the guideline-directed medical therapy; and it is composed of comprehensive patient education, medication reconciliation and titration, disease management, post-discharge care follow-up within 2 weeks, phone call follow-up occurring within 3 days of discharge, and identifying and addressing barriers for care transitions.

TC is an evolving standard of care in HF management. The AHA and TJC established Advanced Certification in Heart Failure (ACHF) for acute care hospitals (TJC, 2016). The availability of inpatient and outpatient services and TC services for HF patients was one of the required certification criteria (TJC, 2016). Care coordination and care transition drove the GDMT in HF management (TJC, 2016). The GWTG-HF was the AHA's (2016) quality improvement program. It has multiple features including data registry, real-time benchmarking on performance measures, and decision support. A follow-up visit or phone call within 72 hours of post-discharge, HF disease management referral, and HF 30-day readmission and mortality rates were the reporting measures of AHA (2016) GWTG-HF. Hence, the continuum of care in HF management covers inpatient and outpatient services; TC ensures the continuity of care across all settings; and the 30-day all-cause readmission is the quality indicator in directing HF management.

### **Systematic Reviews Supporting TC Programs**

I identified two systematic reviews on HF and nurse-led interventions in the Cochrane Database of Systematic Reviews. The 2012 Cochrane systematic review on disease management interventions for patients with HF to reduce readmission and mortality by comparing to usual care included twenty-five Randomized controlled trials (RCTs). The disease management interventions were grouped into three types that included case management interventions (phone call follow-ups and home visits), clinic interventions (HF clinic follow-up), and multidisciplinary interventions (team-approached care from the hospital to home). The authors of the review concluded that among recently hospitalized HF patients, HF nurse-led case management interventions

reduced HF and all-cause readmissions and mortality; multidisciplinary interventions might have effects on the reduction of HF and all-cause readmissions; and clinic interventions had weak evidence to support its efficacy (Takeda, 2012). The 2016 Cochrane systematic review on evaluating Nurse-Led Titration (NLT) of beta-adrenergic blocking agents, Angiotensin Converting Enzyme Inhibitors (ACEIs), and Angiotensin Receptor Blockers (ARBs) in HF patients with Reduced Ejection Fraction (HFrEF) included seven RCTs with a total of 1684 participants (Driscoll, 2016). The RCTs compared the titration and optimization of those evidence-based drug therapies by a nurse to the optimization by other practitioners in patients with HFrEF (Driscoll, 2016). The systematic review demonstrated that NLT of medication titrations reduced all-cause hospital readmissions and mortality in patients with HFrEF (Driscoll, 2016). Those two systematic reviews have demonstrated the importance of nurses' leadership roles and significant clinical impact on HF management. Post-discharge home visiting and telephone follow-up, multidisciplinary team approach, and medication titration are effective measures to reduce readmissions.

### **Peer-Reviewed Publications Supporting TC Programs**

A search of CINAHL and MEDLINE simultaneous search database using combination phrases *transitional care* and *heart failure* and *readmissions* generated 75 peer-reviewed articles. Limiting the search to 5 years (January 2011 to August 2016) and English language yielded 58 peer-reviewed articles. After applying inclusion and exclusion criteria as outlined below, there were 14 articles remained. Of these, seven were systematic reviews, two were retrospective data analysis, and five were mixed-



method studies.

Inclusion criteria were systematic reviews or original research studies in peer-reviewed journals that examined TC programs or interventions to reduce all-cause or HF-specific hospital readmissions in adult patients with HF in the United States (US). Non-US countries were not included because of difference in health care delivery and reimbursement system. Exclusion criteria were studies with a narrow focus on specific populations such as home health care, long-term care, surgery, military services, dementia, and palliative care; studies had targeted interventions including telemonitoring.

TC has improved quality of care and cost containment in HF management. Many peer-reviewed studies indicated that TC programs and interventions were effective to reduce all-cause and/or HF-specific readmissions and mortality, to improve quality of life, and to lower costs of care (Albert, 2015; Albert, 2016; Centeno & Kahveci, 2014; Feltner et al., 2014; Guirguis-Blake, 2016; Monza, Harris, & Shaw, 2015; Russell, Rosati, Sobolewski, Marren, & Rosenfeld, 2011; Stamp, Machado, & Allen, 2014; Stauffer et al., 2011; Vedel & Khanassov, 2015; White & Hill, 2014). In the systematic review of Feltner et al. (2014) with 47 RCTs, home-visiting programs and multidisciplinary HF clinic interventions reduced all-cause readmission and mortality; structure telephone support reduced HF-specific readmission and mortality. However, in the systematic review, only a few RCTs reported 30-day readmission rates (Feltner et al., 2014). Kociol et al. (2012) conducted a national data survey on hospital-based strategies of readmission reduction programs from 100 hospitals participating in AHA's GWTG-HF. Kociol et al. (2012) categorized hospital strategies into three groups including inpatient care, TC, and

quality improvement. Among participated hospitals, their strategies varied in all three groups and the study findings demonstrated that only complete TC strategies were related to 30-day readmission reduction among three groups (Kociol et al., 2012). Thus, TC is an evidence-based practice to reduce the 30-day readmission.

The effective TC programs and strategies include a variety of interventions. The evidence-based TC interventions were home visiting programs with telephone follow-ups and multidisciplinary HF clinics (Albert, 2016; Feltner et al., 2014; Guirguis-Blake, 2016; Stamp et al., 2014; Vedel & Khanassov, 2015). Feltner et al. (2014) concluded that home-visiting programs and multidisciplinary HF clinics reduced all-cause readmissions and mortality and structured telephone support reduced HF-specific readmissions and mortality. Because the mean readmission timing was within 12 days of discharge for the majority hospitalized HF patients (Dharmarajan et al., 2013), early follow-up within 7 days of discharge was a necessary TC intervention (DeVore et al., 2016). Other TC interventions having effects on reduction of readmissions in patients with HF included care collaboration and coordination, patient education, improvement of self-management behaviors (Russell et al., 2011; White & Hill, 2014). Feltner et al. (2014) reported that effective interventions to reduce all-cause readmissions or mortality were delivered in person, high intensity, and multidisciplinary care approach (Albert, 2016; Centeno & Kahveci, 2014; Monza, Harris, & Shaw, 2015). Albert et al. (2016) summarized eight themes of TC programs that included discharge planning, multidisciplinary team-based collaboration and communication, timely and organized critical information; medication reconciliation and adherence, social and community services supports, patient education

including monitoring signs and symptoms of HF after discharge, outpatient clinic follow-up, palliative care and end of life care. In summary, the patient-centered TC programs are intensive and complex. The evidence-based TC interventions involve multidisciplinary care approach, care collaboration, care communication, home visiting, and phone follow-up.

Although many studies showed the effectiveness of TC in reducing readmissions, the evidence of TC has remained undetermined. The efficacy of TCP was disputed because of the inconsistency in TC interventions and study designs (Albert et. 2015; Kociol et al., 2012; Stamp et al., 2014). Kociol et al. (2012) reported that hospital-based readmission reduction strategies varied, and processes of care on inpatient and quality improvement measures had no effect on readmission reduction. Kansagara et al. (2016) conducted a systematic review on TC interventions to identify common strategies and themes. Ten studies revealed the evidence of discharge planning and hospital at home interventions to reduce readmissions and the TC interventions in general were multifaceted and flexible to accommodate individual needs (Kansagara et al., 2016). However, the efficacy of TC interventions was weak because of a variety of study populations, study designs, and clinical settings (Kansagara et al., 2016). Albert et al. (2015) and Stamp et al. (2014) indicated the need for further research to generalize the evidence of TC interventions in HF management. In summary, the standard of TC interventions is not well defined, and the efficacy of TCP for readmission reduction is still under debate.

### **Peer-Reviewed Publications Supporting Nurse-orAPN-led TC Programs**

A search of CINAL and MEDLINE simultaneous search database using combination phrases *nurse-led* or *advanced practice nurse-led* or *nurse practitioner-led* with *heart failure* and *readmission* generated 54 articles, limiting the search within 5 years yield 28 articles. After applying above inclusion and exclusion criteria, there were eight articles remaining. Of those, five were systematic reviews and meta-analysis, and three were mixed-method trials. Four out of five systematic reviewed were discussed in sections of the systematic review and peer-reviewed publications supporting TC programs.

Nurses and APNs play leadership and clinician roles in TC programs for readmission reduction and disease management. APN-, NP-, and nurse-led TC programs were effective to reduce all-cause readmissions and mortality in patients with HF (Monza et al., 2015; Takeda, 2012; Stauffer et al., 2011; Stamp et al., 2014). The 2012 Cochrane systematic review on disease management interventions for patients with HF indicated that HF nurse-led case management interventions (home care visits and telephone follow-up) after one year follow-up reduced HF-specific and all-cause readmissions and all-cause mortality; HF clinic interventions had no significant effects on HF-specific and all-cause readmissions and mortality; and multidisciplinary interventions were associated with reduction of HF-specific and all-cause readmissions (Takeda, 2012). The 2012 systematic review and meta-analysis included 19 RCTs evaluating the effect of nurse-led HF management programs (HF-MPs) before the discharge. The review revealed that the nurse-led HF-MPs were effective to reduce HF-specific and all-cause readmissions

(Lambrinou, Kalogirou, Lamnisos, & Sourtzi, 2012). An APN-led program designated the nurse practitioner as the program leader who assumed the primary responsibility for disease management of specified population such as patients with HF (Lowery et al., 2012). An APN-led clinic provided a holistic, patient-centered, and cost effective care and played a significant role in serving health care needs of low-income and uninsured population in the United States (Campbell, 2016). NPs at an APN-led clinic diagnosed and treated individual responses to health issues and promote population health (Campbell, 2016). Hence, Nurses and APNs play important roles in leading and operating HF management programs to reduce readmissions.

Many studies have demonstrated the efficacy of nurse-led TC programs in reducing readmissions. For example, Stamp et al. (2014) conducted a systematic review on the effects of a nurse-led TC program with 20 studies, and the findings indicated that home visiting and telephone follow-up were effective in reduction of readmissions in HF patients. A nurse-led multidisciplinary team approach study improved care transitions, reduced 30-day readmissions, improved patient education and satisfaction (Thompson, 2014). Smith et al. (2015) conducted an RTC trial on nurse-led multidisciplinary HF group clinic appointments in comparison with usual care. The RTC included 198 HF patients and indicated that a nurse-led multidisciplinary HF group clinic appointments increased patient's HF self-care knowledge and improved self-management behaviors that lead to a reduction in readmissions (Smith et al., 2015). Therefore, those studies had clearly identified the effectiveness of nurse-led HF TC programs for readmission reduction and improvement of self-management skills.

There are no consensus on essential elements of a nurse or APN-led TC programs in HF management. More research studies were needed to support the evidence of a nurse or APN-led TC programs for a broader application (Albert et al., 2015; Lambrinou et al., 2012;). Researchers were not able to determine standardized strategies of a successful nurse-led HF-MPs from a systematic review of 19 RCTs on the effect of HF-MPs before discharge, and further studies were needed to standardize strategies of HF-MPs (Lambrinou et al., 2012). In the review of Feltner et al. (2014), telemonitoring and nurse-led clinic interventions had no effect on the readmission and mortality. However, Mikulich et al. (2013) had a controversial finding that an APN-led HF clinic reduced HF readmissions. The APN-led HF clinic consisted of patient education including disease process, medication, daily weight, diet and activity, sodium and fluid restriction, self-care instructions, titration of drugs and compliance (Mikulich et al., 2013). Accordingly, the efficacy of nurse-led HF clinics is also undetermined due to the lack of standardized interventions and conflicted study results of nurse-led clinics.

Despite the fact that many studies indicated that nurses commonly functioned as leadership roles in TC programs, it is unclear of the educational background of nurses leading HF management programs (see Albert et al., 2015). Furthermore, the evidence of APN-led TC programs in readmission reduction was also not well identified in the literature review (see Albert et al., 2015). For instance, the APN-led TCP program was not well recognized and utilized to reduce 30-day readmissions for patients with HF due to the weak evidence and application (Price, 2012). In the systematic review of Albert et al. (2015), nurses functioned as coordinator roles in TC programs; however, only a few

studies mentioned the APN-led TC programs, and the evidence of APN-led TC programs and interventions was weak because of a variety of study designs and strategies of TC programs (Albert et al., 2015). There was a great need for nurse researchers to conduct RCTs to address gaps in TC models and interventions (Albert et al., 2015; Stamp et al., 2014). In the final analysis, the efficacy of TC and an APN-led TCP in reducing readmissions in patients with HF is not determined, and it remains as a practice gap in HF management.

Hospital strategies are vital for readmission reduction in patients with HF. Bradley et al. (2013) reported a cross-sectional web-based national survey on hospital strategies in reducing HF 30-day readmissions among hospitals participating in national quality initiatives. A total of 599 hospitals responded to the survey, and six strategies were significantly and individually related to lower 30-day HF readmissions (Bradley et al. 2013). Those hospital strategies included building partnerships with community physicians and physician groups, networking with local hospitals, medication reconciliations by nurses, post-discharge appointments arrangement, communicating discharge summaries with a patient's primary care provider, and test results follow-up (Bradley et al., 2013). The combination of strategies had a cumulative effect on reducing readmission rates (Bradley et al., 2013). However, only a few hospitals had implemented multiple strategies (Bradley et al., 2013). Hence, the findings highlighted the need for improvement in adopting those recommended hospital strategies cumulatively to reduce 30-day readmissions for patients at high risk for readmissions.

At my practicum site, two HF nurse practitioners evaluated their practice based on the study of Bradley et al. (2013). They added a discharge planner into the HF program who made phone follow-up call within 72 hours of discharge and assisted in arranging post-discharge follow-up appointments. They also implanted an APN-led TCP in August 2015 to offer an early post-discharge follow-up within one to two weeks of discharge for hospitalized patients with HF.

In this DNP project, I examined the effectiveness of an APN-led TC program in reducing 30-day readmissions in patients with HF. I evaluated HF existing archival data to determine if APN roles in leading a patient-centered TCP with a multidisciplinary team approach to provide care coordination and to promote care continuum for hospitalized patients for HF were an effective method for reducing 30-day readmissions.

### **Literature Review Summary**

Appendix A was a summary of the review of the literature related to this project. Each article had a graded level of evidence identified using the Melnyk and Fineout-Overholt's hierarchy of evidence recommended (2005). Table 1 demonstrated the Melnyk and Fineout-Overholt's hierarchy of evidence (2005).



Table 1

*Hierarchy of Evidence*

Level	Evidence
Level I	Evidence from a systematic review or meta-analysis of RCTs or clinical practice guidelines based on systematic reviews of RCTs
Level II	Evidence from at least one well-designed RCT
Level III	Evidence from well-designed controlled trials without randomization
Level IV	Evidence from well-designed case-control and cohort studies
Level V	Evidence from systematic reviews of descriptive or qualitative studies
Level VI	Evidence from a single descriptive or qualitative study
Level VII	Evidence from authority opinions/reports from experts

*Note.* From “*Evidence-based practice in nursing and healthcare*,” by Melynck, B. & Fineout-Overholt, E. 2005, p.10. Philadelphia, PA: Lippincott Williams & Wilkins. Copyright 2005 by Lippincott Williams & Wilkins. Reprinted with permission.

The summary of the evidence was eleven summaries at Level 1, one summary at Level II, six summaries at Level IV, one summary at Level V, two summaries at Level VI, and one summary at Level VII.

### **Local Background and Context**

The practicum site was a 420-bed acute care urban hospital located in Southern California. The hospital had two HF NPs who oversee inpatient and outpatient HF patients. Two APNs started a pilot TC program as a Quality Improvement (QI) project in August 2015 to reduce 30-day all-cause readmissions and to facilitate a safe transition from hospital to home for hospitalized patients with HF. The members of the APN-led TCP included two HFNPs, one discharge planner, and one licensed vocational nurse (LVN). The services of the APN-led TCP included HFNPs’ inpatient visits, patient and family education, outpatient HF clinic follow-up after discharge, prearranging follow-up appointments within 1 week of discharge, and follow-up phone with 72 hours of

discharge. During the HF clinic visits, HFNPs assessed patient's signs and symptoms of HF including fluid status, weight, compliance; reinforced patient education on medication, diet, activity, and fluid restriction; titrated medication to an optimal dose and collaborated with patients' PCPs and cardiologists for continuing care. In the TCP program, the HFNPs assumed the leadership and clinical roles to oversee and operate the TC program. The discharge planner assisted making post-discharge follow-up phone calls, prearranging follow-up appointments with PCPs, cardiologists, and HFNPs in the TCP. The discharge planner also verified insurance and located a PCP for patients followed up in the HF clinic. The LVN assisted HFNPs performing clinical activities including gathering test results, medication reconciliation, and setting up clinic visits.

The hospital was not yet an HF certified center by The Joint Commission (TJC); however, the facility had been participating in the AHA GWTG-HF program for years. The two HF NPs were planning to apply for HF certification in the near future. The pilot TCP for HF patients was not formally funded by the hospital. It was imperative to conduct an evaluation of the APN-led TCP to examine the effectiveness and efficiency of APN-led care coordination (Camicia et al., 2013).

### **Role of the DNP Student**

I realized that chronic disease management is a challenge after working both primary and acute care settings. I had worked in the Intensive Care Unit (ICU) for many years. I had an internship in a cardiologist office near the practicum site as a post-MSN student. Through working in inpatient and outpatient settings, I had seen the challenges

of chronic disease management due to the fragmented health care delivery system and the lack of collaboration and corporation among health care providers across all settings.

The DNP project was designed to evaluate the efficacy of an APN-led TC program to reduce hospital readmissions in patients with HF at the practicum site. The two HF NPs provided me with a de-identified data of HF program after I obtained the hospital's permission for accessing the data. Then I organized the data into Excel spreadsheets to demonstrate the differences in readmissions and demographic information between patients who followed up in the APN-led TCP and patients who followed up with their community physicians (including their PCPs and cardiologists). I would provide program evaluation findings to nursing leadership in the hospital. Because the APN-led TCP in HF management was a first pilot program in the practicum hospital as a QI project, the program evaluation was essential to help administration and other stakeholders decide whether to adopt and extend an APN-led TC model into other chronic disease management programs such as stroke.

### **Summary**

HF is a common and costly medical condition (Mozaffarian et al., 2016). Hospitalized patients with HF are at high risk for hospital readmissions that are considered preventable by using a multidisciplinary care collaboration approach (Takeda, 2012). The TCP is a newly emerged care model to coordinate patient care safely and more efficiently across different settings and or various levels of care for a continuum of care (Albert et al., 2015). Under the ACA, hospital readmission rate is a quality outcome measure that drives value-based quality care (CMS, 2016). The literature indicated that

nurses played leadership roles in TC programs (Naylor et al., 2011). However, the evidence regarding the efficacy of APN's roles in the TCP is limited, and the standardized strategies of TCP are also not well defined in the literature (see Albert et al., 2015).

The student practicum site had a piloting APN-led TCP to reduce 30-day readmissions in patients with HF. This DNP project was to evaluate the efficacy of the APN-led TCP in reducing hospital readmissions in patients with HF. The program evaluation results potentially helped management stakeholders decide whether to adopt the APN-led TCP as a standardized care model for chronic disease management in the organization and validated the APN -led TC program as a quality improvement approach in HF management.

### Section 3: Collection and Analysis of Evidence

#### **Introduction**

HF is a costly medical condition and serious population health problem. HF affected 5.7 million American adults from 2009 to 2012, and the cost for HF management, including direct and nondirect expense, was \$30.7 billion in 2012 (Mozaffarian et al., 2016). HF has remained the primary discharge diagnosis since 2000 (CDC, 2015a; Mozaffarian et al., 2016). The 28-day and 1-year mortality rates for hospitalized HF patients are 10.4% and 29.5%, respectively (Chang et al., 2014), and the CMS's 30-day all-cause hospital readmission rate for HF is 25% (Roger, 2013). Hospital readmissions for patients with HF are considered preventable (Medicare Payment Advisory Commission, 2007). Under the ACA, the CMS (2016) implemented the HPPS in 2012 to penalize hospitals with excess 30-day readmissions on certain measured conditions including HF. A patient-centered and multidisciplinary collaboration care approach has emerged as a TC model that hospitals can use to reduce the 30-day readmissions of patients with HF (Albert et al., 2015). The 2010 IOM report highlighted the importance of nurses in redesigning the American health care system. However, evidence of the effectiveness of an APN-led TC program in disease management is limited (Albert, et al., 2015). The purposes of the DNP project were to evaluate the efficacy of an APN-led TCP to reduce 30-day all-cause readmissions of patients with HF, and to validate the concept of TCP in HF management.

My practicum site was a 420-bed acute care urban hospital located in Southern California. Two HF APNs oversee inpatient and outpatient HF management. The 30-day

all-cause hospital readmission rate for HF patients was 28.2% in 2014 (HFNPs, personal communication, December 23, 2015). The HF APNs piloted a TCP as their QI project in August 2015, to reduce 30-day all-cause readmissions for HF and to facilitate a safe transition from hospital to home in patients with HF. The TCP consists of HF APNs' inpatient visits and HF clinic follow-up after discharge. The APN-led TC program for HF patients was not formally funded by the hospital. The hospital HF program was not currently certified as an advanced HF center by TJC. However, the hospital has been participating in the AHA GWTG–HF program for years. It was vital to conduct an evaluation of the APN-led TCP to examine the effectiveness and efficiency of APN-led TCP in reducing 30-day readmissions in patients with HF. The logic model (Hodges & Videto, 2011) served as a theoretical framework to guide my evaluation of the APN-led TCP in HF management.

### **Practice-Focused Question**

The HF all-cause-related readmission has been a dominant issue in California and for the practicum site hospital. The Health Services Advisory Group (HSAG, 2016), a CMS contracted quality improvement organization, found that the 30-day all-cause readmission rate was 18.5% in the State of California, compared with 18.6% nationwide. However, the region the hospital belonged to had the highest 30-day all-cause readmission rate in the state, at 21% (HSAG, 2016). For the 30-day all-cause readmissions for HF, the average readmission rate for the state was 25.3%; this was above the national benchmark of around 22% (HSAG, 2016).

The HSAG (2016) indicated that most readmissions were occurring between 8 to 14 days post-discharge. Breaking these statistics down by the source of readmissions shows that skilled nurse facilities (SNF) had a 30-day HF all-cause readmission rate of 27.4%, followed by 26.4% for home health agencies (HHA), and 24.4% for home (HSAG, 2016). The 2014 HSAG report showed that the 30-day HF all-cause readmission rate for the practicum site hospital was 27.5%, which was above the state level at 25.1%, and the national level of about 18% (HFNPs, personal communication, December 23, 2015). The 30-day HF all-cause readmission rate from home was 29%. This was similar to SNF, but slightly higher than HHA at 27.8% (HFNPs, December 23, 2015). The California Department Public Health (CDPH, 2016) data showed that heart disease was the leading cause of deaths in California in 2013. Heart disease has become a threat to population health; it was thus critical for me to identify barriers to HF management.

The primary barriers contributing to high readmission rates in HF patients are issues related to follow-up visits and communication about the discharge information with PCPs or cardiologists after hospital discharge (HFNPs, December 23, 2015; Yancy et al., 2013). Jongsma (2015), working in a nearby hospital, confirmed the challenge of arranging follow-up appointments with PCPs after discharge, and suggested the need to have a nurse-led outpatient clinic for facilitating timely follow-up and physician collaboration after discharge for hospitalized HF patients.

The ACCF and the AHA have recommended the adoption of a TCP system-wide in managing HF patients to reduce readmissions and improve safe and quality care (Yancy et al., 2013). Nurses, including APNs, have assumed leadership roles in leading

TC interventions and programs. However, there was limited evidence supporting the efficacy of TC programs for hospital readmission reduction in patients with HF (Albert et al., 2015). Multiple researchers have used nurses to manage HF TC programs, but only a few have mentioned the use of APN, and the educational background of nurses has not been well defined in the literature (Albert et al., 2015).

The purposes of this DNP project were to evaluate the effectiveness of the APN-led TCP at my practicum site as a QI project to reduce 30-day all-cause readmissions in patients with HF who were discharged to home. I hoped that program evaluation results would help senior managers decide whether to adopt the APNs-led TCP as an emerging care model for chronic disease management in the organization. The practice-focused question was:

For hospital discharged HF patients, was an APN-led TCP effective in reducing hospital 30-day all-cause readmissions?

#### **Source of Evidence**

I used the following sources of evidence to address the practice-focused question.



Table 2

*Sources of Evidence*

Source	Evidence
Professional and Government Organizations	American Heart Association(AHA): Professional Heart Daily/Guideline and Statements, AHA/ Get with the Guideline-HF, Heart Failure Society of American (HFSA), American Association of Heart Failure Nurses (AAHFN), American College of Cardiology (ACC), Agency for Healthcare Research and Quality (AHRQ), Centers for Disease Control and Prevention (CDC), Centers of Medicare and Medicaid Services (CMS), The Joint Commission (TJC), Walden University Library Home.
Online Research Databases	Cochrane Database of Systematic Reviews, MEDLINE with full text, CINAHL plus with full text, Pub Med, Walden University's Scholar Works.
Books	Hodges, B. C., & Videto, D. M. (2011). <i>Assessment and planning in health programs</i> . (2 <sup>nd</sup> ed). Sudbury, MA: Jones & Bartlett Learning.
Expert Opinions	Dr. Whitehead and Dr. Valdez
Practicum Site Data	Cindy Peters RN, HFNP and Pat Long RN, HFNP Hospital Archival Data from Quality Improvement Records

The evidence-based HF guidelines recommended by AHA/ACC and HFSA served as guides to identify the practice gap at the practicum site. The CMS's reimbursement rules and regulations, AHA GWGTG–HF recommendations, and AHA and the TJC joint advanced HF certification statements helped me evaluate organizational adherence to performance and quality measures related to HF management. The online research databases assisted in determining the strength and consistency of evidence-based HF practice. Hodges and Videto's (2011) book provided steps on health care program evaluation. The expert opinions ensured my program evaluation project's design and

method were appropriate for the targeted population. The practicum site data provided information on hospital quality improvement performance in HF management.

Hospital readmissions for patients with HF are costly and preventable. A patient-centered TC program has been a recommended disease management care model, but the evidence of the effectiveness of APN-led TCP in HF management has been limited (Albert et al., 2015; Feltner et al., 2014; Lindenfeld et al., 2010; Yancy et al., 2013). The project's purpose was to evaluate the effectiveness of an APN-led TCP in reducing 30-day all-cause readmissions in patients with HF.

### **Archival and Operational Data**

The practicum site had a variety of data sources for its HF program. The two HFNPs oversaw the HF data and initiated QI measures accordingly. Data were used for quality improvement, so patient consents were waived per hospital policy. Based on the Medicare claims, HSAG provided the hospital quarterly data on 30-day all-cause readmission rates for HF that compared the practicum site hospital levels with regional and state levels. The AHA's GWTG-HF generated a monthly quality measures report on processes and outcomes of HF management such as patient education and hospital mortality. However, those reports from HSAG and GWTG-HF were not disseminated quarterly. The hospital manager of quality improvement and decision support generated monthly discharge lists of HF as the index admission and quarterly HF 30-day all-cause readmission patients' records.

HFNPs have collected data on the TC program in Excel spreadsheets since the QI project HF TCP started in August 2015. The Excel spreadsheet includes the following data: 30-day readmissions, emergency room visits, mortality, compliance with evidence-based medications, Ejection Fraction (EF), age, insurance, physician follow-up appointments, and HF clinic visits. Additionally, HFNPs were authorized to access into the Electronic Medical Records (EMR) for obtaining hospital costs and have receive a utilization summary for hospitalized patients with an index of admission for HF upon their discharge on a daily basis.

The purpose of the project was to assess the efficacy of an APN-led TCP in reducing 30-day readmissions in patients with HF who were recently discharged from the hospital to home for an index admission of HF. The practicum site's HF data were relevant to helping determine the effectiveness of an APN-led TCP from hospital to home to reduce 30-day readmissions. The limitation of the HF TCP data collected by HFNPs was the inability to capture readmissions to other hospitals. The hospital's quarterly 30-day readmission report also did not including readmissions to other hospitals. The HSAG data were only applied to all Medicare fee for service patients. Furthermore, the TCP's data collection occurred in the second month after two HFNPs implemented the TCP; that could have potentially limited the number of patients participating in the APN-led TCP.

Two HF NPs kept all HF data on password-protected computers in the HF office. There was no patient consent required because the data were considered HF quality improvement per the hospital policy. The information technology (IT) department in the

hospital authorized a code of *quality improvement* to HFNPs for accessing and auditing the EMR in patients with HF. For the purpose of the project, I obtained the hospital's permission to access HF program data and analyze the existing data for program evaluation.

### **Evidence Generated for the Doctoral Project**

In this DNP project, I analyzed the existing QI data to evaluate the differences in the 30-day readmissions in patients with HF who were discharged to home and followed up their post-discharge care either with an APN-led TCP or with their community physicians including PCPs and/or cardiologists.

**Participants.** Prior to discharge, hospitalized adult patients with HF were offered to follow up in the HF clinic within one week to two weeks after discharge except for patients who fell into the following categories: hemodialysis, palliative care, and hospice care, Against Medical Advice (AMA), HHA, discharged to Skilled Nurse Facility (SNF), and Long-Term Acute Care (LTAC). Patients enrolled in a locally managed care medical group were also excluded from the APN-led TCP. HFNPs had collected one year of HF data for the TC program. The participants in the QI project were all hospitalized adult patients who were 18 years and older with an index admission of HF. The QI data covered patients with a HF index admission who were discharged to home from September 2015 to August 2016.

**Procedures.** For this project, I accessed and analyzed the existing HF data for one year from September 2015 to August 2016. The data was analyzed to determine the differences in 30-day all-cause readmissions in discharged home patients with an index

admission of HF between patients following up with an APN-led TCP and their community physicians after discharge. In this DNP project, I defined the post-discharge care occurring within 30 days of discharge. I excluded patients who were on hemodialysis, palliative care, hospice care, AMA, HHA, SNF, LTAC, and the stated local managed care group. I assigned patients who only followed their post-discharge care with the APNs in the APN-led TCP group and patients who were scheduled to follow-up their post-discharge care with their physicians in the community physicians' group. I also excluded patients who followed up their post-discharge care with both APNs and physicians from the project and patients, who never showed up for their post-discharge care appointments with the APNs.

I used the basic logic model (Hodges & Videto, 2011) to guide my program evaluation. As shown in Table 3, the explanation of logic model demonstrated processes and elements of program evaluation.

Table 3

*Explanation of the Basic Logic Model*

Logic Model Stages	Components	Purposes/Meanings	Examples
Planned work	Resources/ Inputs	To support the program 'work	Human, financial, organizational, and community resources
	Activities	To generate intended results with available resources	Program implementation: processes, tools, events, technology, and actions
Planned results	Outputs	Direct products of program activities	Types, levels and targets of services generated by the program.
	Outcomes	Specific short-term and long-term changes in program target participants	Behavior, knowledge, skills, status and level of functioning.
	Impact	Long-term effects	Intended or unintended change occurring in organizations, communities or systems

Note from "The logic model development guide," by W.K. Kellogg Foundation, (2004), Copyright 2004 by the W.K. Kellogg Foundation. Reprinted with permission.

**Protections.** All patients were protected under the Health Insurance Portability and Accountability Act of 1996 (HIPAA) privacy rule. A patient's consent was waived by the hospital because the data were being utilized for quality improvement purpose. However, I applied and obtained hospital's permission to access and analyze HF data related to an APN-led TC program evaluation as well as Walden University's IRB approval for conducting data analysis of an APN-led TCP at the practicum site. The Walden University's IRB ensured that all Walden University research met both the university's ethical standards and U.S. federal regulations (Laureate Education, Inc., 2016).

### **Analysis and Synthesis**

I analyzed the existing HF QI data to evaluate the APN-led TCP outcome in 30-day all-cause readmissions. This QI project compared the 30-day all-cause readmissions in hospital-discharged patients with HF who followed up with an APN-led TCP to patients who followed up with their community physicians (including primary care providers and/or cardiologists). The data of HF QI project ranged from September 2015 to August 2016. This QI project included adult HF patients who were 18 years old and older with a recent hospitalization's index admission of HF.

In compliance with the HIPAA privacy rule (U.S. Department of Health & Human Services, 2016), I removed patient's identification (e.g., name, medical number, and birthdates) for data analysis. I organized the de-identified HF QI data in Excel spreadsheets. Patients who followed up in the APN-led TCP had a separated Excel spreadsheet from patients who followed up with their community physicians. However,

each Excel spreadsheet had the same elements including patient de-identification number, age, sex, 30-day readmission, insurance, and HF types. For statistical analysis, each item in the Excel spreadsheet was assigned a number. For example, each patient was coded as a combination of three number (e.g., 001); for the sex, 1 represented female, and 2 represented male; patients with 30-day readmissions was coded as 1 and 0 for patients with no readmissions; for insurance, a different number represented the different insurance for patients (e.g., 5 for Medi-Cal [Medicaid] insurance); and diastolic HF was coded with 1 and 2 was for systolic HF. All data were stored in the password-required computers in the HFNPs' office.

In this DNP project, the chi-square test (Polit, 2010) was used to infer the existing relationship between categorical variables. The two categories included the 30-day readmission status (readmitted and not readmitted) in patients with HF who were discharged from the same hospital from September 2015 to August 2016 and followed up their post-discharge care with two different provider groups (APN-led TCP and a community physicians' group).

### **Summary**

HF is a common and costly population health issue. It continues to be associated with high hospital readmissions and mortality. HF 30-day readmission and mortality are quality measures and associated with reimbursement (CMS, 2015; 2016). A TC program or model is an emerging disease management care model that can reduce readmissions and mortality, improve quality of life, and reduce costs of care in patients with HF (see Albert, 2016; Feltner et al., 2014; Stamp et al., 2014).



## Section 4: Findings and Recommendations

### **Introduction**

HF is a serious public health problem. In 2012, 5.7 million adults in the United States were diagnosed with HF (Mozaffarian et al., 2016). HF is associated with high mortality rates, hospital admissions and readmissions, and costs. Patients with HF usually died within 5 years, and one in nine deaths are related to this disease (Mozaffarian et al., 2016). HF accounts for many patients' primary admission and discharge diagnosis (Mozaffarian et al., 2016).

The 2007 Medicare Payment Advisory Commission report showed that 75% of readmissions in Medicare patients were avoidable. Hence, The CMS (2016) established a HRRP in 2012 to reduce payments to hospitals with excessive amounts of readmissions. The lack of care coordination and care communication across settings, and the unavailability of post-discharge care follow-up have been identified as barriers in transitions of care that contribute to high readmission rates (Albert et al., 2015). As a result, TC has emerged as a disease management approach (Yancy et al., 2013; Albert et al., 2015). However, the efficacy of TCP in readmission reduction has been under debate because of the inconsistency of study designs and the lack of standard of TC interventions (Albert et al., 2015).

Nurses often played leadership roles in disease management programs for post-discharge care, but little was known about the effectiveness of APNs' in leading a TCP to reduce 30-day readmissions in patients with HF (Albert et al., 2015). Therefore, I

conducted this quality improvement project to evaluate the efficacy of an APN-led TCP in reducing 30-day all-cause readmissions in patients with HF.

The practice-focused question was:

For hospital-discharged HF patients, was an APN-led TCP effective in reducing hospital 30-day readmissions?

In this DNP project, I explored existing quality improvement data to evaluate the efficacy of an APN-led TCP in reducing 30-day readmissions in patients with HF. The sources of evidence were the quality improvement data from the practicum site. The HF nurse practitioners authorized my use of the quality improvement data and provided me with four different data reports: the TCP list, quarterly HF readmission lists, monthly HF discharge lists, and HSAG quarterly reports. The project's two variables were readmission status within 30 days of discharge, and post-discharge care providers (e.g., an APN-led TCP and the community physicians' group). Since two variables were categorical data, I used a chi-square test to examine the association between readmission status and post-discharge care providers. De-identified datasets were developed to include patients' age, sex, HF types, and insurance.

### **Findings and Implications**

My Walden University IRB approval number was 03-17-17-0484336. Figure 2 shows the data collection process.




Figure 2. Flowchart for data collection process.

I screened a total of 630 patients from monthly HF discharge lists from September

2015 to August 2016. Exclusions included 251 patients who were less than 18 years old, on hemodialysis, AMA, HHA, Hospice and palliative care, or discharged to SNF, LTAC, and the managed care group. Eighty-one patients participated in the APN-led TCP, and 289 patients had prearranged physician follow-up appointments. Thirty-four patients in the APN-led TCP were eliminated for no-shows and cancellations. I included a total of 345 patients in the final analysis. Forty-seven of these patients were in the APN-led TCP, and 298 patients were in the physicians' group.


Table 4  
*Cross Tabulations and Chi-Square Results for 30-Day Readmissions With Post-Discharge Care Providers*

Providers	Readmission status			$X^2$	$df$	$p$
	Yes	No	Total			
APN-led	7	40	47	0.236	1	0.627
TCP						
Physicians	53	245	298			
Total	60	285	345			

The results of the chi-square analysis showed a nonsignificant association between 30-day readmission status and post-discharge care with different health care providers [ $\chi^2(1, N = 345) = 0.236, p = 0.627$ ]. The critical chi-square statistical value for  $p=0.05$  with 1 degree of freedom was 3.84, and the calculated chi-square value was less than the critical value ( $0.236 < 3.84$ ). There was not a statistically significant association between readmission status and post-discharge care providers. Hence, I concluded that the 30-day

readmission rates were same between the APN-led TCP and the community physicians' group.


I demonstrated patients' mean age and sex distribution for both groups in Figure 3.



*Figure 3.* Bar graph showing the sex distribution for each group.

The Figure 3 bar graph shows patients' mean age and female and male distribution for the APN-led TCP and the physicians' group. Patients in the APN-led TCP were approximately 16 years younger than those who were in the physician's group (49.3% vs. 65.62%). More than two thirds of patients in the APN-led TCP were male, and less than one third of them were female (76.6% vs. 23.3%). In the physician's group, the distribution of male and female patients was similar (56.71% vs. 43.29%).


I displayed patients' HF types based on EF scores in the Figure 4.



*Figure 4.* Bar graph showing HF types for each group.

The Figure 4 bar graph shows the distribution of diastolic and systolic HF for each group. The 89.4% of patients in the APN-led TCP had systolic HF that was known to have a higher mortality rate, and the remaining 10.6% of patients had diastolic HF. On the contrary, the distributions of systolic and diastolic HF were similar (51.34% vs. 48.66%).

I summarized payer sources for each group in the Figure 5.




*Figure 5.* Bar graph showing insurance types for each group.

The Figure 5 bar graph showed patients' insurance in each group. In the APN-led TCP, most patients had Medi-Cal (48.94%), followed by Medicare (21.28%), PPO (19.15%), and HMO (10.64%). On the other hand, more than half of patients (51%) in the physician's group had Medicare, followed by Medi-Cal (38.39%), PPO (5.03%), and HMO (4.70%). There was a high percentage rate of patients in the APN-led TCP who had restricted care access because of their low-income status compared with patients in the physicians' group.

Figure 3, Figure 4, and Figure 5 show patients' demographic characteristics in age, sex, HF type, and insurance between two groups. Because the statistical tools were not used to calculate those demographic differences between two groups, I was not able to conclude whether the population characteristics were statistically significantly different in age, sex, HF types, and insurances between the APN-led TCP and the community physicians' group.

I also included the 30-day all-cause readmission rates from home in Medicare patients with HF before, during, and after the implementation of the APN-led TC in the Figure 6.



*Figure 6.* Bar graph showing comparisons of 30-day all-cause readmission rates from home in Medicare patients with HF from Health Services Advisory Group's Reports.

The Figure 6 bar graph shows the 30-day all-cause readmission rates from home in Medicare patients with HF before and after the implementation of the APN-led TCP. The APN-led TCP was implemented in Q3 2015. The all-cause 30-day HF readmission rates from home were 29% in Q2 2013 to Q1 2014, followed by 26.8% in Q2 2015 to Q1 2016, and 24.4% in Q3 2015 to Q2 2016. Those two HF nurse practitioners' rationale for starting the APN-led TCP was based on the HF all-cause readmission rates from HSAG reports Q2 2013 to Q1 2014. They implemented the APN-led TCP in Q3 2015 as their quality improvement project to reduce the 30-day HF readmissions. At the same time, this group added a discharge planner to the HF program to arrange follow-up



appointments for hospitalized patients with HF before releasing them to home. Additionally, two HF NPs assigned the discharge planner a home phone call list within 72 hours of discharge. In the project, patients who were in SNF, on hospice, and with HHA were not included. Some patients had received pharmacists' consultations on medications before they went home; however, this was not consistent practice for all hospitalized patients with HF. There was a trending down pattern in readmission reduction from home in Medicare patients with HF after the implementation of the APN-led TCP. Although this DNP project did not address the HF readmissions based on payer sources, the HSAG's data on HF readmissions in patients with Medicare supported the value of the APN-led TCP in readmission reduction.

In this DNP project, limitations included the small number of patients in the APN-led TCP compared with the physicians group, the inability to capture 30-day readmissions to other hospitals, and the inability to confirm whether patients in the physicians group followed up their prearranged post-discharge care appointments. At my practicum site, the two HFNPs implemented this quality improvement project (the APN-led TCP) in August, 2015, and they started to collect the program data in September, 2015. Therefore, the short timing might cause the small number of patients participated in the APN-led TCP. As the DNP project was a quality improvement data analysis, I was not authorized to call patients and the community physicians to find out whether patients had been readmitted to other hospitals within 30 days of post-discharge as well as whether they followed up their prearranged post-discharge care appointments with their physicians. Hence, the readmission rates in the project were not representing the overall

readmissions for all discharged patients with HF in the practicum hospital, and the number of patients in the community physicians' group was not the number of patients who actually followed up their post-discharge care appointments with their physicians.

The implications resulting from the findings in terms of individuals, communities, institutions, and systems were that the APNs played important roles in providing TC to patients with HF for readmission reduction. In this DNP project, the chi-square analysis results revealed that the readmission rates were same between the APN-led TCP and the physicians' group in providing TC to hospital-discharged patients with HF. However, 48.94% of patients in the APN-led TCP were Medi-Cal, and most of them had no primary care providers. Without the APN-led TCP, the hospital HF readmission outcome could be worse than the current status because of the lack of post-discharge care follow-up for patients with Medi-Cal. Furthermore, the hospital HSAG's data revealed the trending down HF readmissions from home in Medicare Patients after implementing the APN-led TCP; this result demonstrated the value of APN-led TCP in readmission reduction. Readmission was one of the quality measures that impacted hospital reimbursement. Because of the APN-led TCP's readmission outcome as same as the physicians, providing post-discharge care to a large proportion of Medi-Cal patients, impacting readmissions for Medicare patients from home, the hospitals' administrator should consider supporting the APN-led TCP. And they should continue to collect another 6-month HF QI data, such as also to include readmission rates based on payer sources, to validate the efficacy of the APN-led TCP to reduce 30-day readmissions in patients with HF.

From this DNP project's findings, the potential implications to positive social change were increasing care access and improving population health for low-income patients with HF. Although the DNP project did not validate the efficacy of APN-led TCP to reduce 30-day readmissions in patients with HF, patients were different in terms of payer sources and HF type, between the APN-led TCP and the physicians group. Regarding the insurance type, more than half of patients in the APN-led TCP had Medical insurance while half of patients in the physicians group had Medicare. In general, physicians accepted patients with Medicare, and few physicians took patients with Medical which was Medicaid in California. Primary care physicians were reluctant to take Medicaid patients because of the lower reimbursement rate and longer reimbursement time compared with Medicare and commercial insurance (Long, 2013). Additionally, patients with Medicaid insurance had more social and behavior problems, and transportation and compliance issues; hence, physicians spent more time treating them than patients with private insurance whom had similar medical problems (Long, 2013). The lower reimbursement rates and complex of care accounted for the lack of primary care providers for patients with Medicaid insurance. Furthermore, patients with systolic HF had a higher mortality rate and poorer progress than those who had diastolic HF (Mosterd & Hoes, 2007). In the DNP project, the majority of patients in the APN-led TCP had systolic HF. With the unwillingness of physicians to take care of those sick patients, the APN-led TCP was a great access to patients with low-income and less primary care access. Therefore, the availability of care access could potentially improve the population health, reduce unnecessary readmission, improve mortality associated with

hospitalizations, and reduce medical care spending. However, more data analysis was needed to examine the relationship between payer sources and readmissions, with a large number of patients in the APN-led TCP.

### **Recommendations**

The proposed recommendation that would potentially address the gap-in-practice was to continue the APN-led TCP for hospital-discharged patients with chronic conditions such as HF until another 6-month HF QI data were collected and analyzed. The lack of care coordination and care transition from hospital to home had contributed to high hospital readmissions (Albert et al., 2015). Additionally, the difficulty of making post-discharge follow-up appointments with physicians had been considered one of the barriers for causing the lack of care transition for hospital-discharged patients. This DNP project's findings indicated that there were no statistically significant differences in readmission reduction between the APN-led TCP and the community physicians' group, and the 30-day all-cause readmission rates for HF were same for hospital-discharged patients received their post-discharge care provided either by the APN-led TCP or the physicians' group. In this DNP project, the finding demonstrated that the average ages in the APN-led TCP were 14.5 years younger than the physicians group; more than half of them had Medi-Cal insurance and did not have their primary care physicians. Hence, it was important to support the APN-led TCP as an alternative care approach in overcoming physician shortage and promoting care transitions from hospital to home in order to combating the high hospital readmissions in patients with HF. Meanwhile, more data collection and analysis including the relation between payer sources and readmissions

were needed to validate the efficacy of the APN-led TCP to reduce 30-day readmissions in patients with HF.

The current APN-led TCP included two nurse practitioners, one discharge planner, and one LVN. The discharge planner played a critical role in setting up post-discharge follow-up appointments with the APN-led TCP or the community physicians before discharge, calling patients at home within 72 hours of hospital discharge, verifying patients' insurance, and finding their primary care providers for patients in the APN-led TCP. The part-time LVN assisted two NPs in operating the HF TC clinic, collecting patients' medical records including laboratory and other diagnostic test results, and sending medical records to other providers. Two APNs had overseen the TCP to ensure the safety and quality of care transitions from hospital to home for patients with HF. During hospitalizations, APNs along with the multidisciplinary team provided patient education and assisted with the anticipated discharge processes. For patients in the TCP, two NPs continued to see patients in the TC HF clinic within the first 2 weeks after discharge and made care transitions to their PCPs accordingly. Additionally, hospital discharge pharmacists only provided consultations to patients in the hospital and provided 30-day free discharge medication for patients either could not afford it or had no insurance. A future recommendation was that the APN-led TCP should involve a multidisciplinary team, comprised of social workers and or discharge planners, nurses, nurse practitioners, and pharmacists. Stranges et al. (2015) also reported a similar practice model that a multidisciplinary team approach including medical providers,

clinical pharmacists, and social workers could reduce 30-day all-cause readmissions. Hence, a multidisciplinary team approach was an essential part of an APN-led TCP.

### **Strength and Limitations of the Project**

Although this DNP project did not validate the efficacy of the APN-led TCP in readmission reduction in patients with HF, the project findings still demonstrated the APNs' leadership roles in redesigning health care delivery to promote population health and to reduce health care costs. The project's findings demonstrated that the 30-day readmissions were not associated with post-discharge care providers, and the readmission rates were same between the APN-led TCP and the physicians' group. The HSAG's reports demonstrated the trending down readmission rates in patients with HF who were readmitted from home after HF nurse practitioner started the APN-led TCP. The HFNPs implemented the APN-led TCP in August 2015. The HSAG readmission rate from home in Q 2 2013 to Q1 2014 was 29%, and the readmission rate was reduced to 24.4% in Q3 2015 to Q 2 2016 after starting the TCP. The APN-led TCP and the prearranged physician post-discharge follow-up appointments by the discharge planner might contribute the reduction in readmissions from home for Medicare patients. Two HFNPs piloted the APN-led TCP as a quality improvement project after they saw the alarming readmissions in patients with HF on the HSAG's reports. They also added a discharge planner to the HF program to help make post-discharge follow up appointments, and phone call follow up within 72 hours of discharge. The TC was an emerging disease management model in reducing hospital readmission in patients with HF; however, the evidence of the APN's role in leading the TCP was limited (Albert et al., 2015). This

project's results validated the APN's leadership roles in health care reform to improve quality of care and population health.

This DNP project had several limitations. First, I was unable to verify whether patients in the physicians group followed up their prearranged post-discharge care appointments. I confirmed that all patients in the APN-led TCP followed up their post-discharge care with the APN-led TCP within 2 weeks of discharge. Hence, the uncertainty of physicians' post-discharge follow-up weakens the project findings. Second, there were a disproportionate number of patients between groups. I had 47 patients in the APN-led TCP and 298 patients in the physicians group. As I collected data reports of the APN-led TCP 1 month after the program started, this could cause a small number of patients in the APN-led TCP. Third, the non-randomization project design limited the generalizability of findings. Fourth, there was a lack of standardization on TC strategies between the APN-led TCP and the physicians' group. Patients in the APN-led TCP received a consistent standard of care. However, various physicians provided post-discharge care to patients in the physicians' group. The standard of TC interventions between the two groups was not compared.

## Section 5: Dissemination Plan

### **Dissemination Plan**

To disseminate the DNP project findings to the practicum site, I will present the project results at a quarterly HF multidisciplinary meeting and at a quarterly department cardiologist meeting. My preceptor and I will meet the vice president of the Heart Institute to discuss my project findings and recommendations including formally adopting the APN-led TCP, adding a pharmacist to the APN-led TCP, and engaging physicians in developing the standard of TC strategies. Lastly, I will submit a project poster to regional and national conferences to disseminate the project findings.

The appropriate audiences for dissemination of the doctoral project to the broader nursing profession include HF nurses, nurse practitioners, care managers, and nurse leaders who deal with hospital readmissions. Other audiences include social workers, discharge planners, pharmacists, insurance representatives, quality improvement experts, and physicians. Appropriate venues for dissemination of the project may be the America Heart Failure Nurse annual conference, the American Association of Nurse Practitioners annual conference, and the practicum site annual research conference.

### **Analysis of Self**

The DNP project has enriched my professional and personal experiences as a practitioner, scholar, and project manager. I had been working in the acute care setting for more than 20 years in different roles, and had known that patient education was critical while patients were in the hospital. However, I had not considered how much information they could retain while they were in physical and emotional distress in the



hospital. The clinical experiences helped me realize that post-discharge education, such as information on medication, diet, activities, disease processes, recognizing worsening signs and symptoms, and follow-up care, were extremely important to build patients' self-management skills at home. HF is a chronic condition, and patients' self-management skills are essential for disease management and readmission reduction. The TC was an effective approach to continue reinforcing patients' education and to address their needs from the hospital to home, such as transportation to clinics. Furthermore, from the clinical experience, I learned that many primary care physicians and even some cardiologists were not familiar with the evidence-based HF practice guidelines. Thus, it is pivotal to have HF experts properly manage patients with HF. The health care system is complex, especially in regard to managed care. I have seen patients who had insurance for a long time but were not able to see their assigned doctors because the doctors were far away from where they lived. The APN-led TCP helped them contact their insurance provider and connected them with different doctors they could reach easily. The after-discharge care was a critical window, and many unaddressed patients' needs could cause hospital-discharged patients back to hospitals. From this DNP project, I have developed my long-term professional goal of working on TC for patients with chronic conditions.

The DNP project was a long scholarly journey. Along the way, I have gained tremendous support, guidance, and help from my committee members and the doctoral writing workshops. The challenges I encountered were the project designs as a quality improvement project, the selection of statistical tools, and scholarly writing. The insights

I gained were related to academic processes such as the project design, literature review, IRB application, and scholarly writing strategies.

### **Summary**

The TC model is an emerging disease management practice used to reduce readmissions in patients with HF. However, the literature included limited evidence on the effectiveness of APN-led TCP for readmission reductions. In this DNP project, I extracted and analyzed existing quality improvement data reports, and compared readmission statuses between an APN-led TCP and the physicians' group. There were no statistically significant differences on readmission status between two groups. This DNP project highlighted the APNs' leadership roles in TC programs to reduce readmissions and costs. It also highlighted clinical roles in providing medical care to those patients with low income. The project also clearly showed that a TCP was a patient-centered multidisciplinary team approach that consisted of nurse practitioners, nurses, social workers, and pharmacists. However, there were numerous weaknesses that limited the broad applications of project findings: the uncertainty of follow-up, the lack of TC strategies, a small sample size, a convenient sampling method, and a single project site. Hence, I recommended a future study on payer sources and readmissions to include a randomized research method with multiple study sites to strengthen study findings.

## References

- Albert, N. M., Barnason, S., Deswal, A., Hernandez, A., Kociol, R., Lee, E., White-Williams, C. (2015). Transitions of care in heart failure: a scientific statement from the American Heart Association. *Circulation. Heart Failure*, 8(2), 384-409. doi:10.1161/HHF.0000000000000006
- Albert, N. M. (2016). A systematic review of transitional-care strategies to reduce rehospitalization in patients with heart failure. *Heart & Lung: The Journal of Critical Care*, 45(2), 100-113. doi:10.1016/j.hrtlng.2015.12.001
- American Heart Association. (2016). Get with the guidelines- heart failure. Retrieved from [http://www.heart.org/idc/groups/heart-public/@wcm/@hcm/@gwtg/documents/downloadable/ucm\\_467882.pdf](http://www.heart.org/idc/groups/heart-public/@wcm/@hcm/@gwtg/documents/downloadable/ucm_467882.pdf)
- Bradley, E. H., Curry, L., Horwitz, L. I., Sipsma, H., Wang, Y., Walsh, M. N., ... Krumholz, H. M. (2013). Hospital strategies associated with 30-day readmission rates for patients with heart failure. *Circulation. Cardiovascular Quality and Outcomes*, 6(4), 444-450. doi:10.1161/CIRCOUTCOMES.111.000101
- Burns, P. B., Rohrich, R. J., & Chung, K. C. (2011). The levels of evidence and their role in evidence-based medicine. *Plastic and Reconstructive Surgery*, 128(1), 305-310. doi:10.1097/PRS.0b013e318219c17
- California Department Public Health. (2016). Center for health statistics and informatics death data trend summary. Retrieved from <http://www.cdph.ca.gov/programs/ohir/Pages/DDTSummaryIndex.aspx>
- Camicia, M., Chamberlain, B., Finnie, R. R., Nalle, M., Lindeke, L. L., Lorenz, L., & ...

- McMenamin, P. (2013). The value of nursing care coordination: a white paper of the American Nurses Association. *Nursing Outlook*, 61(6), 490-501. Retrieved from <http://www.nursingworld.org/carecoordinationwhitepaper>
- Campbell, K.S. (2016). Enhancing Interest and knowledge of how to start a nurse practitioner-led clinic (Unpublished doctoral dissertation). Otterbein University, Westerville, Ohio.
- Centeno, M. M., & Kahveci, K. L. (2014). TC models: preventing readmissions for high-risk patient populations. *Critical Care Nursing Clinics of North America*, 26(4), 589-597. doi:10.1016/j.ccell.2014.08.009
- Centers for Disease Control and Prevention. (2010). Logic model. Retrieved from [http://www.cdc.gov/nccdphp/dnpao/hwi/programdesign/logic\\_model.htm](http://www.cdc.gov/nccdphp/dnpao/hwi/programdesign/logic_model.htm)
- Centers of Disease Control and Prevention.(2015). Heart failure fact sheet.Retrieved from [http://www.cdc.gov/dhdsp/data\\_statistics/fact\\_sheets/fs\\_heart\\_failure.htm](http://www.cdc.gov/dhdsp/data_statistics/fact_sheets/fs_heart_failure.htm)
- Centers for Medicare & Medicaid Services. (2015a). 2014 Measure information about the 30-day all cause hospital readmission measure, calculated for the value-based payment modifier program. Retrieved from <https://www.cms.gov/Medicare/Medicare-Fee-for-ServicePayment/PhysicianFeedbackProgram/Downloads/2014-ACR-MIF.pdf>
- Centers for Medicare & Medicaid Services. (2015b). Outcome measures. Retrieved from <https://www.cms.gov/medicare/quality-initiatives-patient-assessment-instruments/hospitalqualityinits/outcomemeasures.html>
- Centers for Medicare & Medicaid Services. (2016). Readmissions reduction program

(HRRP). Retrieved from <https://www.cms.gov/medicare/medicare-fee-for-service-payment/acuteinpatientpps/readmissions-reduction-program.5html>

Chang, P. P., Chambless, L. E., Shahar, E., Bertoni, A. G., Russell, S. D., Ni, H., ...

Rosamond, W. D. (2014). Incidence and survival of hospitalized acute decompensated heart failure in four US communities (from the Atherosclerosis Risk in Communities Study). *American Journal of Cardiology*, *113*(3), 504-510.  
doi:10.1016/j.amjcard.2013.10.032

Davidson, P.M., Cockburn, J., & Newton, P.J. (2008). Unmet needs following hospitalization with heart failure: implications for clinical assessment and program planning. *Journal of Cardiovasc Nurses*, *23*, 541–546.

doi:10.1097/01.JCN.0000338927.43469.35

DeVore, A. D., Cox, M., Eapen, Z. J., Yancy, C. W., Bhatt, D. L., Heidenreich, P. A., ...

Hernandez, A. F. (2016). Temporal trends and variation in early scheduled follow-up after a hospitalization for heart failure: Findings from Get With The Guidelines-Heart Failure. *Circulation: Heart Failure*, *9*(1).

doi:10.1161/CIRCHEARTFAILURE.115.002344

Dharmarajan, K., Hsieh, A.F., Lin Z, Bueno, H., Ross, J.S., Horwitz, L.I., Barreto-Filho,

J.A., ... Krumholz, H.M.(2013). Diagnoses and timing of 30-day readmissions after hospitalization for heart failure, acute myocardial infarction, or pneumonia. *Journal of the American Medical Association*, *309*(4), 355-363.

doi:10.1001/jama.2012.216476

Driscoll, A. (2016). Nurse-led titration of angiotensin converting enzyme inhibitors, beta-

adrenergic blocking agents, and angiotensin receptor blockers for people with heart failure with reduced ejection fraction. *Cochrane Database of Systematic Reviews*, (1), 1-41. doi:10.1002/14651858.CD009889.pub2

- Echeverry, L. M., Lamb, K. V., & Miller, J. (2015). Impact of APN home visits in reducing healthcare costs and improving function in homebound heart failure. *Home Healthcare Now*, 33(10), 532-537. doi:10.1097/NHH.0000000000000304
- Feltner, C., Jones, C. D., Cené, C. W., Zheng, Z., Sueta, C. A., Coker-Schwimmer, E. L., ... Jonas, D. E. (2014). TC interventions to prevent readmissions for persons with heart failure: a systematic review and meta-analysis. *Annals of Internal Medicine*, 160(11), 774-784. doi:10.7326/M14-0083
- Grove, S., Burns, N., & Gray, J. (2013). *The practice of nursing research: Appraisal, synthesis, and generation of evidence* (7<sup>th</sup>ed.). St. Louis, MO: Saunders Elsevier
- Guirguis-Blake, J. (2016). TC Interventions to prevent readmissions for patients with heart failure. *American Family Physician*, 93(5), 401-403. Retrieved from <http://www.aafp.org/afp/2016/0301/p401.html>
- Health Service Advisory Group (2016). Medicare fee-for-service (FFS) hospital readmissions: Q3 2014—Q2 2015 State of California. Retrieved from [http://www.hsag.com/globalassets/providers/nplh-ca/california\\_statewide\\_readmission\\_report.pdf](http://www.hsag.com/globalassets/providers/nplh-ca/california_statewide_readmission_report.pdf)
- Hodges, B. C., & Videto, D. M. (2011). *Assessment and planning in health programs* (2<sup>nd</sup>ed). Sudbury, MA: Jones & Bartlett Learning
- Hall, M.J., Levant, S., & DeFrances, C.J. (2012). Hospitalization for congestive heart

failure: United States, 2000–2010. *NCHS Data Brief, 108*, 1-8. Retrieved from <https://www.cdc.gov/nchs/data/databriefs/db108.pdf>

Institute of Medicine (IOM). (2010). *The future of nursing: Leading change, advancing health*. Retrieved from <https://www.nationalacademies.org/hmd/~/media/Files/Report%20Files/2010/The-Future-of-Nursing/Future%20of%20Nursing%202010%20Recommendations.pdf>.

Institute of Medicine (IOM). (2001). *Crossing the quality chasm: A new health for the 21<sup>st</sup> century*. Retrieved from <https://www.nationalacademies.org/hmd/~/media/Files/Report%20Files/2001/Crossing-the-Quality-Chasm/Quality%20Chasm%202001%20%20report%20brief.pdf>.

Jongsma, M. (2015). *Care transition gaps: Risk identification and intervention* (Unpublished doctoral dissertation). Walden University, Minneapolis, Minnesota.

Kansagara, D., Chiovaro, J. C., Kagen, D., Jencks, S., Rhyne, K., O'Neil, M. . . .

Englander, H. (2016). So many options, where do we start? An overview of the care transitions literature. *Journal of Hospital Medicine, 11*(3), 221-230.  
doi:10.1002/jhm.2502

Kociol, R. D., Peterson, E. D., Hammill, B. G., Flynn, K. E., Heidenreich, P. A., Piña, I. L., ... Hernandez, A. F. (2012). National survey of hospital strategies to reduce heart failure readmissions: Findings from the Get With the Guidelines-Heart Failure registry. *Circulation: Heart Failure, 5*(6), 680-

687.doi:10.1161/CIRCHEARTFAILURE.112.967406

- Krumholz, H.M., Currie, P.M., Riegel, B., Phillips, C.O., Peterson, E.D., Smith, R., ... Faxon, D.P. (2006). A taxonomy for disease management: A scientific statement from the American Heart Association Disease Management Taxonomy Writing Group. *Circulation, 114*(13), 1432-45. doi:10.1161/CIRCULATIONAHA.106.177322
- Ladd, S., Sitaker, M., Patanian, M., & Jernigan, J. (2008). Adapting logic models over time: The Washington State Heart Disease and Stroke Prevention Program Experience. *Preventing Chronic Disease, 5*(2), A60. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2396971/>
- Lambrinou, E., Kalogirou, F., Lamnisis, D., & Sourtzi, P. (2012). Effectiveness of heart failure management programmes with nurse-led discharge planning in reducing re-admissions: A systematic review and meta-analysis. *International Journal of Nursing Studies, 49*(5), 610-624. doi:10.1016/j.ijnurstu.2011.11.002
- Laureate Education, Inc. (2016). Institutional review board for ethical standards in research. Baltimore, MD: Author. Retrieved from <http://academicguides.waldenu.edu/researchcenter/orec>
- Lindenfeld, J., Albert, N., Boehmer, J., Collins, S., Ezekowitz, J., Givertz, M. . . . Walsh, M. (2010). HFSA 2010 Comprehensive heart failure practice guideline. *Journal of Cardiac Failure, 16*(6), e1-194. doi:10.1016/j.cardfail.2010.04.004
- Long, S. K. (2013). Physicians may need more than higher reimbursements to expand Medicaid participation: findings from Washington State. *Health Affairs (Project Hope), 32*(9), 1560-1567. doi:10.1377/hlthaff.2012.1010



- Lowery, J., Hopp, F., Subramanian, U., Wiitala, W., Welsh, D. E., Larkin, A. . . .
- Vaitkevicius, P. (2012). Evaluation of a nurse practitioner disease management model for chronic heart failure: a multi-site implementation study. *Congestive Heart Failure (Greenwich, Conn.)*, 18(1), 64-71. doi:10.1111/j.1751-7133.2011.00228.x
- Medicare Payment Advisory Commission (U.S.). (2007). Report to the Congress: Promoting greater efficiency in Medicare. Retrieved from [http://www.medpac.gov/docs/default-source/reports/Jun07\\_EntireReport.pdf?sfvrsn=0](http://www.medpac.gov/docs/default-source/reports/Jun07_EntireReport.pdf?sfvrsn=0)
- Madigan, E. A., Gordon, N. H., Fortinsky, R. H., Koroukian, S. M., Piña, I., & Riggs, J. S. (2012). Rehospitalization in a national population of home health care patients with heart failure. *Health Services Research*, 47(6), 2316-2338. doi:10.1111/j.1475-6773.2012.01416.x
- Mahramus, T. L., Penoyer, D. A., Sole, M. L., Wilson, D., Chamberlain, L., & Warrington, W. (2013). Clinical nurse specialist assessment of nurses' knowledge of heart failure. *Clinical Nurse Specialist CNS*, 27(4), 198-204. doi:10.1097/NUR.0b013e3182955735
- Melynk, B., & Fineout-Overholt, E. (2005). *Evidence-based practice in nursing and healthcare*. Philadelphia, PA: Lippincott Williams & Wilkins
- Mikulich, S., Parikh, A., Franklin, G., Grelak, S., Echols, G., & Johnson, D. (2013). Decreasing Heart Failure Readmissions with APN led Heart Failure Clinic. *Heart & Lung*, 42(6), e7. doi:10.1016/j.hrtlng.2013.10.008

- Monza, K., Harris, D., & Shaw, C. (2015). The role of the nurse navigator in the management of the heart failure patient. *Critical Care Nursing Clinics of North America*, 27(4), 537-549. doi:10.1016/j.cnc.2015.07.010
- Mosterd, A., & Hoes, A. W. (2007). Clinical epidemiology of heart failure. *Heart*, 93(9), 1137–1146. doi:0.1136/hrt.2003.025270
- Mozaffarian, D., Benjamin, E. J., Go, A. S., Arnett, D. K., Blaha, M. J., Cushman, M., ... Turner, M. B. (2016). Heart disease and stroke statistics-2016 update: A report from the American Heart Association. *Circulation*, 133(4), e38-e360. doi:10.1161/CIR.0000000000000350
- National Quality Measures Clearinghouse (NQMC).(2015). Measure summary: Heart failure (HF): hospital 30-day, all-cause, unplanned risk-standardized readmission rate (RSRR) following HF hospitalization. In: National Quality Measures Clearinghouse (NQMC) [Web site]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ). Retrieved from <https://www.qualitymeasures.ahrq.gov>
- Naylor, M. D., Aiken, L. H., Kurtzman, E. T., Olds, D. M., & Hirschman, K. B. (2011). The care span: The importance of TC in achieving health reform. *Health Affairs (Project Hope)*, 30(4), 746-754. doi:10.1377/hlthaff.2011.0041
- Nelson, J. M & Pulley, A. L. (2015). TC can reduce hospital readmissions. *Am NursToday*.10(4). Retrieved from <http://www.medscape.com/viewarticle/844301>
- Polit, D.K. (2010). *Statistics and data analysis for nursing research* (2<sup>nd</sup>ed). Upper Saddle River, NJ: Pearson Education Inc

- Price, A. (2012). Specialist nurses improve outcomes in heart failure. *Nursing Times*, 108(40), 22-24. Retrieved from <http://www.nursingtimes.net/clinical-subjects/cardiology/specialist-nurses-improve-outcomes-in-heart-failure/5049958.fullarticle>
- Roger, V. L. (2013). Epidemiology of heart failure. *Circulation Research*, 113(6), 646-659. doi:10.1161/CIRCRESAHA.113.300268
- Russell, D., Rosati, R. J., Sobolewski, S., Marren, J., & Rosenfeld, P. (2011). Implementing a TC program for high-risk heart failure patients: findings from a community-based partnership between a certified home healthcare agency and regional hospital. *Journal for Healthcare Quality: Official Publication of the National Association For Healthcare Quality*, 33(6), 17-23. doi:10.1111/j.1945-1474.2011.00167.x
- Smith, C. E., Piamjariyakul, U., Dalton, K. M., Russell, C., Wick, J., & Ellerbeck, E. F. (2015). Nurse-led multidisciplinary heart failure group clinic appointments: Methods, materials, and outcomes used in the clinical trial. *Journal of Cardiovascular Nursing*, 30(4 Suppl 1), S25-S34. doi:10.1097/JCN.0000000000000255
- Stamp, K. D., Machado, M. A., & Allen, N. A. (2014). TC programs improve outcomes for heart failure patients. *Journal of Cardiovascular Nursing*, 29(2), 140-154. doi:10.1097/JCN.0b013e31827db560
- Stauffer, B., Fullerton, C., Fleming, N., Ogola, G., Herrin, J., Stafford, P., & Ballard, D. (2011). Effectiveness and cost of a TC program for heart failure: a prospective

study with concurrent controls. *Archives of Internal Medicine*, 171(14), 1238-1243. doi:10.1001/archinternmed.2011.274

Stranges, P. M., Marshall, V. D., Walker, P. C., Hall, K. E., Griffith, D. K., & Remington, T. (2015). A multidisciplinary intervention for reducing readmissions among older adults in a patient-centered medical home. *American Journal of Managed Care*, 21(2), 106-113

Takeda, A. (2012). Clinical service organisation for heart failure. *Cochrane Database of Systematic Reviews*, (9), doi:10.1002/14651858.CD002752.pub3

The Joint Commission.(2016). Revisions to advanced certification requirements for heart failure. Retrieved from [https://www.jointcommission.org/assets/1/18/Prepub\\_HF\\_Maintenance\\_Jan2017.pdf](https://www.jointcommission.org/assets/1/18/Prepub_HF_Maintenance_Jan2017.pdf)

Thompson, C. (2014). Safe passage: Anurse-led multidisciplinary team approach to improving transitions and reducing readmissions for heart failure patients. *Heart & Lung*, 43(4), 386. doi:10.1016/j.hrtlng.2014.06.021

U.S. Department of Health & Human Services. (2016). Guidance regarding methods for de-identification of protected health information in accordance with the Health Insurance Portability and Accountability Act (HIPAA) privacy rule. Retrieved from <http://www.hhs.gov/hipaa/for-professionals/privacy/special-topics/de-identification/>

Vedel, I., & Khanassov, V. (2015). TC for patients with congestive heart failure: A systematic review and meta-analysis. *Annals of Family Medicine*, 13(6), 562-571.

doi:10.1370/afm.1844

W. K. Kellogg Foundation. (2004). The logic model development guide. Battle Creek, Michigan. Retrieved from

<http://www.smartgivers.org/uploads/logicmodelguidepdf.pdf>

White, S. M., & Hill, A. (2014). A heart failure initiative to reduce the length of stay and readmission rates. *Professional Case Management*, 19(6), 276-284.

doi:10.1097/NCM.0000000000000059

Yancy, C. W., Jessup, M., Bozkurt, B., Butler, J., Casey, D. J., Drazner, M. H., & ...

Wilkoff, B. L. (2013). 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Journal of the American College of Cardiology*, 62(16), e147-e239. doi:10.1016/j.jacc.2013.05.019

## Appendix: Literature Summary with Level of Evidence

Reference	Research Method	Main Findings	Level of Evidence
Albert, N. M., Barnason, S., Deswal, A., Hernandez, A., Kociol, R., Lee, E., & ... White-Williams, C. (2015). Transitions of care in heart failure: a scientific statement from the American Heart Association. <i>Circulation. Heart Failure</i> , 8(2), 384-409. doi:10.1161/HHF.000000000000006	Systematic review of RCTs	TC programs were used to reduce HF 30-day readmission. The evidence is weak due to the variety of study designs, methods, aims, and population. More researches are needed to validate TC interventions.	Level 1
Albert, N. M. (2016). A systematic review of transitional-care strategies to reduce rehospitalization in patients with heart failure. <i>Heart &amp; Lung: The Journal Of Critical Care</i> , 45(2), 100-113. doi:10.1016/j.hrtlng.2015.12.001	Systematic review of RCTs	TC models reduced readmissions and improved quality of life in patients with HF. Eight common themes of TC models included discharge planning, the multidisciplinary team approached collaboration and communication, timely and organized critical information; medication reconciliation and adherence, social and community services support, patient education including monitoring signs and symptoms of HF post discharge, outpatient clinic follow-up, and palliative care and end of life care.	Level 1
Centeno, M. M., & Kahveci, K. L. (2014). TC models: preventing readmissions for high-risk patient populations. <i>Critical Care Nursing Clinics Of North America</i> , 26(4), 589-597. doi:10.1016/j.ccell.2014.08.009	Cohort study	A hospital TC program reduced readmissions	Level IV
Driscoll, A. (2016). Nurse-led titration of angiotensin converting	Cochrane systematic	Nurse -Led Titration(NLT) of medication titration	Level I

<p>enzyme inhibitors, beta-adrenergic blocking agents, and angiotensin receptor blockers for people with heart failure with reduced ejection fraction. <i>Cochrane Database Of Systematic Reviews</i>, (1), doi:10.1002/14651858.CD009889.pub2</p>	<p>review</p>	<p>reduced all-cause hospital readmissions and mortality in patients with HFrEF (seven RCTs with a total of 1684 participants)</p>	
<p>DeVore, A. D., Cox, M., Eapen, Z. J., Yancy, C. W., Bhatt, D. L., Heidenreich, P. A., &amp; ... Hernandez, A. F. (2016). Temporal Trends and Variation in Early Scheduled Follow-Up After a Hospitalization for Heart Failure: Findings from Get With The Guidelines-Heart Failure. <i>Circulation. Heart Failure</i>, 9(1), doi:10.1161/CIRCHEARTFAILURE.115.002344</p>	<p>Descriptive Study</p>	<p>Data sources were Get with the Guidelines-HF and Medicare claims. Individual patients including older age, anemia, DM, CKD, the use of anticoagulation at discharge were more likely for scheduled early follow-up. And there was a small improvement of early follow-up from 26% to 30%. (52,438 patients discharged from 239 hospitals from 2009 to 2012)</p>	<p>Level VI</p>
<p>Feltner, C., Jones, C. D., Cené, C. W., Zheng, Z., Sueta, C. A., Coker-Schwimmer, E. J., &amp; ... Jonas, D. E. (2014). TC Interventions To Prevent Readmissions for People With Heart Failure. <i>Agency for Healthcare Research and Quality</i>. Retrieved from <a href="http://effectivehealthcare.ahrq.gov/index.cfm/search-for-guides-reviews-and-reports/?productid=1911&amp;pageaction=displayproduct">http://effectivehealthcare.ahrq.gov/index.cfm/search-for-guides-reviews-and-reports/?productid=1911&amp;pageaction=displayproduct</a></p>	<p>Systematic review of RCTs</p>	<p>Home-visiting programs and multidisciplinary heart failure (MDS-HF) clinics reduced all-cause readmission and mortality, Structured telephone support (STS) reduced HF-specific readmission and mortality, and telemonitoring, and nurse-led clinic interventions had no effect to reduce readmissions. Effective interventions to reduce all-cause readmissions or mortality were delivered in person, high intensity, and multidisciplinary care approach (47 RCTs)</p>	<p>Level 1</p>

<p>GUIRGUIS-BLAKE, J. (2016). TC Interventions to Prevent Readmissions for Patients with Heart Failure. <i>American Family Physician</i>, 93(5), 401-403.</p>	<p>Reports from experts</p>	<p>The discussion of risks and benefits of TC interventions including multidisciplinary clinic intervention and home visiting for HF patients to reduce mortality and readmission</p>	<p>Level VII</p>
<p>Kansagara, D., Chiovaro, J. C., Kagen, D., Jencks, S., Rhyne, K., O'Neil, M., &amp; ... Englander, H. (2016). So many options, where do we start? An overview of the care transitions literature. <i>Journal Of Hospital Medicine</i>, 11(3), 221-230. doi:10.1002/jhm.2502</p>	<p>Systematic review of RCTs</p>	<p>TC programs featured with complete discharge planning and hospital -at-home interventions reduced readmissions. However, the evidence was weak due to the inconsistency in study interventions and strategies, population, and settings (10 reviews)</p>	<p>Level 1</p>
<p>Kociol, R. D., Peterson, E. D., Hammill, B. G., Flynn, K. E., Heidenreich, P. A., Piña, I. L., &amp; ... Hernandez, A. F. (2012). National survey of hospital strategies to reduce heart failure readmissions: findings from the Get With the Guidelines-Heart Failure registry. <i>Circulation. Heart Failure</i>, 5(6), 680-687. doi:10.1161/CIRCHEARTFAILURE.112.967406</p>	<p>Single descriptive study</p>	<p>The survey showed that the majority of hospital strategies were not associated with the reduction of 30-day readmission. The complete discharge and TC processes were associated with readmissions in patients with HF. Quality improvement and inpatient care processes had no effect on readmission reduction, and hospital strategies varied</p>	<p>Level VI</p>
<p>Lambrinou, E., Kalogirou, F., Lamnisis, D., &amp; Sourtzi, P. (2012). Effectiveness of heart failure management programs with nurse-led discharge planning in reducing re-admissions: A systematic review and meta-analysis. <i>International Journal Of Nursing Studies</i>, 49(5), 610-624. doi:10.1016/j.ijnurstu.2011.11.002</p>	<p>Systematic review and meta-analysis review of RCTs</p>	<p>The nurse-driven HF management programs before discharge potentially reduced readmissions. However, more studies were needed to identify essential interventions of a nurse-driven HF management program before discharge (Nineteen RCTs)</p>	<p>Level 1</p>



<p>Lindenfeld, J., Albert, N., Boehmer, J., Collins, S., Ezekowitz, J., Givertz, M., &amp; ... Walsh, M. (2010). HFSA 2010 Comprehensive Heart Failure Practice Guideline. <i>Journal Of Cardiac Failure, 16</i>(6), e1-194. doi:10.1016/j.cardfail.2010.04.004</p>	<p>Clinical practice guideline based on systematic reviews of RCTs</p>	<p>Comprehensive HF disease management programs included patient and family education and counseling, self-care management, early follow-up post hospital discharge, care access and care coordination, optimization of drug therapy, early detection of signs and symptoms of fluid overload, and addressing social and financial needs</p>	<p>Level I</p>
<p>Mikulich, S., Parikh, A., Franklin, G., Grelak, S., Echols, G., &amp; Johnson, D. (2013). Decreasing Heart Failure Readmissions with APN led Heart Failure Clinic. <i>Heart &amp; Lung, 42</i>(6), e7. doi:10.1016/j.hrtlng.2013.10.008</p>	<p>Cohort study</p>	<p>An APN/APN-led HF clinic decreased readmissions. The activities of HF clinic included patient education (HF disease process, medication, daily weight, sodium and fluid restriction, activity level and self-care instruction), medication titration, and addressing compliance issues. Exclusion criteria included HD, hospice care, nursing home and home care.</p>	<p>Level IV</p>
<p>Monza, K., Harris, D., &amp; Shaw, C. (2015). The Role of the Nurse Navigator in the Management of the Heart Failure Patient. <i>Critical Care Nursing Clinics Of North America, 27</i>(4), 537-549. doi:10.1016/j.cnc.2015.07.010</p>	<p>Review of descriptive studies</p>	<p>A nurse-led TC program reduced 30-day readmission, enhanced self-management behaviors, improved follow-up visits, and facilitated the continuity of care across settings.</p>	<p>Level V</p>

<p>Russell, D., Rosati, R. J., Sobolewski, S., Marren, J., &amp; Rosenfeld, P. (2011). Implementing a TC program for high-risk heart failure patients: findings from a community-based partnership between a certified home healthcare agency and regional hospital. <i>Journal For Healthcare Quality: Official Publication Of The National Association For Healthcare Quality</i>,33(6), 17-23. doi:10.1111/j.1945-1474.2011.00167.x</p>	Cohort study	<p>A joined TC program developed for high-risk HF patients between a hospital and a home health agency. The TC group (223) had fewer readmissions than usual care group (224).</p>	Level IV
<p>Stamp, K. D., Machado, M. A., &amp; Allen, N. A. (2014). TC Programs Improve Outcomes for Heart Failure Patients. <i>Journal Of Cardiovascular Nursing</i>, 29(2), 140-154. doi:10.1097/JCN.0b013e31827db560</p>	Systematic review of RCTs	<p>A nurse - led TC programs for patients with HF improved quality of life and reduced readmissions and costs of care. The successful strategies were home visits alone or in combination with telephone calls. The research gap in TC programs for HF patients needed large RCTs (integrative review n = 20)</p>	Level I
<p>Stauffer, B. D., Fullerton, C., Fleming, N., Ogola, G., Herrin, J., Stafford, P. M., &amp; Ballard, D. J. (2011). Effectiveness and cost of a TC program for heart failure: a prospective study with concurrent controls. <i>Archives Of Internal Medicine</i>,171(14), 1238-1243. doi:10.1001/archinternmed.2011.274</p>	Cohort study	<p>An APN-led TC program for patients with HF reduced 30-day hospital readmissions and had no effect on LOS. The TC program reduced the hospital financial gain about \$227 for each patient in the TC program.</p>	Level IV

<p>Smith, C. E., Piamjariyakul, U., Dalton, K. M., Russell, C., Wick, J., &amp; Ellerbeck, E. F. (2015). Nurse-Led Multidisciplinary Heart Failure Group Clinic Appointments: Methods, Materials, and Outcomes Used in the Clinical Trial. <i>The Journal Of Cardiovascular Nursing</i>, 30(4 Suppl 1), S25-S34. doi:10.1097/JCN.000000000000025</p>	<p>RCT</p>	<p>The nurse-led multidisciplinary HF group clinic appointments improved HF self-care knowledge and self-management skills that reduced readmissions in compared with usual care (192 HF patients)</p>	<p>Level II</p>
<p>Takeda, A. (2012). Clinical service organisation for heart failure. <i>Cochrane Database Of Systematic Reviews</i>, (9), doi:10.1002/14651858.CD002752.pub3</p>	<p>Cochrane systematic review</p>	<p>HF nurse-led case management interventions reduced HF and all-cause readmissions and mortality; multidisciplinary interventions might have effects on the reduction of HF and all-cause readmissions, and clinic interventions had weak evidence to support its efficacy (25 RCTs)</p>	<p>Level I</p>
<p>Thompson, Christine. "Safe Passage: a Nurse-Led Multidisciplinary Team Approach to Improving Transitions and Reducing Readmissions for Heart Failure Patients." <i>Heart &amp; Lung</i> 43, no. 4 (July 2014): 386. <i>CINAHL Plus with Full Text</i>, EBSCOhost(accessed August 31, 2016).</p>	<p>Cohort study</p>	<p>A nurse- multidisciplinary team approach/collaboration reduced 30-day readmission rates, improved patient education, and satisfaction (HACAPS score) using patient engagement, teamwork, and providing timely feedback to frontline staff. Team collaboration improved TC processes for preparing patient's self-management skills at home, which included identification of high readmission risk patients and setting up warming signs in medical charts, identification of readmission risk factors, patient/caregiver</p>	<p>Level IV</p>

		education, medication reconciliation, scheduled follow-up appointments before discharge, and follow-up calls	
Vedel, I., & Khanassov, V. (2015). TC for Patients With Congestive Heart Failure: A Systematic Review and Meta-Analysis. <i>Annals Of Family Medicine</i> , 13(6), 562-571. doi:10.1370/afm.1844	Systematic review	TC interventions reduced readmissions and ED visits. The most effective high-intensity interventions included home visiting and telephone follow-up, and the moderate-intensity interventions needed to be implanted more than six months to see the effect.(41 RCTs)	Level I
White, S. M., & Hill, A. (2014). A Heart Failure Initiative to Reduce the Length of Stay and Readmission Rates. <i>Professional Case Management</i> , 19(6), 276-284. doi:10.1097/NCM.0000000000000059	Cohort study	The implementation of care coordination improved TC, decreased readmissions and LOS and improved self-management behaviors in HF patients (inpatient Medical-surgical unit with 291 patients)	Level IV
Yancy, C. W., Jessup, M., Bozkurt, B., Butler, J., Casey, D. J., Drazner, M. H., & ... Wilkoff, B. L. (2013). 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. <i>Journal Of The American College Of Cardiology</i> , 62(16), e147-e239. doi:10.1016/j.jacc.2013.05.019	Clinical practice guideline based on systematic reviews	HF TC consisted of extensive patient education, medication reconciliation, hand-off updated care summary to follow-up providers, and early post discharge follow-up arrangement. The comprehensive patient and family education involved follow-up visits at the next care setting, worsening signs and symptoms of HF, medications, and activity and diet.	Level I