

Available online at www.sciencedirect.com

Journal of Functional Analysis 247 (2007) 138–168

**JOURNAL OF
Functional
Analysis**

www.elsevier.com/locate/jfa

The restricted Grassmannian, Banach Lie–Poisson spaces, and coadjoint orbits

Daniel Beltiță^{a,*}, Tudor S. Ratiu^b, Alice Barbara Tumpach^c^a *Institute of Mathematics “Simion Stoilow” of the Romanian Academy, PO Box 1-764, RO-014700 Bucharest, Romania*^b *Section de Mathématiques and Centre Bernoulli, École Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland*^c *Section de Mathématiques, École Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland*

Received 14 June 2006; accepted 1 March 2007

Communicated by D. Voiculescu

Abstract

We investigate some basic questions concerning the relationship between the restricted Grassmannian and the theory of Banach Lie–Poisson spaces. By using universal central extensions of Lie algebras, we find that the restricted Grassmannian is symplectomorphic to symplectic leaves in certain Banach Lie–Poisson spaces, and the underlying Banach space can be chosen to be even a Hilbert space. Smoothness of numerous adjoint and coadjoint orbits of the restricted unitary group is also established. Several pathological properties of the restricted algebra are pointed out.

© 2007 Elsevier Inc. All rights reserved.

Keywords: Restricted Grassmannian; Poisson manifold; Coadjoint orbit

1. Introduction

The present paper is devoted to an investigation of the relationship between the restricted Grassmannian and the recently initiated theory of Banach Lie–Poisson spaces.

The restricted Grassmannian (whose definition is recalled after Proposition 2.11 below) is a quite remarkable infinite-dimensional *Kähler* manifold that plays an important role in many areas

* Corresponding author.

E-mail addresses: daniel.beltita@imar.ro (D. Beltiță), tudor.ratiu@epfl.ch (T.S. Ratiu), alice.tumpach@epfl.ch (A.B. Tumpach).

of mathematics and physics. There are many interesting objects related to the restricted Grassmannian, such as: loop groups (see [33, Proposition 8.3.3]), the coadjoint orbits $\text{Diff}^+(S^1)/S^1$ and $\text{Diff}^+(S^1)/\text{PSU}(1, 1)$ of the group of orientation-preserving diffeomorphisms of the circle ([33, Proposition 6.8.2] and [34, Proposition 5.3]). It is related to the integrable system defined by the KP hierarchy (see [35]) and to the fermionic second quantization (see [39]). On the other hand, the notion of a Banach Lie–Poisson space was recently introduced in [28] and is an infinite-dimensional version of the Lie–Poisson spaces, that is, the Poisson manifolds provided by dual spaces of finite-dimensional Lie algebras (see for instance [31] for the finite-dimensional theory). Specifically, a *Banach Lie–Poisson space* is a Banach space \mathfrak{b} whose topological dual \mathfrak{b}^* is endowed with a structure of Banach Lie algebra such that the subspace \mathfrak{b} of $(\mathfrak{b}^*)^*$ is invariant under the corresponding coadjoint action. Equivalently, the Lie bracket of \mathfrak{b}^* is separately weak*-continuous. This new class of infinite-dimensional linear Poisson manifolds is remarkable in several respects: it includes all the preduals of W^* -algebras, thus establishing a bridge between Poisson geometry and the theory of operator algebras, and hence it provides links with algebraic quantum theories; it interacts in a fruitful way with the theory of extensions of Lie algebras (see [29]); and finally, there exist large classes of Banach Lie–Poisson spaces which share with the finite-dimensional Poisson manifolds the fundamental property that the characteristic distribution is integrable, the corresponding integral manifolds being in addition Poisson submanifolds which are symplectic and, in several important situations, are even *Kähler* manifolds (see [7]).

We have mentioned here two types of infinite-dimensional Kähler manifolds: the restricted Grassmannian and certain symplectic leaves in infinite-dimensional Lie–Poisson spaces introduced in [28]. This brings us to the first question addressed in the present paper.

Question 1.1. *Is the restricted Grassmannian a symplectic leaf in a Banach Lie–Poisson space?*

The main result of our paper shows that the answer to this question is essentially affirmative; see Section 5 for the precise statements and a detailed discussion of this problem. Specifically, we shall employ the method of central extensions to construct a certain Banach Lie–Poisson space $\tilde{\mathfrak{u}}_2$ whose characteristic distribution is integrable (Theorem 5.1) and one of the integral manifolds of this distribution is symplectomorphic to the connected component Gr_{res}^0 of the restricted Grassmannian (Theorem 5.3). Using a similar method, we realize the restricted Grassmannian as a symplectic leaf in yet another Banach Lie–Poisson space, which is the predual to a 1-dimensional central extension of the restricted Lie algebra $\mathfrak{u}_{\text{res}}$. See Section 2 for a detailed discussion of the Poisson geometry of this new Banach Lie–Poisson space $(\tilde{\mathfrak{u}}_{\text{res}})_*$.

This second construction is closely related to another area where the theory of restricted groups interacts with the theory initiated in [28]. Specifically, we also address the following question on the predual $(\mathfrak{u}_{\text{res}})_*$ of the restricted Lie algebra.

Question 1.2. *Does the real Banach space $(\mathfrak{u}_{\text{res}})_*$ have a natural structure of Banach Lie–Poisson space and is its characteristic distribution integrable?*

By the very construction of the Banach Lie–Poisson space $(\tilde{\mathfrak{u}}_{\text{res}})_*$, the predual $(\mathfrak{u}_{\text{res}})_*$ appears as a Poisson submanifold of $(\tilde{\mathfrak{u}}_{\text{res}})_*$ and carries a natural structure of Banach Lie–Poisson space. Nonetheless, the answer to the second part of Question 1.2 turns out to be much more difficult to give than the one to Question 1.1 inasmuch as the restricted algebra \mathcal{B}_{res} (see Notation 1.3 below) is a dual Banach $*$ -algebra with many pathological properties (summarized in Section 6):

its unitary group is unbounded, its natural predual is not spanned by its positive cone, and a conjugation theorem for its maximal $*$ -subalgebras fails to be true. Despite these unpleasant properties, we show that the characteristic distribution of $(u_{\text{res}})_*$ has numerous smooth integral manifolds, which are, in particular, *smooth* coadjoint orbits of the restricted unitary group U_{res} (see Section 3). For the sake of completeness, a short section of the paper (Section 4) is devoted to investigating smoothness of adjoint orbits of U_{res} .

Notation 1.3. We conclude this introduction by setting up some notation to be used throughout the paper. In the following, \mathcal{H} will denote a complex Hilbert space, endowed with a decomposition $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$ into the orthogonal sum of two closed subspaces.

It will follow implicitly from the hypotheses of various statements when additional conditions on the Hilbert space \mathcal{H} are imposed. For example, Corollary 3.7 requests the existence of a certain *countable* orthonormal basis, so \mathcal{H} needs to be separable. Also, sometimes it is assumed that both \mathcal{H}_{\pm} are infinite-dimensional. This is the case in Section 6 where we need operators that do *not* belong to the restricted algebra \mathcal{B}_{res} .

The orthogonal projection onto \mathcal{H}_{\pm} will be denoted by p_{\pm} . The Banach ideal of trace class operators on \mathcal{H} will be denoted by $\mathfrak{S}_1(\mathcal{H})$ and $\mathfrak{S}_2(\mathcal{H})$ will denote the Hilbert ideal of Hilbert–Schmidt operators on \mathcal{H} . We let $\mathcal{B}(\mathcal{H})$ be the algebra of all bounded linear operators on \mathcal{H} . We shall also need the Banach Lie group of unitary operators on \mathcal{H} ,

$$U(\mathcal{H}) = \{u \in \mathcal{B}(\mathcal{H}) \mid u^*u = uu^* = \text{id}\},$$

whose Lie algebra is

$$\mathfrak{u}(\mathcal{H}) = \{a \in \mathcal{B}(\mathcal{H}) \mid a^* = -a\}.$$

Now let us define the following skew-Hermitian element:

$$d := i(p_+ - p_-) \in \mathfrak{u}(\mathcal{H}).$$

The restricted Banach algebra and the restricted unitary group are respectively defined as follows:

$$\mathcal{B}_{\text{res}} = \{a \in \mathcal{B}(\mathcal{H}) \mid [d, a] \in \mathfrak{S}_2(\mathcal{H})\} = \{a \in \mathcal{B}(\mathcal{H}) \mid \|a\|_{\text{res}} := \|a\| + \|[d, a]\|_2 < \infty\}, \quad \text{and}$$

$$U_{\text{res}} = \{u \in U(\mathcal{H}) \mid [d, u] \in \mathfrak{S}_2(\mathcal{H})\} = U(\mathcal{H}) \cap \mathcal{B}_{\text{res}}.$$

The Lie algebra of U_{res} is the following Banach Lie algebra:

$$\mathfrak{u}_{\text{res}} = \{a \in \mathfrak{u}(\mathcal{H}) \mid [d, a] \in \mathfrak{S}_2(\mathcal{H})\} = \mathfrak{u}(\mathcal{H}) \cap \mathcal{B}_{\text{res}}.$$

Let us define the following Banach Lie algebra:

$$(u_{\text{res}})_* = \{\rho \in \mathfrak{u}(\mathcal{H}) \mid [d, \rho] \in \mathfrak{S}_2(\mathcal{H}), p_{\pm}\rho|_{\mathcal{H}_{\pm}} \in \mathfrak{S}_1(\mathcal{H}_{\pm})\}.$$

A connected Banach Lie group with Lie algebra $(u_{\text{res}})_*$ is

$$U_{1,2} = \{a \in U(\mathcal{H}) \mid a - \text{id} \in \mathfrak{S}_2(\mathcal{H}), p_{\pm}a|_{\mathcal{H}_{\pm}} \in \text{id} + \mathfrak{S}_1(\mathcal{H}_{\pm})\}.$$

The group U_1 and its Lie algebra \mathfrak{u}_1 are defined as follows:

$$U_1 = \{a \in U(\mathcal{H}) \mid a - \text{id} \in \mathfrak{S}_1(\mathcal{H})\}, \quad \text{and} \quad \mathfrak{u}_1 = \mathfrak{u}(\mathcal{H}) \cap \mathfrak{S}_1(\mathcal{H}).$$

Finally, the Hilbert–Lie group U_2 and its Lie algebra \mathfrak{u}_2 are defined by

$$U_2 = \{a \in U(\mathcal{H}) \mid a - \text{id} \in \mathfrak{S}_2(\mathcal{H})\}, \quad \text{and} \quad \mathfrak{u}_2 = \mathfrak{u}(\mathcal{H}) \cap \mathfrak{S}_2(\mathcal{H}).$$

2. The Banach Lie–Poisson space associated to the universal central extension of $\mathfrak{u}_{\text{res}}$

In this section we construct a Banach Lie–Poisson space $(\tilde{\mathfrak{u}}_{\text{res}})_*$ whose dual is the universal central extension of the restricted algebra $\mathfrak{u}_{\text{res}}$. (See [24] for the definition of universal central extension and Proposition 2.4 below for the justification of this fact.) The Poisson structure of $(\tilde{\mathfrak{u}}_{\text{res}})_*$ is defined by (2.8) in Proposition 2.5. Let us first justify the suggestive notation $(\mathfrak{u}_{\text{res}})_*$.

Proposition 2.1. *The Lie algebra $(\mathfrak{u}_{\text{res}})_*$ is a predual of the unitary restricted algebra $\mathfrak{u}_{\text{res}}$, the duality pairing $\langle \cdot, \cdot \rangle$ being given by*

$$\langle \cdot, \cdot \rangle : (\mathfrak{u}_{\text{res}})_* \times \mathfrak{u}_{\text{res}} \rightarrow \mathbb{R}, \quad (b, c) \mapsto \text{Tr}(bc). \tag{2.1}$$

Proof. Consider two arbitrary elements

$$a = \begin{pmatrix} a_{++} & a_{+-} \\ -a_{+-}^* & a_{--} \end{pmatrix} \in \mathfrak{u}_{\text{res}} \quad \text{and} \quad \rho = \begin{pmatrix} \rho_{++} & -\rho_{-+}^* \\ \rho_{-+} & \rho_{--} \end{pmatrix} \in (\mathfrak{u}_{\text{res}})_*.$$

Then

$$a\rho = \begin{pmatrix} a_{++}\rho_{++} + a_{+-}\rho_{-+} & -a_{++}\rho_{-+}^* + a_{+-}\rho_{--} \\ -a_{+-}^*\rho_{++} + a_{--}\rho_{-+} & a_{+-}^*\rho_{-+}^* + a_{--}\rho_{--} \end{pmatrix}, \tag{2.2}$$

hence

$$\text{Tr}(a\rho) = \text{Tr}(a_{++}\rho_{++}) + 2\Re \text{Tr}(a_{+-}\rho_{-+}) + \text{Tr}(a_{--}\rho_{--}), \tag{2.3}$$

where $\Re z$ denotes the real part of the complex number z . Recall that the bilinear functional

$$\mathcal{B}(\mathcal{H}_{\pm}) \times \mathfrak{S}_1(\mathcal{H}_{\pm}) \rightarrow \mathbb{C}, \quad (b, c) \mapsto \text{Tr}(bc),$$

induces a topological isomorphism of complex Banach spaces $(\mathfrak{S}_1(\mathcal{H}_{\pm}))^* \simeq \mathcal{B}(\mathcal{H}_{\pm})$. It follows that the trace induces a topological isomorphism of real Banach spaces

$$(\mathfrak{u}(\mathcal{H}_{\pm}) \cap \mathfrak{S}_1(\mathcal{H}_{\pm}))^* \simeq \mathfrak{u}(\mathcal{H}_{\pm}). \tag{2.4}$$

Indeed, the \mathbb{C} -linearity of the trace implies that for $b \in \mathcal{B}(\mathcal{H}_{\pm})$ the following conditions are equivalent:

$$(\forall c \in \mathfrak{u}(\mathcal{H}_{\pm}) \cap \mathfrak{S}_1(\mathcal{H}_{\pm})) \quad \text{Tr}(bc) = 0 \quad \iff \quad (\forall c \in \mathfrak{S}_1(\mathcal{H}_{\pm})) \quad \text{Tr}(bc) = 0.$$

Moreover the condition

$$(\forall c \in \mathfrak{u}(\mathcal{H}_\pm) \cap \mathfrak{S}_1(\mathcal{H}_\pm)) \quad \text{Tr}(bc) \in \mathbb{R}$$

implies

$$(\forall c \in \mathfrak{u}(\mathcal{H}_\pm) \cap \mathfrak{S}_1(\mathcal{H}_\pm)) \quad \text{Tr}(b + b^*)c = 0,$$

hence b belongs to $\mathfrak{u}(\mathcal{H}_\pm)$. On the other hand, the duality pairing of complex Hilbert spaces

$$\mathfrak{S}_2(\mathcal{H}_-, \mathcal{H}_+) \times \mathfrak{S}_2(\mathcal{H}_+, \mathcal{H}_-) \rightarrow \mathbb{C}, \quad (b, c) \mapsto \text{Tr}(bc),$$

induces a duality pairing of the underlying real Hilbert spaces by

$$\mathfrak{S}_2(\mathcal{H}_-, \mathcal{H}_+) \times \mathfrak{S}_2(\mathcal{H}_+, \mathcal{H}_-) \rightarrow \mathbb{R}, \quad (b, c) \mapsto \Re \text{Tr}(bc). \tag{2.5}$$

In view of formula (2.3), we conclude that the trace induces a topological isomorphism of real Banach spaces

$$((\mathfrak{u}_{\text{res}})_*)^* \simeq \mathfrak{u}_{\text{res}}.$$

That is, $(\mathfrak{u}_{\text{res}})_*$ is indeed a predual to $\mathfrak{u}_{\text{res}}$, the duality pairing being induced by (2.4) and (2.5). \square

Definition 2.2. We define the Banach Lie algebra $\tilde{\mathfrak{u}}_{\text{res}}$ as the central extension of $\mathfrak{u}_{\text{res}}$ with continuous two-cocycle s given by

$$s(A, B) := \text{Tr}(A[d, B]), \tag{2.6}$$

for all $A, B \in \mathfrak{u}_{\text{res}}$. That is, $\tilde{\mathfrak{u}}_{\text{res}}$ is the Banach Lie algebra $\mathfrak{u}_{\text{res}} \oplus \mathbb{R}$ endowed with the bracket $[\cdot, \cdot]_d$ defined by

$$[(A, a), (B, b)]_d = ([A, B], s(A, B)). \tag{2.7}$$

Remark 2.3. Note that by the very definition of $\mathfrak{u}_{\text{res}}$, one has $[d, \mathfrak{u}_{\text{res}}] \subset (\mathfrak{u}_{\text{res}})_*$. It follows from the duality pairing (2.1), that s is well defined by (2.6). To see that s defines a two-cocycle on $\mathfrak{u}_{\text{res}}$, let us remark that s is $(2i)$ -times the Schwinger term of [39]. It follows from Corollary II.12 in the aforementioned work that s defines a non-trivial element in the second continuous Lie algebra cohomology space $H^2(\mathfrak{u}_{\text{res}}, \mathbb{R})$. The corresponding $U(1)$ -extension of the unitary restricted group U_{res} is isomorphic to the $U(1)$ -extensions U_{res}^\sim and \hat{U}_{res} of U_{res} constructed in [39].

Proposition 2.4. *The cohomology class $[s]$ is a generator of the continuous Lie algebra cohomology space $H^2(\mathfrak{u}_{\text{res}}, \mathbb{R})$.*

Proof. According to [23, Proposition I.11], the second continuous Lie algebra cohomology space $H^2(\mathcal{B}_{\text{res}}, \mathbb{C})$ of the restricted Lie algebra \mathcal{B}_{res} is 1-dimensional. Note that a continuous \mathbb{R} -valued 2-cocycle v on $\mathfrak{u}_{\text{res}}$ extends by \mathbb{C} -linearity to a continuous \mathbb{C} -valued 2-cocycle $v^\mathbb{C}$ on the complex Lie algebra \mathcal{B}_{res} . The cocycle $v^\mathbb{C}$ is a coboundary if and only if there exists

a continuous linear map $\alpha: \mathcal{B}_{\text{res}} \rightarrow \mathbb{C}$ such that $v^{\mathbb{C}}(x, y) = \alpha([x, y])$ for every $x, y \in \mathcal{B}_{\text{res}}$. But since $v^{\mathbb{C}}$ restricts to the \mathbb{R} -valued 2-cocycle v on $\mathfrak{u}_{\text{res}}$, this is the case if and only if there exists $\beta := \mathfrak{R}\alpha: \mathfrak{u}_{\text{res}} \rightarrow \mathbb{R}$ such that $v(x, y) = \beta([x, y])$ for every $x, y \in \mathfrak{u}_{\text{res}}$. It follows that the extension $v^{\mathbb{C}}$ is a coboundary on \mathcal{B}_{res} if and only if v is a coboundary on $\mathfrak{u}_{\text{res}}$. Consequently, there is a natural linear injection of $H^2(\mathfrak{u}_{\text{res}}, \mathbb{R})$ into $H^2(\mathcal{B}_{\text{res}}, \mathbb{C})$. Since s defines a non-trivial element in $H^2(\mathfrak{u}_{\text{res}}, \mathbb{R})$ (see Remark 2.3) and $\dim_{\mathbb{C}} H^2(\mathcal{B}_{\text{res}}, \mathbb{C}) = 1$, it follows that $\dim_{\mathbb{R}} H^2(\mathfrak{u}_{\text{res}}, \mathbb{R}) = 1$ and thus $H^2(\mathfrak{u}_{\text{res}}, \mathbb{R})$ is generated by s . \square

Proposition 2.5. *The Banach space $(\tilde{\mathfrak{u}}_{\text{res}})_*$ is a Banach Lie–Poisson space for the Poisson bracket*

$$\{f, g\}_d(\mu, \gamma) := \langle \mu, [D_{\mu}f(\mu), D_{\mu}g(\mu)] \rangle + \gamma s(D_{\mu}f, D_{\mu}g), \tag{2.8}$$

where $f, g \in C^{\infty}((\tilde{\mathfrak{u}}_{\text{res}})_*)$, (μ, γ) is an arbitrary element in $(\tilde{\mathfrak{u}}_{\text{res}})_*$, and D_{μ} denotes the partial Fréchet derivative with respect to $\mu \in (\mathfrak{u}_{\text{res}})_*$.

The pairing in Eq. (2.8) is the duality pairing defined by (2.1). We will denote by $\langle \cdot, \cdot \rangle_d$ the duality pairing between $(\tilde{\mathfrak{u}}_{\text{res}})_* = (\mathfrak{u}_{\text{res}})_* \oplus \mathbb{R}$ and $\tilde{\mathfrak{u}}_{\text{res}} = \mathfrak{u}_{\text{res}} \oplus \mathbb{R}$ given by

$$\langle (\mu, \gamma), (A, a) \rangle_d = \langle \mu, A \rangle + \gamma a.$$

Proof of Proposition 2.5. By [28, Theorem 4.2], the Banach space $(\tilde{\mathfrak{u}}_{\text{res}})_*$ is a Banach Lie–Poisson space if and only if its dual $\tilde{\mathfrak{u}}_{\text{res}}$ is a Banach Lie algebra satisfying $\text{ad}_x^*(\tilde{\mathfrak{u}}_{\text{res}})_* \subset (\tilde{\mathfrak{u}}_{\text{res}})_* \subset (\tilde{\mathfrak{u}}_{\text{res}})^*$ for all $x \in \tilde{\mathfrak{u}}_{\text{res}}$. The fact that $\tilde{\mathfrak{u}}_{\text{res}}$ is a Banach Lie algebra follows directly from the continuity of s and from the 2-cocycle identity which implies the Jacobi identity of $[\cdot, \cdot]_d$. To see that the coadjoint action of $\tilde{\mathfrak{u}}_{\text{res}}$ preserves the predual $(\tilde{\mathfrak{u}}_{\text{res}})_*$, note that for every $(A, a), (B, b) \in \tilde{\mathfrak{u}}_{\text{res}}$ and every $(\mu, \gamma) \in (\tilde{\mathfrak{u}}_{\text{res}})_*$, one has

$$\begin{aligned} \langle -\text{ad}_{(A,a)}^*(\mu, \gamma), (B, b) \rangle_d &:= \langle (\mu, \gamma), -\text{ad}_{(A,a)}(B, b) \rangle_d = \langle (\mu, \gamma), -[(A, a), (B, b)]_d \rangle_d \\ &= \langle (\mu, \gamma), (-[A, B], -s(A, B)) \rangle_d = -\text{Tr} \mu[A, B] - \gamma \text{Tr} A[d, B] \\ &= -\text{Tr} \mu[A, B] - \gamma \text{Tr}[A, d]B = \langle (-\text{ad}_A^*(\mu) - \gamma[A, d], 0), (B, b) \rangle_d. \end{aligned} \tag{2.9}$$

Since

$$[(\mathfrak{u}_{\text{res}})_*, \mathfrak{u}_{\text{res}}] \subseteq (\mathfrak{u}_{\text{res}})_*, \quad \text{and} \quad [d, \mathfrak{u}_{\text{res}}] \subset (\mathfrak{u}_{\text{res}})_*,$$

we conclude that $-\text{ad}^*(A)(\mu) - \gamma[A, d]$ belongs to $(\mathfrak{u}_{\text{res}})_*$ for every $A \in \mathfrak{u}_{\text{res}}$. Hence the predual $(\tilde{\mathfrak{u}}_{\text{res}})_*$ is preserved by the coadjoint action. Referring again to [28, Theorem 4.2], it follows that the Poisson bracket of $f, g \in C^{\infty}((\tilde{\mathfrak{u}}_{\text{res}})_*)$ is given by

$$\{f, g\}_d(\mu, \gamma) = \langle (\mu, \gamma), [Df(\mu, \gamma), Dg(\mu, \gamma)]_d \rangle_d.$$

Denoting respectively by D_{μ} and D_{γ} the partial Fréchet derivatives with respect to $\mu \in (\mathfrak{u}_{\text{res}})_*$ and $\gamma \in \mathbb{R}$, one has

$$\begin{aligned} \{f, g\}_d(\mu, \gamma) &= \langle (\mu, \gamma), [(D_\mu f, D_\gamma f), (D_\mu g, D_\gamma g)]_d \rangle_d \\ &= \langle (\mu, \gamma), ([D_\mu f, D_\mu g], s(D_\mu f, D_\mu g)) \rangle_d \\ &= \langle \mu, [D_\mu f, D_\mu g] \rangle + \gamma s(D_\mu f, D_\mu g), \end{aligned}$$

and this ends the proof. \square

Remark 2.6. By [28, Theorem 4.2], it follows that the Hamiltonian vector field associated to a smooth function h on $(\mathfrak{u}_{\text{res}})_*$ is given by

$$X_h(\mu, \gamma) = -\text{ad}^*_{(D_\mu h, D_\gamma h)}(\mu, \gamma) = (-\text{ad}^*_{D_\mu h} \mu - \gamma [D_\mu h, d], 0). \tag{2.10}$$

Remark 2.7. Note that, for each $\gamma \in \mathbb{R}$, $(\mathfrak{u}_{\text{res}})_* \oplus \{\gamma\}$ is a Poisson submanifold of $(\tilde{\mathfrak{u}}_{\text{res}})_*$ for the following Poisson bracket on the first factor

$$\{f, g\}_{d, \gamma}(\mu) := \langle \mu, [D_\mu f(\mu), D_\mu g(\mu)] \rangle + \gamma s(D_\mu f, D_\mu g).$$

Remark 2.8. The central extension $(\tilde{\mathfrak{u}}_{\text{res}})_*$ of the Banach Lie–Poisson space $(\mathfrak{u}_{\text{res}})_*$ is a particular example of the extensions of Banach Lie–Poisson spaces constructed in [29]. Indeed formula (2.8) for the bracket of two functions on $(\tilde{\mathfrak{u}}_{\text{res}})_*$ can be alternatively deduced from the general formula (5.6) in [29, Theorem 5.2], with $\mathfrak{c} = \mathbb{R}$, $\mathfrak{a} = (\mathfrak{u}_{\text{res}})_*$, $\varphi = 0$ and $\omega = s$. The pairing in the second term of the right-hand side of (5.6) [29, Theorem 5.2], is, in this special case, just the pairing between the real line and its dual given by multiplication of real numbers (the element $c \in \mathfrak{c}$ is γ), and the bracket of partial derivatives of the functions f and g with respect to c vanishes since \mathbb{R} is commutative.

Proposition 2.9. *The unitary group U_{res} acts on the Poisson manifold $(\mathfrak{u}_{\text{res}})_* \oplus \{\gamma\} \subset (\tilde{\mathfrak{u}}_{\text{res}})_*$ by the affine coadjoint action as follows. For $g \in U_{\text{res}}$,*

$$g \cdot (\mu, \gamma) := (\text{Ad}^*(g^{-1})(\mu) - \gamma \sigma(g), \gamma),$$

where $\mu \in (\mathfrak{u}_{\text{res}})_*$, $\gamma \in \mathbb{R}$, and where

$$\begin{aligned} \sigma : U_{\text{res}} &\rightarrow (\mathfrak{u}_{\text{res}})_*, \\ g &\mapsto gdg^{-1} - d. \end{aligned}$$

Proof. Let us verify that for every $g \in U_{\text{res}}$ we have $gdg^{-1} - d \in (\mathfrak{u}_{\text{res}})_*$. Consider the block decomposition of g with respect to the direct sum $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$

$$g = \begin{pmatrix} g_{++} & g_{+-} \\ g_{-+} & g_{--} \end{pmatrix} \in U_{\text{res}}.$$

One has

$$\begin{aligned} &\begin{pmatrix} g_{++} & g_{+-} \\ g_{-+} & g_{--} \end{pmatrix} \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \begin{pmatrix} g_{++}^* & g_{+-}^* \\ g_{-+}^* & g_{--}^* \end{pmatrix} \\ &= \begin{pmatrix} ig_{++}g_{++}^* - ig_{+-}g_{-+}^* & ig_{++}g_{+-}^* - ig_{+-}g_{--}^* \\ ig_{-+}g_{++}^* - ig_{--}g_{-+}^* & ig_{-+}g_{+-}^* - ig_{--}g_{--}^* \end{pmatrix}. \end{aligned} \tag{2.11}$$

Since $g_{\pm\mp}$ belongs to $\mathfrak{S}_2(\mathcal{H}_{\mp}, \mathcal{H}_{\pm})$, the off-diagonal blocks of the right-hand side are in $\mathfrak{S}_2(\mathcal{H}_{\pm}, \mathcal{H}_{\mp})$. Further, since

$$\begin{pmatrix} g_{++} & g_{+-} \\ g_{-+} & g_{--} \end{pmatrix} \begin{pmatrix} g_{++}^* & g_{-+}^* \\ g_{+-}^* & g_{--}^* \end{pmatrix} = \begin{pmatrix} g_{++}g_{++}^* + g_{+-}g_{+-}^* & g_{++}g_{-+}^* + g_{+-}g_{--}^* \\ g_{-+}g_{++}^* + g_{--}g_{+-}^* & g_{-+}g_{-+}^* + g_{--}g_{--}^* \end{pmatrix} = \begin{pmatrix} \text{id} & 0 \\ 0 & \text{id} \end{pmatrix},$$

and since $\mathfrak{S}_2 \cdot \mathfrak{S}_2 \subset \mathfrak{S}_1$, one has

$$g_{++}g_{++}^* = \text{id} - g_{+-}g_{+-}^* \in \text{id} + \mathfrak{S}_1(\mathcal{H}_+)$$

and

$$g_{--}g_{--}^* = \text{id} - g_{-+}g_{-+}^* \in \text{id} + \mathfrak{S}_1(\mathcal{H}_-).$$

Consequently,

$$g_{++}g_{++}^* - g_{+-}g_{+-}^* \in \text{id} + \mathfrak{S}_1(\mathcal{H}_+)$$

and

$$g_{-+}g_{-+}^* - g_{--}g_{--}^* \in -\text{id} + \mathfrak{S}_1(\mathcal{H}_-).$$

Moreover, it is clear that the result of the multiplication (2.11) is skew-symmetric. Hence for all $g \in U_{\text{res}}$ we have $gdg^{-1} - d \in (\mathfrak{u}_{\text{res}})_*$.

Denoting by $\text{Aff}((\mathfrak{u}_{\text{res}})_* \oplus \{\gamma\})$ the affine group of transformations of $(\mathfrak{u}_{\text{res}})_* \oplus \{\gamma\}$, it remains to show that

$$\begin{aligned} (\text{Ad}^*, -\gamma\sigma) : U_{\text{res}} &\rightarrow \text{Aff}((\mathfrak{u}_{\text{res}})_* \oplus \{\gamma\}) = \text{GL}((\mathfrak{u}_{\text{res}})_* \oplus \{\gamma\}) \rtimes (\mathfrak{u}_{\text{res}})_*, \\ g &\mapsto (\text{Ad}^*(g^{-1}), -\gamma\sigma(g)) \end{aligned}$$

is a group homomorphism. For this, we have to check that $\sigma(g_1g_2) = \text{Ad}^*(g_1^{-1})\sigma(g_2) + \sigma(g_1)$ for all g_1, g_2 in U_{res} (see [22]). In fact

$$\begin{aligned} \sigma(g_1g_2) &= g_1g_2dg_2^{-1}g_1^{-1} - d = g_1(g_2dg_2^{-1} - d)g_1^{-1} + (g_1dg_1^{-1} - d) \\ &= \text{Ad}^*(g_1^{-1})(\sigma(g_2)) + \sigma(g_1), \end{aligned}$$

and this ends the proof. \square

Proposition 2.10. *The isotropy group of $(0, \gamma) \in (\mathfrak{u}_{\text{res}})_* \oplus \{\gamma\}$ for the U_{res} -affine coadjoint action is a Lie subgroup of U_{res} .*

Proof. An element X in the Lie algebra $\mathfrak{u}_{\text{res}}$ of U_{res} induces by the infinitesimal affine coadjoint action on $(\mathfrak{u}_{\text{res}})_* \oplus \{\gamma\}$ the following vector field:

$$\begin{aligned} X \cdot (\mu, \gamma) &:= \frac{d}{dt} [\exp(tX) \cdot (\mu, \gamma)]_{t=0} \\ &= \left(\frac{d}{dt} [\text{Ad}^*(\exp(-tX))(\mu) - \gamma\sigma(\exp(tX))]_{t=0}, 0 \right) \\ &= (-\text{ad}_X^*(\mu) - \gamma[X, d], 0). \end{aligned}$$

By definition, the Lie algebra of the isotropy group of (μ, γ) is

$$\mathfrak{u}_{(\mu, \gamma)} := \{X \in \mathfrak{u}_{\text{res}} \mid -\text{ad}_X^*(\mu) - \gamma[X, d] = 0\}.$$

The proposition is trivial when μ and γ vanish. For $\mu = 0$ and $\gamma \neq 0$, the Lie algebra $\mathfrak{u}_{(0, \gamma)}$ consist of all elements of $\mathfrak{u}_{\text{res}}$ which commute with d . Hence, for $\gamma \neq 0$, $\mathfrak{u}_{(0, \gamma)} = \mathfrak{u}(\mathcal{H}_+) \oplus \mathfrak{u}(\mathcal{H}_-)$. A topological complement to $\mathfrak{u}_{(0, \gamma)}$ in $\mathfrak{u}_{\text{res}}$ is $\mathfrak{m} := \mathfrak{u}(\mathcal{H}) \cap (\mathfrak{S}_2(\mathcal{H}_+, \mathcal{H}_-) \oplus \mathfrak{S}_2(\mathcal{H}_-, \mathcal{H}_+))$. \square

Proposition 2.11. *The affine coadjoint orbits of U_{res} that are smooth are tangent to the characteristic distribution of the Poisson manifold $(\tilde{\mathfrak{u}}_{\text{res}})_*$.*

Proof. By the proof of Proposition 2.10, the image of the differential of the orbit map is

$$\mathfrak{u}_{\text{res}} \cdot (\mu, \gamma) = \{(-\text{ad}_X^*(\mu) - \gamma[X, d], 0) \mid X \in \mathfrak{u}_{\text{res}}\}.$$

By Remark 2.6, the characteristic space at $(\mu, \gamma) \in (\tilde{\mathfrak{u}}_{\text{res}})_*$ is

$$\begin{aligned} P(\mu, \gamma) &= \{X_h(\mu) = (-\text{ad}_{D_\mu h}^* \mu - \gamma[D_\mu h, d], 0) \mid h \in C^\infty((\mathfrak{u}_{\text{res}})_*)\} \\ &= \{(-\text{ad}_X^* \mu - \gamma[X, d], 0) \mid X \in \mathfrak{u}_{\text{res}}\}. \end{aligned}$$

Thus the assertion follows. \square

The *restricted Grassmannian* Gr_{res} is defined as the set of subspaces W of the Hilbert space \mathcal{H} such that the orthogonal projection from W to \mathcal{H}_+ (respectively to \mathcal{H}_-) is a Fredholm operator (respectively a Hilbert–Schmidt operator). It follows from [33, Propositions 7.1.2 and 7.1.3] that Gr_{res} is a Hilbert manifold and a homogeneous space under the natural action of U_{res} . According to [39, Proposition II.2], the connected components of U_{res} are the sets

$$U_{\text{res}}^k = \left\{ \begin{pmatrix} U_{++} & U_{+-} \\ U_{-+} & U_{--} \end{pmatrix} \in U_{\text{res}} \mid \text{index}(U_{++}) = k \right\} \quad \text{for } k \in \mathbb{Z}.$$

The pairwise disjoint sets

$$\text{Gr}_{\text{res}}^k = \{W \in \text{Gr}_{\text{res}} \mid \text{index}(p_+|_W : W \rightarrow H_+) = k\}, \quad k \in \mathbb{Z},$$

are the images of the connected components of U_{res} by the continuous projection $U_{\text{res}} \rightarrow \text{Gr}_{\text{res}} = U_{\text{res}} / (U(\mathcal{H}_+) \times U(\mathcal{H}_-))$, and thus they are the connected components of Gr_{res} . In particular, the

connected component of Gr_{res} containing \mathcal{H}_+ is Gr_{res}^0 . The Kähler structure of the restricted Grassmannian is defined in [33, Section 7.8]. According to the convention in [33], the Kähler form ω_{Gr} of Gr_{res} is the U_{res} -invariant 2-form whose value at \mathcal{H}_+ is given by

$$\omega_{\text{Gr}}(X, Y) = 2\Im \text{Tr}(X^*Y), \tag{2.12}$$

where $X, Y \in \mathfrak{S}_2(\mathcal{H}_+, \mathcal{H}_-) \simeq T_{\mathcal{H}_+} \text{Gr}_{\text{res}}$ and $\Im z$ denotes the imaginary part of $z \in \mathbb{C}$. Equivalently, ω_{Gr} is the quotient of the following real-valued anti-symmetric bilinear form Ω_{Gr} on $\mathfrak{u}_{\text{res}}$ which vanishes on $\mathfrak{u}(\mathcal{H}_+) \oplus \mathfrak{u}(\mathcal{H}_-)$ and is invariant under the $U(\mathcal{H}_+) \times U(\mathcal{H}_-)$ -action (see [39, Corollary III.8]):

$$\Omega_{\text{Gr}}(A, B) = -\frac{1}{2}s(A, B), \tag{2.13}$$

where A and B belong to $\mathfrak{u}_{\text{res}}$. In this correspondence, an element $A = \begin{pmatrix} A_{++} & -A_{-+}^* \\ A_{-+} & A_{--} \end{pmatrix}$ in $\mathfrak{u}_{\text{res}}$ is identified with the vector $X = A_{-+}$ in $\mathfrak{S}_2(\mathcal{H}_+, \mathcal{H}_-) \simeq T_{\mathcal{H}_+}(\text{Gr}_{\text{res}})$.

Proposition 2.12. *For every $\gamma \neq 0$, the connected components of the U_{res} -affine coadjoint orbit $\mathcal{O}_{(0,\gamma)}$ of $(0, \gamma) \in (\mathfrak{u}_{\text{res}})_* \oplus \{\gamma\}$ are strong symplectic leaves in the Banach Lie–Poisson space $(\tilde{\mathfrak{u}}_{\text{res}})_*$.*

Proof. We recall from Proposition 2.5 that $(\tilde{\mathfrak{u}}_{\text{res}})_*$ is a Banach Lie–Poisson space. By Proposition 2.10, the isotropy group $U_{(0,\gamma)}$ of $(0, \gamma)$ for the U_{res} -affine coadjoint action is a Banach Lie subgroup of U_{res} since its Lie algebra $\mathfrak{u}_{(0,\gamma)}$ is complemented in $\mathfrak{u}_{\text{res}}$. Let us denote by \tilde{U}_{res} the central extension of U_{res} with Lie algebra $\tilde{\mathfrak{u}}_{\text{res}}$ and by $p: \tilde{U}_{\text{res}} \rightarrow U_{\text{res}}$ the projection map in the exact sequence $1 \rightarrow S^1 \rightarrow \tilde{U}_{\text{res}} \rightarrow U_{\text{res}} \rightarrow 1$. The group \tilde{U}_{res} is isomorphic to the unitary subgroup of $\text{GL}_{\tilde{\mathfrak{u}}_{\text{res}}}$, the central extension of the group of invertible elements in $\mathfrak{B}_{\text{res}}$ constructed in [33, Section 6.6] (see also [39, Section II.3]). The usual coadjoint action of \tilde{U}_{res} on the dual of its Lie algebra leaves the predual $(\tilde{\mathfrak{u}}_{\text{res}})_*$ invariant since by Eq. (2.9) and the arguments in the proof of Proposition 2.5 following it, one has

$$-\text{ad}_{(A,a)}^*(\mu, \gamma) = (-\text{ad}_A^*(\mu) - \gamma[A, d], 0) \in (\tilde{\mathfrak{u}}_{\text{res}})_*. \tag{2.14}$$

The isotropy group $\tilde{U}_{(0,\gamma)}$ of $(0, \gamma) \in (\tilde{\mathfrak{u}}_{\text{res}})_*$ for the usual coadjoint action of \tilde{U}_{res} is a Banach Lie subgroup of \tilde{U}_{res} since its Lie algebra

$$\tilde{\mathfrak{u}}_{(0,\gamma)} := \{(A, a) \in \mathfrak{u}_{\text{res}} \mid -\text{ad}_{(A,a)}^*(0, \gamma) = 0\} = \mathfrak{u}_{(0,\gamma)} \oplus \mathbb{R}$$

is complemented in $\tilde{\mathfrak{u}}_{\text{res}}$. It follows from [28, Theorem 7.3] that the homogeneous space $\tilde{U}_{\text{res}}/\tilde{U}_{(0,\gamma)}$, which admits a unique smooth Banach manifold structure making the canonical projection $\tilde{\pi}: \tilde{U}_{\text{res}} \rightarrow \tilde{U}_{\text{res}}/\tilde{U}_{(0,\gamma)}$ a surjective submersion, carries a weak symplectic two-form $\omega_{(0,\gamma)}$ given by

$$\omega_{(0,\gamma)}([\tilde{g}]) (T_{\tilde{g}}\tilde{\pi}(T_e L_{\tilde{g}}\xi), T_{\tilde{g}}\tilde{\pi}(T_e L_{\tilde{g}}\eta)) := \langle (0, \gamma), [\xi, \eta]_d \rangle_d, \tag{2.15}$$

where $\xi, \eta \in \tilde{\mathfrak{u}}_{\text{res}}, \tilde{g} \in \tilde{\mathfrak{U}}_{\text{res}}, [\tilde{g}] := \tilde{\pi}(\tilde{g})$. The usual coadjoint action of $\tilde{\mathfrak{U}}_{\text{res}}$ on the predual $(\tilde{\mathfrak{u}}_{\text{res}})_*$ and the affine coadjoint action of $\mathfrak{U}_{\text{res}}$ on $(\tilde{\mathfrak{u}}_{\text{res}})_*$ defined in Proposition 2.9 are related by

$$\text{Ad}^*(\tilde{g}^{-1})(\mu, \gamma) = p(\tilde{g}) \cdot (\mu, \gamma), \tag{2.16}$$

where $(\mu, \gamma) \in (\tilde{\mathfrak{u}}_{\text{res}})_*$. To see this, note that the coadjoint action of the center of the extended group is trivial. Therefore the corresponding action descends to an action of the restricted unitary group. The tangent maps of the group homomorphisms $\tilde{\mathfrak{U}}_{\text{res}} \rightarrow \text{GL}((\tilde{\mathfrak{u}}_{\text{res}})_*)$ defined by the left- and right-hand sides of (2.16) coincide by Eq. (2.14), hence Eq. (2.16) holds for \tilde{g} in the connected component of the unit in $\tilde{\mathfrak{U}}_{\text{res}}$ which is simply connected by [23, Proposition IV.9(i)]. The general case follows by verifying formula (2.16) for the shift operator since by the remark following [39, Definition and Proposition II.23] and [39, Proposition II.27], we have $\tilde{\mathfrak{U}}_{\text{res}} = \tilde{\mathfrak{U}}_{\text{res}}^0 \rtimes \mathbb{Z}$ where the action of $1 \in \mathbb{Z}$ on $\tilde{\mathfrak{U}}_{\text{res}}^0$ projects to the conjugation by the shift on $\mathfrak{U}_{\text{res}}^0$. Consequently, the $\mathfrak{U}_{\text{res}}$ -affine coadjoint orbit $\mathcal{O}_{(0, \gamma)}$ is the coadjoint orbit of $(0, \gamma)$ for the usual coadjoint action of $\tilde{\mathfrak{U}}_{\text{res}}$. It follows from [28, Theorem 7.4] that the map

$$\iota : [\tilde{g}] \in \tilde{\mathfrak{U}}_{\text{res}} / \tilde{\mathfrak{U}}_{(0, \gamma)} \mapsto \text{Ad}_{\tilde{g}^{-1}}^*(0, \gamma) \in (\tilde{\mathfrak{u}}_{\text{res}})_* \tag{2.17}$$

is an injective weak immersion of the quotient manifold $\tilde{\mathfrak{U}}_{\text{res}} / \tilde{\mathfrak{U}}_{(0, \gamma)}$ into the predual space $(\tilde{\mathfrak{u}}_{\text{res}})_*$, and that the connected component of the affine coadjoint orbit $\mathcal{O}_{(0, \gamma)}$ endowed with the smooth manifold structure making ι into a diffeomorphism and the symplectic form given by $\iota_*(\omega_{(0, \gamma)})$ are symplectic leaves of the Banach Lie–Poisson space $(\tilde{\mathfrak{u}}_{\text{res}})_*$. By [28, Theorem 7.5], this symplectic form is in fact strong. \square

Theorem 2.13. *The connected components of the restricted Grassmannian are strong symplectic leaves in the Banach Lie–Poisson space $(\tilde{\mathfrak{u}}_{\text{res}})_*$. More precisely, for every $\gamma \neq 0$, the $\mathfrak{U}_{\text{res}}$ -affine coadjoint orbit $\mathcal{O}_{(0, \gamma)}$ of $(0, \gamma) \in (\mathfrak{u}_{\text{res}})_* \oplus \{\gamma\}$ is isomorphic to the restricted Grassmannian Gr_{res} via the map*

$$\begin{aligned} \Phi_\gamma : \text{Gr}_{\text{res}} &\rightarrow \mathcal{O}_{(0, \gamma)}, \\ W &\mapsto 2i\gamma(p_W - p_+), \end{aligned}$$

where p_W denotes the orthogonal projection on W . The pull-back by Φ_γ of the symplectic form on $\mathcal{O}_{(0, \gamma)}$ is (-2γ) -times the symplectic form ω_{Gr} on Gr_{res} .

Proof. An element in the affine coadjoint orbit $\mathcal{O}_{(0, \gamma)}$ of $(0, \gamma)$ is of the form (ρ, γ) with

$$\rho = \gamma(gdg^{-1} - d) = 2i\gamma(gp_+g^{-1} - p_+),$$

for some $g \in \mathfrak{U}_{\text{res}}$ (where we have used the identity $p_- = \text{id} - p_+$ to simplify the formula for the affine coadjoint action given in Proposition 2.9). By [39, Corollary III.4(ii)], Φ_γ is bijective for $\gamma \neq 0$. Since the manifold structure of the orbit $\mathcal{O}_{(0, \gamma)}$ is induced by the identification $\mathcal{O}_{(0, \gamma)} = \mathfrak{U}_{\text{res}} / (\mathfrak{U}(\mathcal{H}_+) \times \mathfrak{U}(\mathcal{H}_-))$, it follows from [39, Corollary III.4(i)] that Φ_γ is a diffeomorphism. The symplectic form $\omega_{\mathcal{O}}$ on $\mathcal{O}_{(0, \gamma)}$ is the $\mathfrak{U}_{\text{res}}$ -invariant symplectic form whose value at $(0, \gamma) \in \mathcal{O}_{(0, \gamma)}$ is the given by

$$\omega_{\mathcal{O}}(0, \gamma)(X_f(0, \gamma), X_g(0, \gamma)) = \{f, g\}_d(0, \gamma),$$

where f and g are any smooth functions on $(\mathfrak{u}_{\text{res}})_*$. Using formulas (2.10) and (2.8), it then follows that

$$\omega_{\mathcal{O}}(0, \gamma)(\gamma[D_{\mu}f, d], \gamma[D_{\mu}g, d]) = \gamma s(D_{\mu}f, D_{\mu}g).$$

Hence for every $A, B \in \mathfrak{u}_{\text{res}}$, one has

$$\omega_{\mathcal{O}}(0, \gamma)(\gamma[A, d], \gamma[B, d]) = \gamma s(A, B) = -2\gamma \Omega_{\text{Gr}}(A, B).$$

It follows that the real-valued anti-symmetric bilinear form on $\mathfrak{u}_{\text{res}}$ corresponding to the symplectic form $\omega_{\mathcal{O}}$ on $\mathcal{O}_{(0, \gamma)} = \text{U}_{\text{res}}/(\text{U}(\mathcal{H}_+) \times \text{U}(\mathcal{H}_-))$ equals $-2\gamma \Omega_{\text{Gr}}$ (where the latter identification is given by the orbit map), and this ends the proof. \square

Remark 2.14. We refer to the paper [29] for additional information on the relationship between the Banach Lie–Poisson spaces and the theory of Lie algebra extensions.

3. Coadjoint orbits of the restricted unitary group

This section includes some partial answers to Question 1.2. The main difficulty is to show that the isotropy group of an element in the predual $(\mathfrak{u}_{\text{res}})_*$ is a Lie subgroup of U_{res} , or equivalently that its Lie algebra is complemented in $\mathfrak{u}_{\text{res}}$. Using the averaging method developed in [4,6] for constructing closed complements, we will be able to show that the U_{res} -coadjoint orbit of every element $\rho \in (\mathfrak{u}_{\text{res}})_*$ which commutes with d is a smooth manifold and that its connected components are symplectic leaves of the characteristic distribution (see Proposition 3.3). It follows that the same conclusion holds for every element $\rho \in (\mathfrak{u}_{\text{res}})_*$ which is U_{res} -conjugate to an element commuting with d , or equivalently to a diagonal operator with respect to a Hilbert basis compatible with the eigenspaces of d . The set of elements with the latter property is not equal to the whole $(\mathfrak{u}_{\text{res}})_*$; however, it is dense (for more details see the proof of Corollary 3.5). Recall that in finite dimensions, every element in the Lie algebra $\mathfrak{u}(n)$ of the unitary group $\text{U}(n)$ is $\text{U}(n)$ -conjugate to a diagonal matrix with respect to a given basis of \mathbb{C}^n , or, in other words, $\text{U}(n)$ acts transitively on the set of Cartan subalgebras of $\mathfrak{u}(n)$. This is no longer true in the infinite-dimensional case (see Section 6.3). It is a difficult question to decide whether a given operator ρ in $(\mathfrak{u}_{\text{res}})_*$ or $\mathfrak{u}_{\text{res}}$ has the good property of being U_{res} -conjugate to a diagonal operator with respect to a basis adapted to the decomposition $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$. In Propositions 3.5 and 3.7, we give some concrete criteria to check that property.

Conjecture 3.1. *The real Banach space $(\mathfrak{u}_{\text{res}})_*$ has a natural structure of Banach Lie–Poisson space and its characteristic distribution is integrable.*

We refer to [27] for a discussion of integrable distributions on Banach manifolds. The meaning of the integrability of the characteristic distribution in Conjecture 3.1 is that for every $\mu_0 \in (\mathfrak{u}_{\text{res}})_*$ there exist a connected Banach manifold M and a smooth injective mapping $\psi : M \rightarrow (\mathfrak{u}_{\text{res}})_*$ such that $\mu_0 \in \psi(M)$ and for every $x \in M$ the tangent map $T_x \psi : T_x M \rightarrow T_{\psi(x)}((\mathfrak{u}_{\text{res}})_*)$ is also injective and its range is equal to the fiber of the characteristic distribution at the point $\psi(x) \in (\mathfrak{u}_{\text{res}})_*$. Such a pair (M, ψ) (or just the manifold M , for the sake of simplicity) is said to be an integral manifold of the characteristic distribution of $(\mathfrak{u}_{\text{res}})_*$ through the point μ_0 .

Remark 3.2. It is clear that

$$(\mathfrak{u}_{\text{res}})_* \hookrightarrow \mathfrak{u}_{\text{res}}$$

with a continuous inclusion map. On the other hand, it follows at once by the multiplication formula (2.2) that

$$[(\mathfrak{u}_{\text{res}})_*, \mathfrak{u}_{\text{res}}] \subseteq (\mathfrak{u}_{\text{res}})_*, \tag{3.1}$$

which implies that the predual $(\mathfrak{u}_{\text{res}})_*$ is left invariant by the coadjoint representation of the Banach Lie algebra $\mathfrak{u}_{\text{res}}$. Now the results of [28] imply the following two facts:

- The predual Banach space $(\mathfrak{u}_{\text{res}})_*$ has a natural structure of Banach Lie–Poisson space.
- If $\rho \in (\mathfrak{u}_{\text{res}})_*$ has the property that the corresponding isotropy group

$$U_{\text{res}, \rho} := \{u \in U_{\text{res}} \mid u\rho u^{-1} = \rho\}$$

is a Banach Lie subgroup of U_{res} , then the coadjoint orbit \mathcal{O}_ρ is an integral manifold of the characteristic distribution of $(\mathfrak{u}_{\text{res}})_*$. Moreover, \mathcal{O}_ρ is a weakly symplectic manifold when equipped with the orbit symplectic structure.

Thus, the desired conclusion will follow as soon as we prove that the isotropy group $U_{\text{res}, \rho}$ of any $\rho \in (\mathfrak{u}_{\text{res}})_*$ is a Banach Lie subgroup of U_{res} . Throughout the present paper, by *Banach Lie subgroup* we mean the same notion as in [9] or [28]: a subgroup of a Banach Lie group which has a structure of Banach Lie group of its own with respect to the relative topology and has the additional property that the corresponding Lie subalgebra has a closed complement in the Lie algebra of the ambient Banach Lie group.

As an easy consequence of the Harris–Kaup theorem (see for instance [5, Theorem 4.13]) the isotropy group $U_{\text{res}, \rho}$ of any $\rho \in (\mathfrak{u}_{\text{res}})_*$ does have a structure of Banach Lie group of its own with respect to the relative topology, so the only point that remains to be settled is the existence of a closed complement of the isotropy Lie algebra. The Lie algebra of $U_{\text{res}, \rho}$ is given by

$$\mathfrak{u}_{\text{res}, \rho} = \{a \in \mathfrak{u}_{\text{res}} \mid a\rho = \rho a\} = \{a \in \mathfrak{u}_{\text{res}} \mid (\forall t \in \mathbb{R}) \alpha_t(a) = a\},$$

where

$$\alpha : \mathbb{R} \rightarrow \mathcal{B}(\mathfrak{u}_{\text{res}}), \quad \alpha(t)b := \alpha_t(b) := \exp(t\rho) \cdot b \cdot \exp(-t\rho).$$

It is clear that α is a group homomorphism. Moreover, since $\rho \in (\mathfrak{u}_{\text{res}})_* \subseteq \mathfrak{u}_{\text{res}}$ and the adjoint action of the Banach Lie group U_{res} is continuous, it follows that $\alpha : \mathbb{R} \rightarrow \mathcal{B}(\mathfrak{u}_{\text{res}})$ is norm continuous.

On the other hand, it follows by (3.1) that

$$(\forall t \in \mathbb{R}) \quad \alpha_t((\mathfrak{u}_{\text{res}})_*) \subseteq (\mathfrak{u}_{\text{res}})_*, \tag{3.2}$$

since $\rho \in (\mathfrak{u}_{\text{res}})_*$. Then the concrete form of the duality pairing between $(\mathfrak{u}_{\text{res}})_*$ and $\mathfrak{u}_{\text{res}}$ (see (2.3)) shows that

$$(\forall t \in \mathbb{R}) \quad (\alpha_t|_{(\mathfrak{u}_{\text{res}})_*})^* = \alpha_{-t}, \tag{3.3}$$

and in particular each operator $\alpha_t : \mathfrak{u}_{\text{res}} \rightarrow \mathfrak{u}_{\text{res}}$ is weak*-continuous.

Now a complement to $u_{res, \rho}$ in u_{res} can be constructed by the averaging technique over the amenable group $(\mathbb{R}, +)$ provided one has $\sup_{t \in \mathbb{R}} \|\alpha_t\| < \infty$. (Some references for the aforementioned averaging technique are [4], the proof of Proposition 3.4 in [7], and [6].)

Additionally we note that since for every operator $T : X \rightarrow Y$ between the Banach spaces X and Y the norm of T equals the norm of its dual T^* , it is enough to estimate uniformly the norm of α_t restricted to the predual $(u_{res})_*$. This restriction is an adjoint action of the group corresponding to the predual.

Proposition 3.3. *If $\rho \in (u_{res})_*$ and $[d, \rho] = 0$, then the coadjoint isotropy group of ρ is a Banach Lie subgroup of U_{res} and the connected components of the corresponding U_{res} -coadjoint orbit \mathcal{O}_ρ are smooth leaves of the characteristic distribution of $(u_{res})_*$.*

Proof. According to Remark 3.2 it suffices to show that $\sup_{t \in \mathbb{R}} \|\alpha_t\| < \infty$. The hypothesis $[d, \rho] = 0$ shows that ρ preserves \mathcal{H}_+ and \mathcal{H}_- , that is

$$\rho = \begin{pmatrix} \rho_{++} & 0 \\ 0 & \rho_{--} \end{pmatrix} \in (u_{res})_*.$$

An element $b \in (u_{res})_*$ with block decomposition with respect to the direct sum $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$

$$b = \begin{pmatrix} b_{++} & b_{+-} \\ b_{-+} & b_{--} \end{pmatrix}$$

is the sum of an element

$$b_1 = \begin{pmatrix} b_{++} & 0 \\ 0 & b_{--} \end{pmatrix}$$

in the Lie algebra $u_0 := u_1 \cap (u(\mathcal{H}_+) \times u(\mathcal{H}_-))$ and an element

$$b_2 = \begin{pmatrix} 0 & b_{+-} \\ b_{-+} & 0 \end{pmatrix}$$

in the topological complement $\mathfrak{m} = u(\mathcal{H}) \cap (\mathfrak{S}_2(\mathcal{H}_+, \mathcal{H}_-) \oplus \mathfrak{S}_2(\mathcal{H}_-, \mathcal{H}_+))$ of u_0 in $(u_{res})_*$. Accordingly,

$$\begin{aligned} \|\alpha_t(b)\|_{(u_{res})_*} &= \|\exp(t\rho)b \exp(-t\rho)\|_{(u_{res})_*} \\ &= \|\exp(t\rho)b_1 \exp(-t\rho) + \exp(t\rho)b_2 \exp(-t\rho)\|_{(u_{res})_*} \\ &= \|e^{\text{ad}(t\rho)}(b_1) + e^{\text{ad}(t\rho)}(b_2)\|_{(u_{res})_*}. \end{aligned}$$

Since $\text{ad}(t\rho)$ preserves both u_0 and \mathfrak{m} , it follows that

$$e^{\text{ad}(t\rho)}(b_1) \in u_0 \quad \text{and} \quad e^{\text{ad}(t\rho)}(b_2) \in \mathfrak{m}.$$

By the very definition of the norm $\|\cdot\|_{(u_{res})_*}$, one has

$$\|\alpha_t(b)\|_{(u_{res})_*} = \|e^{\text{ad}(t\rho)}(b_1)\|_1 + \|e^{\text{ad}(t\rho)}(b_2)\|_2,$$

where $\|\cdot\|_1$ (respectively $\|\cdot\|_2$) is the usual norm in \mathfrak{S}_1 (respectively \mathfrak{S}_2). Since the conjugation by a unitary element preserves both $\|\cdot\|_1$ and $\|\cdot\|_2$, it follows that α_t acts by isometries on $(u_{\text{res}})_*$, in particular $\sup_{t \in \mathbb{R}} \|\alpha_t\| < \infty$. \square

Remark 3.4. The calculation in the proof of Proposition 3.3 actually shows that for every $u \in U_{\text{res}}$ satisfying $[d, u] = 0$ we have $\|ubu^{-1}\|_{\text{res}} = \|b\|_{\text{res}}$ whenever $b \in \mathcal{B}_{\text{res}}$. In fact

$$\|ubu^{-1}\|_{\text{res}} = \|ubu^{-1}\| + \|[d, ubu^{-1}]\|_2 = \|b\| + \|u[d, b]u^{-1}\|_2 = \|b\| + \|[d, b]\|_2 = \|b\|_{\text{res}},$$

where the second equality follows since $[d, u] = 0$. Note also that

$$\begin{aligned} \|ab\|_{\text{res}} &= \|ab\| + \|[d, a]b + a[d, b]\|_2 \leq \|a\|\|b\| + \|[d, a]\|_2\|b\| + \|a\|\|[d, b]\|_2 \\ &\leq \|a\|_{\text{res}}\|b\|_{\text{res}}. \end{aligned}$$

Corollary 3.5. *If $\rho \in (u_{\text{res}})_*$ is a finite-rank operator, then the coadjoint isotropy group of ρ is a Banach Lie subgroup of U_{res} and the connected components of the corresponding U_{res} -coadjoint orbit \mathcal{O}_ρ are smooth leaves of the characteristic distribution of $(u_{\text{res}})_*$.*

Proof. The set of finite-rank operators \mathcal{F} is a dense subset of the predual $(u_{\text{res}})_*$. For every skew-symmetric finite-rank operator F there exists a unitary operator $u \in \mathbf{1} + \mathcal{F}$, such that uFu^{-1} leaves both \mathcal{H}_- and \mathcal{H}_+ invariant. (This follows since any two finite-rank operators are contained in a certain finite-dimensional Lie algebra of finite-rank operators; see for instance [15, Chapter I, Lemma 1] or [36, Proposition 3.1].) Note that $u \in U_{\text{res}}$, and the isotropy groups of the elements F and uFu^{-1} are conjugated by the element u . Hence the isotropy group at any finite-rank operator is a Banach Lie subgroup of U_{res} , and this shows that the conclusion of Proposition 3.3 is satisfied if we replace the hypothesis $[d, \rho] = 0$ by the condition that ρ is a finite-rank operator. \square

Remark 3.6. An alternative way to prove Corollary 3.5 is to pick a $*$ -invariant d -invariant subalgebra containing the skew-symmetric finite-rank operator F and thus to reduce things to the finite-dimensional setting.

Corollary 3.7. *Assume that $\rho \in (u_{\text{res}})_*$ and that there exist an orthonormal basis $\{e_n\}_{n \geq 1}$ of the Hilbert space \mathcal{H} and the real numbers $t \in (0, 1)$ and $s \in (0, 3(1-t)/100]$ such that the following conditions are satisfied:*

- (i) *We have $\{e_n \mid n \geq 1\} \subseteq \mathcal{H}_+ \cup \mathcal{H}_-$.*
- (ii) *The matrix $(\rho_{mn})_{m,n \geq 1}$ of ρ with respect to the basis $\{e_n\}_{n \geq 1}$ has the properties*

$$|\rho_{m+1,n+1}| \leq t|\rho_{m,n}| \quad \text{whenever } m, n \geq 1,$$

and

$$|\rho_{m,n}|^2 \leq \frac{s^2}{(mn)^2} |\rho_{mm}\rho_{nn}| \quad \text{whenever } m, n \geq 1 \text{ and } m \neq n.$$

Then the coadjoint isotropy group of ρ is a Banach Lie subgroup of U_{res} and the connected components of the corresponding U_{res} -coadjoint orbit \mathcal{O}_ρ are smooth leaves of the characteristic distribution of $(\mathfrak{u}_{\text{res}})_*$.

Proof. It follows at once by [18, Theorem 1] that there exists an operator $a = -a^* \in \mathfrak{S}_2(\mathcal{H})$ such that the operator $u\rho u^{-1}$ is diagonal with respect to the basis $\{e_n\}_{n \geq 1}$, where $u = \exp a$. In particular we have $u \in U_2 \subseteq U_{\text{res}}$ and $[d, u\rho u^{-1}] = 0$, so that we can use Proposition 3.3 to get the desired conclusion. \square

Remark 3.8. Let $\rho \in \mathcal{B}(\mathcal{H})$. In addition to the applications of Proposition 3.3 in the proofs of Corollaries 3.5 and 3.7, we note that each of the following two conditions is equivalent to the existence of an unitary operator $u \in U_{\text{res}}$ such that $[d, u\rho u^{-1}] = 0$:

- (i) There exists $p \in \mathcal{B}(\mathcal{H})$ such that $p = p^* = p^2$, $p - p_+ \in \mathfrak{S}_2(\mathcal{H})$, and $\rho p = p\rho$.
- (ii) There exists an element $\mathcal{W} \in \text{Gr}_{\text{res}}$ such that $\rho(\mathcal{W}) \subseteq \mathcal{W}$.

In fact, our assertion concerning (i) follows at once since

$$\{p \in \mathcal{B}(\mathcal{H}) \mid p = p^* = p^2 \text{ and } p - p_+ \in \mathfrak{S}_2(\mathcal{H})\} = \{up_+u^{-1} \mid u \in U_{\text{res}}\}$$

according to [10, Lemma 3.1].

On the other hand, the assertion on condition (ii) holds since by [33, Proposition 7.1.3] we have

$$\text{Gr}_{\text{res}} = \{u(\mathcal{H}_+) \mid u \in U_{\text{res}}\}$$

and, in addition, if $p \in \mathcal{B}(\mathcal{H})$ is the orthogonal projection onto some closed subspace $\mathcal{W} \subseteq \mathcal{H}$ then $\rho(\mathcal{W}) \subseteq \mathcal{W}$ if and only if $[p, \rho] = 0$. To see this, recall that $\rho^* = -\rho$, hence $\rho(\mathcal{W}) \subseteq \mathcal{W}$ if and only if $\rho(\mathcal{W}^\perp) \subseteq \mathcal{W}^\perp$.

4. Some smooth adjoint orbits of the restricted unitary group

We are going to investigate in this section the smoothness of adjoint orbits of the restricted unitary group and derive some consequences about the smoothness of affine coadjoint orbits of the restricted unitary group. In particular, we shall find sufficiently many smooth adjoint orbits of U_{res} to fill an open subset of the Lie algebra $\mathfrak{u}_{\text{res}}$ (see Proposition 4.2), as well as sufficiently many smooth affine coadjoint orbits of U_{res} to fill an open subset of the Lie algebra $(\mathfrak{u}_{\text{res}})_*$ (see Corollary 4.4).

Lemma 4.1. Assume that the element

$$\rho = \begin{pmatrix} \rho_{++} & \rho_{+-} \\ \rho_{-+} & \rho_{--} \end{pmatrix} \in \mathfrak{u}_{\text{res}}$$

satisfies the conditions

$$\sigma(\rho_{++}) \cap \sigma(\rho_{--}) = \emptyset \tag{4.1}$$

(where $\sigma(\rho_{\pm\pm})$ denotes the spectrum of $\rho_{\pm\pm}$) and

$$\|\rho_{+-}\|_2 < \frac{1}{2} \text{dist}(\sigma(\rho_{++}), \sigma(\rho_{--})). \tag{4.2}$$

Then there exists $u \in U_{\text{res}}$ such that $[d, u^{-1}\rho u] = 0$.

Proof. The hypotheses (4.1) and (4.2) imply that there exists a Hilbert–Schmidt operator $k : \mathcal{H}_+ \rightarrow \mathcal{H}_-$ satisfying the operator Riccati equation

$$k\rho_{+-}k + k\rho_{++} - \rho_{--}k = \rho_{-+}.$$

(This result was obtained in [21]; see also [1, Theorem 4.6 and Remark 4.7], as well as [2].) Then the operator

$$g = \begin{pmatrix} \text{id}_{\mathcal{H}_+} & k^* \\ k & -\text{id}_{\mathcal{H}_-} \end{pmatrix}$$

is invertible and has the properties $[d, g] \in \mathfrak{S}_2(\mathcal{H})$, $g = g^*$, $[d, g^2] = 0$ and

$$[d, g^{-1}\rho g] = 0 \tag{4.3}$$

(see [1, Section 2.3]). Now let $g = us$ be the polar decomposition of the invertible operator $g \in \mathcal{B}(\mathcal{H})$, where $u \in \mathcal{B}(\mathcal{H})$ is unitary and $s = (g^*g)^{1/2}$.

On the other hand, since $d^* = -d$, it follows that the commutant $\{d\}'$ is a von Neumann algebra of operators on \mathcal{H} . Thus, since $g = g^*$ and $g^*g = g^2 \in \{d\}'$, it is straightforward to deduce that $(g^*g)^{1/2} \in \{d\}'$, that is, $[d, s] = 0$. Now recall that $[d, g] \in \mathfrak{S}_2(\mathcal{H})$ to deduce that the unitary operator $u = gs^{-1}$ satisfies $[d, u] \in \mathfrak{S}_2(\mathcal{H})$, that is, $u \in U_{\text{res}}$.

Moreover by (4.3) we have

$$0 = [d, g^{-1}\rho g] = [d, s^{-1}u^{-1}\rho us] = s^{-1}[d, u^{-1}\rho u]s,$$

where the latter equality follows since we have seen that $[d, s] = 0$. Now we get $[d, u^{-1}\rho u] = 0$, as desired. \square

Proposition 4.2. *For any $\gamma \in \mathbb{R} \setminus \{0\}$, there exists an open U_{res} -invariant neighborhood V of $\gamma d \in \mathfrak{u}_{\text{res}}$ such that V is a union of smooth adjoint orbits of the Banach Lie group U_{res} .*

Proof. Denote by V_γ the set of all elements

$$\rho = \begin{pmatrix} \rho_{++} & \rho_{+-} \\ \rho_{-+} & \rho_{--} \end{pmatrix} \in \mathfrak{u}_{\text{res}}$$

satisfying conditions

$$\sigma(\rho_{\pm\pm}) \subseteq \{y \in i\mathbb{R} \mid |y \mp \gamma| < 1/3\}$$

and

$$\|\rho_{\pm\mp}\|_2 < \frac{2}{3}.$$

Recall that $\rho_{\pm\pm} \mp \gamma i$ is skew-Hermitian, hence its spectral radius equals its operator norm and the condition $\sigma(\rho_{\pm\pm}) \subseteq \{y \in i\mathbb{R} \mid |y \mp \gamma i| < 1/3\}$ is equivalent to $\|\rho_{\pm\pm} \mp \gamma i\| < 1/3$. Note that

$$\begin{pmatrix} \rho_{++} & 0 \\ 0 & \rho_{--} \end{pmatrix} = \rho - \frac{1}{2i}[d, \rho],$$

hence the condition on the spectrum of $\rho_{\pm\pm}$ defines an open subset of $\mathfrak{u}_{\text{res}}$. On the other hand, the condition $\|\rho_{\pm\mp}\|_2 < 2/3$ is equivalent to $\|[d, \rho]\|_{\text{res}} < 2\sqrt{2}/3$ (since $\rho_{\pm\mp}^* = \rho_{\mp\pm}$) hence it also describes an open subset of $\mathfrak{u}_{\text{res}}$. It follows that V_γ is an open neighborhood of $\gamma d \in \mathfrak{u}_{\text{res}}$. We are going to show that the set

$$V := \bigcup_{u \in \mathfrak{U}_{\text{res}}} \text{Ad}_{\mathfrak{U}_{\text{res}}}(u)V_\gamma \subseteq \mathfrak{u}_{\text{res}}$$

has the desired properties.

Indeed, V is clearly invariant under the adjoint action of $\mathfrak{U}_{\text{res}}$, it is a union of open sets, and one of these open sets contains γd . Moreover, it follows by Lemma 4.1 along with the construction of V that for every $\rho \in V$ there exists $u \in \mathfrak{U}_{\text{res}}$ such that $[d, u^{-1}\rho u] = 0$. Next denote $\tilde{\rho} = u^{-1}\rho u$, so that $\exp(t\rho) = u \exp(t\tilde{\rho})u^{-1}$ for all $t \in \mathbb{R}$. Then for all $t \in \mathbb{R}$ and $b \in \mathfrak{u}_{\text{res}}$ it follows by means of Remark 3.4 that

$$\begin{aligned} \|\exp(t\rho)b \exp(-t\rho)\|_{\text{res}} &= \|u \exp(t\tilde{\rho})u^{-1}bu \exp(-t\tilde{\rho})u^{-1}\|_{\text{res}} \\ &\leq \|u\|_{\text{res}} \|\exp(t\tilde{\rho})u^{-1}bu \exp(-t\tilde{\rho})\|_{\text{res}} \|u^{-1}\|_{\text{res}} \\ &= \|u\|_{\text{res}} \|u^{-1}bu\|_{\text{res}} \|u^{-1}\|_{\text{res}} \\ &\leq \|u\|_{\text{res}}^2 \|u^{-1}\|_{\text{res}}^2 \|b\|_{\text{res}}. \end{aligned}$$

Consequently the 1-parameter group

$$\alpha : \mathbb{R} \rightarrow \mathcal{B}(\mathfrak{u}_{\text{res}}), \quad \alpha_t(b) = \exp(t\rho)b \exp(-t\rho),$$

satisfies

$$\sup_{t \in \mathbb{R}} \|\alpha_t\| \leq \|u\|_{\text{res}}^2 \|u^{-1}\|_{\text{res}}^2.$$

Now the arguments in Remark 3.2 show that the adjoint isotropy group of ρ is a Lie subgroup of $\mathfrak{U}_{\text{res}}$, and thus the adjoint orbit of ρ is smooth. \square

An alternative way to see that the set V_γ in the previous proof is open follows by the well-known upper continuity of the spectrum as a function of the operator (see, e.g. [12,14,26]). We also note that a shorter argument for the fact that the adjoint isotropy group of ρ is a Lie subgroup of $\mathfrak{U}_{\text{res}}$ consists in an application of Proposition 3.3 for $u^{-1}\rho u$ along with the fact that the stabilizer of $u^{-1}\rho u$ is conjugate to the stabilizer of ρ .

Corollary 4.3. *For any $\gamma \in \mathbb{R} \setminus \{0\}$, there exists an open $\mathfrak{U}_{1,2}$ -invariant neighborhood V of $\gamma d \in \mathfrak{u}_{\text{res}} = \mathfrak{u}_{1,2}^*$ such that V is a union of smooth coadjoint orbits of the Banach Lie group $\mathfrak{U}_{1,2}$.*

Proof. Apply Proposition 4.2 along with the fact that $\mathfrak{U}_{1,2} \hookrightarrow \mathfrak{U}_{\text{res}}$ and the adjoint action of $\mathfrak{U}_{\text{res}}$ restricts to the coadjoint action of $\mathfrak{U}_{1,2}$. \square

Corollary 4.4. *For any $\gamma \in \mathbb{R} \setminus \{0\}$, there exists an open U_{res} -invariant neighborhood W of $(0, \gamma) \in (\tilde{u}_{\text{res}})_*$ such that W is a union of smooth affine coadjoint orbits of the Banach Lie group U_{res} .*

Proof. For any $(\mu, \lambda) \in (\tilde{u}_{\text{res}})_*$, the operator $\rho = \mu - \lambda d$ belongs to $\mathfrak{u}_{\text{res}}$ and

$$\|\mu - \lambda d\|_{\text{res}} \leq |\lambda| + \|\mu\|_{(\tilde{u}_{\text{res}})_*}$$

which implies that the linear map $\theta : (\mu, \lambda) \in (\tilde{u}_{\text{res}})_* \mapsto \mu - \lambda d \in \mathfrak{u}_{\text{res}}$ is continuous. With the notation introduced in the proof of Proposition 4.2, let $W_\gamma := \{(\mu, \lambda) \in (\tilde{u}_{\text{res}})_* \mid \mu - \lambda d \in V_{-\gamma}\}$, that is, $W_\gamma = \theta^{-1}(V_{-\gamma})$ and hence W_γ is open in $(\tilde{u}_{\text{res}})_*$. Note that $(0, \gamma) \in W_\gamma$. Moreover since

$$g \cdot (\mu, \lambda) = (\mu, \lambda) \iff g\mu g^{-1} - \lambda(gdg^{-1} - d) = \mu,$$

the isotropy group of any (μ, λ) for the affine coadjoint action of U_{res} equals the isotropy group of $\mu - \lambda d$ for the adjoint action of U_{res} , hence is a Banach Lie subgroup of U_{res} by Proposition 4.2. Now

$$W := \bigcup_{u \in U_{\text{res}}} u \cdot W_\gamma \subseteq (\tilde{u}_{\text{res}})_*$$

has the desired properties. \square

5. The Banach Lie–Poisson space associated to the central extension of \mathfrak{u}_2

Denote by $\tilde{\mathfrak{u}}_2 := \mathfrak{u}_2 \oplus \mathbb{R}$ the central extension of \mathfrak{u}_2 defined by the restriction of s to $\mathfrak{u}_2 \times \mathfrak{u}_2$, where s is the two-cocycle defined in (2.6). The natural isomorphism $(\tilde{\mathfrak{u}}_2)^* \simeq \tilde{\mathfrak{u}}_2$ implies that $\tilde{\mathfrak{u}}_2$ is a Banach Lie–Poisson space, for the Poisson bracket given by

$$\{f, g\}_d(\mu, \gamma) := \langle \mu, [D_\mu f(\mu), D_\mu g(\mu)] \rangle + \gamma s(D_\mu f, D_\mu g),$$

where $f, g \in C^\infty(\tilde{\mathfrak{u}}_2)$, (μ, γ) is an arbitrary element in $\tilde{\mathfrak{u}}_2$, and D_μ denotes the partial Fréchet derivative with respect to $\mu \in \mathfrak{u}_2$.

Theorem 5.1. *The characteristic distribution of the Banach Lie–Poisson space $\tilde{\mathfrak{u}}_2$ is integrable.*

Proof. In order to prove that the characteristic distribution is integrable, it suffices to check that all of the affine coadjoint isotropy groups are Lie subgroups of the Hilbert Lie group U_2 . For this purpose we note that, for arbitrary $(\mu, \gamma) \in \tilde{\mathfrak{u}}_2$, the corresponding isotropy group of the affine coadjoint action of U_2 on $\tilde{\mathfrak{u}}_2$ is

$$(U_2)_{(\mu, \gamma)} = \{g \in U_2 \mid \mu = g\mu g^{-1} - \gamma g d g^{-1} + \gamma d\},$$

according to the explicit expression of the affine coadjoint action in Proposition 2.9. The previous equality implies that

$$(U_2)_{(\mu, \gamma)} = \{g \in \mathbb{C}\mathbf{1} + \mathfrak{S}_2(\mathcal{H}) \mid g^* g = g g^* = 1 \text{ and } \mu = g\mu g^{-1} - \gamma g d g^{-1} + \gamma d\},$$

and now it is clear that $(U_2)_{(\mu,\gamma)}$ is an algebraic subgroup of degree ≤ 2 of the group of invertible elements in the unital Banach algebra $\mathbb{C}\mathbf{1} + \mathfrak{S}_2(\mathcal{H})$. Then the Harris–Kaup theorem (see for instance [5, Theorem 4.13]) implies that $(U_2)_{(\mu,\gamma)}$ is a Lie group with respect to the topology inherited from $\mathbb{C}\mathbf{1} + \mathfrak{S}_2(\mathcal{H})$. In particular, this topology coincides with the one inherited from U_2 . Since U_2 is a Hilbert Lie group, hence the Lie algebra of $(U_2)_{(\mu,\gamma)}$ has a complement in the Lie algebra of U_2 , it then follows that $(U_2)_{(\mu,\gamma)}$ is a Banach Lie subgroup of U_2 , and this concludes the proof. (Compare Remark 3.2.) \square

The transitivity of the action of the Lie group U_2 on the connected component Gr_{res}^0 of the restricted Grassmannian has been established in [10, Theorem 3.5] and [23, Proposition V.7]. That the action of the subgroup $U_{1,2}$ of U_2 on Gr_{res}^0 is transitive has been proved in [37, Section 1.3.4] with the help of the canonical basis defined in [33, Section 7.3] and associated to any element of the restricted Grassmannian. Below we give a shorter and geometrical proof of the latter fact.

Proposition 5.2. *The connected component Gr_{res}^0 of the restricted Grassmannian is a homogeneous space under the unitary group $U_{1,2} \subset U_2$.*

Proof. The restricted Grassmannian is a symmetric space of the restricted unitary group U_{res} . It follows from the description of geodesics in [3, Proposition 8.8] (see also [11,30] or its infinite-dimensional version as given in [25, Example 3.9] or [38, Proposition 1.9]) that each geodesic of Gr_{res} starting at $W \in Gr_{res}^0$ is given by

$$\beta(t) = (\exp tX) \cdot \mathcal{H}_+, \quad X \in \mathfrak{m}_W, \tag{5.1}$$

where \mathfrak{m}_W is the orthogonal in \mathfrak{u}_{res} to the Lie algebra of the isotropy group of W . For $W = \mathcal{H}_+$ we have $\mathfrak{m} = \mathfrak{u}(\mathcal{H}) \cap (\mathfrak{S}_2(\mathcal{H}_+, \mathcal{H}_-) \oplus \mathfrak{S}_2(\mathcal{H}_-, \mathcal{H}_+))$, and for $W = g \cdot \mathcal{H}_+$ with $g \in U_{res}$, we have $\mathfrak{m}_W = g\mathfrak{m}g^{-1}$. Note that for $X \in \mathfrak{m}$, $\exp tX$ belongs to $U_{1,2} \subset U_2$. Since the Hopf–Rinow theorem is no longer true in the infinite-dimensional case, it is not clear whether every two elements in the complete connected manifold Gr_{res}^0 can be joined by a geodesic. Nevertheless [13, Theorem B] asserts that, for every $W \in Gr_{res}^0$, the set of elements which can be joined to W by a unique minimal geodesic contains a dense G_δ set. Moreover from the properties of the Riemannian exponential map, there exists a neighborhood \mathcal{V} of \mathcal{H}_+ in Gr_{res}^0 such that every element in \mathcal{V} can be joined to \mathcal{H}_+ by a (minimal) geodesic. Hence an arbitrary element $W \in Gr_{res}^0$ can be joined to an element $W' \in \mathcal{V}$ by a geodesic

$$\beta_1(t) = (\exp tX_1) \cdot W', \quad X_1 \in \mathfrak{m}_{W'}, \quad t \in [0, 1],$$

and W' can be joined to \mathcal{H}_+ by a geodesic

$$\beta_2(t) = (\exp tX_2) \cdot \mathcal{H}_+, \quad X_2 \in \mathfrak{m}, \quad t \in [0, 1].$$

Consequently

$$W = \beta_1(1) = (\exp X_1) \cdot W' = (\exp X_1)(\exp X_2) \cdot \mathcal{H}_+.$$

But X_1 belongs to $\mathfrak{m}_{W'} = \exp(X_2)\mathfrak{m}\exp(-X_2)$, hence

$$W = (\exp X_2 \exp X_3) \cdot \mathcal{H}_+,$$

where $X_3 = \text{Ad}(\exp(-X_2))(X_1)$ belongs to \mathfrak{m} . Since $\exp X_3$ and $\exp X_2$ are elements of the unitary group $U_{1,2}$, it follows that their product belongs to $U_{1,2}$. Thus $U_{1,2}$ acts transitively on Gr_{res}^0 . \square

Theorem 5.3. *The connected component Gr_{res}^0 of the restricted Grassmannian is a strong symplectic leaf in the Banach Lie–Poisson space $\tilde{\mathfrak{u}}_2$. More precisely, for every $\gamma \neq 0$, the U_2 -affine coadjoint orbit $\tilde{\mathcal{O}}_{(0,\gamma)}$ of $(0, \gamma) \in \tilde{\mathfrak{u}}_2$ is diffeomorphic to Gr_{res}^0 via the application*

$$\begin{aligned} \Phi_\gamma : \text{Gr}_{\text{res}}^0 &\rightarrow \tilde{\mathcal{O}}_{(0,\gamma)}, \\ W &\mapsto 2i\gamma(p_W - p_+), \end{aligned}$$

where p_W denotes the orthogonal projection on W . The pull-back by Φ_γ of the symplectic form on $\tilde{\mathcal{O}}_{(0,\gamma)}$ is (-2γ) -times the symplectic form ω_{Gr} on Gr_{res}^0 .

Proof. The assertion follows by the method of proof of Theorem 2.13, since Gr_{res}^0 is transitively acted upon by the group U_2 according to Proposition 5.2. \square

Next we shall investigate the existence of invariant complex structures on certain covering spaces of the symplectic leaves of $\tilde{\mathfrak{u}}_2$ (Corollary 5.6 below). To this end we need two facts holding in a more general setting. In connection with the first of these statements, we note that invariant complex structures on certain homogeneous spaces related to derivations of L^* -algebras have been previously obtained by a different method in [22, Theorem IV.5].

Proposition 5.4. *Let \mathfrak{X} be a real Hilbert Lie algebra with a scalar product denoted by $(\cdot | \cdot)$. Assume that there exists a connected Hilbert Lie group $U_{\mathfrak{X}}$ whose Lie algebra is \mathfrak{X} ; we write $\mathbf{L}(U_{\mathfrak{X}}) = \mathfrak{X}$.*

Now let $D : \mathfrak{X} \rightarrow \mathfrak{X}$ be a bounded linear derivation such that

$$(\forall x, y \in \mathfrak{X}) \quad (Dx | y) = -(x | Dy). \tag{5.2}$$

Consider the closed subalgebra $\mathfrak{h}_0 := \text{Ker } D$ of \mathfrak{X} and define

$$H_0 := \langle \exp_{U_{\mathfrak{X}}}(\mathfrak{h}_0) \rangle,$$

that is, the subgroup of $U_{\mathfrak{X}}$ generated by the image of \mathfrak{h}_0 by the exponential map.

If it happens that H_0 has a structure of Banach Lie group with respect to the topology inherited from $U_{\mathfrak{X}}$, then it is actually a Banach Lie subgroup of $U_{\mathfrak{X}}$ and the smooth homogeneous space $U_{\mathfrak{X}}/H_0$ has an invariant complex structure.

Proof. Denote $\mathcal{L} := \mathfrak{X}_{\mathbb{C}}$, that is, the complex Hilbert Lie algebra which is the complexification of \mathfrak{X} and is endowed with the complex scalar product $(\cdot | \cdot)$ extending the scalar product of \mathfrak{X} . We denote the complex linear extension of D to \mathcal{L} again by D .

Then $D^* = -D$ as operators on the complex Hilbert space \mathcal{L} , so that $-iD \in \mathcal{B}(\mathcal{L})$ is a self-adjoint operator. Let us denote its spectral measure by $\delta \mapsto E(\delta)$. Thus $E(\cdot)$ is a spectral measure on \mathbb{R} and we have

$$D = i \int_{\mathbb{R}} t dE(t).$$

Also denote $S = (-\infty, 0]$, which is a closed subsemigroup of \mathbb{R} , and

$$\mathfrak{k} := \text{Ran } E(-S) = \text{Ran } E([0, \infty)) \subseteq \mathcal{L}.$$

Then \mathfrak{k} is a closed subspace of \mathcal{L} since it is the range of an idempotent continuous map. In addition, since D is a derivation of the Hilbert Lie algebra \mathfrak{X} and S is a closed semigroup, it follows by [5, Proposition 6.4] that \mathfrak{k} is a complex subalgebra of \mathcal{L} with the following properties:

- (i) $[\mathfrak{h}_0, \mathfrak{k}] \subseteq \mathfrak{k}$,
- (ii) $\mathfrak{k} \cap \bar{\mathfrak{k}} = \mathfrak{h}_0 + i\mathfrak{h}_0 (= \text{Ker } D)$, and
- (iii) $\mathfrak{k} + \bar{\mathfrak{k}} = \mathcal{L}$.

Moreover, for every $y \in \mathfrak{h}_0$ and all $x \in \mathfrak{X}$ we have

$$D[y, x] = [Dy, x] + [y, Dx] = [y, Dx]$$

since $Dy = 0$. Therefore, we have $D \circ \text{ad}_{\mathfrak{X}} y = \text{ad}_{\mathfrak{X}} y \circ D$ for each $y \in \mathfrak{h}_0$. According to the definition of H_0 , it then follows that for arbitrary $h \in H_0$ we have $\text{Ad}_{U_{\mathfrak{X}}} h \circ D = D \circ \text{Ad}_{U_{\mathfrak{X}}} h$ on \mathfrak{X} . Then the latter equality holds throughout \mathcal{L} , and it then follows that the operator $\text{Ad}_{U_{\mathfrak{X}}} h : \mathcal{L} \rightarrow \mathcal{L}$ commutes with every value of the spectral measure $E(\cdot)$. In particular we have $\text{Ad}_{U_{\mathfrak{X}}}(h) \circ E(-S) = E(-S) \circ \text{Ad}_{U_{\mathfrak{X}}}(h)$, whence

$$(i') \quad (\forall h \in H_0) \text{Ad}_{U_{\mathfrak{X}}}(h)\mathfrak{k} \subseteq \mathfrak{k}.$$

Now [5, Theorem 6.1] shows that the smooth homogeneous space $U_{\mathfrak{X}}/H_0$ has an invariant complex structure. \square

Proposition 5.5. *Let \mathcal{H} be an infinite-dimensional complex Hilbert space and let $a \in \mathcal{B}(\mathcal{H})$ such that $a^* = -a$. Denote by*

$$D = \text{ad}_{u_2} a : u_2 \rightarrow u_2, \quad x \mapsto [a, x],$$

the derivation of the compact L^ -algebra u_2 defined by a , and denote*

$$\mathfrak{h}_0 := \text{Ker } D = \{x \in u_2 \mid [a, x] = 0\}.$$

Next denote

$$H := \{u \in U_2 \mid uau^{-1} = a\}$$

and in addition define

$$H_0 := \langle \exp(\mathfrak{h}_0) \rangle.$$

That is, H_0 is the subgroup of U_2 generated by the image of \mathfrak{h}_0 by the exponential map. Then the following assertions hold:

- (j) Both H and H_0 are Banach Lie subgroups of U_2 .
- (jj) The subgroup H_0 is the connected component of $\mathbf{1} \in H$.
- (jjj) The natural map

$$U_2/H_0 \rightarrow U_2/H, \quad uH_0 \mapsto uH,$$

is an U_2 -equivariant smooth covering map.

Proof. Consider the Banach algebra $\mathcal{A} := \mathbb{C}\mathbf{1} + \mathfrak{S}_2(\mathcal{H})$ and denote by $\varphi: \mathcal{A} \rightarrow \mathbb{C}$ the continuous linear functional uniquely defined by the conditions $\varphi(\mathbf{1}) = 1$ and $\text{Ker } \varphi = \mathfrak{S}_2(\mathcal{H})$. Then we have

$$H = \{u \in \mathcal{A}^\times \mid u^*u = uu^* = \mathbf{1}, ua = au, \text{ and } \varphi(u) = 1\}$$

hence, by the Harris–Kaup theorem (see for instance [5, Theorem 4.13]), H is a subgroup of \mathcal{A}^\times that carries a Banach Lie group structure of its own. In addition, the Lie algebra

$$\mathbf{L}(H) = \{x \in \mathcal{A} \mid x^* = -x \text{ and } xa = ax\} = \mathfrak{h}_0,$$

of H has a closed complement in u_2 since the latter is a real Hilbert space. Thus H is a Banach Lie subgroup of U_2 .

On the other hand, H_0 has the structure of connected Lie group such that the inclusion map $H_0 \hookrightarrow U_2$ is an immersion and $\mathbf{L}(H_0) = \mathfrak{h}_0$. (See for instance [5, Theorem 3.5] and its proof.) Since $H_0 \subseteq H$ and $\mathbf{L}(H_0) = \mathbf{L}(H) = \mathfrak{h}_0$, it then follows that H_0 is the connected component of $\mathbf{1} \in H$. This can be seen directly by Lie theoretic methods; specifically, one just has to use the fact that the exponential map of any Banach Lie group is a local diffeomorphism at 0. An alternative approach is to use the proof of Lie’s second theorem by means of the Frobenius theorem (see for instance [19, Chapter VI, Theorem 5.4]). According to that proof, the connected group H_0 is the integral manifold through $\mathbf{1}$ corresponding to a smooth left-invariant integrable distribution on U_2 whose fiber at $\mathbf{1}$ is (the complemented closed Lie subalgebra) \mathfrak{h}_0 . Now recall the universality property of the integral leaves of integrable distributions according to [19, Chapter VI, Theorem 4.2] or, more generally [27, Theorem 4(iii)], which implies that the inclusion map $H_0 \hookrightarrow H$ is smooth. Then the wished-for property that H_0 is open in H follows since H_0 and H have the same tangent space at $\mathbf{1} \in H_0 \subseteq H$.

By either of these methods it follows that H_0 is an open subgroup of the Banach Lie subgroup H of U_2 , and then H_0 is in turn a Lie subgroup of U_2 . Thus assertions (j) and (jj) are proved. Assertion (jjj) follows since the natural map $U_2/H_0 \rightarrow U_2/H$ is clearly an U_2 -equivariant map whose tangent map at every point is an isomorphism. \square

In the following statement we need the notion of symplectic leaf in a Banach Lie–Poisson space. Let G be a Banach Lie group with Lie algebra \mathfrak{g} . Assume that \mathfrak{g} admits a predual \mathfrak{g}_*

such that the coadjoint action of G on \mathfrak{g}^* preserves the predual space \mathfrak{g}_* . Then, for any $\rho \in \mathfrak{g}_*$ such that the isotropy subgroup $G_\rho := \{g \in G \mid \text{Ad}_g^* \rho = \rho\}$ is a Banach Lie subgroup of G , the coadjoint orbit $\mathcal{O} := \{\text{Ad}_g^* \rho \mid g \in G\} \subset \mathfrak{g}_*$ is a Banach manifold diffeomorphic to the quotient G/G_ρ , weakly immersed in \mathfrak{g}_* , and the Banach Lie–Poisson structure of \mathfrak{g}_* induces on \mathcal{O} a weak symplectic form given by the usual formula (see [28, Theorems 7.3 and 7.4]). Weak immersion means that the derivative of the inclusion is only injective without any assumption on the closedness of the range, let alone splitting assumptions. This statement was also used in Remark 3.2 for $G = U_{\text{res}}$. See also the comments preceding it regarding integrable distributions on Banach manifolds. Several classes of Banach Lie–Poisson spaces that are unions of smooth symplectic leaves are given in [7]. In the corollary below the situation is simpler because we are dealing with a Hilbert Lie–Poisson space.

Corollary 5.6. *Every symplectic leaf of the Hilbert Lie–Poisson space \tilde{u}_2 is transitively acted on by U_2 by means of the affine coadjoint action and is U_2 -equivariantly covered by some complex homogeneous space of U_2 .*

Proof. Let $(\mu, \gamma) \in \tilde{u}_2$ arbitrary and denote $a := \mu - \gamma d \in \mathcal{B}(\mathcal{H})$. With the notation of Proposition 5.5, it is clear that H is equal to the isotropy group of the affine coadjoint action of U_2 . Thus the symplectic leaf $\tilde{\mathcal{O}}_{(\mu, \gamma)}$ through (μ, γ) is U_2 -equivariantly diffeomorphic to U_2/H . Now the conclusion follows since U_2/H is U_2 -equivariantly covered by the complex homogeneous space U_2/H_0 , according to Propositions 5.4 and 5.5. \square

Remark 5.7. It follows by Corollary 5.6 that every simply connected symplectic leaf of the Banach Lie–Poisson space \tilde{u}_2 has an U_2 -invariant complex structure. For instance, this is the case for the connected component Gr_{res}^0 of the restricted Grassmannian viewed as a symplectic leaf of \tilde{u}_2 by means of Theorem 5.3.

6. Some pathological properties of the restricted algebras

6.1. Unbounded unitary groups in the restricted algebra

We are going to point out a property that provides a good illustration for the difference between the Banach $*$ -algebra \mathcal{B}_{res} and a C^* -algebra (Proposition 6.2 below).

Lemma 6.1. *Let $a \in \mathcal{B}(\mathcal{H}_-, \mathcal{H}_+)$ and assume that $a = v|a|$ and $a^* = w|a^*|$ are the polar decompositions of a and a^* , where $|a| \in \mathcal{B}(\mathcal{H}_-)$ and $|a^*| \in \mathcal{B}(\mathcal{H}_+)$, while $v: \mathcal{H}_- \rightarrow \mathcal{H}_+$ and $w: \mathcal{H}_+ \rightarrow \mathcal{H}_-$ are partial isometries. Next, denote*

$$\rho = \begin{pmatrix} 0 & a \\ -a^* & 0 \end{pmatrix} \in \mathcal{B}(\mathcal{H}).$$

Then

$$\exp \rho = \begin{pmatrix} \cos |a^*| & v \sin |a| \\ -w \sin |a^*| & \cos |a| \end{pmatrix}.$$

Proof. We have

$$\rho^2 = \begin{pmatrix} -aa^* & 0 \\ 0 & -a^*a \end{pmatrix} = - \begin{pmatrix} |a^*|^2 & 0 \\ 0 & |a|^2 \end{pmatrix}$$

hence

$$(\forall n \geq 0) \quad \rho^{2n} = (-1)^n \begin{pmatrix} |a^*|^{2n} & 0 \\ 0 & |a|^{2n} \end{pmatrix}.$$

This implies that for every $n \geq 0$ we have

$$\begin{aligned} \rho^{2n+1} &= \rho \cdot \rho^{2n} = (-1)^n \begin{pmatrix} 0 & v|a| \\ -w|a^*| & 0 \end{pmatrix} \begin{pmatrix} |a^*|^{2n} & 0 \\ 0 & |a|^{2n} \end{pmatrix} \\ &= (-1)^n \begin{pmatrix} 0 & v|a|^{2n+1} \\ -w|a^*|^{2n+1} & 0 \end{pmatrix}. \end{aligned}$$

Consequently

$$\exp \rho = \sum_{n=0}^{\infty} \left(\frac{1}{(2n)!} \rho^{2n} + \frac{1}{(2n+1)!} \rho^{2n+1} \right) = \begin{pmatrix} \cos |a^*| & v \sin |a| \\ -w \sin |a^*| & \cos |a| \end{pmatrix}$$

which concludes the proof. \square

Proposition 6.2. *All of the unitary groups $(\mathbf{1} + \mathcal{F}) \cap \mathbf{U}(\mathcal{H})$, $\mathbf{U}_{1,2}$, and \mathbf{U}_{res} are unbounded subsets of the unital associative Banach algebra \mathcal{B}_{res} .*

Proof. We have

$$(\mathbf{1} + \mathcal{F}) \cap \mathbf{U}(\mathcal{H}) \subseteq \mathbf{U}_{1,2} \subseteq \mathbf{U}_{\text{res}}$$

so it suffices to show that

$$\sup \{ \|u\|_{\text{res}} \mid u \in (\mathbf{1} + \mathcal{F}) \cap \mathbf{U}(\mathcal{H}) \} = \infty. \tag{6.1}$$

To this end let $n \geq 1$ be an arbitrary positive integer, pick a projection $q_n = q_n^* = q_n^2 \in \mathcal{B}(\mathcal{H}_-)$ with $\dim(\text{Ran } q_n) = n$ and define $a_n := v_n((\pi/2)q_n) = (\pi/2)v_n \in \mathcal{B}(\mathcal{H}_-, \mathcal{H}_+)$, where $v_n: \mathcal{H}_- \rightarrow \mathcal{H}_+$ is an arbitrary partial isometry such that $v_n^*v_n = q_n$. Then $|a_n| = (\pi/2)q_n$, so that $\sin |a_n| = q_n$ and then $\|(\sin |a_n|)\|_2 = \sqrt{\dim(\text{Ran } q_n)} = \sqrt{n}$. Now Lemma 6.1 shows that the element

$$\rho_n = \begin{pmatrix} 0 & a_n \\ -a_n^* & 0 \end{pmatrix} \in \mathbf{u}(\mathcal{H}) \cap \mathcal{F}$$

satisfies

$$\|\exp(\rho_n)\|_{\text{res}} \geq \|(\sin |a_n|)\|_2 = \sqrt{n}.$$

Now the desired conclusion (6.1) follows since $\exp(\rho_n) \in (\mathbf{1} + \mathcal{F}) \cap \mathbf{U}(\mathcal{H})$ and $n \geq 1$ is arbitrary. \square

6.2. The predual of the restricted algebra is not spanned by its positive cone

It is well known that every self-adjoint normal functional in the predual of a W^* -algebra can be written as the difference of two positive normal functionals. It is also well known and easy to see that a similar property holds for the preduals of numerous operator ideals. More precisely, if \mathfrak{J} and \mathfrak{B} are Banach operator ideals such that the trace pairing

$$(\mathfrak{B}, \mathfrak{J}) \rightarrow \mathbb{C}, \quad (T, S) \mapsto \text{Tr}(TS)$$

is well defined and induces a topological isomorphism of the topological dual \mathfrak{B}^* onto \mathfrak{J} , then for every $T = T^* \in \mathfrak{B}$ there exist $T_1, T_2 \in \mathfrak{B}$ such that $T_1 \geq 0, T_2 \geq 0$ and $T = T_1 - T_2$. In fact, we can take $T_1 = (|T| + T)/2$ and $T_2 = (|T| - T)/2$, and we have $T_1, T_2 \in \mathfrak{B}$ since $|T| \in \mathfrak{B}$. (The latter property follows since if $T = W|T|$ is the polar decomposition of T , then $|T| = W^*T \in \mathfrak{B}$.)

We shall see in Proposition 6.4 below that the predual $(u_{\text{res}})_*$ of the restricted Lie algebra fails to have the similar property of being spanned by its elements ρ with $i\rho \geq 0$. In fact, the linear span of these elements turns out to be the proper subspace u_1 of $(u_{\text{res}})_*$.

Lemma 6.3. *Let \mathcal{H}_\pm be two complex separable Hilbert spaces, $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-, 0 \leq a_\pm \in \mathcal{B}(\mathcal{H}_\pm)$, and $t \in \mathcal{B}(\mathcal{H}_-, \mathcal{H}_+)$. Also denote*

$$a = \begin{pmatrix} a_+ & t \\ t^* & a_- \end{pmatrix} \in \mathcal{B}(\mathcal{H}).$$

Then the following assertions hold:

- (i) We have $a \geq 0$ if and only if the inequality

$$|\langle \xi, t\eta \rangle|^2 \leq \langle \xi, a_+\xi \rangle \cdot \langle \eta, a_-\eta \rangle \tag{6.2}$$

holds for all $\xi \in \mathcal{H}_+$ and $\eta \in \mathcal{H}_-$.

- (ii) If $a \geq 0$ and in addition $a_\pm \in \mathfrak{S}_1(\mathcal{H}_\pm)$ and $t \in \mathfrak{S}_2(\mathcal{H}_-, \mathcal{H}_+)$, then

$$\|t\|_2 \leq (\text{Tr} a) / \sqrt{2}. \tag{6.3}$$

Proof. For assertion (i) see Exercise 3.2 at the end of Chapter 3 in [32].

Next, let $\{\xi_i\}_{i \geq 1}$ and $\{\eta_j\}_{j \geq 1}$ be orthonormal bases in the Hilbert spaces \mathcal{H}_+ and \mathcal{H}_- , respectively. Then (6.2) shows that

$$(\forall i, j \geq 1) \quad |\langle \xi_i, t\eta_j \rangle|^2 \leq \langle \xi_i, a_+\xi_i \rangle \cdot \langle \eta_j, a_-\eta_j \rangle.$$

Now recall that $(\|t\|_2)^2 = \sum_{i,j \geq 1} |\langle \xi_i, t\eta_j \rangle|^2$, $\text{Tr} a_+ = \sum_{i \geq 1} \langle \xi_i, a_+\xi_i \rangle$, and $\text{Tr} a_- = \sum_{j \geq 1} \langle \eta_j, a_-\eta_j \rangle$. Thus, adding the above inequalities, we get

$$(\|t\|_2)^2 \leq (\text{Tr} a_+) \cdot (\text{Tr} a_-) \leq (\text{Tr} a_+ + \text{Tr} a_-)^2 / 2 = (\text{Tr} a)^2 / 2$$

and assertion (ii) follows. \square

Proposition 6.4. *The following assertions hold:*

- (i) *If $a \in (\mathfrak{u}_{\text{res}})_*$ and $ia \geq 0$, then $a \in \mathfrak{S}_1(\mathcal{H})$ and $\|a\|_1 \leq \|a\|_{(\mathfrak{u}_{\text{res}})_*} \leq (1 + \sqrt{2})\|a\|_1$.*
- (ii) *If $\rho \in (\mathfrak{u}_{\text{res}})_* \setminus \mathfrak{u}_1$ then there exist no $\rho_1, \rho_2 \in (\mathfrak{u}_{\text{res}})_*$ such that $i\rho_1 \geq 0$, $i\rho_2 \geq 0$, and $\rho = \rho_1 - \rho_2$.*

Proof. (i) Let $a \in (\mathfrak{u}_{\text{res}})_*$ such that $ia \geq 0$, and denote $ia =: \begin{pmatrix} a_+ & \\ & a_- \end{pmatrix}$. Then

$$\begin{aligned} \|a\|_1 &= \|ia\|_1 = \text{Tr}(ia) = \text{Tr} a_+ + \text{Tr} a_- = \|a_+\|_1 + \|a_-\|_1 \\ &\leq \|ia\|_{(\mathfrak{u}_{\text{res}})_*} = \|a_+\|_1 + \|a_-\|_1 + 2\|t\|_2 \\ &\leq \|a_+\|_1 + \|a_-\|_1 + \sqrt{2} \cdot \text{Tr}(ia) = (1 + \sqrt{2})\|ia\|_1 = (1 + \sqrt{2})\|a\|_1, \end{aligned}$$

where the second inequality follows by Lemma 6.2(ii). Consequently, for all $a \in (\mathfrak{u}_{\text{res}})_*$ with $ia \geq 0$ we have $\|a\|_1 \leq \|a\|_{(\mathfrak{u}_{\text{res}})_*} \leq (1 + \sqrt{2})\|a\|_1$.

(ii) Let $\rho \in (\mathfrak{u}_{\text{res}})_* \setminus \mathfrak{u}_1$ and assume that there exist elements $\rho_1, \rho_2 \in (\mathfrak{u}_{\text{res}})_*$ such that $i\rho_1 \geq 0$, $i\rho_2 \geq 0$, and $\rho = \rho_1 - \rho_2$. Then $i\rho_1, i\rho_2 \in \mathfrak{S}_1(\mathcal{H})$ according to the assertion (i), which we have already proved. Consequently, $\rho_1, \rho_2 \in \mathfrak{u}_1$, whence $\rho = \rho_1 - \rho_2 \in \mathfrak{u}_1$. This is a contradiction with the assumption on ρ , which concludes the proof. \square

6.3. The Cartan subalgebras of $\mathfrak{u}_{\text{res}}$ are not U_{res} -conjugate

For a (finite-dimensional) compact connected semi-simple Lie subgroup G of the unitary group $U(n)$, every element X of the Lie algebra \mathfrak{g} of G is conjugate to a diagonal element with respect to a given basis \mathcal{B} of \mathbb{C}^n by an element of G . This can be seen as follows (see [17, Chapter V, Theorem 6.4] for more general results). Take a diagonal element $H \in \mathfrak{g}$ with respect to \mathcal{B} such that the one-parameter subgroup $\exp tH$ is dense in the torus whose Lie algebra is the set of diagonal matrices belonging to \mathfrak{g} . On G , consider the continuous function $g \mapsto B(H, \text{Ad}(g)(X))$, where B denotes the Killing form of G . By compactness, this function takes a minimum at some g_0 , and for every element Y in \mathfrak{g} one has

$$\left. \frac{d}{dt} B(H, \text{Ad}(\exp tY) \text{Ad}(g_0)(X)) \right|_{t=0} = 0,$$

i.e. $B(H, [Y, \text{Ad}(g_0)(X)]) = 0$. Since the Killing form is $\text{Ad}(G)$ -invariant, one has

$$B(H, [Y, \text{Ad}(g_0)(X)]) = B([\text{Ad}(g_0)(X), H], Y).$$

The non-degeneracy of the Killing form then implies that $[\text{Ad}(g_0)(X), H] = 0$. But H has been chosen such that the centralizer of H is the set of diagonal matrices with respect to \mathcal{B} belonging to \mathfrak{g} . Consequently $\text{Ad}(g_0)(X)$ is a diagonal element in \mathfrak{g} . It follows that the maximal Abelian subalgebras, called *Cartan subalgebras*, of \mathfrak{g} are conjugate under G .

This proof cannot be extended to the infinite-dimensional case since the minimization argument above uses in a crucial manner the compactness of the group. We shall prove below that the conjugacy statement itself does not hold, in general. More precisely, we shall show that not all Cartan subalgebras of $(\mathfrak{u}_{\text{res}})_* \hookrightarrow \mathfrak{u}_{\text{res}}$ are U_{res} -conjugate.

We note that a related fact follows from results in the paper [8]. Specifically, let $\rho_0 \in (\mathfrak{u}_{\text{res}})_*$ such that $[d, \rho_0] = 0$, $\text{Ker } \rho_0 = \{0\}$, and each eigenvalue of ρ_0 has multiplicity 1. Next denote by \mathcal{O}_{ρ_0} the coadjoint U_{res} -orbit of ρ_0 , let $\rho \in (\mathfrak{u}_{\text{res}})_*$, and define

$$f_\rho : \mathcal{O}_{\rho_0} \rightarrow (0, \infty), \quad f_\rho(b) = \|\rho - b\|_2.$$

If the function f_ρ happens to have a critical point $\rho_1 \in \mathcal{O}_{\rho_0}$, then $[\rho_1, \rho] = 0$ according to [8]. Since $\rho_1 \in \mathcal{O}_{\rho_0}$, there exists $u \in U_{\text{res}}$ such that $\rho_1 = u\rho_0u^{-1}$, and then $[\rho_0, u^{-1}\rho u] = 0$. The latter equality implies that $u^{-1}\rho u$ commutes with all of the spectral projections of ρ_0 . Hence $[d, u^{-1}\rho u] = 0$ in view of the spectral assumptions on ρ_0 , and then Proposition 3.3 applied to $u^{-1}\rho u$ shows that the coadjoint isotropy group of ρ is a Banach Lie subgroup of U_{res} and the corresponding U_{res} -coadjoint orbit \mathcal{O}_ρ is a smooth leaf of the characteristic distribution of $(\mathfrak{u}_{\text{res}})_*$.

Proposition 6.5. *The unitary group U_{res} does not act transitively on the set of Cartan subalgebras of its Lie algebra.*

Proof. Endow the Hilbert space \mathcal{H} with an orthonormal basis $\mathcal{B} = \{e_n\}_{n \in \mathbb{Z} \setminus \{0\}}$, such that $\{e_{-n}\}_{n \in \mathbb{N} \setminus \{0\}}$ is an orthonormal basis of \mathcal{H}_+ and $\{e_n\}_{n \in \mathbb{N} \setminus \{0\}}$ an orthonormal basis of \mathcal{H}_- . The set \mathcal{D} of skew-Hermitian bounded diagonal operators with respect to \mathcal{B} form a Cartan subalgebra of $\mathfrak{u}_{\text{res}}$. Now consider the following subset of the set of anti-diagonal elements in $\mathfrak{u}_{\text{res}}$:

$$\mathcal{J} = \{J \in \mathfrak{u}_{\text{res}} \mid J(e_n) \in \mathbb{R}e_{-n} \ \forall n \in \mathbb{Z} \setminus \{0\}\}.$$

Since the coefficients $J_{-k,k}$, $k \in \mathbb{Z} \setminus \{0\}$, of $J \in \mathcal{J}$ satisfy $J_{-k,k} = -J_{k,-k}$, it follows from an easy computation that \mathcal{J} is Abelian. An element $B = (B_{i,j}) \in \mathfrak{u}_{\text{res}}$ commutes with every element $J = (J_{i,j})$ in \mathcal{J} if and only if

$$([B, J]_{i,-k}) = (B_{i,k}J_{k,-k} - J_{i,-i}B_{-i,-k}) \tag{6.4}$$

vanishes for every $J \in \mathcal{J}$. This implies the following conditions:

$$\begin{aligned} B_{i,k} &= 0 && \text{for } i \notin \{k, -k\}; \\ B_{k,k} &= B_{-k,-k} && \text{for } k \in \mathbb{Z} \setminus \{0\}; \\ B_{-k,k} &= -B_{k,-k} && \text{for } k \in \mathbb{Z} \setminus \{0\}. \end{aligned}$$

It follows that the maximal Abelian subalgebra \mathcal{C} of $\mathfrak{u}_{\text{res}}$ which contains \mathcal{J} is $\mathcal{J} + \mathcal{D}_+$, where

$$\mathcal{D}_+ = \{D = (D_{i,j}) \in \mathcal{D} \mid D_{-k,-k} = D_{k,k} \ \forall k \in \mathbb{Z} \setminus \{0\}\}.$$

Let us prove by contradiction that the Cartan subalgebras \mathcal{C} and \mathcal{D} are not conjugate under U_{res} . Suppose that there exists a unitary operator

$$g = \begin{pmatrix} g_{++} & g_{+-} \\ g_{-+} & g_{--} \end{pmatrix} \in U_{\text{res}}$$

such that $g\mathcal{J}g^{-1} \subseteq \mathcal{D}$. Consider an element

$$J = \begin{pmatrix} 0 & J_{+-} \\ J_{-+} & 0 \end{pmatrix} \in \mathcal{J}$$

which is a Hilbert–Schmidt operator that is not trace class. One has

$$\begin{aligned} gJg^{-1} &= \begin{pmatrix} g_{++} & g_{+-} \\ g_{-+} & g_{--} \end{pmatrix} \begin{pmatrix} 0 & J_{+-} \\ J_{-+} & 0 \end{pmatrix} \begin{pmatrix} g_{++}^* & g_{+-}^* \\ g_{-+}^* & g_{--}^* \end{pmatrix} \\ &= \begin{pmatrix} g_{+-}J_{-+}g_{++}^* + g_{++}J_{+-}g_{+-}^* & g_{+-}J_{-+}g_{-+}^* + g_{++}J_{+-}g_{--}^* \\ g_{--}J_{-+}g_{++}^* + g_{-+}J_{+-}g_{+-}^* & g_{--}J_{-+}g_{-+}^* + g_{-+}J_{+-}g_{--}^* \end{pmatrix}. \end{aligned}$$

By hypothesis, gJg^{-1} is a diagonal operator

$$D = \begin{pmatrix} D_{++} & 0 \\ 0 & D_{--} \end{pmatrix}$$

with $D_{++} = g_{+-}J_{-+}g_{++}^* + g_{++}J_{+-}g_{+-}^*$ and $D_{--} = g_{--}J_{-+}g_{-+}^* + g_{-+}J_{+-}g_{--}^*$. Now, since g belongs to U_{res} , g_{+-} and g_{-+} are Hilbert–Schmidt. Since J belongs to $\mathfrak{S}_2(\mathcal{H})$, J_{+-} and J_{-+} are Hilbert–Schmidt as well. From the relation $\mathfrak{S}_2 \cdot \mathfrak{S}_2 \subset \mathfrak{S}_1$, it follows that D_{++} and D_{--} are trace class, hence D belongs to $\mathfrak{S}_1(\mathcal{H})$. But this implies that $J = g^{-1}Dg$ is also trace class, since $\mathfrak{S}_1(\mathcal{H})$ is an ideal of $\mathcal{B}(\mathcal{H})$. This leads to a contradiction by the choice of $J \in \mathcal{J}$. It follows that elements in $\mathcal{J} \setminus \mathfrak{S}_1(\mathcal{H})$ are not U_{res} -conjugate to diagonal elements with respect to \mathcal{B} . Consequently, the Cartan subalgebra \mathcal{C} and \mathcal{D} are not U_{res} -conjugate. \square

Remark 6.6. The proof of Proposition 6.5 implies that the unitary group U_{res} does not act transitively on the set of Cartan subalgebras of $(u_{\text{res}})_*$. Since every compact skew-Hermitian operator admits an orthonormal basis of eigenvectors, the set of conjugacy classes of Cartan subalgebras in $(u_{\text{res}})_*$ is in bijection with $U(\mathcal{H})/U_{\text{res}}$ and is infinite. The conjugacy classes of Cartan subalgebras are related to the conjugacy classes of maximal tori. An infinite number of conjugacy classes of maximal tori has already been encountered in the case of some groups of contactomorphisms (see [20]). Examples of maximal tori of different dimensions were provided in [16] in some groups of symplectomorphisms.

Acknowledgments

The research of the first author was partially supported by Romanian grant CEx05-D11-23/2005 and that of the second by the Swiss National Science Foundation. We thank Alan Weinstein who motivated this investigation by asking us whether the Grassmannians studied in [37] falls within the framework of the theory developed in [7,28]. Our thanks go also to Tilmann Wurzbacher for several discussions that influenced our presentation. We are grateful to the anonymous referee for carefully reading the manuscript and for many valuable suggestions.

References

[1] V. Adamjan, H. Langer, C. Tretter, Existence and uniqueness of contractive solutions of some Riccati equations, *J. Funct. Anal.* 179 (2) (2001) 448–473.

- [2] S. Albeverio, K.A. Makarov, A.K. Motovilov, Graph subspaces and the spectral shift function, *Canad. J. Math.* 55 (3) (2003) 449–503.
- [3] A. Arvanitoyeorgos, *An Introduction to Lie Groups and the Geometry of Homogeneous Spaces*, Stud. Math. Libr., vol. 22, Amer. Math. Soc., Providence, RI, 2003.
- [4] A.G. Baskakov, Diagonalization of operators and complementability of subspaces of Banach spaces, *Ukrain. Mat. Zh.* 42 (7) (1990) 867–873, transl. in: *Ukrainian Math. J.* 42 (7) (1990) 763–768 (1991).
- [5] D. Beltiță, *Smooth Homogeneous Structures in Operator Theory*, Chapman & Hall/CRC Monogr. Surv. Pure Appl. Math., vol. 137, Chapman & Hall/CRC, New York, 2006.
- [6] D. Beltiță, B. Prunaru, Amenability, completely bounded projections, dynamical systems and smooth orbits, *Integral Equations Operator Theory* 57 (1) (2007) 1–17.
- [7] D. Beltiță, T.S. Ratiu, Symplectic leaves in real Banach Lie–Poisson spaces, *Geom. Funct. Anal.* 15 (4) (2005) 753–779.
- [8] R. Bhatia, P. Šemrl, Distance between Hermitian operators in Schatten classes, *Proc. Edinb. Math. Soc.* (2) 39 (2) (1996) 377–380.
- [9] N. Bourbaki, *Groupes et Algèbres de Lie*, Hermann, Paris, 1972.
- [10] A.L. Carey, Some homogeneous spaces and representations of the Hilbert Lie group $U(H)_2$, *Rev. Roumaine Math. Pures Appl.* 30 (7) (1985) 505–520.
- [11] J. Cheeger, D.G. Ebin, *Comparison Theorems in Riemannian Geometry*, North-Holland, Amsterdam, 1975.
- [12] J.B. Conway, B.B. Morrel, Operators that are points of spectral continuity, *Integral Equations Operator Theory* 2 (2) (1979) 174–198.
- [13] I. Ekeland, The Hopf–Rinow theorem in infinite dimension, *J. Differential Geom.* 13 (1978) 287–301.
- [14] P.R. Halmos, *A Hilbert Space Problem Book*, Van Nostrand, Princeton, NJ, 1967.
- [15] P. de la Harpe, *Classical Banach Lie Algebras and Banach Lie Groups of Operators in Hilbert Space*, Lecture Notes in Math., vol. 285, Springer-Verlag, Berlin, 1972.
- [16] J.-C. Hausmann, S. Tolman, Maximal Hamiltonian tori for polygon spaces, *Ann. Inst. Fourier (Grenoble)* 53 (6) (2003) 1925–1939.
- [17] S. Helgason, *Differential Geometry and Symmetric Spaces*, Academic Press, New York, 1962.
- [18] A. Hinkkanen, On the diagonalization of a certain class of operators, *Michigan Math. J.* 32 (3) (1985) 349–359.
- [19] S. Lang, *Fundamentals of Differential Geometry*, corrected second printing, Grad. Texts in Math., vol. 191, Springer-Verlag, New York, 2001.
- [20] E. Lerman, Contact cuts, *Israel J. Math.* 124 (2001) 77–92.
- [21] A.K. Motovilov, Removal of the resolvent-like energy dependence from interactions and invariant subspaces of a total Hamiltonian, *J. Math. Phys.* 36 (12) (1995) 6647–6664.
- [22] K.-H. Neeb, Highest weight representations and infinite-dimensional Kähler manifolds, in: *Recent Advances in Lie Theory*, Vigo, 2000, in: *Res. Exp. Math.*, vol. 25, Heldermann, Lemgo, 2002, pp. 367–392.
- [23] K.-H. Neeb, Classical Hilbert–Lie groups, their extensions and their homotopy groups, in: *Geometry and Analysis on Finite- and Infinite-Dimensional Lie Groups*, Będlewo, 2000, in: *Banach Center Publ.*, vol. 55, Polish Acad. Sci., Warsaw, 2002, pp. 87–151.
- [24] K.-H. Neeb, Universal central extensions of Lie groups, *Acta Appl. Math.* 73 (1–2) (2002) 175–219.
- [25] K.-H. Neeb, A Cartan–Hadamard theorem for Banach–Finsler manifolds, *Geom. Dedicata* 95 (2002) 115–156.
- [26] J.D. Newburgh, The variation of spectra, *Duke Math. J.* 18 (1951) 165–176.
- [27] F. Nübel, On integral manifolds for vector space distributions, *Math. Ann.* 294 (1) (1992) 1–17.
- [28] A. Odziejewicz, T.S. Ratiu, Banach Lie–Poisson spaces and reduction, *Comm. Math. Phys.* 243 (1) (2003) 1–54.
- [29] A. Odziejewicz, T.S. Ratiu, Extensions of Banach Lie–Poisson spaces, *J. Funct. Anal.* 217 (1) (2004) 103–125.
- [30] B. O’Neill, *Semi-Riemannian Geometry with Applications to Relativity*, Academic Press, New York, 1983.
- [31] J.-P. Ortega, T.S. Ratiu, *Momentum Maps and Hamiltonian Reduction*, Progr. Math., vol. 222, Birkhäuser Boston, Boston, MA, 2004.
- [32] V. Paulsen, *Completely Bounded Maps and Operator Algebras*, Cambridge Stud. Adv. Math., vol. 78, Cambridge Univ. Press, Cambridge, 2002.
- [33] A. Pressley, G. Segal, *Loop Groups*, Oxford Math. Monogr., Clarendon/Oxford Univ. Press, Oxford, 1990.
- [34] G. Segal, Unitary representations of some infinite dimensional groups, *Comm. Math. Phys.* 80 (3) (1981) 301–342.
- [35] G. Segal, G. Wilson, Loop groups and equations of KdV type, *Publ. Math. Inst. Hautes Études Sci.* 61 (1985) 5–6.
- [36] I. Stewart, *Lie Algebras Generated by Finite-Dimensional Ideals*, Res. Notes in Math., vol. 2, Pitman, London, 1975.
- [37] A.B. Tumpach, Variétés kählériennes et hyperkähleriennes de dimension infinie, PhD thesis, École Polytechnique, Paris, 2005.

- [38] A.B. Tumpach, Mostow Decomposition Theorem for a L^* -group and applications to affine coadjoint orbits and stable manifolds, preprint, math-ph/0605039, May 2006.
- [39] T. Wurzbacher, Fermionic second quantization and the geometry of the restricted Grassmannian, in: Infinite Dimensional Kähler Manifolds, Oberwolfach, 1995, in: DMV Sem., vol. 31, Birkhäuser, Basel, 2001, pp. 287–375.