

QUADRATIC FORMS OVER COMPLETE LOCAL RINGS

MANUEL OJANGUREN AND RAMAN PARIMALA

1. INTRODUCTION

Let A be a complete excellent local domain of Krull dimension 2 and K its field of fractions. We further assume that 2 is invertible in A and that the residue field of A is algebraically closed. We first show that the unramified Brauer group of K (with respect to all discrete valuations of K) vanishes. Using this result we prove that every rank 4 quadratic form which is isotropic in all completions of K with respect to discrete valuations, is isotropic. For $K = \mathbb{C}((x, y))$ this was announced by P. Jaworski in 1999, at a conference on quadratic forms in Oberwolfach. The results presented here, obtained immediately after the conference, are a consequence of our efforts to give a short proof of Jaworski's announcement. They subsequently led to various generalizations, notably in the case when the residue field of A is real closed (see [1] and, for further developments, [2] and [3]). Jaworski's proof has now appeared as well [6].

2. THE UNRAMIFIED BRAUER GROUP

Let A be as above and m its maximal ideal. Let $\pi : X \rightarrow \text{Spec}(A)$ be a desingularization [8] of $\text{Spec}(A)$. Since X is obtained from $\text{Spec}(A)$ by a sequence of blowing ups and normalizations, the map π is proper. We denote by X_n the fibre of $\text{Spec}(A/m^{n+1})$.

We prove a result (Lemma 2.2) in the spirit of Lemma 3.3 of [5].

Lemma 2.1. *The natural maps*

$$\text{Pic}(X_n) \rightarrow \text{Pic}(X_{n+1})$$

are surjective.

Proof. This follows from the exact sequence of sheaves

$$0 \longrightarrow \frac{m^n \mathcal{O}_X}{m^{n+1} \mathcal{O}_X} \longrightarrow \left(\frac{\mathcal{O}_X}{m^{n+1} \mathcal{O}_X} \right)^* \longrightarrow \left(\frac{\mathcal{O}_X}{m^n \mathcal{O}_X} \right)^* \longrightarrow 1,$$

noting that

$$H^2\left(X, \frac{m^n \mathcal{O}_X}{m^{n+1} \mathcal{O}_X}\right) = H^2\left(X_0, \frac{m^n \mathcal{O}_X}{m^{n+1} \mathcal{O}_X}\right) = 0$$

because X_0 is of dimension 1.

Lemma 2.2. *The canonical homomorphism*

$$\text{Br}(X) \rightarrow \varprojlim \text{Br}(X_n)$$

is injective.

Proof. Let \mathcal{A} be an Azumaya algebra over X . Denote by \mathcal{A}_n the algebra obtained from \mathcal{A} under base change from X to X_n and suppose that it is trivial for each n . Let

$$u_n : \mathcal{A}_n \xrightarrow{\sim} \mathcal{E}nd(V_n)$$

be an isomorphism, where V_n is a locally free sheaf on X_n . The sheaf V_n is determined by \mathcal{A}_n up to a line bundle. By Lemma 2.1 we can successively modify each V_n in such a way that V_n is isomorphic to $V_{n+1} \otimes_{\mathcal{O}_{X_{n+1}}} \mathcal{O}_{X_n}$. In this case the isomorphisms u_n also form a projective system. By [5], 5.1.4, the projective system $(V_n, n \in \mathbb{N})$ gives a locally free \mathcal{O}_X -module V and an isomorphism

$$u : \mathcal{A} \xrightarrow{\sim} \mathcal{E}nd(V).$$

Corollary 2.3. *The Brauer group of X is trivial. In particular, the unramified Brauer group of K is trivial.*

Proof. In fact, since X_n is a curve over $\text{Spec}(A/m^{n+1})$ and A/m is algebraically closed, $Br(X_n) = 0$ (See [5], page 101). By a well-known purity theorem ([4], Proposition 2.3) an unramified element of $Br(K)$ is in the image of $Br(X) \rightarrow Br(K)$ and hence is zero.

3. QUADRATIC FORMS

Theorem. *Let A be a complete excellent local domain of Krull dimension 2 and K its field of fractions. Assume that 2 is invertible in A and that the residue field of A is algebraically closed. Every rank 4 quadratic form q over K which is isotropic over every completion of K at a discrete valuation, is isotropic.*

Proof. After scaling we may assume that $q = \langle 1, a, b, abd \rangle$ with $a, b, d \in K^*$. If d is a square, then q is the norm form of the quaternion algebra $\mathcal{A} = \left(\frac{a,b}{K}\right)$. The condition that q is isotropic at all completions implies that \mathcal{A} is split at all completions of K . In particular \mathcal{A} is unramified in $Br(K)$ and hence, by 2.3, is zero. In particular, q is hyperbolic.

Suppose now that d is not a square. Let $L = K(\sqrt{d})$. The field L satisfies the same assumptions as K . The form q_L over L has trivial discriminant and is isotropic at all completions of L at discrete valuations. By the previous case, q_L is hyperbolic. The form q therefore contains a multiple of $\langle 1, -d \rangle$ ([7], Ch. 7, Lemma 3.1) and, being of discriminant d , also contains a subform of discriminant 1. Hence q is isotropic.

REFERENCES

1. J.-L. Colliot-Thélène, M. Ojanguren and R. Parimala, *Quadratic forms over fraction fields of two-dimensional henselian rings and Brauer groups of related schemes*, Proceedings of the International Colloquium on ‘Algebra, Arithmetic & Geometry’ held at T.I.F.R., Mumbai, 2000.
2. J.-L. Colliot-Thélène, Ph. Gille et R. Parimala, *Arithmétique des groupes algébriques linéaires sur certains corps géométriques de dimension deux*, C. R. Acad. Sci. Paris **333**, *Série I* (2001), 827–832.
3. J.-L. Colliot-Thélène, Ph. Gille and R. Parimala, *Arithmetic of linear algebraic groups over two-dimensional geometric fields*, preprint (2002).
4. A. Grothendieck, *Le groupe de Brauer II*, Dix exposés sur la cohomologie des schémas, North Holland, 1968.
5. A. Grothendieck, *Le groupe de Brauer III*, Dix exposés sur la cohomologie des schémas, North Holland, 1968.
6. P. Jaworski, *On the strong Hasse principle for fields of quotients of power series rings in two variables.*, Math. Z. **236** (2001), 531–566.
7. T.Y. Lam, *Algebraic theory of quadratic forms*, Benjamin, 1973.
8. J. Lipman, *Rational singularities with applications to algebraic surfaces and unique factorization*, Publ. Math. IHES **36** (1969), 195–280.

MANUEL OJANGUREN, INSTITUT DE MATHÉMATIQUES, UNIVERSITY OF LAUSANNE, CH-1015 LAUSANNE, SWITZERLAND

RAMAN PARIMALA, SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, MUMBAI, INDIA