

GLOBAL DYNAMICS OF THE NONRADIAL ENERGY-CRITICAL WAVE EQUATION ABOVE THE GROUND STATE ENERGY

J. KRIEGER, K. NAKANISHI, AND W. SCHLAG

ABSTRACT. In this paper we establish the existence of certain classes of solutions to the energy critical nonlinear wave equation in dimensions 3 and 5 assuming that the energy exceeds the ground state energy only by a small amount. No radial assumption is made. We find that there exist four sets in $\dot{H}^1 \times L^2$ with nonempty interiors which correspond to all possible combinations of finite-time blowup on the one hand, and global existence and scattering to a free wave, on the other hand, as $t \rightarrow \pm\infty$.

CONTENTS

1. Introduction	1
2. The basic setup	6
2.1. The critical wave equation, Hamiltonian formalism	6
2.2. The translation and scaling symmetries	6
2.3. A change of time and the static linearized operator	7
2.4. Energy expansion	8
2.5. Orthogonality conditions near the ground state	9
2.6. Linearized energy	10
2.7. Modulation equations	10
2.8. Hyperbolic drivers	11
3. Distance function, λ dominance, ejection	11
4. The variational structure in the energy critical setting	14
5. The one-pass theorem	16
6. Blow-up after ejection	20
7. Scattering after ejection	23
References	31

Support of the National Science Foundation, DMS-0757278 (JK) and DMS-0617854 (WS) is gratefully acknowledged. The authors thank Thomas Duyckaerts and Carlos Kenig for the interest in this work, as well as for helpful conversations.

1. INTRODUCTION

Consider the energy-critical nonlinear wave equation with real-valued u

$$\ddot{u} - \Delta u = |u|^{p-1}u, \quad u(t, x) : \mathbb{R}^{1+d} \rightarrow \mathbb{R}, \quad p = \frac{d+2}{d-2} = 2^* - 1, \quad d = 3 \text{ or } 5, \quad (1.1)$$

in the energy space

$$\vec{u}(t) := (u(t), \dot{u}(t)) \in \mathcal{H} := \dot{H}^1(\mathbb{R}^d) \times L^2(\mathbb{R}^d), \quad (1.2)$$

which is the real Hilbert space with the inner product

$$\langle \underline{u}, \underline{v} \rangle_{\mathcal{H}} := \langle \nabla u_1 | \nabla v_1 \rangle + \langle u_2 | v_2 \rangle, \quad \langle f | g \rangle := \int_{\mathbb{R}^d} f(x) \cdot g(x) dx. \quad (1.3)$$

Henceforth $\vec{u} = (u, \dot{u})$ denotes the vector derived from a time function $u(t)$, while a general vector is denoted like $\underline{u} = (u_1, u_2)$. The seminorm on any domain $\Omega \subset \mathbb{R}^d$ is defined by

$$\|\underline{u}\|_{\mathcal{H}(\Omega)}^2 := \|\nabla u_1\|_{L^2(\Omega)}^2 + \|u_2\|_{L^2(\Omega)}^2. \quad (1.4)$$

We remark that the dimensional restriction $d = 3$ or 5 is needed only for using the blow-up characterization by Duyckaerts-Kenig-Merle [4].

Equation (1.1) is locally well-posed for $\vec{u}(0) \in \mathcal{H}$, globally for small data, and may blow up in finite time (for example, for data of negative energy). Moreover, $I = [0, T_*)$ is a finite maximal time of existence if and only if

$$\|u\|_{L_t^q(I; L_x^q(\mathbb{R}^d))} = \infty, \quad q := \frac{2(d+1)}{d-2}. \quad (1.5)$$

For a comprehensive review of these basic issues we refer the reader to [11].

In a previous paper [12] the authors studied the global dynamics of radial solutions to (1.1). To state that result as well as the main result of this paper, we recall some of the basic structures associated with the critical equation. First, one has the conserved energy of (1.1)

$$E(\vec{u}) := \int_{\mathbb{R}^d} \left[\frac{|\dot{u}|^2 + |\nabla u|^2}{2} - \frac{|u|^{2^*}}{2^*} \right] dx \quad (1.6)$$

as well as the conserved momentum

$$P(\vec{u}) := \langle \dot{u} | \nabla u \rangle. \quad (1.7)$$

Remarkably, (1.1) admits the static Aubin solutions of the form

$$W_\sigma = S_{-1}^\sigma W, \quad W(x) = \left[1 + \frac{|x|^2}{d(d-2)} \right]^{1-\frac{d}{2}}, \quad (1.8)$$

where S_{-1}^σ denotes the \dot{H}^1 preserving dilation

$$(S_{-1}^\sigma \varphi)(x) = e^{(d/2-1)\sigma} \varphi(e^\sigma x). \quad (1.9)$$

These are positive radial solutions of the static equation

$$-\Delta W - |W|^{2^*-2}W = 0, \quad (1.10)$$

which are unique, up to dilation and translation symmetries, amongst the non-negative, non-zero (not necessarily radial) C^2 solutions, see [2]. They also minimize the static energy

$$J(\varphi) := \int_{\mathbb{R}^d} \left[\frac{1}{2} |\nabla \varphi|^2 - \frac{1}{2^*} |\varphi|^{2^*} \right] dx, \quad (1.11)$$

among all non-trivial static solutions. The work of Kenig, Merle [10, 11] and Duyckaerts, Merle [6, 7] allows for a characterization of the global-in-time behavior of solutions with $E(\vec{u}) \leq J(W)$.

In this paper we study the behavior of solutions with

$$E(\vec{u}) < \sqrt{J(W)^2 + \varepsilon^4 + |P(\vec{u})|^2}, \quad (1.12)$$

for some small $\varepsilon > 0$. Solutions of subcritical focusing NLKG and NLS equations with radial data in \mathbb{R}^3 of energy slightly above that of the ground state were studied by the latter two authors in [18, 19]. The nonradial subcritical Klein-Gordon equation in three dimensions was treated in [20]. The key feature of (1.1) by contrast to NLKG is the scaling invariance of (1.1) manifested by

$$u(t, x) \mapsto e^{\sigma(d/2-1)t} u(e^{\sigma t}, e^{\sigma x}) = S_{-1}^{\sigma} u(e^{\sigma t}), \quad (1.13)$$

which leaves the energy unchanged. In particular, the analogue of the ‘‘one pass theorem’’ proved in [21] needs to be modified, specifically by replacing the discrete set of attractors $\{Q, -Q\}$ there by a $(2d+1)$ -parameter family of solitons. For any $(\sigma, p, q) \in \mathbb{R} \times \mathbb{R}^d \times \mathbb{R}^d$, denote the scaling-Lorentz transform of W by

$$W_{\sigma}(p, q) = W_{\sigma}(x - q + p(\langle p \rangle - 1)|p|^{-2}p \cdot (x - q)) \quad (1.14)$$

where $\langle p \rangle := \sqrt{1 + |p|^2}$. Then for any fixed $(p, q) \in \mathbb{R}^{2d}$,

$$u(t, x) = W_{\sigma}(p, q + tp/\langle p \rangle) \quad (1.15)$$

gives a ground state soliton of (1.1). Hence the ground state soliton family is

$$\mathcal{S} := \{(W_{\sigma}(p, q), -\nabla W_{\sigma}(p, q) \cdot p/\langle p \rangle) \mid (\sigma, p, q) \in \mathbb{R}^{1+2d}\} \subset \mathcal{H}. \quad (1.16)$$

Note that in the subcritical NLS case [20], the scaling parameter σ is essentially fixed or at least bounded from above and below by the L^2 conservation law, but in the critical case there is no factor which a priori prevents the scale from going to 0 or $+\infty$. On the other hand, by using the Lorentz transform, we can reduce the problem to the case of zero momentum, where the soliton family is

$$\mathcal{S}_0 := \{\vec{W}_{\sigma}(x - q) \mid (\sigma, q) \in \mathbb{R}^{1+d}\} \subset \mathcal{H}, \quad \vec{W}_{\sigma} = (W_{\sigma}, 0), \quad (1.17)$$

with the energy constraint $E(\vec{u}) < J(W) + \varepsilon^2$ (slightly changing $\varepsilon > 0$).

Introduce the “virial functional”

$$K(\varphi) := \int_{\mathbb{R}^d} [|\nabla\varphi|^2 - |\varphi|^{2^*}] dx \quad (1.18)$$

and note that $K(W) = 0$. The following positivity is crucial for the variational structure around W

$$\|\nabla\varphi\|_2^2/d = J(\varphi) - K(\varphi)/2^*. \quad (1.19)$$

Note that the derivative of $J(\varphi)$ with respect to any scaling

$$\varphi(x) \mapsto e^{a\sigma}\varphi(e^{b\sigma}x)$$

except for S_{-1}^σ gives a non-zero constant multiple of $K(\varphi)$. This is a special feature of the scaling critical case, which allows us to work with a single K , whereas in the subcritical case [21] we needed two different functionals and their equivalence.

The main result of this paper is summarized as follows.

Theorem 1.1. *There exist a small $\varepsilon > 0$, a neighborhood \mathcal{B} of $\pm\mathcal{S}_0$ within $O(\varepsilon)$ distance in \mathcal{H} , and a continuous functional $\mathfrak{S} : \mathcal{H}^\varepsilon \setminus \mathcal{B} \rightarrow \{\pm 1\}$, where*

$$\mathcal{H}^\varepsilon := \{\varphi \in \mathcal{H} \mid E(\varphi) < J(W) + \varepsilon^2\}, \quad (1.20)$$

such that the following properties hold: For any solution u in \mathcal{H}^ε on the maximal existence interval $I(u)$, let

$$\begin{aligned} I_0(u) &:= \{t \in I(u) \mid \vec{u}(t) \in \mathcal{B}\}, \\ I_\pm(u) &:= \{t \in I(u) \mid \vec{u}(t) \notin \mathcal{B}, \mathfrak{S}(\vec{u}(t)) = \pm 1\}. \end{aligned} \quad (1.21)$$

Then $I_0(u)$ is an interval, $I_+(u)$ consists of at most two infinite intervals, and $I_-(u)$ consists of at most two finite intervals. $u(t)$ scatters to 0 as $t \rightarrow \pm\infty$ if and only if $\pm t \in I_+(u)$ for large $t > 0$. Moreover, there is a uniform bound $M < \infty$ such that

$$\|u\|_{L_{t,x}^q(I_+(u) \times \mathbb{R}^d)} \leq M, \quad q := \frac{2(d+1)}{d-2}. \quad (1.22)$$

For each $\sigma_1, \sigma_2 \in \{\pm\}$, let A_{σ_1, σ_2} be the collection of initial data $\vec{u}(0) \in \mathcal{H}^\varepsilon$, and for some $T_- < 0 < T_+$,

$$(-\infty, T_-) \cap I(u) \subset I_{\sigma_1}(u), \quad (T_+, \infty) \cap I(u) \subset I_{\sigma_2}(u). \quad (1.23)$$

Then each of the four sets $A_{\pm, \pm}$ has non-empty interior, exhibiting all possible combinations of scattering to zero/finite time blowup as $t \rightarrow \pm\infty$, respectively.

The radial version of this exact theorem was proved in [12]. The main difference from that paper is of course the presence of the translation and Lorentz symmetries which need to be taken into account. Actually, the Lorentz symmetry does not play much role under the energy constraint $E(\vec{u}) < J(W) + \varepsilon^2$, where the solution can approach to $\pm\mathcal{S}$ only if $|P(\vec{u})| \lesssim$

ε . In contrast, the translational freedom is not a priori controlled by conserved quantities, and so we instead eliminate it by suitable orthogonality conditions. In other words, the modulation theory here amounts to a system of $d + 1$ ODEs corresponding to the dilation and translation symmetries.

By using the Lorentz transform, we can extend the above result to bigger energy, depending on the size of momentum.

Corollary 1.2. *There exist a small $\varepsilon > 0$, a neighborhood \mathcal{B} of $\pm\mathcal{S}$ in \mathcal{H} , and a continuous functional $\mathfrak{S} : \tilde{\mathcal{H}}^\varepsilon \setminus \tilde{\mathcal{B}} \rightarrow \{\pm 1\}$, where*

$$\tilde{\mathcal{H}}^\varepsilon := \left\{ \underline{\varphi} \in \mathcal{H} \mid E(\underline{\varphi}) < \sqrt{J(W)^2 + \varepsilon^4 + |P(\underline{\varphi})|^2} \right\}, \quad (1.24)$$

such that the same conclusion holds as in Theorem 1.1 if we replace \mathcal{H}^ε and \mathcal{B} by $\tilde{\mathcal{H}}^\varepsilon$ and $\tilde{\mathcal{B}}$ respectively. Moreover, if $|E(\vec{u})| \leq |P(\vec{u})|$ and $u \not\equiv 0$ then the solution blows up both in $t < 0$ and in $t > 0$.

Actually, we can reduce the corollary to the theorem only for $|E(\vec{u})| > |P(\vec{u})|$, where we can find a Lorentz transform from u to another solution w with $E(\vec{w}) = \sqrt{E(\vec{u})^2 - |P(\vec{u})|^2}$ and $P(w) = 0$, see (7.30).

The other case $|E(\vec{u})| \leq |P(\vec{u})|$ is treated separately, which is essentially known. Indeed, for such a solution u , there is a Lorentz transform to another solution w with $E(w) < J(W)$. If the original solution u is global in one direction, then so is w (see Lemma 7.3). Then we have $K(w(0)) \geq 0$, otherwise the classical result of Payne-Sattinger [22] (or more precisely by Kenig-Merle [11] in the current setting) implies that w blows up in both directions. Then

$$0 \leq \|\dot{w}\|_2^2/2 + \|\nabla w\|_2^2/d = E(\vec{w}) - K(w)/2^* \leq E(\vec{w}). \quad (1.25)$$

This is already a contradiction if $|E(\vec{u})| < |P(\vec{u})|$, since then we can make $E(w) < 0$. In the remaining case $|E(\vec{u})| = |P(\vec{u})|$, we can make $E(\vec{w})$ as small as we wish. Hence the above inequality implies that the energy norm can be made arbitrarily small. Then the small data scattering implies that $\|w\|_{L_{t,x}^q(\mathbb{R}^{1+d})} \lesssim \sqrt{E(\vec{w})}$. However, since $\|u\|_{L_{t,x}^q(\mathbb{R}^{1+d})}$ is Lorentz invariant, this implies that the original solution $u \equiv 0$.

The rest of the paper is devoted to prove the main theorem. We differ strongly from [12] in terms of the basic formalism which defines our approach. To be more precise, we perform a change of coordinates in the time variable which allows us to work with a fixed reference Hamiltonian in the perturbative analysis rather than a moving one as in [11]. This leads to some simplifications in the ejection lemma, for example, see Lemma 3.2. We remark that the formalism is also different from the one used in the nonradial subcritical equation [20], where a complex formulation was chosen, and more essentially, in the choice of orthogonality conditions, which also brings some simplification.

One application of Theorem 1.1 is the following corollary, which removes the radial assumption from [5, Corollary 6.3]. The solution W_+ is the one

discovered by Duyckaerts, Merle [7]. It is a radial $\dot{H}^1 \times L^2$ solution, exists globally in forward time and approaches W in \dot{H}^1 , and blows up in finite negative time. As above, the dimension satisfies $d = 3$ or $d = 5$.

Corollary 1.3. *Let u be an energy solution of (1.1) such that $E(\vec{u}) = E(W, 0)$. Denote by (u_0, u_1) the initial conditions of u . Assume that*

$$\int |\nabla u_0|^2 dx > \int |\nabla W|^2 dx$$

Then u blows up in finite time in both time directions or $u = W_+$ up to the symmetries of the equation.

In [7, Theorem 2, (c)] this result is proved (nonradially) under the additional condition that $u_0 \in L^2$. Using [12], this L^2 condition was removed in [5], but only in the radial setting. As noted in [5, Remark 6.5], the removal of the radial assumption in [12] would then complete [7] in the sense that the L^2 -condition can be removed even nonradially. This is what we accomplish in this paper, whence Corollary 1.3. For the proof, we refer the reader to [5].

2. THE BASIC SETUP

2.1. The critical wave equation, Hamiltonian formalism. The Cauchy problem for $\vec{u} = (u, \dot{u})$

$$\vec{u}_t = J\mathcal{D}\vec{u} + (0, |u_1|^{p-1}u_1), \quad J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \mathcal{D} = \begin{pmatrix} -\Delta & 0 \\ 0 & 1 \end{pmatrix} \quad (2.1)$$

is locally wellposed in \mathcal{H} , with the conservation of energy:

$$E(\vec{u}) := \frac{1}{2} \|\vec{u}(t)\|_{\mathcal{H}}^2 - \frac{1}{2^*} \|u(t)\|_{2^*}^{2^*}. \quad (2.2)$$

The equation is the Hamiltonian flow in \mathcal{H} with conserved Hamiltonian E relative to the symplectic form

$$\omega(\underline{u}, \underline{v}) := \langle J\underline{u}, \underline{v} \rangle_{L^2} = \langle u_2 | v_1 \rangle - \langle u_1 | v_2 \rangle. \quad (2.3)$$

2.2. The translation and scaling symmetries. Another feature of equation (1.1) is its invariance with respect to the scaling:

$$\vec{S}^\sigma := S_{-1}^\sigma \otimes S_0^\sigma, \quad S_a^\sigma \varphi(x) := e^{(d/2+a)\sigma} \varphi(e^\sigma x), \quad (2.4)$$

which is a unitary group acting on \mathcal{H} , with the generator

$$\vec{\Lambda} := \vec{S}' = \Lambda_{-1} \otimes \Lambda_0, \quad \Lambda_a := r\partial_r + d/2 + a = S_a'. \quad (2.5)$$

With Λ_a^* denoting the adjoint relative to $L^2(\mathbb{R}^d)$ one has $\Lambda_a^* = -\Lambda_{-a}$ and thus $\vec{\Lambda}^* = -\Lambda_1 \otimes \Lambda_0$. Similarly, the unitary group of translations is denoted by

$$(T^c v)(x) := v(x - c), \quad T' = -\nabla, \quad c \in \mathbb{R}^d. \quad (2.6)$$

Our analysis in this paper is around the static Aubin solution

$$W(x) = \left(1 + \frac{|x|^2}{d(d-2)}\right)^{1-\frac{d}{2}}, \quad -\Delta W = W^p, \quad (2.7)$$

whose vector and scaled versions are denoted by

$$\vec{W} := (W, 0), \quad \vec{W}_\sigma := \vec{S}^\sigma \vec{W} := (W_\sigma, 0), \quad \vec{\Lambda} \vec{W} := (W', 0). \quad (2.8)$$

Let $\vec{u} = \underline{u} = T^c \vec{S}^\sigma (\vec{W} + \underline{v})$ be a solution with $\sigma = \sigma(t)$ and $c = c(t)$. In general, \underline{v} need not have the structure (1.2), which is why we do not write \vec{v} . Noting that

$$\begin{aligned} \nabla S_a^\sigma &= S_a^\sigma e^\sigma \nabla, \quad \nabla S_a' = (S_a' + 1) \nabla, \\ (T^c \underline{u})_t &= T^c (\underline{u}_t - \dot{c} \nabla \underline{u}), \quad (\vec{S}^\sigma \underline{u})_t = \vec{S}^\sigma (\underline{u}_t + \dot{\sigma} \vec{\Lambda} \underline{u}), \end{aligned} \quad (2.9)$$

where $\dot{c} = c_t$, $\dot{\sigma} = \sigma_t$, we obtain the equation of \underline{v} :

$$\underline{v}_t = e^\sigma [J\mathcal{L}\underline{v} + \underline{N}(\underline{v})] + (e^\sigma \dot{c} \cdot \nabla - \dot{\sigma} \vec{\Lambda})(\vec{W} + \underline{v}) \quad (2.10)$$

where the linearized and superlinear operators are defined by

$$\begin{aligned} \mathcal{L} &= \begin{pmatrix} L_+ & 0 \\ 0 & 1 \end{pmatrix}, \quad L_+ = -\Delta - pW^{p-1}, \\ \underline{N}(\underline{v}) &= (0, N(v_1)), \quad N(v) = (W + v)^p - W^p - pW^{p-1}v. \end{aligned} \quad (2.11)$$

The structure of the spectrum of L_+ over $L^2(\mathbb{R}^d)$ is as follows: the discrete spectrum consists of a unique negative eigenvalue of L_+ which we denote by $-k^2$. The associated eigenfunction is the ground state of L_+ , denoted by ρ :

$$L_+ \rho = -k^2 \rho, \quad \rho > 0, \quad \|\rho\|_2 = 1. \quad (2.12)$$

The essential spectrum of L_+ is $[0, \infty)$, and it is purely absolutely continuous. At the threshold 0, one has an eigenvalue of multiplicity d , with eigenfunctions ∇W , and a resonance function $W' = \Lambda_{-1} W$ which is unique.

2.3. A change of time and the static linearized operator. The time-dependent coefficient on the linearized operator is removed by the standard change of time variable from t to τ :

$$\frac{d\tau}{dt} = e^{\sigma(t)}, \quad \underline{v}_\tau = J\mathcal{L}\underline{v} + \underline{N}(\underline{v}) + (e^\sigma c_\tau \cdot \nabla - \sigma_\tau \vec{\Lambda})(\vec{W} + \underline{v}). \quad (2.13)$$

The ‘‘generalized’’ eigenvectors of $J\mathcal{L}$ are

$$\begin{aligned} J\mathcal{L}\nabla \vec{W} &= 0, \quad J\mathcal{L}J\nabla \vec{W} = -\nabla \vec{W}, \quad J\mathcal{L}\vec{\Lambda} \vec{W} = 0, \quad J\mathcal{L}J\vec{\Lambda} \vec{W} = -\vec{\Lambda} \vec{W}, \\ J\mathcal{L}\underline{g}^\pm &= \pm k \underline{g}^\pm, \quad \underline{g}^\pm := (1, \pm k)\rho / \sqrt{2k}, \end{aligned} \quad (2.14)$$

where ρ is the aforementioned ground state of L_+ . The normalization here is such that $\omega(\underline{g}^+, \underline{g}^-) = 1$. Define $\vec{\rho} := (\rho, 0)$. Note that $\vec{\Lambda} \vec{W} \notin L^2$ for $d < 5$. Hence we decompose

$$\begin{aligned} \underline{v} &= \lambda_+ \underline{g}^+ + \lambda_- \underline{g}^- + \mu \cdot \nabla \vec{W} + \underline{\gamma} \\ \lambda_{\pm} &:= \omega(\underline{v}, \pm \underline{g}^{\mp}), \quad \mu := \omega(\underline{v}, J \nabla \vec{\rho}) / a_W = \langle v_1 | \nabla \rho \rangle / a_W, \end{aligned} \quad (2.15)$$

where

$$a_W := \frac{1}{d} \langle -\Delta W | \rho \rangle = \frac{1}{d} \langle W^{2^*-1} | \rho \rangle. \quad (2.16)$$

By construction,

$$\omega(\underline{\gamma}, \underline{g}^{\mp}) = 0, \quad \omega(\underline{\gamma}, J \nabla \vec{\rho}) = 0 \quad (2.17)$$

The more natural $\mu := \omega(\underline{v}, J \nabla \vec{W}) = \langle v_1 | \nabla W \rangle$ is problematic since the latter inner product is not well-defined.

Note that we did not extract the remaining root-mode $J \nabla \vec{W}$ from $\underline{\gamma}$, which corresponds to Lorentz “boosts”, i.e., translations in momentum.

2.4. Energy expansion. Using (2.15) the energy is expanded as

$$E(\vec{u}) - E(\vec{W}) = \frac{1}{2} \langle \mathcal{L} \underline{v} | \underline{v} \rangle - C(\underline{v}) = -k \lambda_+ \lambda_- + \frac{1}{2} \langle \mathcal{L} \underline{\gamma} | \underline{\gamma} \rangle - C(\underline{v}), \quad (2.18)$$

where the superquadratic part is given by

$$C(\underline{v}) := \int_{\mathbb{R}^d} \left[\frac{|W + v_1|^{2^*} - W^{2^*}}{2^*} - W^p v_1 - \frac{p}{2} W^{p-1} |v_1|^2 \right] (x) dx. \quad (2.19)$$

One has the estimate

$$|C(\underline{v})| \lesssim \|W^{2^*-3} v_1^3\|_1 + \|v_1\|_{2^*}^2 \quad (2.20)$$

since $2^* > 3$.

Lemma 2.1. *Let $f \in \dot{H}^1(\mathbb{R}^d)$ with $f \perp \rho$. Then*

$$\langle L_+ f | f \rangle \geq 0 \quad (2.21)$$

and

$$\langle L_+ f | f \rangle + |\langle f | \Lambda_0 \rho \rangle|^2 + |\langle f | \nabla \rho \rangle|^2 \simeq \|\nabla f\|_2^2 \quad (2.22)$$

where the implicit constants in (2.22) only depend on the dimension.

Proof. The first statement follows from the self-adjointness of L_+ and the description of the spectrum of L_+ . For the second, we need to invoke the calculus of variations and the concentration-compactness method. It is clear from Sobolev imbedding and Hölder’s inequality that the left-hand side of (2.22) dominates the right-hand side. Suppose the reverse inequality of (2.22) fails. Then there exists a sequence $\{f_n\} \subset \dot{H}^1(\mathbb{R}^d)$ with $\|\nabla f_n\|_2 = 1$ and $f_n \perp \rho$ which further satisfies

$$\langle L_+ f_n | f_n \rangle \rightarrow 0, \quad \langle f_n | \Lambda_0 \rho \rangle \rightarrow 0, \quad \langle f_n | \nabla \rho \rangle \rightarrow 0 \quad (2.23)$$

After passing to a subsequence we may assume that $f_n \rightharpoonup f_\infty$ in $\dot{H}^1(\mathbb{R}^d)$ and $L^{2^*}(\mathbb{R}^d)$, as well as $f_n \rightarrow f_\infty$ strongly in $L^2_{\text{loc}}(\mathbb{R}^d)$. Then $f_\infty \perp \rho$, and

$$\langle f_\infty | \Lambda_0 \rho \rangle = 0, \quad \langle f_\infty | \nabla \rho \rangle = 0 \quad (2.24)$$

From the local convergence in L^2 we conclude that

$$\int_{\mathbb{R}^d} W^{2^*-2}(x) |f_n(x)|^2 dx \rightarrow \int_{\mathbb{R}^d} W^{2^*-2}(x) |f_\infty(x)|^2 dx \quad (2.25)$$

By the first condition in (2.23) the left-hand side in (2.25) also tends to 1. But then $\langle L_+ f_\infty | f_\infty \rangle \leq 0$, and from (2.21) we conclude that $\langle L_+ f_\infty | f_\infty \rangle = 0$ whence

$$\|\nabla f_n\|_2 \rightarrow \|\nabla f_\infty\|_2 \quad \text{as } n \rightarrow \infty$$

Finally, this means that $f_n \rightarrow f_\infty$ strongly in $(\dot{H}^1 \cap L^{2^*})(\mathbb{R}^d)$. In summary, $L_+ f_\infty = 0$ and so

$$f_\infty = \alpha W' + \vec{\beta} \nabla W$$

Inserting this into (2.24) implies that $\alpha = 0$ and $\vec{\beta} = 0$. To see this, we first note that $\langle W' | \Lambda_0 \rho \rangle \neq 0$ which follows from $\langle W' | \rho \rangle = 0$ and

$$\begin{aligned} b_W &:= \langle W' | \Lambda_0 \rho \rangle = -k^{-2} \langle [L_+, x \cdot \nabla] W' | \rho \rangle \\ &= k^{-2} \langle (2L_+ + p(p-1)W^{p-2}W') W' | \rho \rangle \\ &= k^{-2} p(p-1) \langle W^{p-2} (W')^2 | \rho \rangle > 0 \end{aligned} \quad (2.26)$$

Since $\langle \nabla W | \Lambda_0 \rangle = 0$, we infer from this that $\alpha = 0$. On the other hand,

$$\langle \nabla_j W | \nabla_k \rho \rangle = -\frac{1}{d} \delta_{jk} \langle -\Delta W | \rho \rangle = -\frac{1}{d} \langle W^{2^*-2} | \rho \rangle \neq 0$$

and so $\vec{\beta} = 0$. But this clearly contradicts $\|\nabla f_\infty\|_2 = 1$, whence we have arrived at a contradiction. \square

In what follows we denote

$$\alpha := \langle v_1 | \Lambda_0 \rho \rangle = \langle \gamma_1 | \Lambda_0 \rho \rangle \quad (2.27)$$

so that (2.22) implies the following: for any $\underline{\gamma}$ with $\gamma_1 \perp \rho, \nabla \rho$ we have

$$\|\underline{\gamma}\|_{\mathcal{H}}^2 \simeq \langle \mathcal{L} \underline{\gamma} | \underline{\gamma} \rangle + \alpha^2 \quad (2.28)$$

2.5. Orthogonality conditions near the ground state. Now we introduce the crucial *orthogonality conditions* near the family of static solutions $\pm \mathcal{S}_0$. Indeed, we claim that for any $\underline{u} \in \mathcal{H}$ with

$$\min_{\pm} \text{dist}_{\mathcal{H}}(\underline{u}, \pm \mathcal{S}_0) \ll 1 \quad (2.29)$$

admits the representation $\underline{u} = T^c \vec{S}^\sigma (\pm \vec{W} + \underline{v})$ where we can choose $(\sigma, c) \in \mathbb{R}^{1+d}$ such that

$$0 = \alpha = \langle v_1 | \Lambda_0 \rho \rangle, \quad 0 = \mu = \langle v_1 | \nabla \rho \rangle = \omega(\underline{v}, J \nabla \vec{\rho}). \quad (2.30)$$

(2.30) are the orthogonality conditions which we use in this paper. To verify this claim, take any v_1 small in \dot{H}^1 (and thus small in $L^{2^*}(\mathbb{R}^d)$), and define (taking $+\mathcal{S}_0$ for simplicity)

$$F(\sigma, c) := \langle S_{-1}^{-\sigma} T^{-c}(W + v_1) - W | (\Lambda_0 \rho, \nabla \rho) \rangle \in \mathbb{R}^{1+n}. \quad (2.31)$$

Note that $F(0) = \langle v_1 | (\Lambda_0 \rho, \nabla \rho) \rangle$ is small and

$$F'(0) = \langle W + v_1 | (\Lambda_1, \nabla) \otimes (\Lambda_0, \nabla) \rho \rangle$$

is close to

$$\langle W | (\Lambda_1, \nabla) \otimes (\Lambda_0, \nabla) \rho \rangle = -\text{diag}(b_W, a_W, \dots, a_W)$$

It follows from the inverse function theorem that there exists (σ, c) small with $F(\sigma, c) = 0$. But then \tilde{v}_1 defined by means of

$$W + v_1 = S_{-1}^{\sigma} T^{-c}(W + \tilde{v}_1) \quad (2.32)$$

satisfies $\langle \tilde{v}_1 | \Lambda_0 \rho \rangle = 0$ and $\langle \tilde{v}_1 | \nabla \rho \rangle = 0$.

The orthogonality conditions (2.30) are equivalent to $\alpha = 0$ and $\mu = 0$ in (2.15), which “eliminate” the dilation and translation symmetries, respectively.

2.6. Linearized energy. Change variables from (λ_+, λ_-) to (λ_1, λ_2) as follows:

$$\lambda_1 = \frac{1}{\sqrt{2k}}(\lambda_+ + \lambda_-), \quad \lambda_2 = \sqrt{\frac{k}{2}}(\lambda_+ - \lambda_-), \quad \lambda_{\pm} = \sqrt{\frac{k}{2}}(\lambda_1 \pm k^{-1}\lambda_2). \quad (2.33)$$

Note that in these variables we have $\lambda_j = \langle v_j | \rho \rangle$ ($j = 1, 2$) and

$$v_1 = \lambda_1 \rho + \mu \nabla W + \gamma_1, \quad v_2 = \lambda_2 \rho + \gamma_2. \quad (2.34)$$

Now we define the nonlinear energy distance near $\pm\mathcal{S}_0$ by means of the equations

$$\begin{aligned} \mathcal{E}(\vec{u}) &:= E(\vec{u}) - J(W) + k^2 \lambda_1^2 + \alpha^2 + |\mu|^2 \\ &= \frac{1}{2}(k^2 \lambda_1^2 + \lambda_2^2) + \frac{1}{2} \langle \mathcal{L} \underline{\gamma} | \underline{\gamma} \rangle + \alpha^2 + |\mu|^2 - C(\underline{v}). \end{aligned} \quad (2.35)$$

Here $\underline{u} = T^c S^{\sigma}(\pm \vec{W} + \underline{v})$ with $\|\underline{v}\|_{\mathcal{H}}$ small and some choice of \pm , and we use the decomposition (2.15). Lemma 2.1 together with $C(\underline{v}) = o(\|\underline{v}\|_{\mathcal{H}}^2)$, implies that

$$\mathcal{E}(\vec{u}) \simeq \|\underline{v}\|_{\mathcal{H}}^2. \quad (2.36)$$

Hence it is natural to define the linearized energy norm by

$$\|\underline{v}\|_E^2 := \frac{1}{2}(k^2 \lambda_1^2 + \lambda_2^2) + \frac{1}{2} \langle \mathcal{L} \underline{\gamma} | \underline{\gamma} \rangle + \alpha^2 + |\mu|^2. \quad (2.37)$$

2.7. Modulation equations. Differentiation of the orthogonality conditions (2.30) in τ using the equation (2.13) yields

$$\begin{aligned} 0 &= \partial_\tau \langle v_1 | \Lambda_0 \rho \rangle = \langle v_2 | \Lambda_0 \rho \rangle - c_\tau e^\sigma \langle v_1 | \nabla \Lambda_0 \rho \rangle - \sigma_\tau [b_W - \langle v_1 | \Lambda_1 \Lambda_0 \rho \rangle], \\ 0 &= \partial_\tau \langle v_1 | \nabla \rho \rangle = \langle v_2 | \nabla \rho \rangle + c_\tau e^\sigma [a_W I_d - \langle v_1 | \nabla^2 \rho \rangle] + \sigma_\tau \langle v_1 | \Lambda_1 \nabla \rho \rangle, \end{aligned} \quad (2.38)$$

where $a_W, b_W > 0$ are as in (2.15) and (2.26), I_d is the d -dimensional unit matrix, and $\nabla^2 \rho$ is the Hessian. The modulation equations (2.38) determine the evolution of (σ, c) as long as \underline{v} remains small in \mathcal{H} . For future reference, we remark that in the notation of (3.3)

$$\langle v_2 | \Lambda_0 \rho \rangle = \langle \gamma_2 | \Lambda_0 \rho \rangle, \quad \langle v_2 | \nabla \rho \rangle = \langle \gamma_2 | \nabla \rho \rangle \quad (2.39)$$

whence

$$|\sigma_\tau| + |c_\tau| e^\sigma \lesssim \|\underline{\gamma}\|_{\mathcal{H}}, \quad (2.40)$$

as long as $\|\underline{v}\|_E$ is small.

2.8. Hyperbolic drivers. On the other hand, the unstable/stable modes evolve by the following equations derived from (2.13) with $\lambda_\pm = \omega(\underline{v}, \pm \underline{g}^\mp)$

$$\partial_\tau \lambda_\pm = \pm k \lambda_\pm \mp c_\tau e^\sigma \omega(\underline{v}, \nabla \underline{g}^\mp) \mp \sigma_\tau \omega(\underline{v}, \vec{\Lambda}^* \underline{g}^\mp) \pm \omega(\underline{N}(\underline{v}), \underline{g}^\mp). \quad (2.41)$$

In terms of λ_1 and λ_2 these equations become

$$\begin{aligned} \partial_\tau \lambda_1 &= \lambda_2 - c_\tau e^\sigma \langle v_1 | \nabla \rho \rangle + \sigma_\tau \langle v_1 | \Lambda_0 \rho \rangle \\ \partial_\tau \lambda_2 &= k^2 \lambda_1 - c_\tau e^\sigma \langle v_2 | \nabla \rho \rangle + \sigma_\tau \langle v_2 | \Lambda_1 \rho \rangle + \langle N(v_1) | \rho \rangle. \end{aligned} \quad (2.42)$$

The relation between these systems is given by (2.33). Under the orthogonality conditions (2.30) the first equation in (2.42) simplifies to

$$\partial_\tau \lambda_1 = \lambda_2. \quad (2.43)$$

This will be important to guarantee convexity of the distance function in the ejection lemma. However it should not be confused with the definition $\partial_t u_1 = u_2$, even though $\lambda_j = \langle v_j, \rho \rangle$, as t and τ are different.

3. DISTANCE FUNCTION, λ DOMINANCE, EJECTION

The following lemma establishes the existence of a distance function associated with the soliton manifold $\pm \mathcal{S}_0$ in such a way that near this manifold the distance function is proportional to the unstable mode in a suitable sense.

Lemma 3.1. *There exists $\delta_E > 0$ and $d_W(\underline{u}) : \mathcal{H} \rightarrow [0, \infty)$ continuous such that*

$$d_W(\underline{u}) \simeq \inf_{\pm, \sigma, c} \|\underline{u} \mp \vec{W}_\sigma(\cdot - c)\|_{\mathcal{H}}, \quad (3.1)$$

and so that for any $d_W(\underline{u}) \leq \delta_E$ there exists a unique vector $(\mathfrak{s}, \sigma, c) \in \{\pm 1\} \times \mathbb{R}^{1+d}$ with

$$\underline{u} = T^c \vec{S}^\sigma (\mathfrak{s} \vec{W} + \underline{v}), \quad \langle v_1 | \Lambda_0 \rho \rangle = 0, \quad \langle v_1 | \nabla \rho \rangle = 0. \quad (3.2)$$

Decomposing

$$\underline{v} = \lambda_+ \underline{g}^+ + \lambda_- \underline{g}^- + \underline{\gamma}, \quad \lambda_\pm = \omega(\underline{v}, \pm \underline{g}^\mp), \quad (3.3)$$

we have

$$d_W^2(\underline{u}) \simeq \|\underline{v}\|_E^2 = k^2(\lambda_+^2 + \lambda_-^2) + \frac{1}{2} \langle \mathcal{L} \underline{\gamma} | \underline{\gamma} \rangle \quad (3.4)$$

In addition, if

$$2(E(\underline{u}) - J(W)) < d_W^2(\underline{u}) < \delta_E^2 \quad (3.5)$$

then $d_W(\underline{u}) \simeq |\lambda_1| = |\lambda_+ + \lambda_-|$. Finally,

$$d_W(\underline{u}) \leq \delta_E \implies d_W^2(\underline{u}) = E(\underline{u}) - J(W) + k^2 \lambda_1^2 \quad (3.6)$$

Proof. There are $0 < \delta_A \ll 1$ and $C \geq 1$ such that putting $d_0(\underline{u}) := C \text{dist}_{\mathcal{H}}(\pm \mathcal{S}_0, \underline{u})$, the above arguments starting from (2.29) work in the region $d_0(\underline{u}) \leq \delta_A$, and

$$d_0(\underline{u})^2 \simeq E(\underline{u}) - J(W) + k^2 \lambda_1^2 =: d_1^2(\underline{u}) \leq d_0^2(\underline{u}). \quad (3.7)$$

Hence there exists $\delta_E \in (0, \delta_A)$ such that

$$d_0(\underline{u}) \leq \delta_A \text{ and } d_1(\underline{u}) \leq \delta_E \implies d_0(\underline{u}) \leq \delta_A/2. \quad (3.8)$$

Now we choose a smooth cutoff function $\chi \in C^\infty(\mathbb{R})$ satisfying $\chi(r) = 1$ for $|r| \leq 1$ and $\chi(r) = 0$ for $|r| \geq 2$ and set

$$d_W(\underline{u}) := \chi(2d_0(\underline{u})/\delta_A) d_1(\underline{u}) + (1 - \chi)(2d_0(\underline{u})/\delta_A) d_0(\underline{u}). \quad (3.9)$$

If $d_0(\underline{u}) \geq \delta_A$ then $d_W(\underline{u}) = d_0(\underline{u}) \geq \delta_A > \delta_E$. Hence if $d_W(\underline{u}) \leq \delta_E$ then $d_0(\underline{u}) < \delta_A$ and $d_1(\underline{u}) \leq d_W(\underline{u}) \leq \delta_E$, so $d_0(\underline{u}) \leq \delta_A/2$, $d_W(\underline{u}) = d_1(\underline{u})$. The stated properties now follow easily from the considerations in the previous section. \square

The following lemma is the analogue of the ‘‘ejection lemma’’ in our previous papers, see [21]. As usual, we shall need the Payne-Sattinger functional

$$K(u) = \int_{\mathbb{R}^d} [|\nabla u|^2 - |u|^{2^*}] dx \quad (3.10)$$

in our analysis of the global dynamics, which is why it appears below.

Lemma 3.2. *There exists $\delta_H \in (0, \delta_E)$ with the following properties: Let u be a solution on an open interval I such that for some $t_0 \in I$*

$$\delta_0 := d_W(\vec{u}(t_0)) \leq \delta_H, \quad E(\vec{u}) - J(W) \leq \delta_0^2/2, \quad (3.11)$$

and

$$\partial_t d_W(\vec{u}(t_0)) \geq 0. \quad (3.12)$$

Apply the decomposition from Lemma 3.1. Then for $t > t_0$ in I and as long as $d_W(\vec{u}(t)) \leq \delta_H$, $d_W(\vec{u}(t))$ is increasing, and

$$d_W(\vec{u}(t)) \simeq -\mathfrak{s}\lambda_+(t) \simeq -\mathfrak{s}\lambda_1(t) \simeq e^{k\tau}\delta_0, \quad (3.13)$$

where $\tau(t)$ is the solution of the ODE $\tau'(t) = e^{\sigma(t)}$ with $\tau(t_0) = 0$. Moreover,

$$\begin{aligned} |\sigma(t) - \sigma(t_0)| &\lesssim d_W(\vec{u}(t)), \\ \mathfrak{s}K(u(t)) &\gtrsim d_W(\vec{u}(t)) - C_* d_W(\vec{u}(t_0)), \end{aligned} \quad (3.14)$$

$$|\lambda_-(t)| + \|\underline{\gamma}(t)\|_{\mathcal{H}} \lesssim \delta_0 + d_W^2(\vec{u}(t)),$$

for some absolute constant $C_* > 0$ and $\mathfrak{s} = \pm 1$ is fixed on the time interval.

Proof. By Lemma 3.1 and (3.11), we conclude that $|\lambda_1(t_0)| \simeq \delta_0$. Furthermore, as long as $d_W(\vec{u}(t))$ remains sufficiently small and one has $d_W(\vec{u}(t)) \geq \delta_0$, the relation

$$|\lambda_1(t)| \simeq d_W(\vec{u}(t)) \quad (3.15)$$

is preserved. In particular, if $d_W(\vec{u}(t))$ is increasing this relation is preserved. We shall therefore *assume* (3.15) in our argument that establishes the monotonicity and (3.13). The logic here is that once we have shown these properties to be correct, then the validity of (3.15) follows a posteriori by the method of continuity.

Differentiating (3.6) using (2.42) and (2.38) as well as (3.2) yields,

$$\partial_\tau d_W^2(\vec{u}) = 2k^2 \lambda_1 \partial_\tau \lambda_1 = 2k^2 \lambda_1 \lambda_2 \quad (3.16)$$

and

$$\partial_\tau^2 d_W^2(\vec{u}) = 2k^2(k^2 \lambda_1^2 + \lambda_2^2) + O(\lambda_1^3). \quad (3.17)$$

In conjunction with the previous lemma we conclude from (3.17) that $d_W^2(u)$ is increasing and convex in τ , as long as it remains sufficiently small. Next, we remark that λ_1 and $\partial_\tau \lambda_1 = \lambda_2$ have the same sign, since

$$2k^2 \lambda_1 \lambda_2 = \partial_\tau d_W^2(\vec{u}) \geq \partial_\tau d_W^2(\vec{u})|_{t=t_0} = 2d_W(\vec{u}(t_0))\partial_t d_W(\vec{u}(t_0)) \geq 0. \quad (3.18)$$

This implies that $|\lambda_+| \geq |\lambda_-|$ and thus $|\lambda_1| \simeq |\lambda_+|$.

The evolution of λ_+ in τ is determined by (2.41), which states that

$$\partial_\tau \lambda_\pm = \pm k \lambda_\pm + O(d_W^2(\vec{u})). \quad (3.19)$$

Since $d_W(\vec{u}) \lesssim |\lambda_+|$, we see that $|\lambda_+| \simeq e^{k\tau}\delta_0$, and

$$|\lambda_-| \lesssim \delta_0 + \lambda_+^2 \quad (3.20)$$

as claimed. As for the scaling parameters, (2.38) yields $|\partial_\tau \sigma| \lesssim \|\gamma\|_{\mathcal{H}} \lesssim d_W(\vec{u})$, and hence integrating the exponential bound in τ implies $|\sigma - \sigma(t_0)| \lesssim d_W(\vec{u})$.

The $\underline{\gamma}$ -part is estimated as in Lemma 4.3 of [21]. To this end define

$$\underline{v}_d := \lambda_+ \underline{g}^+ + \lambda_- \underline{g}^- = \underline{v} - \underline{\gamma}. \quad (3.21)$$

Then

$$\begin{aligned} E(\vec{W} + \underline{v}) &= J(W) + \frac{1}{2}(\lambda_2^2 - k^2\lambda_1^2) + \frac{1}{2}\langle \mathcal{L}\underline{\gamma}|\underline{\gamma} \rangle - C(\underline{v}) \\ E(\vec{W} + \underline{v}_d) &= J(W) + \frac{1}{2}(\lambda_2^2 - k^2\lambda_1^2) - C(\underline{v}_d) \end{aligned} \quad (3.22)$$

whence

$$E(\vec{u}) - E(\vec{W} + \underline{v}_d) = \frac{1}{2}\langle \mathcal{L}\underline{\gamma}|\underline{\gamma} \rangle + C(\underline{v}_d) - C(\underline{v}) \quad (3.23)$$

as well as (where $\underline{v}_{d,1}$ denotes the first component of \underline{v}_d)

$$\begin{aligned} \partial_\tau E(\vec{W} + \underline{v}_d) &= \lambda_2 \partial_\tau \lambda_2 - k^2 \lambda_1 \partial_\tau \lambda_1 - \langle N(\underline{v}_{d,1}) | \partial_\tau \underline{v}_{d,1} \rangle \\ &= \lambda_2 (\langle N(\underline{v}_1) | \rho \rangle + \sigma_\tau \langle \underline{v}_2 | \Lambda_1 \rho \rangle - c_\tau e^\sigma \langle \underline{v}_2 | \nabla \rho \rangle - \langle N(\underline{v}_{d,1}) | \rho \rangle) \\ &= O(d_W^2(\vec{u}) \|\underline{\gamma}\|_{\mathcal{H}}) \end{aligned} \quad (3.24)$$

where we used (2.40) to bound σ_τ and $c_\tau e^\sigma$. In view of the preceding,

$$\begin{aligned} |E(\vec{u}) - E(\vec{W} + \underline{v}_d)| &\lesssim |E(\underline{u}(t_0)) - E(\vec{W} + \underline{v}_d(t_0))| + \\ &\quad + |E(\vec{W} + \underline{v}_d) - E(\vec{W} + \underline{v}_d(t_0))| \\ &\lesssim \delta_0^2 + \|\underline{\gamma}\|_{L^\infty([t_0, t], \mathcal{H})} d_W^2(\vec{u}) \end{aligned}$$

where we used the exponential growth of $d_W(\vec{u})$ to pass to the last line. Furthermore,

$$\begin{aligned} \frac{1}{2}\langle \mathcal{L}\underline{\gamma}|\underline{\gamma} \rangle &\leq |E(\vec{u}) - E(\vec{W} + \underline{v}_d)| + |C(\underline{v}_d) - C(\underline{v})| \\ &\lesssim \delta_0^2 + \|\underline{\gamma}\|_{L^\infty([t_0, t], \mathcal{H})} d_W^2(\vec{u}) \end{aligned} \quad (3.25)$$

In conclusion,

$$\|\underline{\gamma}\|_{\mathcal{H}} \lesssim \delta_0 + d_W^2(\vec{u}) \quad (3.26)$$

as claimed.

Finally, expanding the K -functional, one checks that

$$K(W + v_1) = -(2^* - 2)\langle W^{2^*-1} | v_1 \rangle + O(\|v_1\|_{\dot{H}^1}^2) \quad (3.27)$$

Inserting the expansion $v_1 = \lambda_1 \rho + \gamma_1$ into (3.27) and using the bounds on $\lambda_1(t)$ and $\underline{\gamma}(t)$ that we just established implies the desired properties of K . \square

We remark that unlike the subcritical nonradial paper [20] the distance function is convex near a minimum and thus increasing in Lemma 3.2. The difference lies with the choice of orthogonality conditions corresponding to the translational symmetry, which in our case insure that $\partial_\tau \lambda_1 = \lambda_2$. This coincides with the behavior in the radial subcritical case, see [21].

4. THE VARIATIONAL STRUCTURE IN THE ENERGY CRITICAL SETTING

We recall the following characterization of the ground state:

$$\begin{aligned} J(W) &= \inf\{J(\varphi) \mid K(\varphi) = 0, \varphi \in \dot{H}^1(\mathbb{R}^d), \varphi \neq 0\} \\ &= \inf\left\{\frac{1}{d}\|\nabla\varphi\|_2^2 \mid K(\varphi) \leq 0, \varphi \in \dot{H}^1(\mathbb{R}^d), \varphi \neq 0\right\} \end{aligned} \quad (4.1)$$

where $J(\varphi)$ is the static energy defined in (1.11), and $\pm W$ are the unique minimizer up to the dilation (as in W_σ) and translation symmetries. In other words, W is the unique (up to the same symmetries) extremizer of the Sobolev embedding $\dot{H}^1(\mathbb{R}^d) \hookrightarrow L^{2^*}(\mathbb{R}^d)$. We need the following variational structure outside of the soliton tube.

Lemma 4.1. *For each $0 < \delta < 1$ there is $\varepsilon_1 = \varepsilon_1(\delta) > 0$ such that if $\underline{u} \in \mathcal{H}$ satisfies $J(u_1) < J(W) + \varepsilon_1^2(\delta)$ and $d_W(\underline{u}) > \delta$, then we have either*

$$K(u_1) > \min\{\kappa(\delta), c\|\nabla u_1\|_{L^2}^2\} \quad (4.2)$$

or else

$$K(u_1) < -\kappa(\delta) \quad (4.3)$$

for suitable $\kappa(\delta) > 0$ and some absolute constant $c > 0$.

Proof. We first eliminate the u_2 component from \underline{u} : if $\|u_2\|_2 \ll \delta$, then it follows that $d_W(u_1, 0) > \delta/2$. On the other hand, if $\|u_2\| \simeq \delta$, then assuming $\varepsilon_1(\delta) \ll \delta$ as we may, it follows that

$$J(u_1) < J(W) - c\delta^2$$

with some absolute constant c . But then we must have $\|u_1 - W_\sigma(\cdot - c)\|_{\dot{H}^1} \gtrsim \delta$ for all σ, c . Hence

$$\delta \lesssim d_W(u_1, 0) \simeq \text{dist}(u_1, \mathcal{S}_0) \quad (4.4)$$

in all cases. In the rest of proof we regard $u = u_1 \in \dot{H}^1(\mathbb{R}^d)$ with $\text{dist}(u, \mathcal{S}_0) \gtrsim \delta$.

By the critical Sobolev imbedding, the statement holds provided $\|\nabla u\|_2 < c_0$ where $c_0 > 0$ is some absolute constant. Thus, assume the lemma fails and let $\{u_n\}_{n=1}^\infty \subset \dot{H}^1$ be a sequence with

$$\|\nabla u_n\|_2 \rightarrow c \geq c_0, \quad K(u_n) \rightarrow 0, \quad J(u_n) < J(W) + \frac{1}{n} \quad (4.5)$$

as well as $\text{dist}(u_n, \mathcal{S}_0) \gtrsim \delta_0$. Since

$$J(u_n) = \frac{1}{d}\|\nabla u_n\|_2^2 + \frac{1}{2^*}K(u_n) \quad (4.6)$$

we see that $\{u_n\}_{n=1}^\infty$ is bounded in $\dot{H}^1 \cap L^{2^*}$ and so $c < \infty$. Then the latter two conditions of (4.5) implies that $\{u_n\}$ is an extremizing sequence for the critical Sobolev embedding $\dot{H}^1(\mathbb{R}^d) \subset L^{2^*}(\mathbb{R}^d)$, and so, by the celebrated theorem of P.-L. Lions [15, Theorem I.1], it is compact in \dot{H}^1 up to scaling

and translation, hence converging strongly to the unique minimizer W up to scaling and translation. But this clearly contradicts $\text{dist}(u_n, \mathcal{S}_0) \gtrsim \delta_0$. \square

As in the previous works [21], we can define a sign functional by combining the ejection lemma with the variational structure exhibited in the previous lemma.

Corollary 4.2. *Let $\delta_S := \delta_H/(2C_*) > 0$ where $\delta_H > 0$ and $C_* \geq 1$ are the constants from Lemma 3.2. Let $0 < \delta \leq \delta_S$ and*

$$\mathfrak{H}_{(\delta)} := \{\underline{u} \in \mathcal{H} \mid E(\underline{u}) < J(Q) + \min(d_W^2(\underline{u})/2, \varepsilon_1^2(\delta))\}, \quad (4.7)$$

where $\varepsilon_1(\delta)$ is defined in Lemma 4.1. Then there exists a unique continuous function $\mathfrak{S} : \mathfrak{H}_{(\delta)} \rightarrow \{\pm 1\}$ satisfying

$$\begin{cases} \underline{u} \in \mathfrak{H}_{(\delta)}, d_W(\underline{u}) \leq \delta_E & \implies \mathfrak{S}(\underline{u}) = -\text{sign } \lambda_1, \\ \underline{u} \in \mathfrak{H}_{(\delta)}, d_W(\underline{u}) \geq \delta & \implies \mathfrak{S}(\underline{u}) = \text{sign } K(\underline{u}), \end{cases} \quad (4.8)$$

where we set $\text{sign } 0 = +1$.

Proof. The proof is the same as in the subcritical radial case, see [18]. \square

5. THE ONE-PASS THEOREM

A key step in the proof of our main theorem is to show that the sign $\mathfrak{S}(\underline{u}(t))$ can change at most once for any solution of (1.1). This goes by the name of *one-pass theorem*, see [21]. The current section is entirely devoted to this theorem:

Theorem 5.1. *There exist $0 < \varepsilon_* \ll \delta_* \ll \delta_H$ with the following properties: Let $\vec{u} \in C(I; \mathcal{H})$ be a solution of (1.1) on an open interval I , satisfying for some $\varepsilon \in (0, \varepsilon_*]$, $\delta \in (\sqrt{2}\varepsilon, \delta_*]$ and $T_1 < T_2 \in I$*

$$E(\vec{u}) \leq J(W) + \varepsilon^2, \quad d_W(\vec{u}(T_1)) < \delta = d_W(\vec{u}(T_2)). \quad (5.1)$$

Then $d_W(\vec{u}(t)) > \delta$ for all $t > T_2$ in I .

Proof. By increasing T_1 and decreasing T_2 if necessary, we may assume in addition that $\sqrt{2}\varepsilon < d_W(\vec{u}(T_1))$ and $\partial_t d_W(\vec{u}(t))|_{t=T_1} \geq 0$. Then Lemma 3.2 applies for all $t \in [T_1, T_2]$ and so $d_W(\vec{u}(t))$ is increasing for $t > T_1$ until it reaches δ_H (the small absolute scale in the ejection lemma). Arguing by contradiction, we assume that for some $t > T_2$ we have $d_W(\vec{u}(t)) \leq \delta$. Such a t can occur only away from T_2 (this will be made more precise shortly), and after $d_W(\vec{u}(t))$ has increased to size $\delta_H \gg \delta$. Moreover, by applying Lemma 3.2 backward in time, we can find $T_3 > T_2$ such that $d_W(\vec{u}(t))$ decreases from δ_H down to δ as $t \nearrow T_3$, and so that

$$d_W(\vec{u}(t)) > \delta = d_W(\vec{u}(T_3)) = d_W(\vec{u}(T_2))$$

for $T_2 < t < T_3$. We may further assume

$$\sigma(u(T_2)) = 0 \leq \sigma(u(T_3)), \quad (5.2)$$

by rescaling and reversing time, if necessary. Here σ is defined in Lemma 3.1.

We now proceed by combining the proof ideas of the analogous theorem for the critical radial wave equation [12] with that for the subcritical nonradial Klein-Gordon equation, see [20] (with slight improvement). Following the latter reference, we first show that the centers of the ground state as given by the path $c(t)$, diverge from each other between times T_2 and T_3 by an amount $\ll T_3 - T_2$. Once this is done, we shall adapt the virial argument from [12] to the nonradial context, which will then allow us to exclude almost homoclinic orbits. It will be understood that all times t belong to the interval I .

By spatial translation, we may assume that $c(T_2) = 0$. By the ejection we have

$$T_3 - T_2 \gtrsim \log(\delta_E/\delta) \gg 1, \quad (5.3)$$

and by the finite speed of propagation

$$|c(T_3)| \leq T_3 - T_2 + O(1), \quad (5.4)$$

where $c(t) = c(u(t)) \in \mathbb{R}^d$ is defined by Lemma 3.1 as long as $\vec{u}(t)$ is close to \mathcal{S}_0 , which is true when t is close to T_2 or T_3 . Consider a localized center of energy defined by (with $\vec{u} = (u_1, u_2)$)

$$\mathfrak{C}(t) := \langle xw|e(\vec{u}) \rangle, \quad e(\vec{u}) := [|u_2|^2 + |\nabla u_1|^2]/2 - |u_1|^{2^*}/2^*, \quad (5.5)$$

where $w(t, x)$ is the cut-off function onto a light cone defined by

$$w(t, x) = \chi(|x|/(t - T_2 + S)) \quad (5.6)$$

for some $1 \ll S = S(\delta) < \delta^{-2}$ to be determined, and some $\chi \in C^\infty(\mathbb{R})$ satisfying $\chi(r) = 1$ for $|r| \leq 1.5$ and $\chi(r) = 0$ for $|r| \geq 2$. Then using the equation of u , we have

$$\begin{aligned} \dot{\mathfrak{C}}(t) &= \langle \dot{w}x|e(\vec{u}) \rangle - P(\vec{u}) + \langle (1-w)u_2|\nabla u_1 \rangle - \langle xu_2|\nabla w \cdot \nabla u_1 \rangle \\ &= O(E_{\text{ext}}(t)) - P(\vec{u}), \end{aligned} \quad (5.7)$$

where $E_{\text{ext}}(t) := \|\vec{u}(t)\|_{\mathcal{H}(|x|>t-T_2+S)}^2$ denotes the exterior free energy. Hence,

$$|\mathfrak{C}(T_3) - \mathfrak{C}(T_2)| \lesssim (T_3 - T_2) \max_{T_2 \leq t \leq T_3} (E_{\text{ext}}(t) + |P(\vec{u})|). \quad (5.8)$$

The conserved momentum is small because

$$|P(\vec{u})| \leq |P(\vec{u}) - P(T^c \vec{S}^\sigma \vec{W})| \lesssim \|\underline{v}(T_1)\|_{\mathcal{H}} + \|\underline{v}(T_1)\|_{\mathcal{H}}^2 \lesssim \delta. \quad (5.9)$$

Using the finite propagation as in [20] and [12] we have for all $t \geq T_2$

$$E_{\text{ext}}(t) \lesssim E_{\text{ext}}(T_2) \lesssim S^{2-d} + \delta^2 \lesssim S^{-1} \quad (5.10)$$

On the other hand, the radial symmetry and the rate of decay of W imply that

$$|\mathfrak{C}(T_2)| \lesssim \sqrt{S}\|\underline{v}\|_E + S\|\underline{v}\|_E^2 \lesssim \sqrt{S}\delta \quad (5.11)$$

The contribution of $W_{\sigma(T_3)}(x - c(T_3))$ at $t = T_3$ is estimated as follows. Denote $c_3 := c(T_3)$, $\sigma_3 := \sigma(T_3)$. Then

$$\begin{aligned} \mathfrak{C}(T_3) &= \langle xw | e(\vec{W}_{\sigma_3}(\cdot - c_3)) \rangle + \langle xw | e(\vec{u}(T_3)) - e(\vec{W}_{\sigma_3}(\cdot - c_3)) \rangle \\ &=: A + B \end{aligned} \quad (5.12)$$

Now, using (5.2),

$$\begin{aligned} A &= c_3 \langle w | e(\vec{W}_{\sigma_3}(\cdot - c_3)) \rangle + \langle (x - c_3)w | e(\vec{W}_{\sigma_3}(\cdot - c_3)) \rangle \\ &= c_3(E(\vec{W}) + o(1)) + \langle (x - c_3)w | e(\vec{W}_{\sigma_3}(\cdot - c_3)) \rangle, \end{aligned} \quad (5.13)$$

where $o(1)$ is with respect to the limit $S \rightarrow \infty$ (uniformly for the other parameters c_3 , σ_3 , T_2 and T_3). Exploiting (5.4) and the obvious cancellation yields

$$|\langle (x - c_3)w | e(\vec{W}_{\sigma_3}(\cdot - c_3)) \rangle| \lesssim 1 + \log(1 + S^{-1}(T_3 - T_2)). \quad (5.14)$$

On the other hand,

$$B \lesssim (T_3 - T_2 + S)\delta. \quad (5.15)$$

Combining these estimates yields

$$|c(T_3)| \lesssim |\mathfrak{C}(T_3) - c(T_3)(E(\vec{W}) + o(1))| + |\mathfrak{C}(T_3) - \mathfrak{C}(T_2)| + |\mathfrak{C}(T_2)|, \quad (5.16)$$

and therefore

$$|c(T_3)| \lesssim 1 + S^{-1}(T_3 - T_2) + (T_3 - T_2 + S)\delta + \sqrt{S}\delta. \quad (5.17)$$

To obtain the desired contradiction, we use the localized virial identity

$$\begin{aligned} V_w(t) &:= \langle wu_2 | (x\nabla + d/2)u_1 \rangle, \\ \dot{V}_w(t) &= -K(u_1(t)) + O(E_{\text{ext}}(t)) = -K(u_1(t)) + O(S^{-1}) \end{aligned} \quad (5.18)$$

with the same choice of w as above. By similar considerations as above, one has the upper bounds

$$\begin{aligned} |V_w(T_2)| &\lesssim \delta S^{\frac{1}{2}}, \\ |V_w(T_3)| &\lesssim \delta(|c_3| + (T_3 - T_2 + S)^{\frac{1}{2}}) + \delta^2(T_3 - T_2 + S). \end{aligned} \quad (5.19)$$

Setting $S := \delta^{-1}$ in (5.17) implies

$$|c(T_3)| \lesssim 1 + \delta(T_3 - T_2)$$

and

$$|V_w(T_2)| + |V_w(T_3)| \lesssim \delta^{\frac{1}{2}} + \delta^2(T_3 - T_2) + \delta(T_3 - T_2)^{\frac{1}{2}} \quad (5.20)$$

We claim that integrating the differential equation in (5.18) and exploiting the ejection dynamics and the variational structure (cf. [21]) leads to the lower bound

$$\int_{T_2}^{T_3} \mathfrak{s}K(u_1(t)) dt \gtrsim \nu(\delta, \delta_H)\delta(T_3 - T_2) + \delta_H, \quad (5.21)$$

where $0 < \nu(\delta, \delta_H) \rightarrow \infty$ as $\delta \rightarrow +0$ and δ_H fixed. This clearly contradicts (5.20), provided that we choose $\delta_* \ll \delta_H^2$ small enough.

It remains to prove (5.21). Let \mathcal{T} be the set of times at which the distance $d_W(\vec{u}(t))|_{[T_2, T_3]}$ reaches a local minima in $[\delta, \delta_S]$. In particular, $\mathcal{T} \ni T_2, T_3$ by the choice of T_2 and T_3 . For every $t_* \in \mathcal{T}$, we can apply the ejection Lemma 3.2 from $t = t_*$ in both time directions. Then we get an interval $I(t_*) \subset [T_2, T_3]$ such that $t_* \in I(t_*)$, $d_W(\vec{u})$ within $I(t_*)$ is decreasing for $t < t_*$ and increasing for $t > t_*$, and $d_W(\vec{u}) = \delta_H$ on $\partial I(t_*) \setminus \{T_2, T_3\}$. Moreover, imposing

$$0 < \varepsilon_* < \varepsilon_1(\delta_S), \quad (5.22)$$

we can ensure that \vec{u} stays in \mathfrak{H}_{δ_S} for $t \in [T_2, T_3]$, so that the sign \mathfrak{s} in the ejection lemma is the same for all $I(t_*)$ by Corollary 4.2. Furthermore, the exponential behavior allows us to estimate

$$\begin{aligned} |I(t_*)| &= \int_{t \in I(t_*)} e^{-\sigma} d\tau \simeq e^{-\sigma(t_*)} \log(\delta_H/d_W(\vec{u}(t_*))) \leq e^{-\sigma(t_*)} \log(\delta_H/\delta), \\ \int_{I(t_*)} \mathfrak{s} K(u(t)) dt &\gtrsim \int_{t \in I(t_*)} (e^{k\tau} - C_*) d_W(\vec{u}(t_*)) e^{-\sigma} d\tau \simeq e^{-\sigma(t_*)} \delta_H. \end{aligned} \quad (5.23)$$

Summing this over all $t_* \in \mathcal{T}$ including T_2 and T_3 , we get

$$\int_{J_1} \mathfrak{s} K(u(t)) dt \gtrsim \delta_H + \frac{\delta_H/\delta}{\log(\delta_H/\delta)} \delta |J_1|, \quad J_1 := \bigcup_{t_* \in \mathcal{T}} I(t_*). \quad (5.24)$$

For the remaining times, we have

$$\inf_{t \in J_0} d_W(\vec{u}(t)) \geq \delta_S, \quad J_0 := [T_2, T_3] \setminus J_1, \quad (5.25)$$

by the definition of J_1 , so that under (5.22) we can use the variational bound of Lemma 4.1. If $\mathfrak{s} = -1$, then we have

$$\int_{J_0} \mathfrak{s} K(u(t)) dt \geq \kappa(\delta_S) |J_0|. \quad (5.26)$$

Adding (5.26) and (5.24) concludes the $\mathfrak{s} = -1$ case of (5.21).

If $\mathfrak{s} = +1$, then the same argument encounters the difficulty that outside of the δ_H -ball the lower bound of Lemma 4.1 may become degenerate due to smallness of $\|\nabla u\|_2^2$. Indeed, replacing $\kappa(\delta_S)$ in the above argument by $\min(\kappa(\delta_S), c\|\nabla u\|_2^2)$ and using the uniform bound on $\|\vec{u}\|_{\mathcal{H}}$ in the region $\mathfrak{S} = +1$ yields

$$\int_{T_2}^{T_3} K(u_1(t)) dt \gtrsim \frac{\delta_H/\delta}{\log(\delta_H/\delta)} \delta \int_{T_2}^{T_3} \|\nabla u(t)\|_2^2 dt + \delta_H. \quad (5.27)$$

This leads to (5.21) for $\mathfrak{s} = +1$ if

$$\int_{T_2}^{T_3} \|\nabla u(t)\|_2^2 dt \gtrsim T_3 - T_2. \quad (5.28)$$

Therefore assume that

$$\int_{T_2}^{T_3} \|\nabla u(t)\|_2^2 dt \leq \kappa_2(T_3 - T_2) \quad (5.29)$$

with some absolute constant κ_2 . To lead (5.29) to a contradiction, we use the (localized) energy equipartition

$$\partial_t \langle wu_t | u \rangle = \|\dot{u}(t)\|_2^2 - K(u(t)) + O(E_{\text{ext}}(t)) \geq 2E(\vec{u}) - \|\nabla u\|_2^2 + O(\delta), \quad (5.30)$$

where w , E_{ext} and $S = \delta^{-1}$ are as before. Taking $\delta_*, \kappa_2 \ll J(W)$, we obtain

$$[\langle wu_t | u \rangle]_{T_2}^{T_3} \geq E(\vec{u})(T_3 - T_2). \quad (5.31)$$

On the other hand, the same argument as for (5.20) yields

$$|\langle wu_t | u \rangle(T_3) - \langle wu_t | u \rangle(T_2)| \lesssim \delta^{1/2} + \delta^2(T_3 - T_2) + \delta(T_3 - T_2)^{1/2}, \quad (5.32)$$

which contradicts (5.31) since $T_3 - T_2 \gg 1 \gg \delta$. \square

The above result has some important implications for the sign functional from Corollary 4.2. To be specific, let

$$\begin{aligned} \mathcal{H}_* &= \{\underline{\varphi} \in \mathcal{H} \mid E(\underline{\varphi}) \leq J(W) + \varepsilon_*^2\}, \\ \mathcal{H}_X &= \{\underline{\varphi} \in \mathcal{H}_* \mid E(\underline{\varphi}) < J(W) + d_W^2(\underline{\varphi})/2\}. \end{aligned} \quad (5.33)$$

It is easy to see that $\mathcal{H}_* \setminus \mathcal{H}_X$ is a small neighborhood of $\pm \mathcal{S}_0$.

Corollary 5.2. *The sign function \mathfrak{S} in Corollary 4.2 is continuous on \mathcal{H}_X , and has the following properties.*

- (1) *Every solution u in \mathcal{H}_* can change $\mathfrak{S}(\vec{u}(t))$ at most once. Moreover, it can enter or exit the region $d_W(\vec{u}) < \delta_*$ at most once.*
- (2) *The region $\mathfrak{S} = +1$ is bounded in \mathcal{H} , while the region $\mathfrak{S} = -1$ is unbounded.*
- (3) *If $\underline{\varphi} \in \mathcal{H}_X$ and $E(\underline{\varphi}) \leq J(W) + \varepsilon_1^2(d_W(\underline{\varphi}))$, then $\mathfrak{S}(\underline{\varphi}) = \text{sign } K(\varphi_1)$, with the convention $\text{sign } 0 = +1$.*
- (4) *If $\underline{\varphi} \in \mathcal{H}_X$ and $d_W(\underline{\varphi}) \leq \delta_S$, then $\mathfrak{S}(\underline{\varphi}) = -\text{sign } \lambda_1(\varphi_1)$.*

The proof is the same as in the radial case [12], so we omit it. Note that $\mathcal{H}_* \setminus \mathcal{H}_X$ is included in $d_W < \delta_*$, and that (3)–(4) completely determine $\mathfrak{S}(\underline{\varphi})$, since we have chosen $\varepsilon_* < \varepsilon_1(\delta_S)$ in (5.22). Moreover, $\mathfrak{S}(\underline{\varphi})$ depends only on φ_1 .

It remains to determine the fate of the solutions in \mathcal{H}_* with $d_W \geq \delta_*$. We will do this in the following two sections for $\mathfrak{S} = \pm 1$, respectively.

6. BLOW-UP AFTER EJECTION

In analogy to [12], we now prove¹ the following

¹The proof is essentially the same as in [12], but since we employ somewhat different notation, we provide the details for the reader's convenience.

Proposition 6.1. *No solution $\vec{u} \in \mathcal{H}_*$ satisfying the conditions in Theorem 5.1 can stay strongly continuous with respect to the topology of \mathcal{H}_* and satisfy the requirements*

$$d_W(\vec{u}) \geq \delta, \quad \mathfrak{S} = -1, \quad (6.1)$$

for all $t > T_2$.

Proof. Suppose, for the sake of contradiction, that the solution actually exists on $(0, \infty)$. Write $w = \chi(\frac{|x|}{t+R})$ for some $R \gg 1$ to be chosen, and $\chi \in C_0^\infty(\mathbb{R})$ a non-negative cutoff function, with $\chi'(r) \leq 0$ for $r \geq 0$ and $\chi(r) = 1$ on $r \leq 1$. Also introduce

$$y(t) = \langle wu|u \rangle. \quad (6.2)$$

Then we have

$$\dot{y}(t) = \langle \dot{w}u + 2w\dot{u}|u \rangle \geq 2\langle w\dot{u}|u \rangle \quad (6.3)$$

Writing $E_{\text{ext}}(t) := \|\vec{u}(t)\|_{\mathcal{H}(|x|>t+R)}^2$, we find using Hardy's inequality

$$\ddot{y} = \langle 2w|\dot{u}^2 - |\nabla u|^2 + |u|^{2^*} \rangle + \langle \ddot{w}u|u \rangle + \langle 4\dot{w}u|\dot{u} \rangle + 2\langle u\nabla w|\nabla u \rangle \quad (6.4)$$

$$= 2(\|\dot{u}\|_2^2 - K(u)) + O(E_{\text{ext}}(t)) \quad (6.5)$$

It follows from the finite propagation as before that we can choose τ large enough such that $E_{\text{ext}}(t) \ll \varepsilon_*^2$ for all $t > 0$.

We next follow the argument in the proof of Theorem 5.1, especially the part after (5.21). Thus with T_2 as in the proof of Theorem 5.1, we write

$$[T_2, \infty) = J_0 \cup J_1 \quad (6.6)$$

with J_0 and J_1 defined just as in the proof of Theorem 5.1, with T_3 replaced by $+\infty$. Then as before we find $-K(u) > \kappa(\delta_M)$ on J_0 ; on the complement

$$J_1 = \bigcup_{t_* \in \mathcal{T}} I(t_*), \quad (6.7)$$

we also obtain the lower bound

$$-\int_{I(t_*)} K(u(t)) dt \gg \delta|I(t_*)|. \quad (6.8)$$

We conclude that

$$\lim_{t \rightarrow +\infty} \dot{y}(t) = \lim_{t \rightarrow +\infty} y(t) = +\infty \quad (6.9)$$

and $y(t)$ is increasing for large enough t .

Next, write

$$\|\dot{u}\|_2^2 - K(u) = \left(1 + \frac{2^*}{2}\right)\|\dot{u}\|_2^2 + \frac{2^* - 2}{2}\|\nabla u\|_2^2 - 2^*E(\vec{u}) \quad (6.10)$$

If $t \in J_0$, then from the variational characterization of W for $K < 0$, we have

$$\|\nabla u(t)\|_2 > \|\nabla W\|_2 \quad (6.11)$$

and so

$$E(\vec{u}) < J(W) + \varepsilon_*^2 = \frac{2^* - 2}{2 \cdot 2^*} \|\nabla W\|_2^2 + \varepsilon_*^2 < \frac{2^* - 2}{2 \cdot 2^*} \|\nabla u(t)\|_2^2 + \varepsilon_*^2,$$

which in conjunction with (6.10) implies

$$\|\dot{u}\|_2^2 - K(u) > (1 + \frac{2^*}{2}) \|\dot{u}\|_2^2 - 2^* \varepsilon_*^2. \quad (6.12)$$

We also have the bound from Lemma 4.1

$$\|\dot{u}\|_2^2 - K(u) > \|\dot{u}\|_2^2 + \kappa(\delta_S), \quad (6.13)$$

which on account of (5.22) and interpolation with the bound (6.12) implies

$$\ddot{y}(t) > 4(1 + c) \|\dot{u}(t)\|_2^2 + 2\varepsilon_*^2, \quad (6.14)$$

provided $t \in J_0$, where $c > 0$ (e.g. $c = 1/(2(d-1))$).

Next, we consider the case when $t \in J_1$. We use the following general inequality in $\dot{H}^1(\mathbb{R}^d)$, [12, Lemma 5.2]:

$$\|\nabla W\|_2^2 \leq \|\nabla u\|_2^2 + \frac{d-2}{2} K(u) + O(K^2(u)/\|\nabla u\|_2^2). \quad (6.15)$$

As $\|\nabla u(t)\|_2^2 \simeq \|\nabla W\|_2^2$ for $t \in J_1$, it follows that

$$E(\vec{u}) < J(W) + \varepsilon_*^2 \leq \frac{2^* - 2}{2 \cdot 2^*} \|\nabla u\|_2^2 + \frac{d-2}{2d} K(u) + O(K^2(u) + \varepsilon_*^2). \quad (6.16)$$

Then (6.10) implies that if $t \in J_1$,

$$\ddot{y} > 4(1 + c) \|\dot{u}\|_2^2 - 2K(u) - O(K^2(u) + \varepsilon_*^2). \quad (6.17)$$

To obtain a contradiction, we next observe that

$$\langle \dot{w}u | u \rangle = -\langle x \cdot \nabla w | |\tilde{w}u|^2 \rangle / (t + R) = \langle wu | \tilde{w}(r\partial_r + d/2)\tilde{w}u \rangle / (t + R), \quad (6.18)$$

where $\tilde{w} := \tilde{\chi}(|x|/(t+R))$ with some $\tilde{\chi} \in C^\infty(\mathbb{R})$ satisfying $\tilde{\chi} = 1$ on $\text{supp } \chi'$ and $\tilde{\chi}(r) = 0$ for $|r| \leq 1$. Hence

$$|\dot{y}| \leq \|wu\|_2 \|2\dot{u} + \tilde{w}(r\partial_r + d/2)\tilde{w}u/(t+R)\|_2 \quad (6.19)$$

and so

$$|\dot{y}|^2/y \leq \|2\dot{u} + \tilde{w}(r\partial_r + d/2)\tilde{w}u/(t+R)\|_2^2 \leq 4\|\dot{u}\|_2^2 + O(E_{\text{ext}}(t)). \quad (6.20)$$

We then infer from (6.14) that for $t \in J_0$,

$$\ddot{y}(t) \geq (1 + c) \frac{\dot{y}(t)^2}{y(t)} + \varepsilon_*^2, \quad (6.21)$$

and from (6.17) that for $t \in J_1$,

$$\ddot{y}(t) \geq (1 + c) \frac{\dot{y}(t)^2}{y(t)} - K(u(t)) - O(K^2(u(t)) + \varepsilon_*^2). \quad (6.22)$$

Now consider

$$\partial_t^2 y^{-c} = -cy^{-1-c} \left[\ddot{y} - (1 + c) \frac{\dot{y}(t)^2}{y(t)} \right]. \quad (6.23)$$

Again from the asymptotic behavior of $K(u)$ on each $I(t_*)$ in J_1 given by Lemma 3.2, (6.22), and the fact that y^{-1-c} is decreasing for large enough t imply that

$$\int_{J(t_*) \cap (-\infty, T)} y(t)^{-1-c} [-K(u(t)) - O(K^2(u(t)) + \varepsilon_*^2)] dt < 0, \quad (6.24)$$

for large $t_* > T_2$ and any $T > t_*$. In particular, we infer that

$$-\partial_t y^{-c}(t_*) \geq \inf_{t \in I(t_*)} \partial_t y^{-c}(t), \quad (6.25)$$

while $\partial_t y^{-c}(t)$ is decreasing in J_0 . Hence for some $t_* \in \mathcal{T}$ and for all $t > t_*$ we have

$$\partial_t y^{-c}(t) \leq \partial_t y^{-c}(t_*) < 0. \quad (6.26)$$

This implies finite time blow up, contradicting the earlier assumption. \square

7. SCATTERING AFTER EJECTION

Here we essentially repeat the argument given in [12] for the reader's convenience, with the small changes necessitated by the presence of space and momentum translations. In the region $\mathfrak{S} = +1$, we already know that all solutions are uniformly bounded in \mathcal{H} , but it is not sufficient for global existence of strongly continuous solution in the critical case. Now we resort to the recent result by Duyckaerts-Kenig-Merle [4] to preclude concentration (type II) blow-up. This is the only place where we have to restrict the dimensions² to 3 or 5

Proposition 7.1. *No solution as in Theorem 5.1 blows up in \mathcal{H}_X with $\mathfrak{S} = +1$ in the region $t \geq T_2$.*

Proof. First, Lemma 3.2 precludes blow-up in the hyperbolic region, since the scaling parameter is a priori bounded during the ejection process, which is valid when reversing the time direction. Hence a blow-up may happen only when $d_W(\vec{u}(t)) > \delta_H$, where $K(u(t)) \geq 0$ and so the energy assumption in Theorem 5.1 implies

$$\frac{\|\dot{u}(t)\|_2^2}{2} + \frac{\|\nabla u(t)\|_2^2}{d} = E(\vec{u}) - \frac{K(u(t))}{2^*} < J(W) + \varepsilon_*^2 = \frac{\|\nabla W\|_2^2}{d} + \varepsilon_*^2. \quad (7.1)$$

This allows us to employ the main result in [4], after reducing ε_* if necessary. Suppose u is a solution on $[0, T_+)$ in \mathcal{H}_X with $\mathfrak{S} = +1$ and $d_W(\vec{u}(t)) > \delta_H$ with the blow-up time $T_+ < \infty$. According to their result, we can then write for t sufficiently near T_+

$$\vec{u}(t) = \vec{W}_{\sigma(t)}(x - c(t)) + \varphi + o(1) \quad \text{in } \mathcal{H}, \quad (7.2)$$

²Strictly speaking, the long-time perturbation argument should be also modified for $d > 6$ in the scattering proof of Proposition 7.2, but it is a minor issue. See [17, 9] for the solution.

for some $\sigma(t) \rightarrow \infty$, $c(t) \in \mathbb{R}^d$ and some fixed $\underline{\varphi} \in \mathcal{H}$. It is then easily checked that as $t \rightarrow T_+ - 0$ we have

$$K(u(t)) = K(W_{\sigma(t)}) + K(\varphi_1) + o(1) = K(\varphi_1) + o(1), \quad (7.3)$$

from which we infer in particular that $K(\varphi_1) \geq 0$. Similarly, we obtain

$$J(W) + \varepsilon_*^2 > E(\vec{u}) = J(W) + E(\underline{\varphi}), \quad (7.4)$$

which implies via $K(\varphi_1) \geq 0$,

$$\|\varphi_2\|_2^2/2 + \|\nabla\varphi_1\|_2^2/d = E(\underline{\varphi}) - K(\varphi_1)/2^* < \varepsilon_*^2. \quad (7.5)$$

This however contradicts $d_W(\vec{u}(t)) > \delta_H \gg \varepsilon_*$ near T_+ . \square

Next we employ the Kenig-Merle scheme from [10, 11] to improve the above result. The one-pass theorem will be incorporated in the same way as in the subcritical case [21]. Extinction of the critical element requires a little extra work due to the possibility of concentration, which will be however reduced to the above proposition.

Proposition 7.2. *Every solution staying in \mathcal{H}_X with $\mathfrak{S} = +1$ and $d_W \geq \delta_*$ for $t > 0$ scatters to 0 as $t \rightarrow +\infty$ with uniformly bounded Strichartz norms on $[0, \infty)$.*

The restriction $d_W \geq \delta_*$ is essential for the uniform Strichartz bound, since the latter does not hold for all scattering solutions, even for $E(\vec{u}) < J(W)$.

Proof. We argue by contradiction. Let u_n be solutions on $[0, \infty)$ in \mathcal{H}_X satisfying

$$\begin{aligned} E(\vec{u}_n) &\rightarrow E_* \leq J(W) + \varepsilon_*^2, & \|u_n\|_{L_{t,x}^q(0,\infty)} &\rightarrow \infty, \\ d_W(\vec{u}_n(t)) &\geq \delta_*, & \mathfrak{S}(\vec{u}_n(t)) &= +1, \quad (t > 0) \end{aligned} \quad (7.6)$$

where we choose $q = 2(d+1)/(d-2)$ so that $L_{t,x}^q$ is an admissible Strichartz norm for the wave equation on \mathbb{R}^d . Here and after, $X(I)$ denotes the restriction to $I \times \mathbb{R}^d$ of the Banach function space X on $\mathbb{R} \times \mathbb{R}^d$. It is well-known that $L_{t,x}^q$ and the energy norm are sufficient to control all the other Strichartz norms, such as $L_t^p \dot{B}_{p,2}^{1/2}$ with $p = 2(d+1)/(d-1)$, as well as the nonlinear term in some dual admissible norm such as in $L_t^{p'} \dot{B}_{p',2}^{1/2}$ (see, for example, [8]).

We may assume that E_* is the minimum for the above property. Following the Kenig-Merle argument, the proof consists of two parts: construction and exclusion of a critical element.

Part I: Construction of a critical element.

Assuming the existence of (7.6), we are going to show that there is a critical element u_* , that is a solution on $[0, \infty)$ in \mathcal{H}_X satisfying

$$E(\vec{u}_*) = E_*, \quad \|u_*\|_{L_{t,x}^q(0,\infty)} = \infty, \quad d_W(\vec{u}_*(t)) \geq \delta_*, \quad \mathfrak{S}(\vec{u}_*(t)) = +1, \quad (7.7)$$

and that its trajectory is precompact modulo dilations and translations in \mathcal{H} .

If $d_W(\vec{u}_n(0)) < \delta_H$, then by Lemma 3.2, we have $d_W(\vec{u}_n(t)) \geq \delta_H$ at some later $t > 0$. Since the Strichartz norm on the ejection time interval is uniformly bounded, we may time-translate each u_n so that

$$d_W(\vec{u}_n(0)) \geq \delta_H, \quad (7.8)$$

without losing (7.6).

Since we chose $\varepsilon_* < \varepsilon_1(\delta_S) \leq \varepsilon_1(\delta_H)$, Lemma 4.1 implies

$$K(u_n(0)) \geq \min(\kappa(\delta_H), c\|\nabla u_n(0)\|_2^2). \quad (7.9)$$

Now apply³ the Bahouri-Gérard decomposition from [1], see also Lemma 4.3 in [11], to $\{\vec{u}_n(0)\}_{n \geq 1}$. Let $U(t)$ denotes the free wave propagator. We conclude that there exist $\lambda_n^j > 0$, $t_n^j \in \mathbb{R}$, $x_n^j \in \mathbb{R}^d$, $\underline{\varphi}^j \in \mathcal{H}$ and free waves w_n^J such that for any $J \geq 1$

$$U(t)\vec{u}_n(0) = \sum_{j=1}^J \vec{V}_n^j(t) + \vec{w}_n^J(t), \quad \vec{V}_n^j(t) := U(t + t_n^j)T_n^j \underline{\varphi}^j, \quad (7.10)$$

where $T_n^j := T^{-x_n^j} \mathcal{S}^{\log \lambda_n^j}$, such that

$$|\log(\lambda_n^j / \lambda_n^k)| + \lambda_n^j |t_n^j - t_n^k| + \lambda_n^j |x_n^j - x_n^k| \rightarrow \infty \quad (7.11)$$

for each $j \neq k$,

$$\begin{aligned} \lim_{n \rightarrow \infty} \left[\|\vec{u}_n(0)\|_{\mathcal{H}}^2 - \sum_{j=1}^J \|\vec{V}_n^j(0)\|_{\mathcal{H}}^2 - \|\vec{w}_n^J(0)\|_{\mathcal{H}}^2 \right] &= 0, \\ \lim_{n \rightarrow \infty} \left[E(\vec{u}_n(0)) - \sum_{j=1}^J E(\vec{V}_n^j(0)) - E(\vec{w}_n^J(0)) \right] &= 0 \end{aligned} \quad (7.12)$$

for each J , and

$$\lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \|w_n^J\|_{L_t^\infty L_x^{2^*}(\mathbb{R}) \cap L_{t,x}^q(\mathbb{R})} = 0. \quad (7.13)$$

The last property applies to any other non-sharp Strichartz norm by interpolation, since those free waves are all uniformly bounded.

³In what follows, we will pass to subsequences without any further mention.

First we check that all components retain the property $K \geq 0$ at $t = 0$. Indeed, one has

$$E(\vec{u}_n) - \frac{1}{2^*} K(u_n(0)) \geq \frac{1}{d} \|\vec{u}_n(0)\|_{\mathcal{H}}^2 = \sum_{j=1}^J \frac{1}{d} \|\vec{V}_n^j(0)\|_{\mathcal{H}}^2 + \frac{1}{d} \|\vec{w}_n^J(0)\|_{\mathcal{H}}^2 + o(1), \quad (7.14)$$

where $o(1) \rightarrow 0$ as $n \rightarrow \infty$. Hence if $\|\nabla u_n(0)\|_2^2 \lesssim \varepsilon_*^2$, then $\|\nabla V_n^j(0)\|_2^2 \lesssim \varepsilon_*^2 \ll 1$, and so $K(V_n^j(0)) \geq 0$. Otherwise, the lower bound in (7.9) is much bigger than ε_*^2 , so for large n , we conclude from the above inequality that

$$\frac{\|\vec{V}_n^j(0)\|_{\mathcal{H}}^2}{d} < J(W), \quad (7.15)$$

which implies $K(V_n^j(0)) \geq 0$, by the variational property of W . The same argument implies $K(w_n^J(0)) \geq 0$ as well. Thus, each component has non-negative energy E .

Now let U^j be the nonlinear profile associated with V_n^j , that is the nonlinear solution satisfying as $n \rightarrow \infty$,

$$\|\vec{U}^j(s_n^j) - U(s_n^j)\varphi^j\|_{\mathcal{H}} \rightarrow 0, \quad s_n^j := \lambda_n^j t_n^j, \quad (7.16)$$

defined uniquely around $t = s_\infty^j := \lim_{n \rightarrow \infty} s_n^j$, such that

$$\|\vec{U}_n^j(0) - \vec{V}_n^j(0)\|_{\mathcal{H}} \rightarrow 0 \quad \vec{U}_n^j(t) := (T_n^j \vec{U}^j)(\lambda_n^j(t + t_n^j)). \quad (7.17)$$

By the scaling invariance of the equation, each U_n^j is also a solution, defined locally around $t = 0$. Hence the above property of $\vec{V}_n^j(0)$ is transferred to U_n^j :

$$K(U_n^j(0)) \geq 0, \quad 0 \leq E(\vec{U}_n^j) = E(\vec{U}^j) \simeq \|\vec{U}_n^j(0)\|_{\mathcal{H}}^2, \quad (7.18)$$

$$\sum_{j=1}^J E(\vec{U}^j) + \lim_{n \rightarrow \infty} E(\vec{w}_n^J(0)) = E_*.$$

We may assume that $j = 1$ gives the maximum among $E(\vec{U}^j)$, then by [11], each U^j for $j > 1$ exists globally and scatters with

$$\sum_{j=2}^J \|U^j\|_{L_{t,x}^q(\mathbb{R})}^2 \lesssim \sum_{j=2}^J E(\vec{U}^j) \leq \frac{2}{3} J(W). \quad (7.19)$$

Now assume $\|U^1\|_{L_{t,x}^q(\mathbb{R})} < \infty$, which is the case if $E(U^1) < J(W)$. Then from the long-time perturbation theory, cf. Theorem 2.20 in [11], one obtains the *nonlinear profile decomposition* for the solutions $u_n(t)$, provided J is

large and fixed, and $n \geq n_0(J)$ is sufficiently large:

$$u_n = \sum_{j=1}^J U_n^j + w_n^J + R_n^J, \quad (7.20)$$

$$\lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \left[\|\vec{R}_n^J\|_{L_t^\infty \mathcal{H}(\mathbb{R})} + \|R_n^J\|_{L_{t,x}^q(\mathbb{R})} \right] = 0,$$

which implies u_n is bounded in $L_{t,x}^q$, contradicting (7.6). Thus we have obtained

$$\|U^1\|_{L_{t,x}^q(T_-, T_+)} = \infty, \quad J(W) \leq E(\vec{U}^1) \leq E_*, \quad \sum_{j=2}^J E(\vec{U}^j) + \|\vec{w}_n^J\|_{\mathcal{H}}^2 \lesssim \varepsilon_*^2, \quad (7.21)$$

where (T_-, T_+) is the maximal existence interval of U^1 .

We now distinguish three cases (a)–(c) by $s_\infty^1 = \lim_{n \rightarrow \infty} \lambda_n^1 t_n^1$:

(a) $s_\infty^1 = \infty$. Then by definition (7.16), U^1 is a local solution around $t = \infty$ with finite Strichartz norms, and

$$\|U_n^1\|_{L_{t,x}^q(0, \infty)} = \|U^1\|_{L_{t,x}^q(s_n^1, \infty)} \rightarrow 0. \quad (7.22)$$

Hence we can use the long-time perturbation argument on $(0, \infty)$, which gives a contradiction via (7.20) as above.

(b) $s_\infty^1 = -\infty$. In this case U^1 scatters at $t = -\infty$ by definition.

If $d_W(\vec{U}^1(t)) > \delta_*/2$ for all $t < T_+$, then $\vec{U}^1(t)$ remains in the region $\mathfrak{S} = +1$ from $t = -\infty$. Hence $T_+ = \infty$ by Proposition 7.1, and $\|U^1\|_{L_{t,x}^q(0, \infty)} = \infty$. Moreover, Theorem 5.1 together with Lemma 3.2 implies that $d_W(\vec{U}^1(t)) \geq \delta_*$ for large t . Hence U^1 is a critical element after some time translation.

Otherwise, $d_W(\vec{U}^1(t_*)) = \delta_*/2$ at some minimal $t_* < T_+$, until which time U^1 remains in $\mathfrak{S} = +1$, and $\|U^1\|_{L_{t,x}^q(-\infty, t_*)} < \infty$. Hence one can apply the nonlinear profile decomposition on the interval $\lambda_n^1(t + t_n^1) \leq t_*$ as in (7.20), which yields in particular that, upon choosing J sufficiently large,

$$d_W(\vec{u}_n((t_* - s_n^1)/\lambda_n^1)) \leq d_W(\vec{U}^1(t_*)) + O(\varepsilon_*) + o(1) \leq \frac{2}{3}\delta_* + o(1), \quad (7.23)$$

as $n \rightarrow \infty$. However, since $t_* - s_n^1 \rightarrow \infty$, this contradicts our assumption

$$\inf_{t \geq 0} d_W(\vec{u}_n(t)) \geq \delta_*. \quad (7.24)$$

(c) $s_\infty^1 \in \mathbb{R}$. Then by the same perturbative arguments as above, the nonlinear profile decomposition (7.20) holds on any compact interval in $(T_-, T_+)/\lambda_n^1 - t_n^1$. Thus, as in the case (b), we deduce from $\inf_{t \geq 0} d_W(\vec{u}_n(t)) \geq \delta_*$ that

$$\inf_{s_\infty^1 \leq t < T_+} d_W(\vec{U}^1(t)) \geq \delta_*/2. \quad (7.25)$$

Then the same argument as in (b) implies that $T_+ = \infty$ and $\|U^1\|_{L_{t,x}^q(s_\infty^1, \infty)} = \infty$, since otherwise U^1 scatters and the nonlinear profile decomposition holds on $[0, \infty)$, contradicting (7.6).

Thus we arrive at the conclusion that $s_\infty^1 < \infty$ and U^1 is a critical element after time translation. This implies $E(U^1) = E_*$ by the minimality, which extinguishes the other profiles U^j ($j > 1$) as well as the remainder w_n^j as $n \rightarrow \infty$, through the nonlinear energy decomposition.

Having constructed a critical element u_* , we apply the above argument to the sequence

$$u_n(t) = u_*(t - t_n), \quad t_n \rightarrow \infty. \quad (7.26)$$

Then the vanishing of all but one (free) profile implies that for some continuous $\sigma(t) \in \mathbb{R}$, $x(t) \in \mathbb{R}^d$,

$$\{T^{x(t)} \vec{S}^{\sigma(t)} \vec{u}_*(t)\}_{t \geq 0} \subset \mathcal{H} \quad (7.27)$$

is precompact, concluding the first part of the proof.

Before proceeding to the extinction, we show that

$$|P(\vec{u})| \lesssim \varepsilon_*. \quad (7.28)$$

Suppose towards a contradiction that $|P(\vec{u}_*)| = |P_1(\vec{u}_*)| \gg \varepsilon_*$. Then, since $J(W) \leq E(\vec{u}_*) < J(W) + \varepsilon_*^2$, we can use the Lorentz transform to reduce the energy below $J(W)$. Indeed, let w be any global strong energy solution of (1.1) and Lorentz transform it as follows: with a parameter $\nu \in \mathbb{R}$,

$$w(t, x) \mapsto w_\nu(t, x) := w(t \cosh \nu + x_1 \sinh \nu, x_1 \cosh \nu + t \sinh \nu, x_2, x_3). \quad (7.29)$$

Then one checks that w_ν is again a strong energy solution of (1.1) which satisfies

$$\begin{aligned} E(w_\nu) &= E(w) \cosh \nu + P_1(w) \sinh \nu, \\ P_1(w_\nu) &= P_1(w) \cosh \nu + E(w) \sinh \nu, \quad P_\alpha(w_\nu) = P_\alpha(w), \quad (\alpha > 1). \end{aligned} \quad (7.30)$$

Now we claim that we can apply the above transform to the forward global solution u_* , and then with some $\nu = O(\varepsilon_*)$ we can construct another forward global solution u_* with $E(\vec{u}_*) < J(W)$ and $\|u_*\|_{L_{t,x}^q(0, \infty)} = \infty$. This contradicts Kenig-Merle's result [11] for $E < J(W)$. In order to transform a solution with infinite Strichartz norm, we argue in the same way as in the subcritical case using the finite propagation speed:

Lemma 7.3. *Let u be a solution of (1.1) in $C(I; \mathcal{H}) \cap L_{t,x}^q(I)$ on a time interval $I \ni T$. Then there is an open neighborhood O of the identity in the Lorentz group, such that the transform of u by any $g \in O$ extends to a solution (with finite energy and L^q) in a space-time region including a time slab which contains T . If u is a solution in $C([T, \infty); \mathcal{H}) \cap L_{loc}^q((T, \infty); L_x^q)$, then for any Lorentz transform u' of u , there exists $T' \in \mathbb{R}$ such that u'*

extends to a solution in $C([T', \infty); \mathcal{H}) \cap L_{loc}^q((T', \infty); L_x^q)$. Moreover, if $\|u'\|_{L_{t,x}^q(T', \infty)} < \infty$ then $\|u\|_{L_{t,x}^q(T, \infty)} < \infty$.

The proof is also the same as for [20, Lemmas 6.1 and 6.2], so we omit it.

Part II: Exclusion of a critical element.

Let u_* be a critical element (7.7), hence

$$\vec{w}_*(t) := \rho(t)^{d/2} \vec{u}_*(t, \rho(t)(x - x(t))), \quad \rho(t) = e^{-\sigma(t)} \quad (7.31)$$

for $t \geq 0$ is precompact in \mathcal{H} . We proceed in three steps.

Step 1: $\limsup_{t \rightarrow \infty} \rho(t)/t < \infty$. To see this, note that by finite propagation speed, we have

$$\lim_{R \rightarrow \infty} \sup_{t \geq 0} \|\vec{u}_*(t)\|_{\mathcal{H}(|x| > t+R)} = 0, \quad (7.32)$$

whence we have

$$\lim_{R \rightarrow \infty} \sup_{t \geq 0} \|\vec{w}_*(t)\|_{\mathcal{H}(|x-x(t)| > (t+R)/\rho(t))} = 0. \quad (7.33)$$

If for some sequence of times $\{s_n\}_{n \geq 1}$ we had $\rho(s_n)/s_n \rightarrow \infty$, then by precompactness of $\{\vec{w}_*(t)\}_{t \geq 0}$, we get $\|\vec{w}_*(s_n)\|_{\mathcal{H}} \rightarrow 0$, whence also $\|\vec{u}_*(s_n)\|_{\mathcal{H}} \rightarrow 0$, which would force $E_* = 0$, a contradiction.

Step 2: $\liminf_{t \rightarrow \infty} \rho(t)/t > 0$. This follows from the localized virial identity (5.18), together with the control on the energy center as well as the energy equipartition, as in the proof of Theorem 5.1. By the precompactness, there is $R > 0$, depending on u_* , such that for all $t \geq 0$

$$\|\vec{u}_*(t)\|_{\mathcal{H}(|x+\rho(t)x(t)| > R\rho(t))} < \varepsilon_*. \quad (7.34)$$

Suppose for contradiction that $\liminf_{t \rightarrow \infty} \rho(t)/t = 0$. Choose $T_3 \gg T_2 \gg 1$ and $R_2, R_3 > 0$ such that

$$\rho(T_j) \ll \varepsilon_* T_j / R, \quad R_j := R\rho(T_j). \quad (7.35)$$

Let $c_j := -\rho(T_j)x(T_j)$ with $c(T_2) = 0$ by space translation. Then (7.34) implies in particular that $\|\vec{u}_*(T_j)\|_{\mathcal{H}(|x-c_j| > R_j)} < \varepsilon_*$, hence by the finite propagation speed we have

$$|c_3| = |c_3 - c_2| \leq |T_3 - T_2| + R_2 + R_3 = |T_3 - T_2| + O(\varepsilon_* T_3), \quad (7.36)$$

where we used (7.35). Let

$$w(t, x) = \chi(|x|/(t - T_2 + R_2)), \quad E_{\text{ext}}(t) = \|\vec{u}_*(t)\|_{\mathcal{H}(|x| > t - T_2 + R_2)}^2, \quad (7.37)$$

with the same χ as in (5.6). Using the finite propagation speed as before, one has

$$\sup_{t > T_2} E_{\text{ext}}(t) \lesssim \varepsilon_*^2. \quad (7.38)$$

For the localized center of energy $\mathfrak{C}(t) = \langle xw | e(\vec{u}) \rangle$, we infer as before

$$\begin{aligned} \dot{\mathfrak{C}}(t) &= O(E_{\text{ext}}(t)) - P(\vec{u}) = O(\varepsilon_*), \quad |\mathfrak{C}(T_2)| \lesssim R_2 \ll \varepsilon_* T_2, \\ \mathfrak{C}(T_3) &= \int_{|x-c_3| < R_3} x e(\vec{u}_*(T_3)) dx + O(\varepsilon_*^2 T_3) = c_3 E(u_*) + O(\varepsilon_* T_3). \end{aligned} \quad (7.39)$$

Thus we obtain upon integrating $\mathfrak{C}(t)$ on (T_2, T_3) ,

$$|c_3| \lesssim \varepsilon_* T_3. \quad (7.40)$$

For the localized virial $V_w(t) = \langle w \dot{u}_* | (r \partial_r + d/2) u_* \rangle$, one has as before

$$\begin{aligned} \dot{V}_w(t) &= -K(u_*(t)) + O(E_{\text{ext}}(t)) \lesssim -K(u_*(t)) + O(\varepsilon_*^2), \\ |V_w(T_2)| &\lesssim R_2 \ll \varepsilon_* T_2, \quad |V_w(T_3)| \lesssim |c_3| + R_3 \lesssim \varepsilon_* T_3, \end{aligned} \quad (7.41)$$

Integrating \dot{V}_w on (T_2, T_3) , and then arguing as for (5.27), we obtain

$$\delta_* \int_{T_2}^{T_3} \|\nabla u_*(t)\|_2^2 dt - O(\delta_S) \lesssim \int_{T_2}^{T_3} K(u_*(t)) dt \lesssim \varepsilon_* T_3, \quad (7.42)$$

where the negative term $-O(\delta_S)$ arises in case⁴ a hyperbolic interval $I(t_*) \not\subset (T_2, T_3)$ has only its negative part inside (T_2, T_3) .

Similarly, using Hardy, we have as before

$$\partial_t \langle w \dot{u}_* | u_* \rangle \geq 2E(\vec{u}_*) - \|\nabla u_*(t)\|_2^2 + O(\varepsilon_*^2), \quad |[\langle w \dot{u}_* | u_* \rangle]_{T_2}^{T_3}| \lesssim \varepsilon_* T_3. \quad (7.43)$$

Integrating the left inequality and combining it with the above estimates, we obtain

$$E(\vec{u}_*) T_3 \lesssim \varepsilon_* \delta_*^{-1} T_3 + \varepsilon_* \delta_S / \delta_*. \quad (7.44)$$

Thus we arrive at $J(W) \leq E(\vec{u}_*) \lesssim \varepsilon_* / \delta_* \ll J(W)$, a contradiction.

It now also follows that we may put $x(t) = 0$ for all $t \geq 0$, since the assumption

$$\limsup_{t \rightarrow \infty} |x(t)| = \infty \quad (7.45)$$

contradicts the compactness property of \vec{w}_* , see (7.33).

Step 3: Construction of a blow up solution via a re-scaling of u_* . Pick a sequence $s_n \rightarrow \infty$ with $\lim_{n \rightarrow \infty} \rho(s_n)/s_n = c \in (0, \infty)$, as well as $\vec{w}_*(s_n) \rightarrow \exists \varphi$ in \mathcal{H} . Define a sequence of solutions

$$u_n(t, x) := s_n^{d/2-1} u_*(s_n t, s_n x) \quad (7.46)$$

whence we have $\vec{u}_n(1) \rightarrow c^{-d/2} \varphi(x/c)$ in \mathcal{H} .

The above two steps imply that \vec{u}_n is precompact in $C([\tau, 1]; \mathcal{H})$ for any $0 < \tau < 1$, and so, after replacement by a subsequence, it converges to some \vec{u}_∞ in $C((0, 1]; \mathcal{H})$. By the local wellposedness theory, it has finite Strichartz norms locally in time, and so u_∞ is the unique strong solution on $(0, 1]$ with

⁴This could be avoided by taking $\varepsilon_* < \varepsilon_1(\delta_*)$ instead of $< \varepsilon_1(\delta_S)$.

the initial condition $\vec{u}_\infty(1) = \underline{\varphi}$. Clearly we also have $d_W(\vec{u}_\infty(t)) \geq \delta_*$ and $\mathfrak{S}(\vec{u}_\infty(t)) = +1$ for $0 < t \leq 1$.

We now show that u_∞ is a solution blowing up at $t = 0$, which contradicts Proposition 7.1. The fact that u_∞ blows up at $t = 0$ follows from the next

Claim: $u_\infty(t, x) = 0$ on $|x| > t$. To see this, pick $0 < \varepsilon \ll 1$ arbitrary, let m large enough such that $\|\vec{w}_*(s_m) - \underline{\varphi}\|_{\mathcal{H}} \ll \varepsilon$ and further pick $R > 0$ such that $\|\underline{\varphi}\|_{\mathcal{H}(|x|>R)} \ll \varepsilon$. Then for $n > m$, we have

$$\|\vec{u}_n(s_m/s_n)\|_{\mathcal{H}(|x|>R\rho(s_m)/s_n)} = \|\vec{w}_*(s_m)\|_{\mathcal{H}(|x|>R)} \ll \varepsilon. \quad (7.47)$$

From this and the finite propagation speed, we deduce that for $s_m/s_n \leq t \leq 1$

$$\|\vec{u}_n(t)\|_{\mathcal{H}(|x|>R\rho(s_m)/s_n+t-s_m/s_n)} \ll \varepsilon. \quad (7.48)$$

Letting $n \rightarrow \infty$, we infer that for $0 < t \leq 1$

$$\|\vec{u}_n(t)\|_{\mathcal{H}(|x|>t)} \ll \varepsilon. \quad (7.49)$$

Since $\varepsilon > 0$ is arbitrary, this implies that u_∞ is supported on $|x| \leq t$, as claimed. This completes the proof of Proposition 7.2. \square

In order to complete the proof of Theorem 1.1, we now exhibit open data sets at time $t = 0$ such that we have blow up/scattering at $t = \pm\infty$, four possibilities in all. However, this has been done in the radial case [12] by producing four solutions starting from the neighborhood $\mathcal{H}_* \setminus \mathcal{H}_X$ and exiting from it in finite time in both time directions, for all four combinations of \mathfrak{S} at the exiting times. Since such behavior is obviously stable in the energy space \mathcal{H} , we get an open set around each solution by the local wellposedness.

In fact, the initial data for such solutions can be given explicitly by

$$\vec{u}(0) = \vec{W} + \varepsilon \underline{a} \rho, \quad \underline{a} = \begin{pmatrix} \pm 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ \pm 1 \end{pmatrix}, \quad 0 < \varepsilon \ll \varepsilon_*. \quad (7.50)$$

For any solution u in the region $d_W(\vec{u}(t)) < \delta_E$, Lemma 3.1 yields

$$\vec{u}(t) = T^{c(t)} \vec{S}^{\sigma(t)} (\mathfrak{s} \vec{W} + \underline{v}(t)), \quad \underline{v}(t) = \underline{\lambda}(t) \rho + \underline{\gamma}(t), \quad (7.51)$$

with $d_W(t) := d_W(\vec{u}(t)) \sim |\underline{\lambda}| + \|\underline{\gamma}\|_E$, and from the proof of Lemma 3.2,

$$\begin{aligned} \partial_\tau \lambda_1 &= \lambda_2, & \partial_\tau \lambda_2 &= k^2 \lambda_1 + O(d_W^2), \\ \partial_\tau [\|\underline{\gamma}\|_E^2 + O(\|\underline{\gamma}\|_E d_W^2)] &= O(\|\underline{\gamma}\|_E d_W^2). \end{aligned} \quad (7.52)$$

In particular, if $\gamma(0) = 0$ and the linearized solution $\underline{\lambda}^0$ for $\underline{\lambda}$ satisfies for $0 \leq \tau' \leq \tau$

$$\int_0^{\tau'} |\underline{\lambda}^0(\tau'')| d\tau'' \lesssim |\underline{\lambda}^0(\tau')| < \delta_E, \quad (7.53)$$

then we deduce that (see [18, 21] for more detail in the subcritical radial case)

$$|\underline{\lambda} - \underline{\lambda}^0| + \|\underline{\gamma}\|_E \lesssim |\underline{\lambda}^0|^2 \simeq d_W^2. \quad (7.54)$$

This is the case for (7.50), with

$$\begin{cases} \underline{a} = (\pm 1, 0) \implies \underline{\lambda}^0 = \pm \varepsilon (\cosh(k\tau), k \sinh(k\tau)), \\ \underline{a} = (0, \pm 1) \implies \underline{\lambda}^0 = \pm \varepsilon (\sinh(k\tau)/k, \cosh(k\tau)). \end{cases} \quad (7.55)$$

Moreover, when $\varepsilon \ll |\underline{\lambda}^0| \ll \delta_S$, the solution is in the region \mathcal{H}_X with $\mathfrak{S}(\vec{u}) = -\text{sign } \lambda_1^0$. Hence the solution for $\underline{a} = (\pm 1, 0)$ respectively blows up and scatters both in $t < 0$ and $t > 0$, while the solutions for $\underline{a} = (0, \pm 1)$ blows up in $\pm t > 0$ and scatters for $t \rightarrow \mp \infty$. One can easily check that the former case $\underline{a} = (\pm 1, 0)$ is actually in the Kenig-Merle (or Payne-Sattinger) criterion $E(\vec{u}) < J(W)$ and $\pm K(u(0)) < 0$, while the latter case $\underline{a} = (0, \pm 1)$ satisfies $E(\vec{u}) > J(W)$.

REFERENCES

- [1] Bahouri, H., Gérard, P. *High frequency approximation of solutions to critical nonlinear wave equations*. Amer. J. Math. **121** (1999), no. 1, 131–175.
- [2] Caffarelli, L., Gidas, B., Spruck, J. *Asymptotic symmetry and local behavior of semi-linear elliptic equations with critical Sobolev growth*. Comm. Pure Appl. Math. **42** (1989), no. 3, 271–297.
- [3] Duyckaerts, T., Kenig, C., Merle, F. *Universality of blow-up profile for small radial type II blow-up solutions of energy-critical wave equation*, preprint, arXiv:0910.2594.
- [4] Duyckaerts, T., Kenig, C., Merle, F. *Universality of the blow-up profile for small type II blow-up solutions of energy-critical wave equation: the non-radial case*, preprint, arXiv:1003.0625.
- [5] Duyckaerts, T., Kenig, C., Merle, F. *Profiles of bounded radial solutions of the focusing, energy-critical wave equation*, preprint 2011.
- [6] Duyckaerts, T., Merle, F. *Dynamic of threshold solutions for energy-critical NLS*. Geom. Funct. Anal. **18** (2009), no. 6, 1787–1840.
- [7] Duyckaerts, T., Merle, F. *Dynamic of threshold solutions for energy-critical wave equation*. Int. Math. Res. Pap. IMRP **2008**.
- [8] Ginibre, J., Soffer, A., Velo, G. *The global Cauchy problem for the critical non-linear wave equation*, J. Funct. Anal. **110** (1992), 96–130.
- [9] Ibrahim, S., Masmoudi, N., Nakanishi, K. *Scattering threshold for the focusing non-linear Klein-Gordon equation*, preprint, arXiv:1001.1474, to appear in Analysis & PDE.
- [10] Kenig, C., Merle, F. *Global well-posedness, scattering, and blow-up for the energy-critical focusing nonlinear Schrödinger equation in the radial case*, Invent. Math. **166** (2006), no. 3, pp. 645–675.
- [11] Kenig, C., Merle, F. *Global well-posedness, scattering and blow-up for the energy-critical focusing non-linear wave equation*. Acta Math. **201** (2008), no. 2, 147–212.
- [12] Krieger, J., Nakanishi, K., Schlag, W. *Global dynamics away from the ground state for the energy-critical nonlinear wave equation*, to appear in Amer. Journal Math.
- [13] Krieger, J., Schlag, W. *On the focusing critical semi-linear wave equation*. Amer. J. Math. **129** (2007), no. 3, 843–913.
- [14] Krieger, J., Schlag, W., Tataru, D. *Slow blow-up solutions for the $H^1(\mathbb{R}^3)$ critical focusing semilinear wave equation*. Duke Math. J. **147** (2009), no. 1, 1–53.
- [15] Lions, P.-L. *The concentration-compactness principle in the calculus of variations. The limit case. I*. Rev. Mat. Iberoamericana **1** (1985), no. 1, 145–201.
- [16] Merle, F., Vega, L. *Compactness at blow-up time for L^2 solutions of the critical nonlinear Schrödinger equation in 2D*. Internat. Math. Res. Notices 1998, no. 8, 399–425.

- [17] Nakanishi, K. *Scattering theory for the nonlinear Klein-Gordon equation with Sobolev critical power*. Internat. Math. Res. Notices **1999**, no. 1, 31–60.
- [18] Nakanishi, K., Schlag, W. *Global dynamics above the ground state energy for the focusing nonlinear Klein-Gordon equation*, Journal Diff. Eq. 250 (2011), 2299–2233.
- [19] Nakanishi, K., Schlag, W. *Global dynamics above the ground state energy for the cubic NLS equation in 3D*, to appear in Calc. Var. and PDE.
- [20] Nakanishi, K., Schlag, W. *Global dynamics above the ground state for the nonlinear Klein-Gordon equation without a radial assumption*. To appear in Arch. Rational Mech. Analysis.
- [21] Nakanishi, K., Schlag, W. *Invariant manifolds and dispersive Hamiltonian evolution equations*, Zürich Lectures in Advanced Mathematics, EMS, 2011.
- [22] Payne, L. E., Sattinger, D. H. *Saddle points and instability of nonlinear hyperbolic equations*. Israel J. Math. **22** (1975), no. 3-4, 273–303.
- [23] Shatah, J., Struwe, M. *Geometric Wave Equations*, Courant Lecture Notes, AMS, 1998.

DEPARTMENT OF MATHEMATICS, EPFL, LAUSANNE, SWITZERLAND

DEPARTMENT OF MATHEMATICS, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF CHICAGO, 5734 SOUTH UNIVERSITY AVENUE, CHICAGO, IL 60615, U.S.A.